

Exposures to different SARS-CoV-2 spike variants elicit neutralizing antibody responses with differential specificity towards established and emerging strains

Running title: Immunity elicited by SARS-2 variants

Matthew T Laurie¹, Jamin Liu^{1,2}, Sara Sunshine¹, James Peng³, Douglas Black³, Anthea M Mitchell^{1,4}, Sabrina A Mann^{1,4}, Genay Pilarowski^{5,6}, Kelsey C Zorn¹, Luis Rubio³, Sara Bravo⁶, Carina Marquez³, Maya Petersen⁷, Diane Havlir³, Joseph DeRisi^{1,4*}

¹Department of Biochemistry and Biophysics, University of California San Francisco, CA 94143, USA

²University of California, Berkeley-University of California, San Francisco Graduate Program in Bioengineering, Berkeley, CA 94720, USA

³Department of Medicine, University of California, San Francisco, San Francisco, CA 94143, USA

⁴Chan Zuckerberg Biohub, San Francisco, CA 94158, USA

⁵The Public Health Company, Oakland, CA 94610, USA

⁶Unidos en Salud, San Francisco, CA, 94143, USA

⁷Division of Biostatistics, University of California, Berkeley, Berkeley, CA 94720, USA

*Corresponding author:

Joseph DeRisi
1700 4th St., San Francisco, CA 94158
Phone:415-418-3647
Email: joe@derisilab.ucsf.edu

Keywords: SARS-CoV-2; COVID-19; neutralization; variant; B.1.617.2 (Delta); vaccination; natural infection; antibody escape; resistance; immune exposure

ABSTRACT

The wide spectrum of SARS-CoV-2 variants with phenotypes impacting transmission and antibody sensitivity necessitates investigation of the immune response to different spike protein versions. Here, we compare the neutralization of variants of concern, including B.1.617.2 (Delta) in sera from individuals exposed to variant infection, vaccination, or both. We demonstrate that neutralizing antibody responses are strongest against variants sharing one or more spike mutations with the immunizing exposure. We also observe that exposure to multiple spike

variants increases the breadth of variant cross-neutralization. These findings contribute to understanding the relationship between exposures and antibody responses and may inform booster vaccination strategies.

INTRODUCTION

Genomic surveillance of SARS-CoV-2 continues to identify a diverse spectrum of emerging variants possessing mutations in the spike gene, the main viral determinant of cellular entry and primary target of neutralizing antibodies [1]. Many spike mutations likely result from selective pressure which improves viral fitness through increased transmissibility or evasion of host immunity [2,3]. Studies have demonstrated that sera from vaccinated and naturally infected individuals yield diminished neutralizing activity against certain variants, including the recently ascendant Delta variant [4,5]. Because serum neutralization titer is an important correlate of real-world protective immunity, these findings suggest that antibody responses elicited by exposure to ancestral spike versions (Wuhan or D614G) will be less effective at preventing future infection by certain variants [6]. However, the diversity and prevalence of variants have fluctuated greatly throughout the pandemic, creating a complex population of individuals that may have inherently different capacity to neutralize certain variants depending on the specific genotype of their previous exposures, including vaccination [7].

In this study, we address the question of variant-elicited immune specificity by determining the breadth of neutralizing activity elicited by exposure to specific SARS-CoV-2 variants, vaccines, or both. To accomplish this, we collected serum from subjects with prior infections by variants B.1 (D614G mutation only), B.1.429 (Epsilon), P.2 (Zeta), and B.1.1.519, which were identified by viral sequencing. We also collected serum from mRNA vaccine recipients who were infected with the B.1 ancestral spike lineage prior to vaccination, infected with B.1.429 prior to vaccination, or had no prior infection. We measured and compared the

neutralization titer of each serum cohort against a panel of pseudoviruses representing each different exposure variant plus the variants of concern B.1.351 (Beta), P.1 (Gamma), B.1.617, and B.1.617.2 (Delta), which have one or more spike mutations of interest in common with one of the exposure variants. Our results provide a quantitative comparison of the degree of neutralization specificity produced by different exposures. We also demonstrate the effect of serial exposure to different spike versions in broadening the cross-reactivity of neutralizing antibody responses. Together, these findings describe correlates of protective immunity within the rapidly evolving landscape of SARS-CoV-2 variants and are highly relevant to the design of future vaccination strategies targeting spike antigens.

METHODS

Serum collection

Samples for laboratory studies were obtained under informed consent from participants in an ongoing community based, SARS-CoV-2 testing, genomic surveillance, and vaccination program “Unidos en Salud”, which serves a predominantly Latino community in the Mission neighborhood in San Francisco, California [8,9]. Subjects with and without symptoms of COVID-19 were screened with the BinaxNOW rapid antigen assay (supplied by California Department of Public Health). Positive rapid tests were followed by immediate disclosure and outreach to household members for testing, supportive community services, and academic partnership for research studies. All samples were sequenced using ARTIC Network V3 primers on an Illumina NovaSeq platform and consensus genomes generated from the resulting raw .fastq files using IDseq [10].

Convalescent serum donors were selected based on sequence-confirmed infection with the following variants of interest: B.1 (D614G mutation only; n=10 donors), B.1.429 (Epsilon; n=15), B.1.1.519 (n=6), P.2 (Zeta; n=1), B.1.526 (Iota; n=1), D614G infection followed by

BNT162b2 vaccination (n=8), B.1.429 infection followed by BNT162b2 vaccination (n=17). Serum was also collected from healthy recipients of BNT162b2 (n=5) and mRNA-1273 (n=6) vaccines (vaccines based on the Wuhan spike sequence), which were confirmed to have no prior SARS-CoV-2 infection by anti-SARS-CoV-2 nucleocapsid IgG assay [11]. All serum was collected from donors at least 14 days (mean 34 days, standard deviation 16 days) after exposure to either SARS-CoV-2 or the second dose of mRNA vaccine. For pooled serum experiments, samples from the same exposure group were pooled at equal volumes. Serum samples from the closely related exposures P.2 and B.1.526 were pooled together for the E484K exposure pool, and samples from BNT162b2 and mRNA-1273 exposures were pooled together for the vaccine exposure pool because of the very similar neutralization specificity observed in individual tests of these sera. Serum samples were heat inactivated at 56°C for 30 minutes prior to experimentation. Subject demographic information and time from positive test/vaccination to serum collection are summarized in Table S1 and precise exposure of each subject is detailed in Table S2.

Pseudovirus production

SARS-CoV-2 pseudoviruses bearing spike proteins of variants of interest were generated using a recombinant vesicular stomatitis virus expressing GFP in place of the VSV glycoprotein (rVSV Δ G-GFP) described previously [12]. The following mutations representative of specific spike variants were cloned in a CMV-driven expression vector and used to produce SARS-CoV-2 spike pseudoviruses: B.1 (D614G), B.1.429/Epsilon (S13I, W152C, L452R, D614G), P.2/Zeta (E484K, D614G), B.1.351/Beta (D80A, D215G, Δ 242-244, K417N, E484K, N501Y, D614G, A701V), P.1/Gamma (L18F, T20N, P26S, D138Y, R190S, K417T, E484K, N501Y, D614G, H655Y, T1027I, V1176F), B.1.1.519 (T478K, D614G, P681H, T732A), B.1.617 (L452R, E484Q, D614G, P681R), and B.1.617.2/Delta (T19R, T95I, G142D, Δ 157-158, L452R, T478K, P681R, D614G, D950N). Pseudoviruses were titered on Huh7.5.1 cells overexpressing

ACE2 and TMPRSS2 (gift of Andreas Puschnik) using GFP expression to measure the concentration of focus forming units (ffu) [13].

Pseudovirus neutralization experiments

Huh7.5.1-ACE2-TMPRSS2 cells were seeded in 96-well plates at a density of 7000 cells/well one day prior to pseudovirus inoculation. Serum samples were diluted into complete culture media (DMEM with 10% FBS, 10mM HEPES, 1x Pen-Strep-Glutamine) using the LabCyte Echo 525 liquid handler and 1500 ffu of each pseudovirus was added to the diluted serum to reach final dilutions ranging from 1:40-1:5120, including no-serum and no-pseudovirus controls. Serum/pseudovirus mixtures were incubated at 37°C for 1h before being added directly to cells. Cells inoculated with serum/pseudovirus mixtures were incubated at 37°C and 5% CO₂ for 24h, resuspended using 10x TrypLE Select (Gibco), and cells were assessed with the BD Celesta flow cytometer. All neutralization assays were repeated in a total of three independent experiments with each experiment containing two technical replicates for each condition. Cells were verified to be free of mycoplasma contamination with the MycoAlert Mycoplasma detection kit (Lonza).

Data analysis

Pseudovirus flow cytometry data was analyzed with FlowJo to determine the percentage of GFP-positive cells, indicating pseudovirus transduction. Percent neutralization for each condition was calculated by normalizing GFP-positive cell percentage to no-serum control wells. Neutralization titers (NT₅₀ and NT₉₀) were calculated from eight-point response curves generated in GraphPad Prism 7 using four-parameter logistic regression. The fold-change in pseudovirus neutralization titer in each serum group was calculated by normalizing each variant NT₅₀ and NT₉₀ value to D614G pseudovirus NT₅₀ and NT₉₀ values in the same serum group. To

compare neutralization titer across a panel of different pseudoviruses and serum groups, the Log₂ fold-change compared to D614G pseudovirus was reported.

RESULTS

We compared the 50% and 90% neutralization titers (NT₅₀ and NT₉₀) of D614G and B.1.429 (Epsilon) pseudoviruses in individual serum samples from subjects exposed to D614G infection, B.1.429 infection, mRNA vaccination, D614G infection followed by mRNA vaccination, and B.1.429 infection followed by mRNA vaccination (Figure 1). Fold-changes in both NT₅₀ and NT₉₀ are reported since these values often differ in magnitude due to differences in neutralization curve slope between different variants and sera. In D614G-exposed and vaccine-exposed serum, we observed approximately 2 to 3-fold decreases in average neutralization titer against B.1.429 pseudovirus compared to D614G pseudovirus. As expected, B.1.429-exposed serum neutralized B.1.429 pseudovirus more efficiently than D614G pseudovirus. Of note, previous infection with either D614G or B.1.429 followed by vaccination led to substantially higher neutralization titers against both pseudoviruses. In contrast to other exposure groups, serum from vaccine recipients previously infected by B.1.429 neutralized D614G and B.1.429 at similar titers, with only a 1.3-fold difference in NT₉₀, indicating that exposure to multiple spike variants elicits a potent response with specificity toward the breadth of prior exposures.

We next investigated how exposure impacts neutralization specificity by crossing a panel of eight different spike variants against serum pools elicited by seven different prior exposures. (Figure 2). A range of reductions in neutralization titer relative to D614G pseudovirus were observed, with B.1.351 (Beta) and B.1.617.2 (Delta) exhibiting the greatest resistance to neutralization in most serum types with up to 12-fold and 4-fold reductions in NT₉₀, respectively. However, reductions in neutralization titer were considerably smaller or absent in serum from subjects previously exposed to a variant bearing some or all of the same spike mutations as the

variant being tested. Specifically, prior exposure to the E484K mutation in the spike receptor binding domain (RBD) produced the greatest neutralization for the four tested variants with mutations at the E484 position: P.1 (Gamma), P.2 (Zeta), B.1.351 (Beta), and B.1.617. Similarly, B.1.617.2 (Delta) was neutralized more effectively by B.1.429-exposed serum with and without subsequent vaccination, and most effectively by B.1.1.519-exposed serum. These effects are presumably due to the shared L452R RBD mutation in both B.1.429 and B.1.617.2, and the shared T478K RBD mutation and P681 furin cleavage site mutation found in both B.1.1.519 and B.1.617.2.

DISCUSSION

In this study, we observe that vaccination and natural SARS-CoV-2 infection elicit neutralizing antibody responses that are most potent against variants that bear spike mutations present in the immunizing exposure. This trend is exemplified by variants with mutations at the spike E484 position, which were all neutralized more effectively by E484K-exposed serum than other serum types. Importantly, we also show that B.1.617.2 (Delta) is neutralized more effectively by serum elicited by prior exposure to two different variants — B.1.429 and B.1.1.519 — which have separate subsets of spike mutations overlapping with mutations in B.1.617.2. Given that different regions throughout the world have experienced variable transmission of different variants prior to the dominance of B.1.617.2, these results suggest that acquired immunity in the population will differ significantly depending on the previous prevalence of variants in each region. Furthermore, these results demonstrate that specificity is strongest for serum neutralizing variants fully homologous to the exposure, but even single shared spike mutations, particularly those in highly antigenic regions such as the RBD, can enhance cross-neutralization.

This study also demonstrates the effect of serial exposure to two different versions of spike on neutralizing antibody response. Infection with B.1.429 (Epsilon) followed by vaccination led to greater cross-neutralization of B.1.429 and B.1.617.2 (Delta) compared to vaccination alone or D614G infection plus vaccination, supporting the notion that exposure to multiple spike variants expands neutralization specificity. A limitation of this study is the relatively small number of serum samples, however the shift in neutralization titer between D614G and variant pseudoviruses shows strong consistency between samples. These serology data leverage human exposures to an array of naturally occurring spike mutations, including those relevant to B.1.617.2, providing a real-world complement to previous animal studies investigating heterologous boosting or multivalent vaccination strategies [14,15]. Together, these results contribute additional evidence suggesting that heterologous boosting strategies may be an important and effective measure to address newly emergent variants with the potential to evade existing acquired immunity. Future studies investigating immune responses to additional emerging variants in vaccinated and unvaccinated individuals will contribute to identifying spike antigen versions that elicit broadly neutralizing antibody responses.

Funding

This work was supported by the University of California San Francisco COVID fund [to J.D., M.L., J.L., and S.S.]; the National Institutes of Health [grant number UM1AI069496 to D.H.; and grant number F31AI150007 to S.S.]; the Chan Zuckerberg Biohub [to J.D. and D.H.]; and the Chan Zuckerberg Initiative [to J.D. and D.H.].

Acknowledgements

We would like to thank Dr. Chuka Didigu, Dorothy Park CRNA, Salu Ribeiro, and Bay Area Phlebotomy and Laboratory services for performing blood draws of study subjects. We thank Dr. Andreas Puschnik for providing the engineered cell line used in this study. We thank Susana Elledge and Dr. James Wells for providing reagents and advice on antibody assays. We thank Drs. Peter Kim, Don Ganem, Sandy Schmidt, and Cori Bargmann for technical assistance and discussion.

Potential conflicts

Dr. DeRisi is a member of the scientific advisory board of The Public Health Company, Inc., and is scientific advisor for Allen & Co. Dr. DeRisi also reports options granted for service on the Scientific Advisory Board of The Public Health Company. None of the other authors have any potential conflicts.

REFERENCES

1. CDC. Coronavirus Disease 2019 (COVID-19) [Internet]. Centers for Disease Control and Prevention. 2020 [cited 2021 Sep 1]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html>
2. Teysou E, Delagrèverie H, Visseaux B, et al. The Delta SARS-CoV-2 variant has a higher viral load than the Beta and the historical variants in nasopharyngeal samples from newly diagnosed COVID-19 patients. *J Infect.* **2021**; :S0163-4453(21)00416–3.
3. Greaney AJ, Starr TN, Barnes CO, et al. Mapping mutations to the SARS-CoV-2 RBD that escape binding by different classes of antibodies. *Nat Commun.* **2021**; 12(1):4196.
4. Shen X, Tang H, Pajon R, et al. Neutralization of SARS-CoV-2 Variants B.1.429 and B.1.351. *New England Journal of Medicine.* Massachusetts Medical Society; **2021**; 384(24):2352–2354.
5. Liu J, Liu Y, Xia H, et al. BNT162b2-elicited neutralization of B.1.617 and other SARS-CoV-2 variants. *Nature.* **2021**; 596(7871):273–275.
6. Corbett KS, Nason MC, Flach B, et al. Immune correlates of protection by mRNA-1273 vaccine against SARS-CoV-2 in nonhuman primates. *Science* [Internet]. American Association for the Advancement of Science; **2021**
7. Liu C, Ginn HM, Dejnirattisai W, et al. Reduced neutralization of SARS-CoV-2 B.1.617 by vaccine and convalescent serum. *Cell.* **2021**; 184(16):4220-4236.e13.
8. Peng J, Liu J, Mann SA, et al. Estimation of secondary household attack rates for emergent spike L452R SARS-CoV-2 variants detected by genomic surveillance at a community-based testing site in San Francisco. *Clin Infect Dis.* **2021**; :ciab283.

9. Chamie G, Marquez C, Crawford E, et al. SARS-CoV-2 Community Transmission disproportionately affects Latinx population during Shelter-in-Place in San Francisco. *Clin Infect Dis*. **2020**; :ciaa1234.
10. Kalantar KL, Carvalho T, de Bourcy CFA, et al. IDseq—An open source cloud-based pipeline and analysis service for metagenomic pathogen detection and monitoring. *GigaScience* [Internet]. **2020** [cited 2021 Jul 14]; 9(10).
11. Elledge SK, Zhou XX, Byrnes JR, et al. Engineering luminescent biosensors for point-of-care SARS-CoV-2 antibody detection. *Nat Biotechnol*. **2021**; :1–8.
12. Hoffmann M, Kleine-Weber H, Pöhlmann S. A Multibasic Cleavage Site in the Spike Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. *Mol Cell*. **2020**; 78(4):779-784.e5.
13. Wang R, Simoneau CR, Kulsuptrakul J, et al. Genetic Screens Identify Host Factors for SARS-CoV-2 and Common Cold Coronaviruses. *Cell*. **2021**; 184(1):106-119.e14.
14. Wu K, Choi A, Koch M, et al. Variant SARS-CoV-2 mRNA vaccines confer broad neutralization as primary or booster series in mice. *bioRxiv*. **2021**; :2021.04.13.439482.
15. Corbett KS, Gagne M, Wagner DA, et al. Protection against SARS-CoV-2 Beta Variant in mRNA-1273 Boosted Nonhuman Primates. *bioRxiv*. **2021**; :2021.08.11.456015.

Figure 1

Neutralization of D614G and B.1.429 pseudoviruses by serum from individuals with different exposures




Figure 1. Plot of 50% and 90% pseudovirus neutralization titers (NT₅₀ and NT₉₀) of serum samples obtained from donors with the indicated infection and/or vaccination exposures. Grey lines connect neutralization titer values for D614G (black dots) and B.1.429 (blue dots) pseudoviruses within each individual serum sample. Geometric mean neutralization titers for each serum group are marked with red lines and fold-change in NT₅₀ and NT₉₀ between D614G and B.1.429 pseudoviruses is shown along with P-value. Dark grey shading marks the interquartile range of titer values in each group and light grey shading marks the 10th-90th percentile of the range. P-values were calculated with a Wilcoxon matched-pairs signed-rank test.

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Figure 2

Change in variant pseudovirus neutralization titer relative to D614G




Figure 2. Matrix of normalized neutralization titers for seven different variant pseudoviruses (rows) neutralized by seven different pools of individual sera grouped by exposure (columns). Data is represented as a heat map of the Log₂ fold-change in NT₅₀ (top left of each box) and NT₉₀ (bottom right of each box) of each variant relative to D614G pseudovirus. All serum samples were collected at least 14 days after the date of the subject’s positive COVID-19 test or date of second vaccine dose. All titer measurements are the mean of at least three independent experiments, each performed with two technical replicates. Positive Log₂ fold-change (blue) indicates an increase in neutralization titer for that variant relative to D614G pseudovirus, while negative Log₂ fold-change (red) indicates a decrease relative to D614G. Statistical significance was determined with unpaired t-tests. All values are statistically significant (P-value < 0.05) except where noted with “ns” to indicate the difference in variant neutralization titer is not significantly different from D614G titer in that serum pool.