

1 **Long-term air pollution exposure and diabetes risk in American older adults: a national**  
2 **cohort study.**

3  
4 **Running title: air pollution and diabetes**

5  
6 Maayan Yitshak Sade, Ph.D. <sup>1</sup>, Liuhua Shi, Sc.D. <sup>2</sup>, Elena Colicino, Ph.D. <sup>1</sup>, Heresh Amini,  
7 Ph.D. <sup>3</sup>, Joel D Schwartz, Ph.D. <sup>4</sup>, Qian Di, Ph.D. <sup>5</sup>, and Robert O Wright, MD MPH. <sup>1</sup>.

8  
9 <sup>1</sup> Icahn School of Medicine at Mount Sinai, Department of Environmental Medicine and Public  
10 Health, New York, NY, USA.

11 <sup>2</sup> Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory  
12 University, Atlanta, Georgia, USA.

13 <sup>3</sup> Department of Public Health, University of Copenhagen, Copenhagen, Denmark

14 <sup>4</sup> Exposure, Epidemiology, and Risk Program, Department of Environmental Health, Harvard  
15 T.H. Chan School of Public Health, Boston, MA, USA.

16 <sup>5</sup> Vanke School of Public Health, Tsinghua University, Beijing, China.

17  
18  
19 **Corresponding author:**

20 Maayan Yitshak Sade

21 Department of Environmental Medicine and Public Health

22 Icahn School of Medicine at Mount Sinai

23 New York, New York, United States

24 [maayan.yitshak-sade@mssm.edu](mailto:maayan.yitshak-sade@mssm.edu)

29 **ABSTRACT**

30 **Background:** Type 2 diabetes is a major public health concern. Studies show that both genetics  
31 and environmental exposures play a significant role in developing the disease. We assessed the  
32 association between air pollution and first documented diabetes occurrence in a national U.S.  
33 cohort of older adults to estimate diabetes risk.

34 **Methods:** We included all Medicare enrollees 65 years and older in the fee-for-service program,  
35 part A and part B, in the contiguous United States (2000-2016). Participants were followed  
36 annually until the first recorded diabetes diagnosis, end of enrollment, or death (264,869,458  
37 person-years). We obtained annual estimates of fine particulate matter (PM<sub>2.5</sub>), nitrogen dioxide  
38 (NO<sub>2</sub>), and warm-months ozone (O<sub>3</sub>) exposures from highly spatiotemporally resolved prediction  
39 models. We assessed the simultaneous effects of the pollutants on diabetes risk using survival  
40 analyses. We repeated the models in cohorts restricted to ZIP codes with air pollution levels not  
41 exceeding the national ambient air quality standards (NAAQS) during the study period.

42 **Results:** We identified 10,024,879 diabetes cases of 41,780,637 people (3.8% of person-years).  
43 The hazard ratio (HR) for first diabetes occurrence was 1.074 (95% CI 1.058; 1.089) for 5 µg/m<sup>3</sup>  
44 increase in PM<sub>2.5</sub>, 1.055 (95% CI 1.050; 1.060) for 5 ppb increase in NO<sub>2</sub>, and 0.999 (95% CI  
45 0.993; 1.004) for 5 ppb increase in O<sub>3</sub>. Both for NO<sub>2</sub> and PM<sub>2.5</sub> there was evidence of non-linear  
46 exposure-response curves with stronger associations at lower levels (NO<sub>2</sub> ≤ 40 ppb, PM<sub>2.5</sub> ≤ 9  
47 µg/m<sup>3</sup>). Furthermore, associations remained in the restricted low-level cohorts. The O<sub>3</sub>-diabetes  
48 exposure-response relationship differed greatly between models and require further investigation.

49 **Conclusion:** Exposures to PM<sub>2.5</sub> and NO<sub>2</sub> are associated with increased diabetes risk, even when  
50 restricting the exposure to levels below the NAAQS set by the U.S. EPA.

51 **Keywords:** air pollution; diabetes; PM<sub>2.5</sub>, NO<sub>2</sub>, O<sub>3</sub>

## 52 INTRODUCTION

53 Type 2 diabetes (T2D) is a major public health concern rising rapidly, with the number of people  
54 diagnosed with the disease worldwide more than doubling in the past 20 years <sup>1</sup>. T2D may cause  
55 major complications, including blindness, cardiovascular damage, and premature mortality <sup>2</sup>, it is  
56 therefore important to identify determinants of the disease.

57 Like many chronic conditions, type I diabetes (T1D) and T2D have a genetic component, but  
58 genetics alone explains only a small portion of the variance. Studies show that in T2D, which  
59 represents over 99% of diabetes cases globally <sup>3</sup>, genetics and environmental exposures play a  
60 significant role <sup>4</sup>. In the last few decades, environmental research focused primarily on  
61 behavioral factors, such as inactivity and diet. However, in recent years, cardiometabolic risk  
62 was linked to various environmental components, including the social environment <sup>5,6</sup>,  
63 environmental pollutants <sup>3,7</sup>, and, more specifically, air pollution <sup>8-13</sup>. A recent meta-analysis  
64 concluded that each ten  $\mu\text{g}/\text{m}^3$  increase in long-term fine particulate matter (PM<sub>2.5</sub>) exposure was  
65 associated with an 8% increase in T2D prevalence risk, and each ten  $\mu\text{g}/\text{m}^3$  increase in long-term  
66 nitrogen dioxide (NO<sub>2</sub>) exposure was associated with a 7% increase in T2D prevalence. O<sub>3</sub> was  
67 not included in that analysis since the number of studies was limited <sup>14</sup>.

68 These positive associations were observed in other systematic reviews as well <sup>15,16</sup>. However,  
69 since it is challenging to investigate the effects of the environment on chronic conditions, current  
70 research has many limitations. These limitations include potential bias in the outcome and  
71 exposure assessment, inability to separate incidence and prevalence cases, healthy survivor bias,  
72 and the inability to account for cardiometabolic risk factors and co-morbidities <sup>15</sup>.

73 Our study addresses many of these limitations by investigating the association between  
74 particulate and gaseous air pollution and first documented diabetes. Our use of highly  
75 spatiotemporally resolved exposure models reduces bias due to exposure misclassification. In  
76 addition, unlike most national studies or chronic air pollution effects, we use comprehensive  
77 chronic conditions data on Medicare enrollees across the U.S. to reduce misclassification of  
78 diabetes and approximate incident cases.

79 In this study, we aim to assess the association between long-term air pollution exposure and first  
80 documented diabetes occurrence, focusing on the effect of air pollution exposure below the U.S.  
81 Environmental Protection Agency (EPA) national ambient air quality standards (NAAQS).

## 82 **METHODS**

### 83 **Study population**

84 We included all Medicare enrollees who were 65 years and older in the fee-for-service (FFS)  
85 program, part A (hospital insurance) and part B (medical insurance), in the contiguous U.S.  
86 between the years 2000-2016. We limited the data to person-years included in these programs  
87 because the algorithm used to identify chronic conditions utilizes claims covered by these three  
88 programs. We entered participants into the cohort on January 1 after they became Medicare  
89 participants and followed participants for each calendar year until the first recorded diabetes  
90 diagnosis, end of enrollment in either of the mentioned Medicare programs, or death – whichever  
91 came first. To avoid gaps in follow-up, once enrollment in the FFS, Medicare part A or B  
92 programs was terminated, those participants were no longer included in the cohort even if they  
93 renewed their enrollment in later years.

94 To better approximate incident cases, we excluded individuals diagnosed with diabetes before or  
95 in their first year of enrollment. We obtained information on diabetes status from the chronic  
96 conditions warehouse database. Diabetes was identified using an algorithm that incorporates  
97 claims indicating that an individual received a healthcare service for diabetes. The algorithm  
98 combines inpatient, outpatient, skilled nursing facilities, home health claims, or carrier claims  
99 (primarily doctor visits)<sup>17</sup>. To be diagnosed with diabetes, beneficiaries must have at least one  
100 claim with the international classification of disease codes of diabetes (ICD-9 or ICD-10) from  
101 either inpatient, skilled nursing facility, home health, or part B services within two years.

102 This study was approved by the Centers for Medicare & Medicaid Services (CMS) under the  
103 data use agreement (#RSCH-2020-55733), the Institutional Review Board of Emory University  
104 (#STUDY00000316), and the Institutional Review Board of Mount Sinai (STUDY 20-01344),  
105 and a waiver of informed consent was granted. The Medicare dataset was stored and analyzed in  
106 the Rollins High-Performance Computing (HPC) Cluster at Emory University, in compliance  
107 with Health Insurance Portability and Accountability Act (HIPAA).

## 108 **Exposures**

109 Our study was focused on three principal air pollutants linked to cardiometabolic health: PM<sub>2.5</sub>,  
110 O<sub>3</sub>, and NO<sub>2</sub>. PM<sub>2.5</sub> are fine inhalable particles, smaller than 2.5 micrometers, comprising a  
111 mixture of solids and liquids. NO<sub>2</sub> is an air pollutant that originates mostly from traffic and high-  
112 temperature combustion processes<sup>18</sup>. O<sub>3</sub> at the ground level is formed naturally and following a  
113 chemical reaction - where air pollutants emitted from sources such as traffic, industry, and  
114 wildfires (i.e., nitrogen oxides and volatile organic compounds) react with sunlight and organic  
115 gases, principally from vegetation<sup>19</sup>.

116 We obtained predictions of PM<sub>2.5</sub> (24-hour average,  $\mu\text{g}/\text{m}^3$ ), NO<sub>2</sub> (daily 1-hour maximum, ppb)  
117 and ozone (daily maximum of 8-hour average, ppb) exposures from validated prediction models  
118 calibrated to measurements at approximately 2000 monitoring stations using an ensemble of  
119 three machine learners (neural network, random forest, and gradient boosting) that provided  
120 daily estimates for a 1 km<sup>2</sup> grid of the contiguous U.S.<sup>20-22</sup>. In brief, each machine learning  
121 algorithm incorporated more than 100 predictor variables from satellite data, land-use  
122 information, weather data, and chemical transport model simulations. The predictions of the  
123 three learners were then integrated using a generalized additive model-based geographically  
124 weighted-averaging technique. The model was calibrated using daily pollutants concentrations  
125 measured at EPA monitoring sites and demonstrated excellent model performance (average  
126 cross-validation R<sup>2</sup>=0.89, 0.84, and 0.86 for annual predictions of PM<sub>2.5</sub>, NO<sub>2</sub>, and O<sub>3</sub>,  
127 respectively). To align with the Medicare data, we aggregated gridded exposures to ZIP codes by  
128 averaging predictions of grid cells within each ZIP code annually. For ozone, we averaged the  
129 exposure only during the warmer months of each year (May – October), a commonly specified  
130 time window to examine associations with health outcomes<sup>23</sup>. We selected this window because  
131 ozone formation is increased in warmer months due to its reaction with sunlight.

## 132 **Covariates**

133 The Medicare data is a dynamic cohort that includes individual-level information on the  
134 participants' sex, race, age, Medicaid eligibility, and date of death. We additionally obtained the  
135 following ZIP code level covariates from the U.S. Census: median household income, population  
136 density, percent home renters, percent of residents with no high school education, and percent of  
137 the population self-identified as black. We obtained data from available U.S. Census years and  
138 the annual American Community Survey, and extrapolated values for missing years. We

139 aggregated annual summer, and winter temperature ZIP code means from gridded Daymet 1- km  
140 models<sup>24</sup>.

## 141 **Statistical analysis**

142 We investigated the effect of the three air pollutants simultaneously using a cox-equivalent re-  
143 parameterized Poisson survival approach<sup>25</sup>. To allow for strata-specific baseline hazard  
144 functions, we stratified the models by follow-up year, calendar year, ZIP code, sex, race (white,  
145 black, other), age, and Medicaid insurance. We calculated diabetes counts in each follow-up  
146 year, calendar year, and ZIP code within strata specified by these individual characteristics, using  
147 the log of the corresponding total person-time as an offset. This approach is proven to be  
148 equivalent to a time-varying cox survival model using the Anderson-Gill formulation<sup>25</sup>. We used  
149 the m-out-n bootstrap method to account for spatial autocorrelation of neighboring ZIP codes.

150 This methods samples ZIP codes randomly for each bootstrap replicate, and estimates  
151 statistically robust confidence intervals<sup>26</sup>. First, we fit a multi-pollutant model using linear terms  
152 for the three pollutants. Then, to allow for nonlinear exposure-response curves, we used  
153 penalized spline functions for each of the pollutants. We fit three multipollutant models in which  
154 we used a penalized spline to estimate the exposure-response curve for one pollutant, and linear  
155 terms for the adjusted pollutants. We adjusted the models for annual summer and winter mean  
156 temperatures, population density, median household income, percent home renters, percent of  
157 residents without a high school diploma, and percent black population.

158 To assess the effect of air pollution concentrations below the NAAQS, we created three  
159 restricted low-exposure subsets comprised of individuals who were always exposed to exposure  
160 levels lower than the national standards within the study period (i.e.,  $PM_{2.5} < 12 \mu\text{g}/\text{m}^3$ ,  $NO_2 < 53$   
161 ppb). As there is no annual standard for  $O_3$ , we selected 50 ppb as a threshold to approximate the

162 current World Health Organization global interim target 1 for peak-season average  
163 O<sub>3</sub> concentration <sup>27</sup>.

#### 164 *Sensitivity analyses*

165 The exclusion of individuals who are not enrolled in the FFS, part A, and Part B programs can  
166 potentially induce a selection bias in the study. Biased results can also occur due to competing  
167 mortality risks. To avoid selection bias, we used inverse probability weights. Probabilities of  
168 enrollment in the cohort and probability of not dying were modeled, accounting for the subjects'  
169 age, sex, race, Medicaid eligibility, and the ZIP code population density, percent population  
170 under the poverty line, percent population without a high school diploma, percent black and  
171 Hispanic population. We calculated these probabilities using pooled logistic regressions. We  
172 then calculated the weight by multiplying the inverse probability of enrolling in the three  
173 programs with the inverse probability of being alive and calculated the averaged weight within  
174 each stratum of follow up year, calendar year, ZIP code, and individual characteristics. We then  
175 repeated the models incorporating the weights and compared the exposure-response curves with  
176 and without weights.

## 177 **RESULTS**

178 We have included 264,869,458 person-years of 41,780,637 people. The mean age was  
179 approximately 76 years, 60% were women, and 90% were white. We observed 10,024,879  
180 diabetes cases (3.8% of person-years) (**Table 1**). The mean and IQR values of the air pollutants  
181 were as follows: PM<sub>2.5</sub> 10.1 µg/m<sup>3</sup> (4.2 µg/m<sup>3</sup>), O<sub>3</sub> 43.2 ppb (7.0 ppb), and NO<sub>2</sub> 18.9 ppb (13.7  
182 ppb). The correlation between the exposures was low to moderate (**Supplementary Table 1**),  
183 with the highest correlation observed between PM<sub>2.5</sub> and NO<sub>2</sub> (r=0.44).



184 Using linear terms, we observed an increased diabetes risk associated with 5  $\mu\text{g}/\text{m}^3$  increase in  
185  $\text{PM}_{2.5}$  (HR=1.074, 95% CI 1.058; 1.089) and 5 ppb increase in  $\text{NO}_2$  (HR=1.055, 95% CI 1.050;  
186 1.060). No linear association was observed with  $\text{O}_3$  (HR 0.999, 95% CI 0.993; 1.004). We then  
187 used penalized spline functions to allow for nonlinear exposure-response curves and found  
188 evidence of nonlinear associations with the three pollutants. Both for  $\text{NO}_2$  and  $\text{PM}_{2.5}$  we observed  
189 stronger associations at the lower ends of the pollutants' distributions. For  $\text{PM}_{2.5}$  the associations  
190 were stronger for levels  $< 8.2 \mu\text{g}/\text{m}^3$  (the 25<sup>th</sup> percentile), followed by a plateaued association. For  
191  $\text{NO}_2$  the associations were stronger for levels  $< 36$  ppb (94<sup>th</sup> percentile), followed by attenuated  
192 associations. The  $\text{O}_3$  exposure-response curve was highly nonlinear with mostly negative  
193 associations (**Figure 1**). The associations with  $\text{NO}_2$  and  $\text{PM}_{2.5}$  were similar in the single pollutant  
194 models and the multivariate model. The  $\text{O}_3$  exposure-response curve differed greatly between the  
195 single and multipollutant models. Unlike the negative associations seen in the multivariate  
196 model, there was no clear trend in the association with the pollutant in the single pollutant model  
197 (**Supplementary Figure 1**).

198 Of the analytical dataset, 33.7% (89,519,417 person-years), 96.4% (255,575,750 person-years),  
199 and 54.4% (144,225,079 person-years) were only exposed to low-level annual  $\text{PM}_{2.5}$ , annual  
200  $\text{NO}_2$ , and warm-season  $\text{O}_3$  during the study period, respectively. High-pollution ZIP codes  
201 excluded from the restricted datasets were mostly located in the mid-west, southeast, and the  
202 mid-Atlantic U.S. regions. We estimated the nonlinear exposure-response curves for the three  
203 pollutants in multivariate regressions in these restricted cohorts, and the associations with  $\text{PM}_{2.5}$   
204 and  $\text{NO}_2$  remained similar and significant. For  $\text{O}_3$ , there was no clear association with diabetes  
205 and the direction of the associations differed across the distribution of the pollutant (**Figure 2**).

206 **Supplementary Table 2** shows the baseline characteristics of those included and excluded from  
207 the analytical dataset. Person-years included in the analytical dataset were of older people. In  
208 addition, the proportion of whites and women was larger, and the proportion of death was much  
209 smaller. We conducted a sensitivity analysis and repeated the models including inverse  
210 probability weights accounting for the probability of being enrolled in the cohort and alive. The  
211 results of the two models were very similar (**Supplementary Figure 2**).

## 212 **DISCUSSION**

213 In a large-scale, national cohort, we observed higher risks for diabetes associated with increases  
214 in NO<sub>2</sub> and PM<sub>2.5</sub> exposures, even when restricting the data to exposure levels below the  
215 NAAQS set by the U.S. EPA. Results regarding the association with O<sub>3</sub> were inconclusive.

216 T2D is characterized by high blood glucose levels and increased insulin resistance leading to  
217 vascular damage and metabolic dysfunctions<sup>29</sup>. A growing body of literature links ambient air  
218 pollution exposure to diabetes risk, as concluded in many recent systematic reviews<sup>15,16</sup>. Studies  
219 also found associations between air pollution exposure and markers of potential underlying  
220 pathways of this association: lower insulin sensitivity<sup>30</sup>, and higher fasting glucose<sup>30,31</sup>.

221 Different mechanisms were proposed to explain the links between air pollution exposure and  
222 T2D. Oxidative stress is considered a major pathway to this association<sup>3</sup>. PM exposure is  
223 associated with oxidative stress, which can lead to lipid peroxidation, reduction of antioxidants,  
224 and activation of pro-inflammatory processes<sup>9,32</sup>. These, in turn, have a major role in the  
225 development and progression of metabolic syndrome and diabetes in particular<sup>33,34</sup>. Another  
226 suggested pathway is PM-induced mitochondrial dysfunction, which decreases brown adipose  
227 tissue<sup>35</sup>. Since stimulated brown adipose tissue regulates glucose homeostasis and insulin  
228 resistance using glucose and free fatty acids during thermogenesis<sup>36,37</sup>, this may systematically

229 interfere with glucose metabolism and insulin sensitivity. Studies also found indications of  
230 oxidative stress and mitochondrial dysfunction following NO<sub>2</sub> and O<sub>3</sub> exposure<sup>38-41</sup>.

231 Our study shows evidence of increased diabetes risk associated with long-term PM<sub>2.5</sub> and NO<sub>2</sub>  
232 exposures. Like our findings, a 2019 meta-analysis concluded that the odds for T2D were 1.03  
233 and 1.05 times higher for incremental increases of 10 µg/m<sup>3</sup> in PM<sub>2.5</sub> and NO<sub>2</sub> exposures,  
234 respectively<sup>16</sup>. Studies assessing the association between O<sub>3</sub> and diabetes are much more  
235 limited, and the direction of the observed associations is inconsistent and varies greatly  
236 depending on the model covariates<sup>42</sup>. For example, a longitudinal cohort of 13,548 individuals  
237 in China found a 5.7% increase in diabetes incidence hazard associated with a ten µg/m<sup>3</sup> increase  
238 in annual O<sub>3</sub> exposure<sup>43</sup>. However, two recent long-term studies did not find a significantly  
239 increased diabetes risk associated with O<sub>3</sub> exposure in multipollutant models accounting for  
240 particulate matter<sup>44,45</sup> and NO<sub>2</sub><sup>44</sup> exposures. Moreover, a 2021 review concluded that evidence  
241 on the association between O<sub>3</sub> exposure and diabetes is insufficient to infer causality<sup>42</sup>. Our  
242 study also showed inconclusive results, possibly related to residual confounding, exposure  
243 measurement error, or the complexities of simultaneously estimating the effects of multiple air  
244 pollutants.

245 The Clean Air Act was last amended in 1990 and requires the U.S. EPA to set NAAQS that  
246 mitigate any harmful consequences of air pollution to human health and the environment<sup>28</sup>.

247 However, recent studies suggest that these thresholds are insufficient to protect human health.  
248 There is evidence of increased mortality risk associated with air pollution levels below the  
249 NAAQS<sup>46-51</sup>. In some analyses, the associations observed at the lower exposure distribution  
250 range were stronger than the higher range of the exposure distribution<sup>50,51</sup>. This may be  
251 attributed to larger measurement error at the rarer higher exposure concentrations that is likely to

252 attenuate the overall effect observed when considering the full ranges of the pollutant exposures  
253 in the analysis<sup>52</sup>. Regarding PM<sub>2.5</sub>, it is also possible that the composition of the particles is  
254 different for low and high pollution days<sup>46</sup>.

255 We limited our study sample to ZIP codes with lower air pollution levels to simulate a scenario  
256 where air pollution levels never exceeded thresholds considered safe by the U.S. EPA. The major  
257 findings of our study are the nonlinear associations and the harmful air pollution effects observed  
258 even from levels below the NAAQS set by the EPA. These results suggest that NO<sub>2</sub>, and PM<sub>2.5</sub>  
259 exposures below the NAAQS are associated with increased cardiometabolic risk. Like our  
260 findings, Paul et al.<sup>53</sup> also observed associations between moderate exposure concentrations of  
261 PM<sub>2.5</sub> and NO<sub>2</sub> and diabetes mortality and incidence risks. Moreover, they report greater PM<sub>2.5</sub>  
262 effects on diabetes incidence risk in lower pollution concentrations.

263 Our study has several limitations. First, there are differences between people included and  
264 excluded from the cohort. This is a limitation of all studies that analyze claims data of the  
265 Medicare cohort. However, our sensitivity analysis suggests that our results were not biased due  
266 to differential probabilities of enrollment or death. Second, the Medicare data does not provide  
267 information on subtypes of diabetes or individual confounders such as BMI, smoking, physical  
268 activity, or lifestyle. However, since exposure is assigned on a ZIP code level, neighborhood  
269 factors are more likely to confound the associations in our study. These neighborhood factors  
270 were accounted for in our models. Finally, we might have had exposure misclassification errors  
271 like other air pollution studies. However, the use of highly spatiotemporally resolved exposure  
272 models reduces this error.

273 In conclusion, assessing the simultaneous effects of particulate and gaseous air pollutants in a  
274 national cohort, we found increased diabetes risk associated with PM<sub>2.5</sub> and NO<sub>2</sub> exposure. The

275 observed effects remained when restricting the data to exposure levels below the NAAQS. For  
276 ozone, the effects were inconclusive and require further investigation. Since current studies of  
277 the link between air pollution and diabetes are scarce and often limited in quality or sample size,  
278 this national study may add robust evidence important for inferring the causal link between air  
279 pollution exposure and the development of diabetes.

## 280 **ACKNOWLEDGEMENTS**

281 This study was supported by the HERCULES Center (P30 ES019776), the Mount Sinai  
282 transdisciplinary center on early environmental exposures (P30 ES023515), the National Institute  
283 on Aging (NIA/NIH R01 AG074357), the National Institute of Environmental Health Sciences  
284 (R21 ES032606, R01 ES032242, 5U2CES026555-03, and R01 ES013744, P30 ES000002, R01  
285 ES032418), and the United States Environmental Protection Agency (US EPA) (RD-83587201).  
286 Its contents are solely the responsibility of the grantee and do not necessarily represent the  
287 official views of the US EPA. Furthermore, the US EPA does not endorse the purchase of any  
288 commercial products or services mentioned in the publication. Finally, H.A. is supported by  
289 Novo Nordisk Foundation Challenge Programme: Harnessing the Power of Big Data to Address  
290 the Societal Challenge of Aging (NNF17OC0027812).

## 291 **CONFLICT OF INTEREST STATEMENT**

292 The authors declare that they have no known competing financial interests or personal  
293 relationships that could have influenced the work reported in this paper.

294

295

296 **REFERENCES**

297 1. Zimmet PZ, Magliano DJ, Herman WH, Shaw JE. Diabetes: a 21st century challenge.  
 298 *The lancet Diabetes & endocrinology*. 2014;2(1):56-64.

299 2. Papatheodorou K, Papanas N, Banach M, Papazoglou D, Edmonds M. Complications of  
 300 Diabetes 2016. *J Diabetes Res*. 2016;2016:6989453.

301 3. Rajagopalan S, Brook RD. Air pollution and type 2 diabetes: mechanistic insights.  
 302 *Diabetes*. 2012;61(12):3037-3045.

303 4. Kaprio J, Tuomilehto J, Koskenvuo M, et al. Concordance for type 1 (insulin-dependent)  
 304 and type 2 (non-insulin-dependent) diabetes mellitus in a population-based cohort of  
 305 twins in Finland. *Diabetologia*. 1992;35(11):1060-1067.

306 5. Khan SU, Javed Z, Lone AN, et al. Social Vulnerability and Premature Cardiovascular  
 307 Mortality Among US Counties, 2014 to 2018. *Circulation*. 2021;144(16):1272-1279.

308 6. Basile Ibrahim B, Barcelona V, Condon EM, Crusto CA, Taylor JY. The Association  
 309 Between Neighborhood Social Vulnerability and Cardiovascular Health Risk Among  
 310 Black/African American Women in the InterGEN Study. *Nurs Res*. 2021;70(5S Suppl  
 311 1):S3-s12.

312 7. Dendup T, Feng X, Clingan S, Astell-Burt T. Environmental Risk Factors for Developing  
 313 Type 2 Diabetes Mellitus: A Systematic Review. *Int J Environ Res Public Health*.  
 314 2018;15(1).

315 8. Brook RD. Cardiovascular effects of air pollution. *Clinical Science*. 2008;115(5-6):175-  
 316 187.

317 9. Brook RD, Rajagopalan S, Pope CA, III, et al. Particulate Matter Air Pollution and  
 318 Cardiovascular Disease An Update to the Scientific Statement From the American Heart  
 319 Association. *Circulation*. 2010;121(21):2331-2378.

320 10. Andersen ZJ, Raaschou-Nielsen O, Ketzler M, et al. Diabetes incidence and long-term  
 321 exposure to air pollution: a cohort study. *Diabetes care*. 2012;35(1):92-98.

322 11. Chandrabose M, Rachele JN, Gunn L, et al. Built environment and cardio-metabolic  
 323 health: systematic review and meta-analysis of longitudinal studies. *Obes Rev*.  
 324 2019;20(1):41-54.

325 12. Li Y, Xu L, Shan Z, Teng W, Han C. Association between air pollution and type 2  
 326 diabetes: an updated review of the literature. *Ther Adv Endocrinol Metab*.  
 327 2019;10:2042018819897046.

328 13. Park SK, Wang W. Ambient Air Pollution and Type 2 Diabetes Mellitus: A Systematic  
 329 Review of Epidemiologic Research. *Curr Envir Health Rpt*. 2014;1:275-286.

330 14. Yang BY, Fan S, Thiering E, et al. Ambient air pollution and diabetes: A systematic  
 331 review and meta-analysis. *Environ Res*. 2020;180:108817.

332 15. Eze CI, Hemkens GL, Bucher CH, et al. Association between Ambient Air Pollution and  
 333 Diabetes Mellitus in Europe and North America: Systematic Review and Meta-Analysis.  
 334 *Environmental Health Perspectives*. 2015.

335 16. Liu F, Chen G, Huo W, et al. Associations between long-term exposure to ambient air  
 336 pollution and risk of type 2 diabetes mellitus: A systematic review and meta-analysis.  
 337 *Environ Pollut*. 2019;252(Pt B):1235-1245.

338 17. Center for Medicare & Medicaid services. Chronic condition algorithms.  
 339 <https://www2.ccwdata.org/web/guest/condition-categories>. Published 2015. Accessed  
 340 Sep 15, 2020.

- 341 18. Lim SS, Vos T, Flaxman AD. A comparative risk assessment of burden of disease and  
342 injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a  
343 systematic analysis for the Global Burden of Disease Study 2010 (vol 380, pg 2224,  
344 2012). *Lancet*. 2013;381(9874):1276-1276.
- 345 19. Lange SS, Mulholland SE, Honeycutt ME. What Are the Net Benefits of Reducing the  
346 Ozone Standard to 65 ppb? An Alternative Analysis. *Int J Environ Res Public Health*.  
347 2018;15(8).
- 348 20. Di Q, Amini H, Shi L, et al. Assessing NO<sub>2</sub> Concentration and Model Uncertainty with  
349 High Spatiotemporal Resolution across the Contiguous United States Using Ensemble  
350 Model Averaging. *Environ Sci Technol*. 2020;54(3):1372-1384.
- 351 21. Requia WJ, Di Q, Silvern R, et al. An Ensemble Learning Approach for Estimating High  
352 Spatiotemporal Resolution of Ground-Level Ozone in the Contiguous United States.  
353 *Environ Sci Technol*. 2020;54(18):11037-11047.
- 354 22. Di Q, Amini H, Shi L, et al. An ensemble-based model of PM<sub>2.5</sub> concentration across  
355 the contiguous United States with high spatiotemporal resolution. *Environ Int*.  
356 2019;130:104909.
- 357 23. Wei Y, Yazdi MD, Di Q, et al. Emulating causal dose-response relations between air  
358 pollutants and mortality in the Medicare population. *Environ Health*. 2021;20(1):53.
- 359 24. NASA Distributed Active Archive Center (DAAC) at NSIDC (DAAC). Daymet: Daily  
360 Surface Weather Data on a 1-km Grid for North America, Version 3.  
361 [https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds\\_id=1328](https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1328). Published 2020. Accessed August  
362 19, 2021.
- 363 25. Shi L, Wu X, Danesh Yazdi M, et al. Long-term effects of PM<sub>2.5</sub> on neurological  
364 disorders in the American Medicare population: a longitudinal cohort study. *Lancet*  
365 *Planet Health*. 2020;4(12):e557-e565.
- 366 26. Bickel P, Götze F. Selected Works of Willem van Zwet. 2012.
- 367 27. Organization WH. *WHO Air quality guidelines for particulate matter, ozone, nitrogen*  
368 *dioxide and sulfur dioxide*. World Health Organization;2005.
- 369 28. United States Environmental Protection Agency. National ambient air quality standards  
370 (NAAQS) table. <https://www.epa.gov/criteria-air-pollutants/naaqs-table#3>. Published  
371 2021. Accessed July 15, 2021.
- 372 29. Cooper-DeHoff RM, Pepine CJ. Metabolic syndrome and cardiovascular disease:  
373 challenges and opportunities. *Clin Cardiol*. 2007;30(12):593-597.
- 374 30. Wolf K, Popp A, Schneider A, et al. Association Between Long-term Exposure to Air  
375 Pollution and Biomarkers Related to Insulin Resistance, Subclinical Inflammation, and  
376 Adipokines. *Diabetes*. 2016;65(11):3314-3326.
- 377 31. Yitshak Sade M, Kloog I, Liberty I, Schwartz J, Novack V. The Association Between Air  
378 Pollution Exposure and Glucose and Lipids Levels. *The Journal of Clinical*  
379 *Endocrinology & Metabolism* 2016.
- 380 32. Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O. Oxidative stress and  
381 antioxidant defense. *World Allergy Organ J*. 2012;5(1):9-19.
- 382 33. Hutcheson R, Rocic P. The metabolic syndrome, oxidative stress, environment, and  
383 cardiovascular disease: the great exploration. *Exp Diabetes Res*. 2012;2012:271028.
- 384 34. Lim CC, Thurston GD. Air Pollution, Oxidative Stress, and Diabetes: a Life Course  
385 Epidemiologic Perspective. *Curr Diab Rep*. 2019;19(8):58.

- 386 35. Xu Z, Xu X, Zhong M, et al. Ambient particulate air pollution induces oxidative stress  
387 and alterations of mitochondria and gene expression in brown and white adipose tissues.  
388 *Particle and fibre toxicology*. 2011;8(1):1-14.
- 389 36. Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance.  
390 *Physiol Rev*. 2004;84(1):277-359.
- 391 37. Bartelt A, Bruns OT, Reimer R, et al. Brown adipose tissue activity controls triglyceride  
392 clearance. *Nat Med*. 2011;17(2):200-205.
- 393 38. Sourij H, Meinitzer A, Pilz S, et al. Arginine bioavailability ratios are associated with  
394 cardiovascular mortality in patients referred to coronary angiography. *Atherosclerosis*.  
395 2011;218(1):220-225.
- 396 39. Schooneman MG, Vaz FM, Houten SM, Soeters MR. Acylcarnitines: reflecting or  
397 inflicting insulin resistance? *Diabetes*. 2013;62(1):1-8.
- 398 40. Zhao YY, Wang HL, Cheng XL, et al. Metabolomics analysis reveals the association  
399 between lipid abnormalities and oxidative stress, inflammation, fibrosis, and Nrf2  
400 dysfunction in aristolochic acid-induced nephropathy. *Sci Rep*. 2015;5:12936.
- 401 41. Alderete TL, Chen Z, Toledo-Corral CM, et al. Ambient and Traffic-Related Air  
402 Pollution Exposures as Novel Risk Factors for Metabolic Dysfunction and Type 2  
403 Diabetes. *Curr Epidemiol Rep*. 2018;5(2):79-91.
- 404 42. LaKind JS, Burns CJ, Pottenger LH, Naiman DQ, Goodman JE, Marchitti SA. Does  
405 ozone inhalation cause adverse metabolic effects in humans? A systematic review. *Crit*  
406 *Rev Toxicol*. 2021;51(6):467-508.
- 407 43. Wang Y, Cao R, Xu Z, et al. Long-term exposure to ozone and diabetes incidence: A  
408 longitudinal cohort study in China. *Sci Total Environ*. 2022;816:151634.
- 409 44. Jerrett M, Brook R, White LF, et al. Ambient ozone and incident diabetes: a prospective  
410 analysis in a large cohort of African American women. *Environment international*.  
411 2017;102:42-47.
- 412 45. Li H, Duan D, Xu J, et al. Ambient air pollution and risk of type 2 diabetes in the  
413 Chinese. *Environ Sci Pollut Res Int*. 2019;26(16):16261-16273.
- 414 46. Shi L, Rosenberg A, Wang Y, et al. Low-Concentration Air Pollution and Mortality in  
415 American Older Adults: A National Cohort Analysis (2001-2017). *Environ Sci Technol*.  
416 2021.
- 417 47. Danesh Yazdi M, Wang Y, Di Q, et al. Long-Term Association of Air Pollution and  
418 Hospital Admissions Among Medicare Participants Using a Doubly Robust Additive  
419 Model. *Circulation*. 2021;143(16):1584-1596.
- 420 48. Shi L, Zanobetti A, Kloog I, et al. Low-Concentration PM<sub>2.5</sub> and Mortality: Estimating  
421 Acute and Chronic Effects in a Population-Based Study. *Environ Health Perspect*.  
422 2016;124(1):46-52.
- 423 49. Yitshak-Sade M, Lane KJ, Fabian MP, et al. Race or racial segregation? Modification of  
424 the PM<sub>2.5</sub> and cardiovascular mortality association. *PloS one*. 2020;15(7):e0236479.
- 425 50. Wei Y, Wang Y, Wu X, et al. Causal Effects of Air Pollution on Mortality Rate in  
426 Massachusetts. *Am J Epidemiol*. 2020;189(11):1316-1323.
- 427 51. Di Q, Dai L, Wang Y, et al. Association of Short-term Exposure to Air Pollution With  
428 Mortality in Older Adults. *Jama*. 2017;318(24):2446-2456.
- 429 52. Steenland K, Karnes C, Darrow L, Barry V. Attenuation of exposure-response rate ratios  
430 at higher exposures: a simulation study focusing on frailty and measurement error.  
431 *Epidemiology*. 2015;26(3):395-401.



- 432 53. Paul LA, Burnett RT, Kwong JC, et al. The impact of air pollution on the incidence of  
433 diabetes and survival among prevalent diabetes cases. *Environ Int.* 2020;134:105333.  
434

435 **Table 1.** Population characteristics (264,869,458 person years)  
 436

<b>Population characteristics</b>	<b>Summary statistics</b>
<b><u>Individual characteristics</u></b>	
Age, Mean (S.D.)	75.97 (7.7)
Female sex, n (%)	158,138,886 (59.7)
Race, n (%)	
White	238,995,206 (90.2)
Black	14,986,036 (5.7)
Other	10,888,216 (4.1)
Medicaid insurance, n (%)	24,986,082 (9.4)
Diabetes, n (%)	10,024,879 (3.8)
Hypertension, n (%)	96,411,807 (36.4)
Ischemic heart disease, n (%)	57,842,331 (21.8)
Death, n (%)	1,125,608 (0.4)
<b><u>ZIP code characteristics</u></b>	
Percent poverty, Mean (SD)	0.1 (0.1)
Population density, Mean (SD)	2,650.1 (7,854.1)
Median house value, Mean (SD)	195,301.5 (159,465.5)
Percent black population, Mean (SD)	0.1 (0.2)
Median household income, Mean (SD)	53,618.0 (22,608.6)
Percent black population, Mean (SD)	0.3 (0.1)

437

438 **FIGURE LEGENDS**

439

440 **Figure 1. The association between PM<sub>2.5</sub>, NO<sub>2</sub>, and warm-months O<sub>3</sub> exposure and diabetes**  
441 **occurrence.**

442 Figure 1 shows the exposure-response curves for the association between PM<sub>2.5</sub>, NO<sub>2</sub>, and warm-  
443 months O<sub>3</sub> and diabetes occurrence. We show the curve from the 0.5<sup>th</sup> percentile of PM<sub>2.5</sub>, i.e.,  
444 with 0.5% poorly constrained extreme values excluded. Results are obtained from a multivariate  
445 model, adjusted for age, race, sex, Medicaid insurance, annual ZIP code means of summer and  
446 winter temperature, and annual ZIP code level sociodemographic variables. The model also  
447 includes a random intercept for each ZIP code and a spline function of year.

448

449

450 **Figure 2. The association between low-levels air pollutants exposure and diabetes**  
451 **occurrence.**

452 Figure 2 shows the exposure-response curves for the association between PM<sub>2.5</sub>, NO<sub>2</sub>, and warm-  
453 months O<sub>3</sub> and diabetes occurrence, in subsets, restricted to ZIP codes in which the annual  
454 exposures levels have not exceeded the national ambient air quality standards during the study  
455 period (i.e., PM<sub>2.5</sub><12 µg/m<sup>3</sup>, NO<sub>2</sub><35 ppb, or O<sub>3</sub><50 ppb). We show the curves from the  
456 0.5<sup>th</sup> percentile of NO<sub>2</sub>, i.e., with 0.5% poorly constrained extreme values excluded. Results are  
457 obtained from a multivariate model, adjusted for age, race, sex, Medicaid insurance, annual ZIP  
458 code means of summer and winter temperature, and annual ZIP code level sociodemographic  
459 variables. The model also includes a random intercept for each ZIP code and a spline function of  
460 year.

461



