1 **Running Head:** Psychological factors and disability
2
3 **Title:** Association between psychological factors related to pain and instrumental
4 activities of daily living disability in older adults with knee osteoarthritis
5

6 **Author names:** Keigo Nanjo, P.T, M.S. [a, b, *]; Takashi Ikeda, P.T, Ph.D. [a, c, d]; Naoko
7 Nagashio, P.T., B.S.[b]; Tomoko Sakai, M.D., Ph.D. [a]; and Tetsuya Jinno, M.D., Ph.D. [e]
8
9 **Affiliations**
10
[a] *Department of Rehabilitation Medicine, Tokyo Medical and Dental University*
Graduate School, Tokyo, Japan

[b] *Department of Rehabilitation, Shonan Kamakura General Hospital, Kanagawa, Japan*

[c] *School of Nursing and Rehabilitation Sciences, Showa University, Kanagawa, Japan*

[d] *Research Institute for Sport and Exercise Sciences, Showa University, Kanagawa,*
Japan

[e] *Department of Orthopaedic Surgery, Dokkyo Medical University Saitama Medical*
Center, Saitama, Japan
19

20 **Acknowledgements:** The authors thank PTs. D. Kurihara, K. Imahira and K. Suda for
21 their cooperation with data collection.
22
23 **Declaration of Interest:** The authors report no conflicts of interest.
24
25 **Funding:** This research did not receive any specific grant from funding agencies in the
26 public, commercial, or not-for-profit sectors.
27
28

1 **Device Status Statement:** The manuscript submitted does not contain information

2 about medical device.

4 **Data availability statement:** The data that support the findings of this study are

5 available from the corresponding author, Nanjo, K., upon reasonable request.

6
7

*Corresponding author:
Keigo Nanjo, P.T., M.S.
Tokyo Medical and Dental University Graduate School, Department of Rehabilitation
Medicine
1-5-24, Yusima, Bunkyo-ku, Tokyo 113-8510, Japan
E-mail: nanjo.reh@tmd.ac.jp
Tel: +81-03-5803-5648

15
16
17
18

Abstract

Objective: To investigate the association between psychological factors related to pain

and instrumental activities of daily living (IADL) disability in older adults with knee

osteoarthritis (OA).

Design: A cross-sectional study.

Setting: Outpatients at a general hospital in Japan.

Participants: One hundred seventy-nine (N=179) patients with knee OA and

aged≥65years.

Interventions: Not applicable.

Main Outcome Measure: Six-item short form of the pain catastrophizing scale (PCS-

6) and four-item short form of the pain self-efficacy questionnaire (PSEQ-4) were used

to assess the psychological status related to pain. IADL status was assessed using eight

activity items. Participants selected "able," "need help," or "unable" depending on their

abilities to perform these IADL activities. If they chose "need help" or "unable" to

perform, for at least one item, we defined them as "disabled."

Results: Of all participants, 88 (49.1%) showed disability in conducting IADL. Binary

logistic regression analysis with the dependent variable as IADL disabled or not and

PCS-6 and PSEQ-4 as independent variables were performed. Age, sex, pain intensity,

depressive symptoms, usual gait speed (UGS), and isometric knee extension strength

were included as covariates in the logistic regression model. Only PSEQ-4 (odds ratio

[OR]=0.90, 95% confidence interval [95%CI]=0.82–0.99, p=0.02) was a significant

independent variable for psychological factors related to pain. PCS-6 was not a

significant variable (OR=1.06, 95%CI=0.94–1.19, p=0.13). Sex (OR=0.38,

95%CI=0.15–0.96, p=0.04) and UGS (OR=0.13, 95%CI=0.02–0.72, p=0.02) were

significantly independent variables.

Conclusion: Our study showed the importance of assessing self-efficacy related to pain

using the PSEQ-4 to determine the presence of IADL disability in older adults with

knee OA.

Keywords: activities of daily living, gait speed, knee, osteoarthritis, pain, self-efficacy

Abbreviations: BADL, basic activities of daily living; CCI, Charlson comorbidity

index; CI, confidence interval; GDS-15, fifteen-item geriatric depression scale ; IADL,

instrumental activities of daily living; IKES, isometric knee extension strength; KL-

score, Kellgen-Lawllence score; KOOS-pain, pain subscale of the knee injury and

osteoarthritis outcome score; OA, osteoarthritis; OR, odds ratio; PCS-6, six-item short

form of the pain catastrophizing scale; PSEQ-4, four-item short form of the pain self-

efficacy questionnaire; ROM, range of motion; TKA, total knee arthroplasty; UGS,

usual gait speed; UKA, unicompartmental knee arthroplasty

1   Knee osteoarthritis (OA) is a prevalent chronic condition related to aging, and one of

2   the common causes of disabilities limiting daily activities among the elderly.[1] Since the

3   population of older adults with knee OA-related disabilities is increasing globally,[2]

4   health care professionals need effective countermeasures to address this concern.

6   Instrumental activities of daily living (IADL) are essential elements of the

independent lives of older adults. It is marked as independent functioning in a given

environment and includes activities such as meal preparation, shopping, commutation,

financial management, and performance of other household chores.[3] Since IADL

disability in older adults influences health-related quality of life,[4] all-cause mortality,[5]

and decline of cognitive function,[6] assessments and interventions for IADL disability

are important in older adults.

14   Disabilities in the daily lives of patients with knee OA are affected by several

15   factors such as gait speed, knee muscle strength, and pain.[7,8] Pain-related psychological

16   factors such as pain catastrophizing and self-efficacy[10,11,12] have also been reported in

1 disabled patients with knee OA. Regarding psychological interventions for patients with

2 knee OA, a systematic review showed effectiveness in improving functional

3 limitation.[13] A randomized controlled trial showed that a combined intervention of

4 exercise and cognitive therapy for pain was more effective in dealing with the disability

5 than exercise or cognitive therapy each alone. [14]

7  In most studies regarding psychological factors related to pain among patients

with knee OA,[9-12,14] disability was treated as a concept that combines basic activities of

daily living (BADL) and IADL. It is well known that disabilities in older adults

progress hierarchically,[15] IADL disability occurs before BADL disability.[16] To prevent

an incident or deterioration of disabilities in older adults with knee OA,

countermeasures should be taken at an early stage of the disability such as IADL

disability. Therefore, identifying specific factors related to IADL disability is needed to

develop effective interventions for treating it. To our knowledge, there are few studies

on IADL disability in older adults with knee OA.[17,18] Psychological interventions

focused on pain management are recommended for disabilities in older adults with knee

1   OA; [13] however, the association between psychological factors related to pain and IADL

2   disability is unclear.

4   The aim of this study was to investigate the association between

5   psychological factors related to pain and IADL disability in older adults with knee OA.

6   We hypothesized that psychological factors such as pain catastrophizing and pain-

7   related self-efficacy were independently associated with IADL disability in older adults

8   with knee OA, even when adjusted for covariates.

11   **Methods**

14   *Participants*

15   Patients with either knee OA scheduled for unilateral total knee arthroplasty (TKA) or

16   unicompartmental knee arthroplasty (UKA) were eligible for this study. The inclusion

criteria were age ≥65 years and diagnosis of bilateral knee OA based on the clinical

guidelines: knee pain for >3 months and Kellgren–Lawrence score (KL-score) of ≥2.[19]

Conversely, the exclusion criteria were: rheumatoid arthritis; knee OA after trauma;

dementia (mini-mental state examination score <23); inability to walk with a cane by

oneself; history of surgery on the lower extremity; a plan for TKA or UKA on the other

side; serious pathologies (e.g., cancer during treatment) and neurological findings (e.g.,

muscle weakness) that could influence the test performance. A total of 357 patients

were identified at the start of the study. Of these, 166 patients were excluded due to a

history of TKA or UKA on the other side (n=106), history of lower extremity surgery

except knee joint (n=32), rheumatoid arthritis (n=13), mini-mental state examination

score <23 (n=3), knee OA after trauma (n=3), and inability to walk by themselves with

a cane (n=9). A total of 191 patients were invited to participate in this study; 12 refused

to participate. A total of 179 patients were included in the study.

Study design

This study used a cross-sectional design. The conducting and reporting of this study

were guided by the STROBE guidelines.[20] Recruitment was conducted at Shonan

Kamakura General Hospital from May 17, 2019, to May 30, 2021. All participants

provided informed consent before the study began. The study was conducted in

accordance with the Declaration of Helsinki and was approved by the research ethics

boards of the Tokushukai Group Ethics Committee (No. TGE01198-024). All

measurements were evaluated by physical therapists one month before the surgery. Age,

sex, body mass index, Charlson comorbidity index [21], and KL-score [22] of both knees

were obtained as confounders from clinical records.

Outcome measures

IADL status

IADL status was assessed on the basis of eight activity items (food preparation,

shopping, housekeeping, doing laundry, taking medication, using transportation, using a

telephone, and handling finances) using the IADL scale proposed by Lawton and

Brody.[3] Participants were asked to answer their abilities to perform these IADL

activities as "able," "need help," or "unable." Based on previous cross-sectional

reports,[23,24] we defined those participants as "disabled" who opted for "need help" or

"unable" to perform for at least one item; otherwise, they were defined as "non-

disabled."

Pain status

Pain catastrophizing and self-efficacy were assessed as psychological factors related to

pain. Pain catastrophizing was assessed using Japanese version of six-item short form of

the pain catastrophizing scale (PCS-6).[25] Pain catastrophizing scale consists of subscales

related magnification, rumination, and helplessness.[26] It is reported that PCS-6 has the

same property as the original version, correlates to pain intensity assessed by numerical

rating scales (r=0.30, p<0.001),[27] and has good internal consistency (Cronbach's

alpha=0.90).[25] Participants were questioned the degree to which they experienced each

of 6 thoughts and feelings when experiencing pain, on a scale from 0 ("not at all") to 4

("all the time"). The total PCS-6 score ranges from 0 to 24, with higher scores

indicating higher levels of pain catastrophizing.

Self-efficacy was assessed using Japanese version of four-item short form of

the pain self-efficacy questionnaire (PSEQ-4).[28] The pain self-efficacy questionnaire is a

10-item self-report questionnaire used to assess self-efficacy in individuals with chronic

pain.[29] It is reported that PSEQ-4 has the same property as the original version,[27]

correlates pain intensity assessed by numerical rating scales (r=-0.35, $p<0.001$), and has

good internal consistency (Cronbach's alpha=0.90).[28] The PSEQ-4 consists of four

questions, and participants were questioned how confident they were on a scale of 0

("not at all confident") to 6 ("completely confident") to perform the given activities

despite pain. The PSEQ-4 scores range from 0 to 24, with higher scores indicating more

confidence in performing the given activities despite pain.

The Japanese version of pain subscale of the knee injury and osteoarthritis

outcome score (KOOS-pain) was used for the representative index of pain intensity.[30]

KOOS-pain consists of nine questions. Participants were questioned about their

condition one week before the evaluation date. Standardized answer choices were

provided, and each question was assigned a score ranging from 0 to 4. A normalized

total score of 0 to 100 was calculated, and higher scores indicated that patients reported

less pain. KOOS-pain correlates with the body pain subscale of the short form-36 health

survey ($r=0.67$, $p<0.01$) and has good internal consistency (Cronbach's alpha=0.90).[30]

Confounders

Gait speed and knee joint functions

Based on previous studies that reported the relationship between IADL disability and

gait speed,[18, 23, 31] usual gait speed (UGS) was measured using a 5-meter gait test.[32] To

test knee joint function, isometric knee extension strength (IKES) was measured using a

handheld dynamometer [a].[33] Knee extension and flexion range of motion (ROM) were

measured using a goniometer.[34] The knee scheduled for surgery was defined as the

affected side, and opposite side was defined as the unaffected side. IKES and knee

ROM measurements were conducted on both knees.

Depressive symptoms

Depressive symptoms were assessed using Japanese version of the fifteen-item geriatric

depression scale (GDS-15)[35] as the prevalence of depressive symptoms in patients with

knee OA is higher than in the general population,[36] and depressive symptoms have been

reported as a factor affecting IADL disability in older adults.[37] GDS-15 consists of 15

questions. The participants answered with a "yes" or "no" response. The total score was

calculated from 0 to 15, with higher scores indicating more depressive symptoms.

Statistical analysis

The participants were divided into two groups according to their IADL status: those

who answered "need help" or "unable" to perform on at least one item were allocated as

the IADL disabled group, and those who answered "able" in all items were assigned as

the IADL non-disabled group. All outcome measures and confounders were compared

between the two groups. The chi-square test was used for categorical variables, and

student's *t*-test and Mann–Whitney *U* test were used for standard and nonstandard

distribution variables, respectively. In the IADL disabled group, the proportion of each

item of IADL disability was shown.

To confirm the relationships between confounders and pain statuses,

correlations between continuous variables (p-values <0.05), while comparing the two

groups for the pain status, were calculated using Pearson's correlation coefficient and

Spearman's rank correlation. Regarding sex differences, the values of pain status

between men and women were compared.

The association of psychological factors related to pain with IADL disability

was analyzed using binary logistic regression models. Previous reports have shown that

age, sex,[38] depressive symptoms [37] and gait speed [18, 23,33] were factors related to IADL

disability in older adults. Three logistic regression models with dependent variable as

IADL status (disabled= 1 or not= 0) were deployed hierarchically. The first model

included pain status (e.g., KOOS-pain, PCS-6, PSEQ-4), age, and sex (men=1) as

independent variables (Model 1). The second model added GDS-15 to Model 1 (Model

2). The final model included Model 2 with UGS and other confounders that had p-

values <0.05 in the two groups comparison (Model 3). All analyses were performed

using R version 4.0.3. [b] for all tests, a p-value <0.05, was considered statistically

significant.

To create a sample size in a logistic regression analysis, the number of cases

(N) = 10 k/p are needed, where k is number of independent variables as covariates and p

is a defined ratio of responders to non-responders at the follow-up points.[39] To avoid

overfitting and comply with the recommendations given by Vittinghoff & McCulloch,[40]

we assumed to arrive at nine factors out of all variables in measurements and

demographics. In a previous study involving older adults with joint pain (n=407),

60.9% of participants had IADL disability.[23] Assuming the ratio p of disabled to non-

disabled was 1:1.5, the minimum sample size in this study was calculated as

$10 \times 9/0.6 = 150$.

Results

All participants completed all assessments, and all confounders were collected from

their medical records. Of all participants, 88 (49.1%) had IADL disabilities. Sixty-one

(69.3%) showed disability in shopping, and 47 (53.4%) showed disability in using

transportation. No participant showed disability in taking medication and using a

telephone (Table 1).

10   The IADL non-disabled group was significantly younger (p=0.001) and had a

greater number of men (p=0.009) than the IADL disabled group. The IADL non-

disabled group showed significantly higher values of UGS (1.02 ± 0.27 [mean ± SD] vs.

0.83 ± 0.27; p<0.001), IKES on the affected side (1.23 [0.96–1.59] (median

[interquartile range]) vs. 0.99 [0.73–1.31]; p<0.001), IKES on the unaffected side (1.26

[0.96–1.61] vs. 0.94 [0.75–1.30]; p<0.001), KOOS-pain (48.6 ± 17.8 vs. 42.5 ± 15.1;

p=0.01) and PSEQ-4 (15.0 [12.0–18.5] vs. 13.0 [11.0–15.0]; p<0.001). Conversely, the

IADL non-disabled group showed significantly lower values of PCS-6 (14.0 [7.5–17.0]

vs. 16.0 [12.0–19.0]; p=0.003) and GDS-15 (3.0 [1.0–6.0] vs. 4.0 [2.0–7.3]; p<0.001)

(Table 2).

The results of Pearson's correlation coefficient and Spearman's rank

correlation coefficients are presented in Table 3. PCS-6 significantly correlated with

UGS (r=-0.26), IKES on the affected side (r=-0.17), and GDS-15 (r=0.36). PEQ-4

significantly correlated with UGS (r=0.20), IKES on the affected side (r=0.27), and

GDS-15 (r=-0.42). KOOS-pain significantly correlated with UGS (r=0.37), IKES on the

affected side (r=0.23), IKES on the unaffected side (r=0.21) and GDS-15 (r=-0.19).

In the comparison of pain status between men and women, women showed

significantly lower values of KOOS-pain (44.1 ± 15.9 vs. 51.0 ± 9.0; p=0.02) and

higher values of PCS-6 (15.0 [11.0–19.0] vs. 13.5 [8.5–16.0]; p=0.03). The values of

PSEQ-4 (13.0 [12.0–17.0] vs. 14.5 [11.3–19.5]; p=0.24) were not significantly

different.

2    In binary logistic regression analysis with IADL disability status as a

3 dependent variable, PCS-6 (odds ratio [OR]=1.07, 95% confidence interval [CI] =1.00–

4 1.14, p=0.004), PSEQ-4 (OR=0.88, 95%CI=0.81–0.95, p=0.002), age (OR=1.12,

5 95%CI=1.05–1.12, p=0.01) and sex (OR=0.33, 95%CI=0.14–0.79, p=0.002) were

6 significant independent variables in Model 1. In Model 2, PSEQ-4 (OR=0.90,

7 95%CI=0.82–0.98, p=0.01), age (OR=1.12, 95%CI=1.04–1.19, p=0.001) and sex

8 (OR=0.33, 95%CI=0.14–0.79, p=0.002) were significantly independent variables. IKES

9 on both sides showed significant differences between the two groups, and the effect size

10 of IKES on the unaffected side was greater than that on the affected side. Therefore,

11 only IKES on the unaffected side was included in the final model to avoid

12 multicollinearity. In the final model, PSEQ-4 (OR=0.90, 95%CI=0.82–0.99, p=0.02),

13 UGS (OR=0.13, 95%CI=0.02–0.72, p=0.02) and sex (OR=0.38, 95%CI=0.15–0.96,

14 p=0.04) were significantly independent variables (Table 4).

Discussion

The present study aimed to investigate the association between psychological factors

related to pain and IADL disability in older adults with knee OA. The values of PCS-6

and PSEQ-4 were significantly different between the IADL non-disabled and IADL

disabled groups. However, in the logistic regression analysis adjusted for covariates,

PSEQ-4 was the only significant psychological factor related to pain associated with

IADL disability.

In the final logistic regression model, sex and UGS were significantly

independent variables. Alexandre et al. have shown that sex is a factor related to IADL

disability in older adults.[38] Our results supported this finding. In particular, a reduction

in gait speed has been reported as a strong factor related to IADL disability in older

adults. [23,33,41] The result in this study is consistent with our previous study that showed

gait speed is a discriminatory factor for IADL disability in older adults with knee OA.[18]

In this study, 49.1% of all participants had IADL disability. This proportion was lower

than that in a previous study involving older adults with joint pain.[23] In our study

population, main items pertaining to IADL disability were outdoor activities such as

shopping (68.6%) and using transportation (50.6%) that required gait speed.

Furthermore, because the effect size of UGS was the greatest of all measurements when

compared between the two groups, it could be a crucial factor associated with IADL

disability in older adults with knee OA.

9   Regarding pain status, a previous report involving patients with low back pain

reported that the minimal importance change of PSEQ-4 was 1.5 points.[42] In our study

results, the median difference in PSEQ-4 between the two groups was 2.0 points. In

addition, the correlation between PSEQ-4 and UGS was weak, and PSEQ-4 was not

significantly different for men and women. Therefore, these findings indicated that

PSEQ-4 was hardly affected by demographics and motor functions and that it had a

specific relationship with IADL disability in older adults with knee OA.

With respect to the relationship between pain self-efficacy and disability, two

cross-sectional studies showed that self-efficacy related to pain was a factor related to

disability in patients with chronic low back pain.[43,44] To our knowledge, however, few

studies have assessed the relationship between disability and self-efficacy using a pain

self-efficacy questionnaire in patients with knee OA. Some cross-sectional studies have

shown that self-efficacy, which partially includes pain elements, is a factor related to

disability in patients with knee OA.[11,12] A cohort study in older adults with knee pain

reported that self-efficacy is a predictor of self-reported disability.[10] With respect to the

relationship between IADL disability and self-efficacy, a previous cross-sectional study

in older adults with chronic joint pain showed that self-efficacy is associated with IADL

disability in logistic regression analysis that included demographics and pain intensity

as covariates.[24] The results of the above studies are consistent with those of our study.

Self-efficacy describes the confidence that a person has in one's ability to achieve a

desired outcome.[45] In patients with knee OA, those with high self-efficacy for

controlling arthritis pain have been found to have higher pain thresholds than those with

low self-efficacy.[46] Therefore, older adults with knee OA who had high self-efficacy

might have the tendency not to voluntarily inhibit activities included in IADL even if

they had knee pain. Consequently, we considered that PSEQ-4 was associated with

IADL disability, even after adjusting for other covariates.

Study limitations

This study had several limitations. First, since this was a cross-sectional study, it was

difficult to determine causal relationships between factors and IADL disability

incidence. Second, because this study included patients with knee OA scheduled for

TKA or UKA, treatment situations were different from general older adults with knee

OA. Therefore, it might have been difficult to adapt the results to older adults with

early-stage knee OA or low pain status. In particular, awaiting surgery could affect pain

intensity and psychological factors related to the pain. Nonetheless, the average KOOS-

pain score in this study population was not inferior to those in other studies.[47] The

median PCS-6 and PSEQ-4 scores were also not inferior to those in previous studies in

patients with chronic pain.[27,28] Third, because IADL disability was treated with

categorical variables as the dependent variable in our logistic regression analyses, we

could not identify the impact of each factor on IADL disability. However, when

comparing the two groups, the effect sizes of the variables that were significantly

related to IADL disability in logistic regression models were roughly the same.

Conclusion

In psychological factors related to pain, self-efficacy assessed using the PSEQ-4 was

found to be associated with IADL disability even after adjusting for covariates. Sex and

gait speed were also factors associated with IADL disability in older adults with knee

OA. Our study showed the importance of assessing self-efficacy related to pain using

the PSEQ-4 to determine the presence of IADL disability in older adults with knee OA.

Future studies are necessary to elucidate the mechanism by which self-efficacy is

related to IADL disability in older adults with knee OA. Furthermore, longitudinal

studies are needed to identify predictors of the development of IADL disability in older

adults with knee OA.

Suppliers

a: μ-tus F-1, Anima Corporation, 3-65-1, Shimoishihara, Chofu, Tokyo, 182-0034,

Japan

b: R version 4.0.3, R Foundation for Statistical Computing, Welthandelsplatz 1, 1020

Vienna, Austria

References

1.   Bijlsma JW, Berenbaum F, Lafeber FP. Osteoarthritis: an update with relevance for

clinical practice. Lancet 2011;377(9783):2115-26.

2.   Global, regional, and national incidence, prevalence, and years lived with disability

for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global

Burden of Disease Study 2015. Lancet 2016;388(10053):1545-602.

3.   Lawton MP, Brody EM. Assessment of older people: self-maintaining and

instrumental activities of daily living. Gerontologist 1969;9(3):179-86.

4.   Fried LP, Kronmal RA, Newman AB, Bild DE, Mittelmark MB, Polak JF et al.

Risk factors for 5-year mortality in older adults: the Cardiovascular Health Study.

JAMA 1998;279(8):585-92.

5.   Scott WK, Macera CA, Cornman CB, Sharpe PA. Functional health status as a

predictor of mortality in men and women over 65. J Clin Epidemiol

1997;50(3):291-6.

6. Rajan KB, Hebert LE, Scherr PA, Mendes de Leon CF, Evans DA. Disability in

basic and instrumental activities of daily living is associated with faster rate of

decline in cognitive function of older adults. J Gerontol A Biol Sci Med Sci

2013;68(5):624-30.

7. Pereira D, Peleteiro B, Araújo J, Branco J, Santos RA, Ramos E. The effect of

osteoarthritis definition on prevalence and incidence estimates: a systematic review.

Osteoarthritis Cartilage 2011;19(11):1270-85.

8. van Dijk GM, Veenhof C, Lankhorst GJ, Dekker J. Limitations in activities in

patients with osteoarthritis of the hip or knee: the relationship with body functions,

comorbidity and cognitive functioning. Disabil Rehabil 2009;31(20):1685-91.

9. Zambon S, Siviero P, Denkinger M, Limongi F, Victoria Castell M, van der Pas S et

al. Role of osteoarthritis, comorbidity, and pain in determining functional

limitations in older populations: European project on osteoarthritis. Arthritis Care

Res (Hoboken) 2016;68(6):801-10.

10. Rejeski WJ, Miller ME, Foy C, Messier S, Rapp S. Self-efficacy and the

progression of functional limitations and self-reported disability in older adults with

knee pain. J Gerontol B Psychol Sci Soc Sci 2001;56(5):S261-5.

11. Maly MR, Costigan PA, Olney SJ. Determinants of self-report outcome measures in

people with knee osteoarthritis. Arch Phys Med Rehabil 2006;87(1):96-104.

12. Creamer P, Lethbridge-Cejku M, Hochberg MC. Factors associated with functional

impairment in symptomatic knee osteoarthritis. Rheumatology (Oxford)

2000;39(5):490-6.

13. Dixon KE, Keefe FJ, Scipio CD, Perri LM, Abernethy AP. Psychological

interventions for arthritis pain management in adults: a meta-analysis. Health

Psychol 2007;26(3):241-50.

14. Bennell KL, Ahamed Y, Jull G, Bryant C, Hunt MA, Forbes AB et al. Physical

therapist-delivered pain coping skills training and exercise for knee osteoarthritis:

randomized controlled trial. Arthritis Care Res (Hoboken) 2016;68(5):590-602.

15. Barberger-Gateau P, Rainville C, Letenneur L, Dartigues JF. A hierarchical model

of domains of disablement in the elderly: a longitudinal approach. Disabil Rehabil

2000;22(7):308-17.

16. Edjolo A, Proust-Lima C, Delva F, Dartigues JF, Pérès K. Natural history of

dependency in the elderly: A 24-year population-based study using a longitudinal

item response theory model. Am J Epidemiol 2016;183(4):277-85.

17. George LK, Ruiz D, Sloan FA. The effects of total knee arthroplasty on physical

functioning in the older population. Arthritis Rheum 2008;58(10):3166-71.

18. Nanjo K, Ikeda T, Nagashio N, Masuda T, Sakai T, Okawa A et al. Gait speed and

pain status as discriminatory factors for instrumental activities of daily living

disability in older adults with knee osteoarthritis. Geriatr. Gerontol. Int 2021;1–6.

https://doi.org/10.1111/ggi.14229

19. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K et al. Development of

criteria for the classification and reporting of osteoarthritis. Classification of

osteoarthritis of the knee. Diagnostic and therapeutic criteria committee of the

American Rheumatism Association. Arthritis Rheum 1986;29(8):1039-49.

20. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP.

The Strengthening the Reporting of Observational Studies in Epidemiology

(STROBE) statement: guidelines for reporting observational studies. J Clin

Epidemiol. 2008;61(4):344-9.

21. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying

prognostic comorbidity in longitudinal studies: development and validation. J

Chronic Dis 1987;40(5):373-83.

22. Kellgren JH, Lawrence JS. Radiological assessment of osteoarthrosis. Ann Rheum

Dis 1957;16(4):494-502.

23. Lee MC, Hsu CC, Tsai YF, Chen CY, Lin CC, Wang CY. Criterion-referenced

values of grip strength and usual gait speed using instrumental activities of daily

living disability as the criterion. J Geriatr Phys Ther 2018;41(1):14-9.

24. Hermsen LA, Leone SS, Smalbrugge M, Dekker J, van der Horst HE. Frequency,

severity and determinants of functional limitations in older adults with joint pain

and comorbidity: results of a cross-sectional study. Arch Gerontol Geriatr

2014;59(1):98-106.

25. Nishigami T, Mibu A, Tanaka K, Yamashita Y, Watanabe A, Tanabe A.

Psychometric properties of the Japanese version of short forms of the pain

catastrophizing scale in participants with musculoskeletal pain: A cross-sectional

study. J Orthop Sci 2017;22(2):351-6.

26. Sullivan MJ, Bishop SR, Pivik J. The pain catastrophizing scale: development and

validation. Psychol Assess 1995;7(4): 524-532.

27. McWilliams LA, Kowal J, Wilson KG. Development and evaluation of short forms

of the pain catastrophizing scale and the pain self-efficacy questionnaire. Eur J Pain

2015;19(9):1342-9.

28. Adachi T, Enomoto K, Yamada K, Inoue D, Nakanishi M, Takahashi N, et al.

Evaluating the psychometric properties of two-item and four-item short forms of

the Japanese pain self-efficacy questionnaire: a cross-sectional study. J Anesth

2019;33(1):58-66.

29. Nicholas MK. The pain self-efficacy questionnaire: Taking pain into account. Eur J

Pain 2007;11(2):153-63.

30. Nakamura N, Takeuchi R, Sawaguchi T, Ishikawa H, Saito T, Goldhahn S. Cross-

cultural adaptation and validation of the Japanese knee injury and osteoarthritis

outcome score (KOOS). J Orthop Sci 2011;16(5):516-23.

31. Zhang L, Guo L, Wu H, Gong X, Lv J, Yang Y. Role of physical performance

measures for identifying functional disability among Chinese older adults: Data

from the China Health and Retirement Longitudinal Study. PLoS One

2019;14(4):e0215693.

32. Shinkai S, Watanabe S, Kumagai S, Fujiwara Y, Amano H, Yoshida H et al.

Walking speed as a good predictor for the onset of functional dependence in a

Japanese rural community population. Age Ageing 2000;29(5):441-6.

33. Katoh M, Yamasaki H. Comparison of reliability of isometric leg muscle strength

measurements made using a hand-held dynamometer with and without a restraining

belt. J Phys. Ther. Sci. 2009;21(1):37-42.

34. Norkin CC, White DJ. Measurement of joint motion: a guide to goniometry. 2nd ed.

Philadelphia: F.A. Davis; 1995.

35. Sugishita K, Sugishita M, Hemmi I, Asada T, Tanigawa T. A validity and

reliability study of the Japanese version of the geriatric depression scale 15 (GDS-

15-J). Clin Gerontol 2017;40(4):233-40.

36. Stubbs B, Aluko Y, Myint PK, Smith TO. Prevalence of depressive symptoms and

anxiety in osteoarthritis: a systematic review and meta-analysis. Age

Ageing. 2016;45(2):228–235.

37. Steffens DC, Hays JC, Krishnan KRR. 1999. Disability in geriatric depression. Am

J Geriatr Psychiatry. 1999;7:34-40.

38. Alexandre TaS, Corona LP, Nunes DP, Santos JL, Duarte YA, Lebrão ML.

Disability in instrumental activities of daily living among older adults: gender

differences. Rev Saude Publica 2014;48(3):379-89.

39. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of

the number of events per variable in logistic regression analysis. J Clin Epidemiol.

1996;49(12):1373–9.

40. Vittinghoff E, McCulloch CE. Relaxing the rule of ten events per variable in

logistic and cox regression. Am J Epidemiol. 2007;165(6):710–8. https://doi.

org/10.1093/aje/kwk052.

41. Donoghue OA, Savva GM, Cronin H, Kenny RA, Horgan NF. Using timed up and

go and usual gait speed to predict incident disability in daily activities among

community-dwelling adults aged 65 and older. Arch Phys Med Rehabil

2014;95(10):1954-61.

42. Chiarotto A, Vanti C, Cedraschi C, Ferrari S, de Lima E Sà Resende F, Ostelo RW

et al. Responsiveness and minimal important change of the pain self-efficacy

questionnaire and short forms in patients with chronic low back pain. J Pain

2016;17(6):707-18.

43. Costa LaC, Maher CG, McAuley JH, Hancock MJ, Smeets RJ. Self-efficacy is

more important than fear of movement in mediating the relationship between pain

and disability in chronic low back pain. Eur J Pain 2011;15(2):213-9.

44. Ayre M, Tyson GA. The role of self-efficacy and fear-avoidance beliefs in the

prediction of disability. Aust Psychol 2001;36(3):250-253.

45. Bandura A. Self-efficacy: toward a unifying theory of behavioral change. Psychol

Rev 1977;84(2):191-215.

46. Keefe FJ, Lefebvre JC, Maixner W, Salley AN, Caldwell DS. Self-efficacy for

arthritis pain: relationship to perception of thermal laboratory pain stimuli. Arthritis

Care Res 1997;10(3):177-84.

47. Collins NJ, Prinsen CA, Christensen R, Bartels EM, Terwee CB, Roos EM. Knee

injury and osteoarthritis outcome score (KOOS): systematic review and meta-

analysis of measurement properties. Osteoarthritis Cartilage 2016;24(8):1317-29.

48. Field A. Discovering statistics using IBM SPSS Statistics. 4th ed. London: Sage

Publications Ltd; 2012.

Table 1. Content of IADL disability in IADL disabled group (n=88)

item	
Preparing food	33 (37.5)
Shopping	61 (69.3)
Housekeeping	40 (45.5)
Doing laundry	6 (6.8)
Taking medication	0 (0)
Using transportation	47 (53.4)
Using a telephone	0 (0)
Handling finances	15 (17.0)

Number (%) was shown.

IADL: Instrumental activity of daily living

Table 2. Baseline characteristics of the study population and comparison between IADL non-disabled and disabled groups

variable	ALL (n= 179)	IADL non-disabled (n=91)	IADL disabled (n=88)	p-value	effect size
Women; number (%)	141 (78.8)	64 (70.6)	77 (87.5)	0.009	0.2
Age; mean±SD	76.5 ± 5.4	75.2 ± 5.2	77.8 ± 5.2	0.001	0.24
Body mass Index(kg/m^2); median (IQR)	23.1 (21.1–26.2)	22.9 (21.4–26.1)	23.1 (21.4–26.1)	0.74	0.02
Family status (living alone); number (%)	36 (20.1)	22 (24.2)	14 (15.9)	0.23	0.09
CCI; median (IQR)	0 (0–1.0)	0 (0–1.0)	0 (0–1.0)	0.63	0.04
KL score on affected side					
II; number (%)	19 (10.6)	9 (9.9)	10 (11.4)	0.44	0.1
III; number (%)	101 (56.4)	48 (52.7)	53 (60.2)		
IV; number (%)	59 (33.0)	34 (37.4)	25 (28.4)		
KL score on unaffected side					
II; number (%)	62 (34.6)	33 (36.3)	29 (33.0)	0.5	0.09
III; number (%)	82 (45.8)	38 (41.8)	44 (50.0)		
IV; number (%)	35 (19.6)	20 (22.0)	15 (17.0)		
Pain status					
KOOS-pain; mean±SD	45.6 ± 16.8	48.6 ± 17.8	42.5 ± 15.1	0.01	0.18
PCS-6; median (IQR)	15.0 (10.0-18.0)	14.0 (7.5-17.0)	16.0 (12.0-19.0)	0.003	0.23
PSEQ-4; median (IQR)	14.0 (12.0-17.0)	15.0 (12.0-18.5)	13.0 (11.0-15.0)	<0.001	0.29

UGS (m/sec); mean±SD		0.93 ± 0.27	1.02 ± 0.27	0.83 ± 0.27	<0.001	0.37
IKES（Nm/kg）; median (IQR)	affected side	1.09 (0.81–1.43)	1.23 (0.96–1.59)	0.99 (0.73–1.31)	<0.001	0.27
	unaffected side	1.10 (0.78–1.47)	1.26 (0.96–1.61)	0.94 (0.75–1.30)	<0.001	0.29
Knee flexion ROM（°）; median (IQR)	affected side	130 (115–135)	130 (115–135)	130 (115–140)	0.86	0.03
	unaffected side	125 (120–140)	125 (120–137.5)	130 (115–140)	0.63	0.03
Knee extension ROM（°）; median (IQR)	affected side	-5 (-10 to 0)	-5 (-10 to 0)	-5 (-10 to 0)	0.51	0.05
	unaffected side	-5 (-10 to 0)	-5 (-5 to 0)	-5 (-10 to 0)	0.25	0.07
GDS-15; median (IQR)		3.0 (1.5–6.0)	3.0 (1.0–6.0)	4.0 (2.0-7.3)	<0.001	0.25

Effect size was estimated using r for continuous variables ($r=\sqrt{t^2/(t^2+df)}$, for standard distribution variables; $r=z/\sqrt{N}$, for nonstandard distribution variables) and Cramér 's V for categorical variables.[48]

CCI, Charlson comorbidity index; GDS-15, fifteen-item geriatric depression scale; IADL, instrumental activities of daily living; IKES, isometric knee extension strength; KL-score, Kellgen-Lawllence score; KOOS-pain, pain subscale of the knee injury and osteoarthritis outcome score; OA, osteoarthritis; PCS-6, six-item short form of the pain catastrophizing scale; PSEQ-4, four-item short form of the pain self-efficacy questionnaire; ROM, range of motion; UGS, usual gait speed

Table 3. Correlation between variables of pain status and covariates

variable	PCS-6	PSEQ-4	KOOS-pain
Age	0.07	-0.04	-0.09
UGS	-0.26[†]	0.20[†]	0.37[‡]
IKES on affected side	-0.17*	0.15	0.23[†]
IKES on unaffected side	-0.10	0.27[‡]	0.21[†]
GDS-15	0.36[‡]	-0.42[‡]	-0.19[†]

*: $p<0.05$, [†]: $p<0.01$, [‡]: $p<0.001$

Pearson's correlation coefficient and Spearman's rank correlation coefficients are presented.

GDS-15, fifteen-item geriatric depression scale; IKES, isometric knee extension strength; KOOS-pain, pain subscale of the knee injury and osteoarthritis outcome score; PCS-6, six-item short form of the pain catastrophizing scale; PSEQ-4, four-item short form of the pain self-efficacy questionnaire; UGS, usual gait speed

Table 4. logistic regression model with the dependent variable as IADL disable or not.

Variable	Crude OR	Mode1		Mode2		Model3	
		Adjusted OR	95%CI	Adjusted OR	95%CI	Adjusted OR	95%CI
PCS-6	1.07	1.07*	1.00–1.14	1.06	0.99–1.13	1.06	0.94–1.19
PSEQ-4	0.86	0.88†	0.81–0.95	0.90*	0.82–0.98	0.90*	0.82–0.99
KOOS pain	0.98	1.00	0.98–1.03	1.00	0.98–1.03	1.01	0.99–1.04
Age	1.10	1.12†	1.05–1.19	1.12†	1.04–1.19	1.07	1.00–1.15
Sex	0.33	0.33*	0.14-0.79	0.33*	0.14-0.79	0.38*	0.15–0.96
GDS-15	1.17			1.08	0.96–1.21	1.06	0.94–1.19
UGS	0.05					0.13*	0.02–0.72
IKES unaffected side	0.35					0.73	0.32–1.64

Model1; Hosmer-Lemeshow test = 0.71, area under the curve (95% CI) = 0.77 (0.70–0.84)

Model2; Hosmer-Lemeshow test = 0.12, area under the curve (95% CI) = 0.77 (0.70–0.84)

Model3; Hosmer-Lemeshow test = 0.47, area under the curve (95% CI) = 0.80 (0.73–0.86)

*: $p < 0.05$, †: $p < 0.01$

CI, confidence interval; GDS-15, fifteen-item geriatric depression scale; IADL, instrumental activities of daily living; IKES, isometric knee extension strength; KOOS-pain, Pain subscale of the knee injury and osteoarthritis outcome score; OR, odds ratio; PCS-6, six-item short form of the pain catastrophizing scale; PSEQ-4, four-item short form of the pain self-efficacy questionnaire; ROM, range of motion; UGS, usual gait speed;