Age-specific effects of childhood BMI on multiple sclerosis risk: a Mendelian Randomisation study

Luke Hone[1]*, Benjamin M Jacobs[1,2]*, Charles Marshall[1,2], Gavin Giovannoni[1,2,3], Alastair Noyce[1,2], Ruth Dobson[1,2]
* Equal contribution

1: Preventive Neurology Unit, Wolfson Institute of Population Health, Queen Mary University London
2: Department of Neurology, Royal London Hospital
3: Blizard Institute, Queen Mary University London

Corresponding author: Dr Benjamin M Jacobs
Preventive Neurology Unit, Wolfson Institute of Population Health, Charterhouse Square, London EC1M 6BQ
Email: b.jacobs@qmul.ac.uk

Abstract
Words
Figures
Tables
References

Objective
Higher childhood Body Mass Index (BMI) during early life is thought to be a causal risk factor for Multiple Sclerosis (MS). We used longitudinal mendelian randomisation (MR) to determine whether there is a critical window during which BMI influences MS risk.

Methods
Summary statistics for childhood BMI and for MS susceptibility were obtained from recent large GWAS. We generated exposure instruments for BMI during four non-overlapping epochs (< 3 months, 3 months - 1.5 years, 2 - 5 years, and 7 - 8 years) and performed MR using the inverse-variance weighted method with standard sensitivity analyses.

Results

At all time epochs other than birth, genetically-elevated BMI was associated with an increased liability to MS: Birth (OR 0.81, 95%CI 0.50-1.31, N_{SNPs}=7, p=0.39), Infancy (OR 1.18, 95%CI 1.04-1.33, N_{SNPs}=18, p=0.01), Early childhood (OR 1.31, 95%CI 1.03-1.66, N_{SNPs}=4, p=0.03), Later childhood (OR 1.34, 95% CI 1.08-1.66, N_{SNPs}=4, p=0.01). There was no evidence that horizontal pleiotropy was biasing the IVW estimates at any of these points. There was evidence of an upwards trend in MS risk from birth to 8 years (Mann-Kendall test, tau = 0.636, 2-sided p-value =0.005).

Conclusion

We provide novel evidence using longitudinal MR that genetically-determined higher BMI during early life (from 3 months) increases MS risk, and that the magnitude of this effect increases towards late childhood and adolescence.

Introduction

Multiple Sclerosis (MS) is an immune-mediated demyelinating disease of the Central Nervous System (CNS), with both genetic and environmental risk factors[1]. Environmental factors including obesity, low serum vitamin D, EBV infection, smoking, air pollution and solvent exposure have all been associated with increased MS risk, although the mechanisms through which they act remain uncertain[1]. Traditional epidemiological study designs used to study environmental risk factors - such as case-control and cohort studies - are liable to unmeasured confounding and reverse causation and may yield associations which are not truly causal[2].

Mendelian Randomisation (MR) is a type of instrumental variable (IV) analysis that uses genetic variants as proxies to examine whether observed associations between an exposure and an outcome are likely to be causal[2,3]. Childhood obesity has been identified as a potential causal MS risk factor in both epidemiological[4] and MR studies[5,6]. A consistent finding is that birth weight is not associated with MS risk, whereas higher BMI during later childhood and adolescence appears to be a risk factor[4,7-9]. However, BMI during early life is a dynamic trait; it is unknown if there is a crucial point during development when BMI influences subsequent MS risk. Understanding the precise nature and timing of this association may shed light on the biological pathways leading to the clinical development of MS, and may inform targeted obesity prevention programmes in early life. Defining this critical window of effect is almost impossible using MR to instrument BMI at a single time point., and combining GWAS of BMI at different time points from different cohorts may introduce confounding from population stratification.

Longitudinal MR analysis of the effect of BMI during development is now possible due to the recent publication of longitudinal BMI GWAS data from the Norwegian Mother, Father, and Child Cohort Study (MoBa) cohort[10]. We set out to explore the causal effect of BMI changes during early life on MS risk and examine the dynamics of this effect over time.

Methods

Genetic datasets:
Exposures dataset
The MoBa cohort is an open-ended cohort study that recruited pregnant women in Norway from 1999 to 2008, with anthropometric measurements of the children by trained nurses across childhood. This cohort was used to perform GWAS of BMI at various ages using data from between 11,095 and 28,681 children across 12 time points from birth to 8 years (birth, 6 weeks, 3 months, 6 months, 8 months, 1 year, 1.5 years, 2 years, 3 years, 5 years, 7 years, and 8 years). 46 distinct genetic loci were associated with raised childhood BMI at various ages[10]. The outcome of this GWAS was standardized BMI, and therefore the units are not directly comparable with published estimates from MR studies using other BMI GWAS for genetic instruments.

We used variants reaching a genome-wide p-value threshold of $<5\times10^{-8}$ at each time point after conditional and joint analysis as instrumental variables. In the primary analysis, time points were amalgamated into four distinct epochs: Birth to 6 weeks, 3 months to 1.5 years, 2 to 5 years, and 7 to 8 years, reflecting the developmental stages of birth, infant, toddler and child. If a variant was associated at $p<5\times10^{-8}$ with >1 time point within an epoch, the association statistics were taken from the time point of statistically-strongest association (i.e. the smallest p value). A secondary analysis then used data at each individual time point. These epochs were defined according to the shared genetics of BMI at each time point, and loosely correspond to the four phases of BMI genetics suggested by the authors[10]. Data are available from the MOBA authors on request.

Outcome dataset
The International Multiple Sclerosis Genetics Consortium (IMSGC) 2019 discovery phase GWAS (47,429 MS cases, 68,374 controls) was used as the outcome dataset[11]. We used summary statistics from the discovery stage meta-analysis (14,802 persons with MS, 26,703 controls). This GWAS discovered 233 independent genome-wide significant signals associated with MS, of which 32 variants lay within the Major Histocompatibility Complex (MHC). Data are available on request from https://imsgc.net/.

Other datasets
GWAS datasets for adult BMI and birthweight were obtained from the GIANT and EGG consortia meta-analyses (respectively) including UK Biobank participants[12,13]. Data on birth weight has been contributed by the EGG Consortium using the UK Biobank Resource and has been downloaded from www.egg-consortium.org. Data on adult BMI can be downloaded from https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files.

Mendelian randomisation & statistical analyses
Mendelian Randomisation (MR) was performed using the TwoSampleMR (version 0.5.5) package in R (version 3.6.1)[14].

Genetic instruments were generated by the following steps:
1. Exposure (BMI) Single Nucleotide Polymorphisms (SNPs) associated with childhood BMI at $p<5\times10^{-8}$ were retained.
2. SNPs were clumped to ensure independence using the PLINK clumping method and a European 1000 Genomes reference dataset. SNPs in linkage disequilibrium (LD) ($R^2>$

0.001) or within a window of 10,000 kb were pruned. The SNP with the lowest p-value in the window or in LD was retained for analysis.

3. MHC SNPs (chr6:25,000,000 - chr6:35,000,000 in hg19) were excluded.
4. Exposure SNPs were extracted from the outcome dataset (IMSGC MS GWAS).
5. Exposure and outcome SNPs were harmonised so that their effects corresponded to the same allele and palindromic SNPs with a minor allele frequency (MAF) > 0.42 were discarded.

MR analysis was performed using a random-effects inverse-variance-weighted method (IVW). Secondary analyses included a test for heterogeneity, tests for horizontal pleiotropy through calculation of an MR-Egger intercept and MR-PRESSO[15], and a Leave-One-Out analysis. In secondary analyses we excluded SNPs explaining more variance in the outcome than exposure (Steiger filtering) as these variants are likely to violate the exclusion-restriction assumption of MR[16]. Results are presented as Odds Ratios (OR) with 95% Confidence Intervals (CIs) and associated p-values. Results are visualised as scatter, forest, funnel and leave-one-out plots. F-statistics for instrument strength were calculated using an online power calculator (https://shiny.cnsgenomics.com/mRnd/)[17] based on R^2 values and sample sizes from the exposure GWAS. To ensure that childhood BMI instrument SNPs were not likely to influence MS susceptibility via their effect on adult BMI, we excluded SNPs which were strongly associated with adult BMI ($P<1x10^{-5}$) in the largest GWAS meta-analysis of adult BMI to date[12]. To provide further confirmation of the birthweight MR result, we repeated the analysis using an independent and larger GWAS of birthweight including UKB participants[13].

Data Availability
We thank MoBa, Prof Stefan Johansson and Marc Vaudel for providing summary statistics for childhood BMI. We thank the IMSGC for providing MS GWAS data. We thank MR-Base for making the TwoSampleMR package available publicly. Download links for datasets used in this study are provided above. All code used in this study is available at https://github.com/benjacobs123456/MR_MOBA_MS.

Consent and Approval
This study was performed using publicly available data sources. There were no experiments performed on human subjects and no direct patient contact, therefore written consent was not required.

Results

Age-epoch analysis
We collapsed BMI during childhood into four windows: birth (birth - 6 weeks inclusive), infancy (3 months - 1.5 years inclusive), early childhood (2 years - 5 years inclusive), and later childhood (7 - 8 years). At all time epochs other than birth, genetically-elevated BMI was associated with an increased liability to MS (tables 1 & 2, figure 1, supplementary figures 1&2): birth (OR 0.81, 95% CI 0.50 - 1.31, N_{SNPs}=7, p=0.39), infancy (OR 1.18, 95% CI 1.04 - 1.33, N_{SNPs}=18, p = 0.01), early childhood (OR 1.31, 95% CI 1.03 - 1.66, N_{SNPs}=4, p=0.03), and later childhood (OR 1.34, 95% CI 1.08 - 1.66, N_{SNPs}=4, p=0.01).

The MR-Egger intercepts did not suggest that unbalanced horizontal pleiotropy was biasing the IVW estimate at any of these epochs (supplementary table 1). Cochran's test of heterogeneity suggested heterogeneity in the IVW estimators during the birth epoch, but not during any other time window (supplementary table 1). Using the MR-PRESSO method, we

found evidence of global pleiotropy for the birth epoch (p<0.001) but no other epochs (infancy p=0.68, early childhood p=0.84, later childhood p=0.43). For the birth epoch, a single outlier (rs11187129) was found to distort the overall MR estimate ($P_{Outlier}$< 0.05), and removal of this outlier led to a point estimate closer to the null ($beta_{corrected}$= -0.06, p=0.69).

To determine whether these results could be driven by SNPs acting in the opposite causal direction (i.e. influencing childhood BMI via susceptibility to MS), we excluded any SNPs explaining more variance in the outcome than the exposure (Steiger filtering)[16]. No SNPs were filtered out using this approach, and thus the causal estimates were unchanged. Leave-one-out sensitivity analysis, in which SNPs are removed in turn from the genetic instrument to ensure no one SNP is driving the result, did not lead to an appreciable change in the MR estimates, although there was some loss of precision (supplementary table 2).

Next, to determine whether the observed MR effects of childhood BMI during infancy, early and late childhood on MS risk could be driven by the effects of these genetic variants on BMI in adulthood, we excluded all variants with nominal evidence of association with adult BMI from a large GWAS of ~700,000 adults[12]. Exclusion of these SNPs did not produce a substantial change in the MR estimates during any epoch, although due to smaller numbers of variants the precision was diminished (supplementary tables 3 & 4; supplementary figure 3). For the later childhood epoch, only one SNP was retained and so the Wald Ratio method was the only method available to carry out the MR analysis.

Individual time point analysis
Secondary analysis at each time point demonstrated results consistent with the epoch-based analysis, albeit with less precision at each individual time point due to the smaller number of variants used for the genetic instruments (supplementary figure 4, supplementary table 5). A Mann-Kendall trend test of the 12 individual time points suggested evidence of a trend in the MR effect estimates (IVW or Wald Ratio where only one SNP was available for analysis; tau = 0.636, 2-sided p-value =0.005)

Although there was no evidence of unbalanced horizontal pleiotropy at any of these time points where there were sufficient SNPs in the instrument to test, there was again evidence of notable heterogeneity at the birth BMI time point (supplementary table 6).

To replicate our negative finding that birth weight does not causally impact MS risk using an external dataset, we repeated the analysis using summary statistics from a larger GWAS of birthweight[13]. This yielded a convincing null (OR 1.20, 95% CI 0.92 - 1.55, N_{SNP}=136, p=0.17) with no evidence to suggest unbalanced horizontal pleiotropy (Egger-intercept = -0.01, p=0.18).

Discussion

We provide novel evidence from longitudinal two-sample mendelian randomisation that higher BMI from 3 months onwards, but not at birth, may be a causal risk factor for MS. For the first time, we demonstrate an increase in magnitude of the effect size of raised BMI across the lifecourse from early to late childhood, suggesting that the effect of BMI on MS risk increases during the transition from childhood to adolescence. By removing SNPs associated with adult BMI, we provide evidence that these effects cannot be fully explained by effects of the genetic instruments on later-life BMI. These results provide evidence that

BMI may influence MS risk earlier in life than previously thought, and support the view that early interventions to prevent childhood obesity may help to prevent some cases of MS.

The key novelty here is the use of MR with longitudinal GWAS from the same cohort to explore the time-varying effects of an exposure, in our case childhood BMI, on MS risk. We use a variety of methods to quantify and account for pleiotropy, and demonstrate using the MR-PRESSO method that pleiotropy is unlikely to influence the positive results we observe at the 3 months - 1.5 years, 2 years - 5 years, and 7 years - 8 years epochs. We demonstrate the value of this approach for unpicking the role of time-varying factors on disease risk, demonstrating a clear trend from a null effect of birthweight to a clear effect of BMI in late childhood, consistent with previous epidemiological findings in MS[7,9].

An important caveat is that we cannot exclude the possibility that the observed effects of genetic instruments for childhood BMI on MS risk may be accounted for by the effects of those variants on adult or adolescent BMI. We aim to overcome this concern by removing all SNPs from the genetic instruments weakly associated with adult BMI from a large and therefore well-powered GWAS of ~700,000 individuals which would be expected to detect strong effects. Although this sensitivity analysis diminished our power, it is striking that we still observe a strong effect of BMI from 3 months - 1.5 years on MS risk. This question could potentially be further addressed with multivariable mendelian randomisation, however given the small number of variants for each genetic instrument and the high collinearity of childhood and adult BMI, we have not pursued this line of analysis in this study. As larger GWAS and therefore more genetic instruments become available for childhood BMI at different early life stages, this will be the crucial test. Preliminary analyses we have performed suggest that much of the effect we observe may be mediated by persistence of elevated BMI into later life, however given the instability of these estimates we do not have high confidence in these results. Replication in other longitudinal early-life cohorts, such as the Avon Longitudinal Study of Parents and Children, and discovery of more loci which exert age-specific effects on BMI will help to reinforce or refute our findings.

Our results support and extend previous observations regarding the effects of early life BMI on subsequent MS risk. Observational data supports the concept that BMI during adolescence, particularly late teenage years, is a risk factor for MS[4,7,9,18]. Mendelian randomisation studies have strengthened the notion that this may be a causal relationship[5,6]. It remains debatable whether childhood BMI is a causal risk factor for MS independently of its persistence into adolescence/adulthood, or whether it purely affects MS risk via its strong correlation with BMI in early adult life. In favour of the latter, a recent study used multivariable MR to regress out the effects of genetic instruments for childhood BMI mediated via adult BMI: controlling for the effect of adult BMI abolished the observed MR effect of childhood BMI on MS risk, suggesting that persistently raised BMI into adulthood, rather than transiently increased BMI during childhood, is the causal risk factor[6]. These results, obtained with larger datasets for early life BMI, suggest that what we observe may be driven by the same phenomenon - pleiotropic SNPs acting on MS risk via their effect on BMI in later life rather than in childhood *per se*. Although we address this concern to an extent by excluding SNPs associated with adult BMI, we cannot exclude this possibility.

A weakness of our study is the relatively small number of SNPs included in the genetic instruments at each time point. This motivated our approach of collapsing SNPs into time epochs based on shared biology and kinetics of BMI during these time windows, however we recognise that this represents an arbitrary simplification of the data. Although at individual

time points our MR instruments are under-powered, our epoch instruments all have good power (F statistics > 10) to detect true effects. The clear null effect of birthweight is confirmed using an external, and better-powered, GWAS[13].

A further possible concern is that population stratification may influence our results, as the exposure GWAS (Norwegian) and outcome GWAS (European) are drawn from slightly different populations. There are three reasons why this is not a major problem for our results. First, the MOBA GWAS were performed on a post-Quality Control subset of individuals who cluster tightly with reference EUR samples from the CEU population (correspondence with authors). Second, these GWAS results show a strong genetic correlation with the 'comparative body size at age 10' trait from UK Biobank. Third, allele frequencies for included SNPs are very similar between this GWAS and reference European datasets, such as UKB.

In summary we have provided evidence from longitudinal mendelian randomisation that the relationship between early life BMI and MS seems to follow a gradient, with no effect at birth and a gradually increasing influence on later-life MS risk. The window during which BMI influences MS risk may stretch earlier than previously thought. Further work is required to determine whether these effects are solely mediated via their effect on BMI during early adulthood. Our findings extend previous MR findings in this field and demonstrate the value of studying time-varying risk factors with MR.

Funding

This work was performed at the Preventive Neurology Unit, which is funded by the Barts Charity. BMJ is supported by an MRC Clinical Research Training Fellowship.

Disclosures

Author contributions

BMJ and RD conceived the idea of this work

The primary analysis was performed by LH with support from BMJ

The initial manuscript was drafted by LH with input from BMJ and RD

All authors provided input into critical revisions of the manuscript prior to submission

References

1. Olsson, T., Barcellos, L. F. & Alfredsson, L. Interactions between genetic, lifestyle and environmental risk factors for multiple sclerosis. *Nat. Rev. Neurol.* **13**, 25–36 (2017).

2. Davey Smith, G. & Ebrahim, S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *International journal of* (2003).

3. Davies, N. M., Holmes, M. V. & Davey Smith, G. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. *BMJ* **362**, k601 (2018).

4. Munger, K. L. *et al.* Childhood body mass index and multiple sclerosis risk: a long-term cohort study. *Mult. Scler.* **19**, 1323–1329 (2013).

5. Jacobs, B. M., Noyce, A. J., Giovannoni, G. & Dobson, R. BMI and low vitamin D are causal factors for multiple sclerosis: A Mendelian Randomization study. *Neurol Neuroimmunol Neuroinflamm* **7**, (2020).

6. Harroud, A. *et al.* Childhood obesity and multiple sclerosis: A Mendelian randomization study. *Mult. Scler.* 13524585211001781 (2021).

7. Jacobs, B. M. *et al.* Gene-Environment Interactions in Multiple Sclerosis: A UK Biobank Study. *Neurol Neuroimmunol Neuroinflamm* **8**, (2021).

8. Munger, K. L., Chitnis, T. & Ascherio, A. Body size and risk of MS in two cohorts of US women. *Neurology* **73**, 1543–1550 (2009).

9. Hedström, A. K., Olsson, T. & Alfredsson, L. High body mass index before age 20 is associated with increased risk for multiple sclerosis in both men and women. *Mult. Scler.* **18**, 1334–1336 (2012).

10. Helgeland, Ø. *et al.* Characterization of the genetic architecture of BMI in infancy and early childhood reveals age-specific effects and implicates pathways involved in Mendelian obesity. *bioRxiv* (2021) doi:10.1101/2021.05.04.21256508.

11. International Multiple Sclerosis Genetics Consortium. Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science* **365**, (2019).

12. Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body mass index in 700000 individuals of European ancestry. *Hum. Mol. Genet.* **27**, 3641–3649 (2018).

13. Warrington, N. M. *et al.* Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic risk factors. *Nat. Genet.* **51**, 804–814 (2019).

14. Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the human phenome. *Elife* **7**, (2018).

15. Verbanck, M., Chen, C.-Y., Neale, B. & Do, R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat. Genet.* **50**, 693–698 (2018).

16. Hemani, G., Tilling, K. & Smith, G. D. Orienting the causal relationship between imprecisely measured traits using genetic instruments. bioRxiv. (2017).

17. Brion, M.-J. A., Shakhbazov, K. & Visscher, P. M. Calculating statistical power in Mendelian randomization studies. *International Journal of Epidemiology* vol. 42 1497–1501 (2013).

18. Hedström, A. K. *et al.* Interaction between adolescent obesity and HLA risk genes in the etiology of multiple sclerosis. *Neurology* **82**, 865–872 (2014).

						Exposure (BMI)			Outcome (MS)		
Epoch	SNP	Nearest Gene	Effect allele	Non-effect allele	EAF in MOBA	Beta	SE	P value	Beta	SE	P value
Birth - 6 weeks	rs11187129	*HHEX*	T	C	0.54108	-0.0471726	0.008338	2.10E-08	0.094401	0.01739	5.68E-08
Birth - 6 weeks	rs11708067	*ADCY5*	A	G	0.765083	-0.0789424	0.009755	5.20E-16	-1.00E-04	0.029553	0.9973
Birth - 6 weeks	rs1482853	*CCNL1*	C	A	0.598464	0.0988178	0.008435	5.90E-32	-0.01511	0.016958	0.3728
Birth - 6 weeks	rs2298615	*EHBP1L1*	C	T	0.76651	-0.0712846	0.012196	5.40E-09	0.045929	0.02142	0.03202
Birth - 6 weeks	rs28642213	*GPSM1*	A	G	0.267403	0.0623516	0.009521	4.70E-11	0.02245	0.028404	0.4293
Birth - 6 weeks	rs7310615	*SH2B3*	C	G	0.45479	-0.0498278	0.008508	6.50E-09	0.067752	0.016702	4.98E-05
Birth - 6 weeks	rs739669	*DLG4*	G	A	0.376996	-0.0715704	0.008601	4.70E-17	-0.02358	0.01775	0.1841
Birth - 6 weeks	rs7772579	*ESR1*	A	C	0.699604	0.0651087	0.009067	5.90E-13	0.002297	0.018227	0.8997
3 months - 1.5 years	rs1032296	*HHIP*	T	C	0.376232	0.0524932	0.009097	1.10E-08	0.025831	0.017131	0.1316
3 months - 1.5 years	rs10493544	*TNNI3K*	T	C	0.430744	0.0538075	0.009404	7.10E-09	-0.00638	0.01639	0.6971
3 months - 1.5 years	rs11676272	*ADCY3*	A	G	0.51411	-0.0887022	0.009299	2.80E-22	-0.0203	0.016619	0.2218
3 months - 1.5 years	rs13038017	*EFCAB8*	T	C	0.47431	-0.0535926	0.009372	1.20E-08	0.001601	0.016545	0.9229
3 months - 1.5 years	rs17145750	*MLXIPL*	C	T	0.840536	0.0697872	0.011991	6.80E-09	-0.02706	0.025971	0.2974
3 months - 1.5 years	rs1772945	*OPRM1*	G	A	0.441547	-0.0562618	0.00938	3.20E-09	-0.0031	0.016224	0.8487
3 months - 1.5 years	rs1820721	*GLPR1*	C	A	0.511425	-0.060729	0.008743	7.20E-12	0.016333	0.016275	0.3156
3 months - 1.5 years	rs1985927	*SCGB1A!*	T	C	0.270127	-0.0597242	0.010637	6.80E-09	-0.0244	0.019192	0.2036
3 months - 1.5 years	rs209421	*UBE3D*	G	T	0.264271	0.0727381	0.009951	5.40E-13	0.006521	0.018044	0.7178
3 months - 1.5 years	rs2585058	*SH3GL3*	G	A	0.525095	0.0630845	0.009321	8.60E-12	0.007373	0.016718	0.6592
3 months - 1.5 years	rs2728641	RP11-405A12.2	C	T	0.477989	0.0501517	0.008951	1.90E-08	0.0244	0.016829	0.1471
3 months - 1.5 years	rs2767486	*LEPR*	A	G	0.838584	-0.14325	0.011859	6.40E-34	-0.03816	0.021062	0.07
3 months - 1.5 years	rs2816985	*NR5A2*	A	G	0.551004	-0.0591627	0.009057	5.40E-11	-0.02762	0.016419	0.09259
3 months - 1.5 years	rs28457693	*PTCH1*	A	G	0.865473	-0.0726875	0.01297	2.40E-08	-0.03449	0.029121	0.2363
3 months - 1.5 years	rs287621	*KLF14*	T	C	0.260505	0.0636843	0.010075	3.70E-10	-0.0141	0.018164	0.4376
3 months - 1.5 years	rs3741508	*NCOR2*	T	G	0.856761	0.0826905	0.013414	1.20E-09	0.009356	0.025142	0.7098
3 months - 1.5 years	rs6538845	*RP11-690J15.1*	T	C	0.515215	-0.0549436	0.008957	1.50E-09	-0.02567	0.016303	0.1154
3 months - 1.5 years	rs6899303	PCSK1	A	C	0.368755	-0.057154	0.009025	5.30E-11	-0.00409	0.016591	0.8052
2 years - 5 years	rs11676272	ADCY3	A	G	0.515086	-0.0750482	0.010748	2.70E-12	-0.0203	0.016619	0.2218
2 years - 5 years	rs12672489	*KLF14*	C	T	0.745792	0.0726685	0.012849	1.10E-08	0.015775	0.018633	0.3972
2 years - 5 years	rs1830890	*PLCE1*	A	G	0.678614	-0.0665976	0.011646	1.30E-08	-0.00733	0.017525	0.6759
2 years - 5 years	rs2767486	*LEPR*	A	G	0.837712	-0.0853349	0.014622	4.60E-09	-0.03816	0.021062	0.07
7 years - 8 years	rs10493544	*TNNI3K*	T	C	0.43154	0.0651591	0.01198	4.80E-08	-0.00638	0.01639	0.6971
7 years - 8 years	rs11676272	*ADCY3*	A	G	0.515866	-0.078168	0.013323	2.90E-09	-0.0203	0.016619	0.2218
7 years - 8 years	rs17817288	*FTO*	A	G	0.513158	-0.0952937	0.013447	1.30E-12	-0.03325	0.016187	0.03999
7 years - 8 years	rs545608	*SEC16B*	G	C	0.766209	-0.0877785	0.015795	3.20E-08	-0.01114	0.020384	0.5848
7 years - 8 years	rs7132908	*FAIM2*	G	A	0.598895	-0.0813492	0.01358	3.30E-09	-0.03999	0.016912	0.01805

Table 1: Single Nucleotide Polymorphisms (SNPs) associated with BMI at various time epochs during early life. 'Epoch' refers to the time window in question. For each SNP, the nearest gene, and the beta, standard error (SE) and associated P value are given for the association with standardized BMI (at each time point) and with MS (i.e. the beta is the log odds ratio per effect allele copy). EAF – effect allele frequency. BMI – body mass index. SNP – single nucleotide polymorphism. SE – standard error. MS – multiple sclerosis.

Table 2: MR estimates from the primary analysis of BMI during each time epoch on MS. For each method, the number of included SNPs is included. The IVW (inverse variance weighted) method was used as the primary analytic method. MR – mendelian randomisation.

Exposure epoch	MR method	Number of SNPs	Beta	SE	P value
Birth - 6 weeks	Inverse variance weighted	7	-0.20891	0.243413	0.390749
3 months - 1.5 years	Inverse variance weighted	18	0.162189	0.062848	0.009861
2 years - 5 years	Inverse variance weighted	4	0.267068	0.122666	0.029467
7 years - 8 years	Inverse variance weighted	4	0.289007	0.109746	0.008453
Birth - 6 weeks	MR Egger	7	1.184225	0.981039	0.281374
3 months - 1.5 years	MR Egger	18	0.259406	0.223306	0.262409
2 years - 5 years	MR Egger	4	1.652619	1.447985	0.371971
7 years - 8 years	MR Egger	4	1.340814	0.760078	0.219773
Birth - 6 weeks	Simple mode	7	0.091193	0.189569	0.647517
3 months - 1.5 years	Simple mode	18	0.10474	0.14939	0.492715
2 years - 5 years	Simple mode	4	0.219738	0.197409	0.346823
7 years - 8 years	Simple mode	4	0.349886	0.181151	0.148952
Birth - 6 weeks	Weighted median	7	-0.0515	0.145521	0.723423
3 months - 1.5 years	Weighted median	18	0.179452	0.086761	0.038606
2 years - 5 years	Weighted median	4	0.248544	0.140316	0.076509
7 years - 8 years	Weighted median	4	0.320545	0.127328	0.01182
Birth - 6 weeks	Weighted mode	7	-0.03037	0.149958	0.846181
3 months - 1.5 years	Weighted mode	18	0.196361	0.116829	0.11109
2 years - 5 years	Weighted mode	4	0.231029	0.189487	0.309874
7 years - 8 years	Weighted mode	4	0.347357	0.165152	0.126174