Misdiagnosis prevents accurate monitoring of transmission and burden for sub-critical

pathogens: a case study of *Plasmodium knowlesi* malaria

John H. Huber

Department of Biological Sciences and Eck Institute for Global Health, University of Notre

Dame, Notre Dame, IN, USA

Corresponding Author: huber.j.h@wustl.edu (JHH)

ABSTRACT

Maintaining surveillance of emerging infectious diseases presents challenges for monitoring their

transmission and burden. Incomplete observation of infections and imperfect diagnosis reduce

the observed sizes of transmission chains relative to their true sizes. Previous studies have

examined the effect of incomplete observation on estimates of pathogen transmission and

burden. However, each study assumed that, if observed, each infection was correctly diagnosed.

Here, I leveraged principles from branching process theory to examine how misdiagnosis could

contribute to bias in estimates of transmission and burden for emerging infectious diseases.

Using the zoonotic *Plasmodium knowlesi* malaria as a case study, I found that, even when

assuming complete observation of infections, the number of misdiagnosed cases within a

transmission chain for every correctly diagnosed case could range from 0 $(0 - 4)$ when R_0 was

0.1 to 86 $(0 - 837)$ when R_0 was 0.9. Data on transmission chain sizes obtained using an

imperfect diagnostic could consistently lead to underestimates of R_0, the basic reproduction

number, and simulations revealed that such data on up to 1,000 observed transmission chains

was not powered to detect changes in transmission. My results demonstrate that misdiagnosis

may hinder effective monitoring of emerging infectious diseases and that sensitivity of

diagnostics should be considered in evaluations of surveillance systems.

INTRODUCTION

For pathogens with sub-critical transmission (i.e., $R_0 < 1$), a robust surveillance system that identifies and correctly diagnoses infections is necessary to monitor changes in pathogen transmission and burden (1). Such pathogen surveillance is important both for measuring progress towards elimination of diseases with immediate public health importance, such as measles (2–4) and malaria (5), and for assessing the future threat of emerging infectious diseases (6), such as avian influenza (7), human monkeypox (1,8), and Middle East respiratory syndrome coronavirus (2,9).

Considerable work has been devoted to advance a mathematical framework that leverages the data collected by surveillance systems to obtain estimates of transmission and burden for pathogens with sub-critical dynamics (1,2,4,10,11). These studies have improved our understanding of a wide range of emerging infectious diseases and have critically evaluated the sensitivity of these estimates to the quality of data from the surveillance system. Crucially, each study modeled variation in surveillance quality through variation in the ascertainment fraction (i.e., the proportion of infections that are detected) and assumed that, once detected, all infections were correctly diagnosed. In reality, however, non-specific clinical and biological features are likely to limit the sensitivity of clinical diagnosis, particularly for emerging infectious diseases (12,13). The extent to which misdiagnosis affects estimates of transmission and burden for pathogens with sub-critical dynamics remains largely unaddressed.

The zoonotic *Plasmodium knowlesi* malaria offers a natural case study to examine the impact of misdiagnosis on estimates of transmission and burden. Endemic to Southeast Asia (14), *P. knowlesi* is a vector-borne disease with most or all infections in humans caused by spillover transmission from the long- and pig-tailed macaque reservoir (15,16). The extent of

transmission between humans is currently unknown (17). Due to morphological similarities with

other *Plasmodium* spp., *P. knowlesi* is routinely misdiagnosed by light microscopy (18). A recent

systematic review and meta-analysis estimated that the sensitivity of light microscopy for

diagnosing *P. knowlesi* infections was less than 1% (19). This high rate of misdiagnosis greatly

affects the quality of surveillance data on *P. knowlesi*, potentially biasing estimates of

transmission and burden.

In this study, I aimed to evaluate the extent to which misdiagnosis of a pathogen affected

the ability to monitor its change in transmission and burden. Using *P. knowlesi* as a case study, I

leveraged an established framework based upon branching process theory to first quantify the

potential magnitude of underestimation of pathogen burden on account of misdiagnosis. Next, I

considered how underestimates of pathogen burden could lead to bias in estimates of R_0, the

basic reproduction number. Finally, I quantified the degree to which misdiagnosis reduced the

statistical power to detect changes in transmission from surveillance data for emerging infectious

diseases, such as *P. knowlesi*.

METHODS

Branching Process Framework of Sub-Critical Transmission

To explore the effects of misdiagnosis on the monitoring of sub-critical transmission (i.e., $R_0 <$

1) of *P. knowlesi*, I extended a framework that uses branching process theory to estimate a

pathogen's R_0 from its size distribution of stuttering transmission chains. Here, I followed

Blumberg and Lloyd-Smith (1,11) and defined a transmission chain as a primary infection (i.e., a

spillover infection from a zoonotic reservoir) and all secondary infections arising from that

primary infection through at least one generation of pathogen transmission.

Assuming that the number of secondary infections caused through one generation of

pathogen transmission followed a negative binomial distribution with mean R_0 and dispersion

parameter κ, I used the branching framework to calculate summary statistics of the transmission

chains. Specifically, I solved for the probability that a transmission chain was truly of size j

infections, r_j, and the mean size of transmission chains, μ.

Following Blumberg and Lloyd-Smith (11), I considered two models of observation of

infections: (i) independent observation and (ii) size-dependent observation. The model of

independent observation assumes that each infection is subject to an independent probability of

observation and correct diagnosis, p_{ind}, that is equal to the product of the observation

probability, p_{det}, and the diagnostic sensitivity, se. By comparison, the model of size-dependent

observation assumes that observation of transmission chains occurs through sentinel infections.

Each infection within a transmission chain is a sentinel infection with probability, p_{sent}, and, if

there is at least sentinel infection within the transmission chain, then all infections within the

transmission chain are observed. Diagnosis of each infection occurs independently and is subject

to sensitivity, se.

I then computed the mean observed transmission chain size, μ^*, as a function of the

transmission parameters (R_0 and κ), the observation model (p_{det} or p_{sent}), and the diagnostic

accuracy (se). This allowed me to relate the distribution of observed transmission chain sizes to

the distribution of true transmission chain sizes and quantify bias in the maximum-likelihood

estimates of transmission, $\widehat{R}_0 = 1 - \frac{1}{\mu^*}$. If all infections are observed and correctly diagnosed,

then $\mu^* = \mu$ and thus $\widehat{R}_0 = R_0$. Violations of this assumption, either through incomplete

observation or misdiagnosis, introduce bias into transmission estimates. A full description of the

branching process framework can be found in the Supplement.

Analyses

Quantifying the Bounds of Total Burden

To first demonstrate how misdiagnosis, in addition to incomplete observation, may lead to an

underestimate of *P. knowlesi* burden, I computed the probability distribution of the true size of a

transmission chain conditional upon the observed size of a transmission chain. That is, given that

I observed a transmission chain of size $\hat{j}$, the probability that the transmission chain is truly of

size j is equal to

$$\Pr(j|\hat{j}) = \frac{\Pr(\hat{j}|j)\,\Pr(j)}{\Pr(\hat{j})}. \quad (1)$$

In eq. (1), Pr (j) is the probability that a transmission chain is of size j, r_j, computed using eq.

(S1), and Pr $(\hat{j})$ is the probability that a transmission chain is of observed size j, s'_j, computed

using eq. (S2) for the model of independent observation and using the numerator of eq. (S7) for

the model of size-dependent observation. For the model of independent observation, the

probability of observing a transmission chain of size $\hat{j}$ given that the transmission chain is truly

of size j is

$$\Pr(\hat{j}|j) = \binom{j}{\hat{j}} \cdot p_{ind}^{\hat{j}} \cdot (1 - p_{ind})^{j-\hat{j}}, \quad (2)$$

where p_{ind} is equal to the product of the probability of detection, p_{det}, and the sensitivity of

diagnosis, se. By comparison, for the model of size-dependent observation, this quantity is

computed as

$$\Pr(\hat{j}|j) = \left(1 - (1 - p_{sent})^j\right) \cdot \binom{j}{\hat{j}} \cdot se^{\hat{j}} \cdot (1 - se)^{j-\hat{j}}. \quad (3)$$

Substituting the respective terms into eq. (1), for the model of independent observation, I

computed the probability that a transmission chain is of true size j given that it is observed to be

of size $\hat{j}$ as

$$\Pr(j|\hat{j}) = \frac{\binom{j}{\hat{j}} \cdot p_{ind}^{\hat{j}} \cdot (1 - p_{ind})^{j-\hat{j}} r_j}{s'_j}. \quad (4)$$

For the model of size-dependent observation, I computed this quantity as

$$\Pr(j|\hat{j}) = \frac{\left(1 - (1 - p_{sent})^j\right) \cdot \binom{j}{\hat{j}} \cdot se^{\hat{j}} \cdot (1 - se)^{j-\hat{j}} r_j}{s'_j}. \quad (5)$$

I used eqs. (4-5) to compute the expected true transmission chain sizes given observed

transmission chains of one, two, or three cases while varying the probability of observation, p_{det}

or p_{sent}, from 0.1 to 1.0 in increments of 0.1. I sampled the sensitivity of the diagnostic method

from the posterior estimate of sensitivity of light microscopy for *P. knowlesi* with mean equal to

1.19 x 10^{-3} (19). For each combination of observed chain size and probability of observation, I

calculated the expected true transmission chain size, assuming that the true value of R_0 was

equal to 0.1, 0.5, or 0.9. These values of R_0 represent low, medium, and high values of sub-

critical transmission and fall within the plausible range of human-to-human transmission of *P.*

knowlesi (20). The dispersion parameter κ was assumed to be 0.1 in all scenarios, though a

supplementary analysis was performed where $\kappa \to \infty$.

Effect of Misdiagnosis on Estimates of Transmission

On account of incomplete observation and misdiagnosis, the observed burden of *P. knowlesi* may

not reflect the true burden. It follows that the mean observed transmission chain size will not

equal the true mean transmission chain size, biasing our estimates of R_0. To explore the extent of

this bias in scenarios where R_0 was equal to 0.1, 0.5, or 0.9, I calculated the mean observed

transmission chain sizes while varying the probabilities of observation, p_{det} or p_{sent}, from 0.1 to

1.0 in increments of 0.1 and while using posterior samples of sensitivity of light microscopy for

P. knowlesi (19). I then compared the maximum-likelihood estimates of $\hat{R}_0$ to the true values of

R_0 for both the models of independent observation and size-dependent observation. In all

scenarios, I assumed that the dispersion parameter κ was 0.1, and a supplementary analysis was

performed were $\kappa \to \infty$.

Effect of Misdiagnosis on Statistical Power to Detect Changes in Transmission

Bias in R_0 estimates on account of misdiagnosis could reduce the statistical power to detect

changes in R_0 over time using data on the size of transmission chains. To measure statistical

power as a function of the number of observed transmission chains, I followed an approach taken

by Blumberg *et al*. (2). I assumed that R_0 was historically equal to 0.1 and then increased to

$R_0 + \Delta R_0$, where ΔR_0 was set to 0.1, 0.5, or 0.9. I then simulated N observed transmission chains

and estimated $\hat{R}_0$ while varying N from 1 to 1,000. I then compared the model in which R_0 was

estimated to have changed to the null hypothesis that there was no change in transmission (i.e.,

$\Delta R_0 = 0$) using the Akaike Information Criterion (AIC) (21). For each number of observed

transmission chains N, I repeated this procedure 1,000 times and computed statistical power as

the proportion of simulations in which I detected a change in transmission on the basis of AIC.

To measure the minimum effect of misdiagnosis on statistical power, I set p_{det} and p_{sent} equal

to 1. Because all infections are observed under this assumption and diagnosis is performed

independently across infections in both models, the models of independent and size-dependent

observation yield identical results. In all scenarios, I assumed that the dispersion parameter κ

was 0.1, and a supplementary analysis was performed were $\kappa \to \infty$.

RESULTS

Assuming complete observation of infections (i.e., p_{det} and p_{sent} equal to 1), misdiagnosis of *P.*

knowlesi infections would underestimate the true *P. knowlesi* burden, with the magnitude of this

effect depending upon R_0 (Fig. 1). For a scenario in which R_0 was 0.1, the expected true size of a

transmission chain is one infection (95% CI: $1 - 5$) if the observed size is one case, four

infections ($2 - 12$) if the observed size is two cases, and seven infections ($3 - 17$) if the observed

size is three cases. Under an alternative scenario in which R_0 was 0.9, the expected true size of

the transmission chains increased to 87 ($1 - 838$), 461 ($49 - 965$), and 650 ($157 - 983$)

infections, respectively.

The effect of incomplete observation (i.e., p_{det} or $p_{sent} < 1$) on the expected burden was

most apparent at an intermediate R_0 of 0.5 and with the model of size-dependent observation

(Fig. 1D). Under this scenario, given a transmission chain of size one, the expected true

transmission chain was 18 infections (1 – 107) if p_{sent} was equal to 0.1, compared to 3 infections

(1 – 80) if p_{sent} was equal to 1. In all other scenarios, the expected burden did not change

significantly with p_{det} or p_{sent}. This occurred because, even with complete observation (i.e.,

p_{det} and p_{sent} equal to 1), 99.881% of *P. knowlesi* cases were expected to be misdiagnosed,

given a sensitivity of 0.119%. Therefore, irrespective of the observation probability, only a

subset of true transmission chain sizes is consistent with the sizes of the observed transmission

chains, given this high percentage of false negatives. If I instead assumed perfect sensitivity of

the method, I observed a greater effect of the observation probability on the expected burden

(Fig. S1).

Figure 1. Effect of misdiagnosis and imperfect observation on the expected burden. The mean

true transmission chain size (dots) and 95% CI (segments) are shown conditional upon on an

observed transmission chain size of one (yellow), two (orange), or three (red) cases and an R_0 of

0.1 (A,B), 0.5 (C,D), and 0.9 (E,F). The horizontal axis is the observation probability,

representing p_{det} for the Model of Independent Observation (A, C, E) and p_{sent} for the Model of

Size-Dependent Observation (B, D, F).

Given that misdiagnosis underestimated the burden of *P. knowlesi* in this analysis, I assessed its

effect on my estimates of transmission, $\hat{R}_0$. Because misdiagnosis caused the average observed

size of transmission chains to be less than the average true size of transmission chains, I

consistently underestimated $\hat{R}_0$, with the effect being more severe at lower R_0 (Fig. 2). For

example, with an R_0 of 0.1 and assuming perfect observation of infections (i.e., p_{det} and p_{sent}

equal to 1), my median $\hat{R}_0$ estimate was 1.9 x 10^{-4} (95% PPI: 2.0 x 10^{-5} – 2.9 x 10^{-3}),

corresponding to a 520-fold (34 – 4900) underestimate of transmission. Under an alternative

scenario in which R_0 was 0.9, my median $\hat{R}_0$ was 0.26 (0.046 – 0.66), corresponding to a 3.5-

fold (1.4 – 19.4) underestimate in transmission.

My estimates of transmission were sensitive to the simulated observation probability,

though the direction of the effect depended upon the assumed model of observation. For the

model of independent observation, $\hat{R}_0$ estimates increased with increasing p_{det}, because the

average observed size of transmission chains increased as more infections were observed (Fig. 2,

left column). By contrast, $\hat{R}_0$ estimates decreased with increasing p_{sent} for the model of size-

dependent observation (Fig. 2, right column). This counterintuitive effect can be explained by the

observation that, if p_{sent} is low, larger transmission chains have a greater probability that at least

one infection is a sentinel infection. This causes a bias in the size of the transmission chains that

are observed at low values of p_{sent}, increasing the mean observed transmission chain size

relative to that at higher values of p_{sent} and thus inflating the $\widehat{R}_0$ estimate.

Figure 2. Effect of misdiagnosis and imperfect observation on estimates of transmission. *The*

posterior median (blue line) and 95% posterior prediction interval (blue shaded region) of

maximum-likelihood estimates of R_0 are shown as a function of the observation probability. The

observation probability represents p_{det} for the Model of Independent Observation (A, C, E) and

p_{sent} for the Model of Size-Dependent Observation (B, D, F). The solid black denotes the true R_0

in each panel, and the grey lines denote two-to-five-fold underestimates of R_0 in each panel.

The underestimates of burden (Fig. 1) and transmission (Fig. 2) indicated that

misdiagnosis of *P. knowlesi* may affect the statistical power to detect changes in transmission

based on the size of observed transmission chains. To test this, I simulated changes in

transmission and measured the statistical power to detect that change. I observed that, under

scenarios in which R_0 increased by 0.9, data on 1,000 observed transmission chains provided

only 10.3% power using an imperfect diagnostic method, compared to 100% if using a perfect

diagnostic method (Fig. 3). At smaller increases in R_0, data on observed transmission chain sizes

obtained using an imperfect diagnostic method had effectively no power to detect a change in

transmission.

Figure 3. Effect of misdiagnosis on the statistical power to detect changes in transmission. *The*

statistical power (%) to detect an increase in transmission is shown as a function of the number

of observed chains for a transmission increase (ΔR_0) of 0.1 (yellow), 0.5 (orange), and 0.9 (red).

Solid lines and points represent an imperfect diagnostic method (i.e., LM) and the dotted lines

represent perfect diagnosis (i.e., PCR).

DISCUSSION

Obtaining accurate estimates of transmission and burden is important for monitoring the

emergence of infectious diseases. Previous studies have explored the extent to which incomplete

observation of infections affects the estimates of transmission for such pathogens (1,2,7,11). In

this work, I built upon these studies by considering the effect of misdiagnosis on estimates of

pathogen transmission and burden. Using the zoonotic *P. knowlesi* malaria as a case study, I

found that misdiagnosis—independent of incomplete observation of infections—may cause us to

underestimate the transmission and burden of pathogens with sub-critical dynamics and hinders

effective, prospective monitoring of changes in transmission.

My results demonstrate that, for pathogens with sub-critical transmission, misdiagnosis

leads to an underestimate of overall pathogen burden. Depending upon the R_0 simulated, I found

that there could be as many as 86 misdiagnosed cases, on average, for each correctly diagnosed

case of *P. knowlesi*, even when assuming complete observation of infections. This effect

increased under select settings when the assumption of complete observation of infections was

relaxed. The underestimation of burden due to misdiagnosis has the potential to shape our

epidemiological understanding of an emerging pathogen. For instance, singleton cases of a

zoonotic pathogen, such as *P. knowlesi*, are commonly assessed as dead-end spillover events

from the zoonotic reservoir (17). However, my simulations suggest that such singleton cases

could instead represent a broad range of epidemiological outcomes, spanning dead-end spillover

events to larger transmission chains.

Due to its effect on observed pathogen burden, misdiagnosis contributed a downward

bias in estimates of transmission. Except for scenarios in which the true simulated R_0 was close

to one, my maximum-likelihood estimates of R_0 approached zero, representing situations in

which we would incorrectly conclude that human-to-human transmission of the pathogen was

unlikely to be occurring. For every scenario considered, the estimate of R_0 was less than the true

value, indicating that bias due to misdiagnosis exceeds the competing positive bias from

incomplete observation when assuming size-dependent observation (1). For pathogens such as *P.*

knowlesi, these simulation results suggest that, in settings where misdiagnosis is common, the

extent of human-to-human transmission could be greater than previously thought. To date, it has

been believed that nearly all cases of *P. knowlesi* in humans are caused by spillover from long-

tailed and pig-tailed macaques, the zoonotic reservoir (17). The lack of observed human-to-

human transmission may be explained by multiple factors, including low parasite densities in

humans (16) and restricted vector habitat preference (15), and is supported by a lack of genetic

diversity across human *P. knowlesi* infections (22). Nevertheless, human-to-human transmission

of *P. knowlesi* has been demonstrated experimentally (23), and these results suggest that, if or

when human-to-human transmission occurs, misdiagnosis could cause us to underestimate its

magnitude.

Finally, I demonstrated that data on the sizes of transmission chains diagnosed using a

diagnostic with realistic sensitivity would be insufficient to monitor changes in transmission.

Even with 1,000 observed transmission chains, I calculated a power of only 10% to detect an

increase in R_0 from 0.1 to 1. This empirical power calculation assumed complete observation of

infections, so it represents an upper bound on the statistical power that we might expect if a

diagnostic with realistic sensitivity was used. Therefore, more sensitive diagnostics, such as

polymerase chain reaction, may be needed to detect changes in transmission that could result

from pathogen evolution (24), among other factors (25–27).

This analysis is subject to a number of limitations. First, the conclusions that I reached

were based upon simulated data only. I used simulations representative of *P. knowlesi* to

illustrate possible outcomes that may occur due to misdiagnosis (19,20), yet I lacked empirical

data on the distribution of transmission chain sizes for *P. knowlesi*. As such, this analysis is not

estimating the true extent of human-to-human transmission of *P. knowlesi*. Second, methods

exist to account for incomplete observation of infections in estimates of R_0 (11). However, as

noted by Blumberg *et al*. (2), it is challenging to estimate the proportion of infections that are

captured by the surveillance system. Consequently, these calculations were conditioned upon the

assumption of complete observation and perfect diagnoses, so violations therein should be

interpreted as the upper bound on the bias that would likely be observed. Finally, I considered a

single pathogen in isolation, though misdiagnosis is commonly due to co-circulation of related

pathogens (12,13,18). Accounting for the upward bias due to false-positive diagnoses from other

pathogens and exploring the magnitude of this effect across epidemiological settings could be

important directions for future work.

ACKNOWLEDGMENTS

I acknowledge funding from a Graduate Research Fellowship from the National Science

Foundation and a Richard and Peggy Notebaert Premier Fellowship from the University of Notre

Dame. The funders had no role in the study design, in the collection, analysis, and interpretation

of data, in the writing of the report, or in the design to submit the article for publication. I thank

Alex Perkins for helpful comments on this manuscript.

SUPPLEMENT

Methods

Mean Transmission Chain Size

Complete Observation and Correct Diagnosis

For a pathogen with sub-critical transmission dynamics, I modeled the number of offspring

caused by a single infection through one generation of transmission as a negative binomial

distribution with mean R_0 and dispersion parameter κ (1,11). Therefore, it follows that

transmission chains of size j occur with probability,

323
$$r_j = \frac{\Gamma(\kappa j + j - 1)}{\Gamma(\kappa j)\Gamma(j+1)} \frac{\left(R_0/\kappa\right)^{j-1}}{\left(1 + \left(R_0/\kappa\right)\right)^{\kappa j + j - 1}}. \quad (S1)$$

In eq. (S1), $\Gamma(\cdot)$ is the gamma function. Because $R_0 < 1$, the mean transmission chain size μ can

be calculated as the mean of a geometric series with common ratio R_0 and is equal to $\frac{1}{1-R_0}$.

Incomplete Observation and Imperfect Diagnosis

In the case of *P. knowlesi* and many other pathogens, the size of transmission chains that are

identified by a surveillance system will be affected by two factors. First, infections in a

transmission chain may not present within the health system, due to a lack of symptoms or

access to treatment. Second, infections in the transmission chain that do present within the health

system may be misdiagnosed and thus inaccurately recorded within the surveillance system.

Both factors will make the observed transmission chain appear smaller in size than the true

transmission chain. Previous work by Blumberg and Lloyd-Smith (1) has quantified the effect of

two models of incomplete observation on estimates of transmission and burden. Here, I build

upon this work by integrating the effect of (mis)diagnosis of infections that occurs secondary to

the observation of infections within the health system.

<u>Model of Independent Observation</u>

The first model of incomplete observation and diagnosis assumes that each individual is subject

to an independent probability p_{ind} equal to the product of observation probability, p_{det}, and the

sensitivity of the diagnostic method, se. Therefore, the probability that we observe and correctly

diagnose j cases from a transmission chain is equal to

346
$$s_j' = \sum_{k=j}^{\infty} r_k \cdot \binom{k}{j} \cdot p_{ind}^j \cdot (1 - p_{ind})^{k-j}, \quad (S2)$$

348   where r_k is the probability that a transmission chain is of true size k, calculated using eq. (S1).

349   The probability that a transmission chain is of observed size j is equal to the normalized

350   probability of s_j', computed as

352
$$r_j' = \frac{s_j'}{1 - s_0'}. \quad (S3)$$

In eq. (S3), s_0' is the probability that a transmission chain is neither observed nor correctly

diagnosed. Due to incomplete observation and misdiagnosis, the probability that a transmission

chain is of observed size j, r_j', is not equal to the probability that a transmission chain is of true

size j, r_j. Finally, because each infection within the transmission chain is subject to an

independent probability of observation and correct diagnosis, the probability p_{obs} that a

randomly sampled infection is observed and correctly diagnosed is equal to p_{ind}.

Model of Size-Dependent Observation

The alternative model assumes that transmission chains are observed through a sentinel

infection, such that, if the sentinel infection is observed, then all other infections in the

transmission chain will be observed. Incorporating the effect of imperfect diagnosis, the

probability that we do not observe a transmission chain of size j is equal to

367
$$v_j = (1 - p_{sent})^j + \left(1 - (1 - p_{sent})^j\right)(1 - se)^j. \quad (S4)$$

In eq. (S4), the first term in the summation is the probability that none of the j infections of the

transmission chain are a sentinel infection, and the second term in the summation is the product

of the probability that at least one infection is a sentinel infection (i.e., the probability that we

observe the transmission chain) and the probability that all j infections are misdiagnosed. Using

eq. (S4), I calculated the probability that a transmission chain is neither observed nor correctly

diagnosed as

376
$$s_0' = \sum_{k=1}^{\infty} r_k \cdot v_k. \quad (S5)$$

378 The probability that a transmission chain is of observed size j is then equal to

380
$$r_j' = \frac{\sum_{k=j}^{\infty} r_k \cdot (1 - (1 - p_{sent})^k) \cdot \binom{k}{j} \cdot se^j \cdot (1 - se)^{k-j}}{1 - s_0'}, \quad (S6)$$

382   and the probability that a randomly chosen infection is observed and correctly diagnosed is equal

383   to

385
$$p_{obs} = \frac{\sum_{j=1}^{\infty} j \cdot r_j \cdot \left(1 - (1 - p_{sent})^j\right) \cdot se}{\mu}. \quad (S7)$$

387   Eq. (S7) accounts for the probability that non-sentinel infections are detected, a quantity that

388   increases as a function of the transmission chain size.

390   <u>Mean Transmission Chain Size</u>

391   For both the model of independent observation and the model of size-dependent observation, the

392   mean observed size of transmission chains is calculated as

394
$$\mu^* = \sum_{j=1}^{\infty} j \cdot r_j' = \frac{p_{obs} \cdot \mu}{1 - s_0'}. \quad (S8)$$

396   **Results**

Figure S1. Effect of misdiagnosis and imperfect observation on the expected true pathogen burden assuming perfect diagnosis. *The mean true transmission chain size (dots) and 95% CI (segments) are shown conditional upon on an observed transmission chain size of one (yellow), two (orange), or three (red) cases and an R_0 of 0.1 (A,B), 0.5 (C,D), and 0.9 (E,F). The horizontal axis is the observation probability, representing p_{det} for the Model of Independent Observation (A, C, E) and p_{sent} for the Model of Size-Dependent Observation (B, D, F).*

Figure S2. Effect of misdiagnosis and imperfect observation on the expected true pathogen burden when κ → ∞. *The mean true transmission chain size (dots) and 95% CI (segments) are shown conditional upon on an observed transmission chain size of one (yellow), two (orange), or three (red) cases and an R_0 of 0.1 (A,B), 0.5 (C,D), and 0.9 (E,F). The horizontal axis is the observation probability, representing p_{det} for the Model of Independent Observation (A, C, E) and p_{sent} for the Model of Size-Dependent Observation (B, D, F).*

413

Figure S3. Effect of misdiagnosis and imperfect observation on estimates of transmission

when κ → ∞. *The posterior median (blue line) and 95% posterior prediction interval (blue*

shaded region) of maximum-level estimates of R_0 are shown as a function of the observation

probability. The observation probability represents p_{det} for the Model of Independent

Observation (A, C, E) and p_{sent} for the Model of Size-Dependent Observation (B, D, F). The solid

black denotes the true R_0 in each panel, and the grey lines denote two-to-five-fold underestimates

of R_0 in each panel.

421

Figure S4. Effect of misdiagnosis on the statistical power to detect changes in transmission

when $\kappa \to \infty$. The statistical power (%) to detect an increase in transmission is shown as a

function of the number of observed chains for a transmission increase (ΔR_0) of 0.1 (yellow), 0.5

(orange), and 0.9 (red). Solid lines and points represent an imperfect diagnostic method (i.e.,

LM) and the dotted lines represent perfect diagnosis (i.e., PCR).

REFERENCES

1. Blumberg S, Lloyd-Smith JO. Inference of R(0) and transmission heterogeneity from the size distribution of stuttering chains. PLoS Comput Biol. 2013;9(5):e1002993.

2. Blumberg S, Funk S, Pulliam JRC. Detecting Differential Transmissibilities That Affect the
Size of Self-Limited Outbreaks. Wilke CO, editor. PLoS Pathog. 2014 Oct
30;10(10):e1004452.

3. Blumberg S, Worden L, Enanoria W, Ackley S, Deiner M, Liu F, et al. Assessing Measles
Transmission in the United States Following a Large Outbreak in California. PLoS Curr
[Internet]. 2015 [cited 2021 Jul 27]; Available from:
https://currents.plos.org/outbreaks/article/assessing-measles-transmission-in-the-united-
states-following-a-large-outbreak-in-california/

4. Farrington CP. Branching process models for surveillance of infectious diseases controlled
by mass vaccination. Biostatistics. 2003 Apr 1;4(2):279–95.

5. Churcher TS, Cohen JM, Novotny J, Ntshalintshali N, Kunene S, Cauchemez S. Measuring
the path toward malaria elimination. Science. 2014 Jun 13;344(6189):1230–2.

6. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to
zoonotic spillover. Nat Rev Microbiol. 2017 Aug;15(8):502–10.

7. Ferguson NM. PUBLIC HEALTH: Enhanced: Public Health Risk from the Avian H5N1
Influenza Epidemic. Science. 2004 May 14;304(5673):968–9.

8. Ambrose MR, Kucharski AJ, Formenty P, Muyembe-Tamfum J-J, Rimoin AW, Lloyd-Smith
JO. Quantifying transmission of emerging zoonoses: Using mathematical models to
maximize the value of surveillance data [Internet]. Epidemiology; 2019 Jun [cited 2021 Jul
27]. Available from: http://biorxiv.org/lookup/doi/10.1101/677021

9. Cauchemez S, Nouvellet P, Cori A, Jombart T, Garske T, Clapham H, et al. Unraveling the
drivers of MERS-CoV transmission. Proc Natl Acad Sci USA. 2016 Aug 9;113(32):9081–6.

10. De Serres G, Gay NJ, Farrington CP. Epidemiology of Transmissible Diseases after
Elimination. American Journal of Epidemiology. 2000 Jun 1;151(11):1039–48.

11. Blumberg S, Lloyd-Smith JO. Comparing methods for estimating R0 from the size
distribution of subcritical transmission chains. Epidemics. 2013 Sep;5(3):131–45.

12. Oidtman RJ, España G, Perkins TA. Co-circulation and misdiagnosis led to underestimation
of the 2015–2017 Zika epidemic in the Americas. Lacerda MVG, editor. PLoS Negl Trop
Dis. 2021 Mar 1;15(3):e0009208.

13. Glennon EE, Jephcott FL, Oti A, Carlson CJ, Bustos Carillo FA, Hranac CR, et al.
Syndromic detectability of haemorrhagic fever outbreaks [Internet]. Epidemiology; 2020
Mar [cited 2021 Jul 27]. Available from:
http://medrxiv.org/lookup/doi/10.1101/2020.03.28.20019463

14. Shearer FM, Huang Z, Weiss DJ, Wiebe A, Gibson HS, Battle KE, et al. Estimating
Geographical Variation in the Risk of Zoonotic Plasmodium knowlesi Infection in Countries

Eliminating Malaria. Churcher TS, editor. PLoS Negl Trop Dis. 2016 Aug 5;10(8):e0004915.

15. Collins WE. *Plasmodium knowlesi* : A Malaria Parasite of Monkeys and Humans. Annu Rev Entomol. 2012 Jan 7;57(1):107–21.

16. Kantele A, Jokiranta TS. Review of cases with the emerging fifth human malaria parasite, Plasmodium knowlesi. Clin Infect Dis. 2011 Jun;52(11):1356–62.

17. Feachem RGA, Chen I, Akbari O, Bertozzi-Villa A, Bhatt S, Binka F, et al. Malaria eradication within a generation: ambitious, achievable, and necessary. The Lancet. 2019 Sep;394(10203):1056–112.

18. Barber BE, William T, Grigg MJ, Yeo TW, Anstey NM. Limitations of microscopy to differentiate Plasmodium species in a region co-endemic for Plasmodium falciparum, Plasmodium vivax and Plasmodium knowlesi. Malar J. 2013 Jan 8;12:8.

19. Huber JH, Elliott M, Koepfli C, Perkins A. The impact of emerging Plasmodium knowlesi on accurate diagnosis by light microscopy: a systematic review and modelling analysis [Internet]. Epidemiology; 2021 Sep [cited 2021 Sep 13]. Available from: http://medrxiv.org/lookup/doi/10.1101/2021.09.08.21263294

20. Imai N, White MT, Ghani AC, Drakeley CJ. Transmission and Control of Plasmodium knowlesi: A Mathematical Modelling Study. Churcher TS, editor. PLoS Negl Trop Dis. 2014 Jul 24;8(7):e2978.

21. Akaike H. A new look at the statistical model identification. IEEE Trans Automat Contr. 1974 Dec;19(6):716–23.

22. Lee K-S, Divis PCS, Zakaria SK, Matusop A, Julin RA, Conway DJ, et al. Plasmodium knowlesi: Reservoir Hosts and Tracking the Emergence in Humans and Macaques. Kazura JW, editor. PLoS Pathog. 2011 Apr 7;7(4):e1002015.

23. Alpert E, Collins WE, Jeter MH, Chin W, Contacos PG. Experimental Mosquito-Transmission of Plasmodium Knowlesi to Man and Monkey. The American Journal of Tropical Medicine and Hygiene. 1968 May 1;17(3):355–8.

24. Geoghegan JL, Senior AM, Di Giallonardo F, Holmes EC. Virological factors that increase the transmissibility of emerging human viruses. Proc Natl Acad Sci USA. 2016 Apr 12;113(15):4170–5.

25. Faust CL, McCallum HI, Bloomfield LSP, Gottdenker NL, Gillespie TR, Torney CJ, et al. Pathogen spillover during land conversion. Ostfeld R, editor. Ecol Lett. 2018 Apr;21(4):471–83.

26. Baeza A, Santos-Vega M, Dobson AP, Pascual M. The rise and fall of malaria under land-use change in frontier regions. Nat Ecol Evol. 2017 May;1(5):0108.

502   27. Morse SS. Factors in the Emergence of Infectious Diseases. Emerg Infect Dis. 1995
503   Mar;1(1):7–15.