Safety and Immunogenicity of an Inactivated Recombinant Newcastle Disease Virus Vaccine

Expressing SARS-CoV-2 Spike: Interim Results of a Randomised, Placebo-Controlled, Phase 1/2 Trial

Authors

Prof Punnee Pitisuttithum, MD[1], Viravarn Luvira, MD[1], Saranath Lawpoolsri, MD[1], Sant

Muangnoicharoen, MD[1], Supitcha Kamolratanakul, MD[1], Chaisith Sivakorn, MD[1], Piengthong Narakorn,

MS[2], Somchaiya Surichan, PhD[2], Sumalee Prangpratanporn, MS[2], Suttida Puksuriwong, PhD[2], Steven

Lamola, MD[3], Laina D Mercer, PhD[3], Rama Raghunandan, PhD[3], Weina Sun, PhD[4], Yonghong Liu, PhD[4],

Juan Manuel Carreño. PhD[4], Rami Scharf, DSc[3], Weerapong Phumratanaprapin, MD[1], Fatima Amanat,

PhD[4], Luc Gagnon, PhD[9], Ching-Lin Hsieh, PhD[10], Ruangchai Kaweepornpoj, MS[2], Sarwat Khan, MS[9],

Manjari Lal, PhD[3], Stephen McCroskery, MS[4], Prof Jason McLellan, PhD[10], Ignacio Mena, PhD[4,5], Marcia

Meseck, MS[7], Benjaluck Phonrat, MS[1], Yupa Sabmee, MS[1], Ratsamikorn Singchareon, MS[2], Stefan

Slamanig, MS[4], Nava Suthepakul, MS[2], Johnstone Tcheou, BS[4], Narumon Thantamnu, BS[1], Sompone

Theerasurakarn, MS[2], Steven Tran, MS[9], Thanakrit Vilasmongkolchai, MS[2], Jessica A White, PhD[3], Prof

Adolfo Garcia-Sastre, PhD[4,5,6,7], Prof Peter Palese, PhD[4,6], Prof Florian Krammer, PhD[4,8], Kittisak

Poopipatpol, BS[2], Ponthip Wirachwong, PhD[2], Richard Hjorth, PhD[3], Bruce L Innis, MD[3]

Author Affiliations:

[1] Faculty of Tropical Medicine, Mahidol University, Faculty of Tropical Medicine, Mahidol University

420/6 Ratchawithi Road, Ratchathewi, Bangkok 10400. Thailand

[2] The Government Pharmaceutical Organization, 1 Thanon Rama VI, Thung Phaya Thai, Ratchathewi,

Bangkok 10400, Thailand

[3] PATH, 2201 Westlake Avenue, Suite 200, Seattle, WA 98121, USA

[4] Department of Microbiology, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Pl, New York,

NY 10029, USA

[5] Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai, 1 Gustave L.

Levy Pl, New York, NY 10029, USA

[6] Department of Medicine, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Pl, New York, NY

10029, USA

[7] The Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Pl, New York, NY

10029, USA

[8] Department of Pathology, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Pl, New York, NY

10029, USA

[9] Nexelis, 525 Bd Cartier O, Laval, QC H7V 3S8, Canada

[10] College of Natural Sciences, The University of Texas at Austin, 120 Inner Campus Dr Stop G2500,

Austin, TX 78712, USA

Correspondence to:

Bruce L Innis, MD, PATH, 2201 Westlake Avenue, Suite 200, Seattle, WA 98121, USA; binnis@path·org

ClinicalTrials·gov Identifier: NCT04764422

Summary

Background

Production of affordable coronavirus disease 2019 (COVID-19) vaccines in low- and middle-income

countries is needed. NDV-HXP-S is an inactivated egg-based Newcastle disease virus vaccine expressing

the spike protein of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It's being developed

in Thailand, Vietnam, and Brazil; herein are initial results from Thailand.

Methods

This phase 1 stage of a randomised, dose-escalation, observer-blind, placebo-controlled, phase 1/2 trial

was conducted at the Vaccine Trial Centre, Mahidol University (Bangkok). Healthy adults aged 18-59

years, non-pregnant and negative for SARS-CoV-2 antibodies were eligible. Participants were block

randomised to receive one of six treatments by intramuscular injection twice, 28 days apart: 1

µg±CpG1018 (a toll-like receptor 9 agonist), 3 µg±CpG1018, 10 µg, or placebo. Participants and

personnel assessing outcomes were masked to treatment. The primary outcomes were solicited and

spontaneously reported adverse events (AEs) during 7 and 28 days after each vaccination, respectively.

Secondary outcomes were immunogenicity measures (anti-S IgG and pseudotyped virus neutralisation).

An interim analysis assessed safety at day 57 in treatment-exposed individuals and immunogenicity

through day 43 per protocol. ClinicalTrials.gov (NCT04764422).

Findings

Between March 20 and April 23, 2021, 377 individuals were screened and 210 were enrolled (35 per

group); all received dose one; five missed dose two. The most common solicited AEs among vaccinees,

all predominantly mild, were injection site pain (<63%), fatigue (<35%), headache (<32%), and myalgia

(<32%). The proportion reporting a vaccine-related AE ranged from 5·7% to 17·1% among vaccine

groups and was 2·9% in controls; there was no vaccine-related serious adverse event. The 10 µg

formulation's immunogenicity ranked best, followed by 3 µg+CpG1018, 3 µg, 1 µg+CpG1018, and 1 µg

formulations. On day 43, the geometric mean concentrations of 50% neutralising antibody ranged from

122·23 IU/mL (1 µg, 95% CI 86·40-172·91) to 474·35 IU/mL (10 µg, 95% CI 320·90-701·19), with 93·9% to

100% of vaccine groups attaining a ≥4-fold increase over baseline.

Interpretation

NDV-HXP-S had an acceptable safety profile and potent immunogenicity. The 3 µg and 3 µg+CpG1018

formulations advanced to phase 2.

Funding

National Vaccine Institute (Thailand), National Research Council (Thailand), Bill & Melinda Gates

Foundation, National Institutes of Health (USA)

Word count = 353

Introduction

There remains a shocking imbalance in the global distribution of coronavirus disease 2019 (COVID-19) vaccines.[1] To achieve control of the COVID-19 pandemic in low- and middle-income countries (LMICs) where most of the global population resides, there must be a great increase in sustainable supply of affordable vaccines. The manufacturing capacity for egg-based inactivated influenza vaccines (IIV) is among the largest in the world; these facilities, some in middle-income countries and operating for less than six months per year, use locally produced embryonated eggs to make more than a billion doses annually of affordable human vaccines.[2] To enable these manufacturers to respond to the COVID-19 pandemic, we developed a COVID-19 vaccine for production in eggs, based on a Newcastle disease virus (NDV) expressing the ectodomain of a novel membrane-anchored, prefusion-stabilized severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spike protein construct, wherein virions are purified and inactivated (NDV-HXP-S).[3-5]

From September to November 2020, manufacturers in Thailand, Vietnam, and Brazil modified their IIV manufacturing process to optimize production of beta-propiolactone (BPL)-inactivated NDV-HXP-S, achieving high yields at pilot scale; the result was three similar processes. A preclinical evaluation of their vaccine candidates, formulated with and without CpG1018, a toll-like receptor 9 agonist adjuvant (Dynavax Technologies)[6] confirmed that they were highly immunogenic and protective in hamsters[5] with no sign of toxicity in rats at the maximum human doses planned for evaluation (3 μg S protein+1·5 mg CpG1018; 10 μg S protein). These results enabled all three manufacturers to initiate clinical development of their vaccine candidates. Herein, we report interim safety and immunogenicity data generated in the phase 1 portion of an adaptive phase 1/2 clinical trial evaluating the NDV-HXP-S vaccine candidate developed by The Government Pharmaceutical Organization of Thailand (GPO). These results provide the first evidence in humans that the NDV vector technology expressing a six-proline

prefusion-stabilized spike protein construct offers a unique platform for affordable manufacturing of a

well-tolerated and highly immunogenic COVID-19 vaccine.

Methods

Study design and participants

The phase 1 segment of a randomised, observer-blind, placebo-controlled, phase 1/2 trial was

conducted at the Vaccine Trial Centre, Faculty of Tropical Medicine, Mahidol University (Bangkok,

Thailand). Participants were recruited from individuals known to the Centre and through

advertisements. Healthy adults 18–59 years of age with body mass index 17 to 40 kg/m^2,-negative

hepatitis B surface antigen and SARS-CoV-2, HIV, and hepatitis C antibody tests were eligible to

participate. A negative urinary pregnancy test was required of women having reproductive capacity

prior to administration of each study vaccine dose. Complete eligibility criteria are described in the trial

protocol provided in the supplementary materials.

Written informed consent was obtained from all participants. The trial complied with the Declaration of

Helsinki and Good Clinical Practice. This study was approved by the Ethics Committee of the Faculty of

Tropical Medicine, Mahidol University (TMEC 21-005) and authorized by the Thailand Food and Drug

Administration (FDA-21-018).

Randomisation and masking

Enrolled subjects were randomly assigned in sequence to one of 6 equal groups (vaccine containing 1 µg

S ± 1·5mg CpG1018 adjuvant, 3 µg S ± 1·5mg CpG1018 adjuvant, 10 µg S, or saline placebo). Subjects

were enrolled in stages, each including active treatment and placebo groups, using a computer-

generated block randomisation sequence prepared by an independent statistician; an unblinded

pharmacist team dispensed each treatment according to the randomization sequence. The first 18

subjects (sentinel cohort) were enrolled to three sequential sentinel groups; 3:1, 1 µg and placebo;

3:3:1, 3 µg or 1 µg+CpG1018 and placebo; and 3:3:1, 10 µg or 3 µg+CpG1018 and placebo, After safety

data were reviewed for the sentinel groups, the next 192 subjects were randomised in 5-dose-cohorts;

32:6, 1 µg and placebo; 32:6, 3 µg and placebo; 32:6, 1 µg+CpG1018 and placebo; 32:7, 10 µg and

placebo; and 32:7, 3 µg+CpG1018 and placebo. All participants and personnel other than the unmasked

pharmacy team and vaccinators were masked to treatment.

Procedures

The recombinant NDV-HXP-S vaccine was manufactured according to current Good Manufacturing

Practice by the GPO in their Influenza Vaccine Plant (Saraburi, Thailand) using locally procured

embryonated eggs inoculated with a master virus seed made and extensively tested for adventitious

agents by the Icahn School of Medicine at Mount Sinai (New York, USA). After incubation for 72 hours at

37°C, eggs were chilled overnight at 4°C, then the allantoic fluids were harvested, clarified, and

concentrated. Recombinant virus particles were purified from the concentrated harvest by two

sequential continuous flow sucrose gradient centrifugations, diafiltered against phosphate-buffered

saline (PBS), inactivated by treatment with 1:4000 BPL for 24 hours at 4°C, and 0.2 micron filter-

sterilized. Vaccine potency was measured by direct enzyme-linked immunosorbent assay (ELISA) using a

human monoclonal antibody (CR3022[7]) to SARS-CoV-2 spike glycoprotein S1 (LakePharma Inc) and an

NDV-HXP-S standard that had been calibrated to a purified HXP-S reference[8] by sodium dodecyl sulphate

polyacrylamide gel electrophoresis (SDS-PAGE) densitometry.

Unmasked staff administered study treatments by intramuscular injection of 0·5 mL on study days 1 and

29. Blood samples were drawn and clinical assessments were done for safety and immunogenicity

endpoints before vaccination on days 1 (dose one), 8, 29 (dose two), 36, and 43; a clinical assessment

for safety only on day 57 was the last timepoint considered for this interim analysis of the phase 1

cohort, although there will be additional immunogenicity and safety assessments on study days 197 and

365. Subjects were observed in the clinic for 30 minutes after each vaccination and were asked to record

any adverse events using paper diary cards during the 7-days after each vaccination. Subjects randomly

allocated to a cell-mediated immunity subset (N=12 per 10 µg, 3 µg+CpG1018, and placebo groups) had

additional blood collected on days 1 and 43 for isolation of peripheral blood mononuclear cells (PBMCs);

these were stored in liquid nitrogen until analysed.

Solicited injection site reactions (pain, tenderness, swelling, induration, erythema) and systemic

symptoms (headache, fatigue, malaise, myalgia, arthralgia, nausea, vomiting, and fever defined as oral

temperature ≥38°C) were recorded by subjects in a diary card that included intensity, then reported by

the investigators; these events were not assessed for causality. Subjects also recorded spontaneously

reported adverse events (AEs) for 28 days; the investigator reported these after grading them for

intensity and categorizing them as serious or not. The investigator also identified the following AEs of

special interest: potential immune-mediated medical conditions (PIMMCs), and AEs of special interest

associated with COVID-19. Intensity of AEs was graded 1-4 as follows: 1 or mild (minimal interference

with daily activities), 2 or moderate (interferes with but does not prevent daily activities), 3 or severe

(prevents daily activities, intervention required), and 4 or very severe (medical intervention required to

prevent disability or death). Investigators assessed unsolicited adverse events for causality (related to

vaccination or not). AEs were graded according to U.S. Department of Health and Human Services

severity grading tables (Food and Drug Administration, Center for Biologics Evaluation and Research

[September 2007] and National Institutes of Health, Division of AIDS [version 2.1, July 2017]). A protocol

safety review committee regularly reviewed blinded safety data; a Data Safety Monitoring Board

monitored unblinded safety data and recommended two formulations for advancing to phase 2.

We measured total anti-SARS-CoV-2 spike (S) IgG using a validated indirect ELISA at Nexelis (Laval,

Canada). Purified recombinant SARS-CoV-2 pre-fusion spike (Nexelis) at 1µg/ml in phosphate buffered

saline (PBS, Wisent Bioproducts) was adsorbed to 96 well Nunc Maxisorb microplates (Thermo Fischer

Scientific) and blocked with 5% skim milk in PBS, containing 0·05% Tween 20. Serial dilutions of test

samples and the assay standard plus controls were added in the plates and incubated for 60 minutes at

room temperature (15-30°C). After washing, horseradish peroxidase (HRP) enzyme-conjugated goat

anti-human IgG-Fc (Jackson ImmunoResearch Laboratories) was added for 60 minutes at room

temperature (15-30°C), then washed. Bound secondary antibody was reacted with 3,3',5,5'-

tetramethylbenzidine (TMB) ELISA peroxidase substrate (Bio-Rad Laboratories) and incubated for 30

minutes at room temperature (15-30°C) before the reaction was stopped with 2N H_2SO_4. Plates were

read at 450 nm with a correction at 620 nm to assess the level of anti-S IgG bound in the microtiter

plate. A reference standard on each plate determined the quantity of anti-S IgG in arbitrary units

(ELU/mL). Concentrations were transformed to binding antibody units per mL (BAU/mL), based on the

WHO International Standard for anti-SARS-CoV-2 immunoglobulin,[9] using a conversion factor

determined during assay validation (1/7·9815). The assay's cut-off and lower limit of quantitation (LLOQ)

was 6·3 BAU/mL.

We measured serum neutralising activity against the Wuhan strain of SARS-CoV-2 in a validated

pseudotyped virus neutralisation assay (PNA) that assessed particle entry-inhibition.[10] Briefly,

pseudotyped virus particles containing a luciferase reporter for detection were made from a modified

vesicular stomatitis virus (VSVΔG) backbone expressing the full-length spike glycoprotein of SARS-CoV-2

(MN908947, Wuhan-Hu-1) from which the last 19 amino acids of the cytoplasmic tail were removed.[11]

Seven two-fold serial dilutions of heat-inactivated serum samples were prepared in 96-well round-

bottom transfer plates (Corning). Pseudotyped virus was added to the serum dilutions at a target

working dilution and incubated at 37°C with 5% CO_2 for 60 ± 5 minutes. Serum-virus complexes were

then transferred onto 96 well white flat-bottom plates (Corning), previously seeded overnight with Vero

E6 cells (Nexelis) and incubated at 37°C and 5% CO_2 for 20 ± 2 hours. Following this incubation,

luciferase substrate from ONE Glo^{TM} Ex luciferase assay system (Promega) was added to the cells. Plates

were then read on a SpectraMax® i3x plate reader (Molecular Devices) to quantify relative luminescence

units (RLU), inversely proportional to the level of neutralising antibodies present in the serum. The

neutralising titre of a serum sample was calculated as the reciprocal serum dilution corresponding to the

50% neutralisation antibody titre (NT_{50}) for that sample; the NT_{50} titres were transformed to

international units per mL (IU/mL), based on the WHO international standard for anti-SARS-CoV-2

immunoglobulin, using a conversion factor determined during assay validation (1/1,872). The assay's

cut-off and LLOQ were 5·3 IU/mL (10 as NT_{50}) and 5.9 IU/mL, respectively. To benchmark vaccine

immunogenicity assessed in BAU/mL and IU/mL, we used a panel of human convalescent serum samples

(HCS) collected 14 days after symptom onset from consecutive cases of mild to moderate COVID-19

illness among health care personnel seen as outpatients in Quebec, Canada during mid-2020. We also

calculated 80% neutralisation titres (NT_{80}); nevertheless, as the PNA was not validated for this

measurement, these results are not presented. We used the same PNA assay to measure NT_{50} (reported

as titres) against pseudotyped virus particles generated for SARS-CoV-2 variants of concern B.1.315[12]

and P.1[13]. In the absence of positive controls for the variant strains of SARS-CoV-2, we used control sera

for the Wuhan-Hu-1 strain.

To assess cellular immunity, we quantified interferon-γ (IFN-γ) and IL-5 producing cells in PBMCs

stimulated with SARS-CoV-2 spike peptide pools (vial 1 158 overlapping peptides, vial 2 157 overlapping

peptides; JPT Peptide) using a human IFN-γ/IL-5 double-colour ELISpot kit (Cellular Technologies) in a

qualified assay. Briefly, activated 96-well plates were coated with anti-human IFN-γ/IL-5 capture

antibodies at 2-8°C. Following overnight (>16h) coating, plates were washed with PBS, and stimulation

media containing SARS-CoV-2 peptide pool 1 or peptide pool 2 or control media was added to wells,

followed by the addition of PBMCs at $2x10^5$ cells/well. After an approximately 44-hour incubation at

$37^{\circ}C\pm1^{\circ}C$ with 5% CO_2, plates were washed to remove cells from the wells. Anti-human IFN-γ/IL-5

detection solution (containing anti-human IFN-γ fluorescein isothiocyanate [FITC] and anti-human IL-5

[biotin] detection antibodies) was then added to the wells and incubated at room temperature (15-30°C)

for 2 h ± 10min to detect IFN-γ and IL-5 cytokine captured on the bottom of the well. Plates were

washed, followed by the addition of a tertiary solution (containing FITC-HRP and streptavidin-alkaline

phosphatase). Following incubation with the tertiary solution, plates were washed, and blue and red

developer solutions were added in sequence (with washes in between), resulting in the appearance of

blue (for IL-5) and red (for IFN-γ) spot forming units (SFUs) in proportion to T cell activity. SFUs were

counted by an ImmunoSpot CTL Analyzer (using CTL ImmunoCapture Software (v7·0·14·0) and CTL

ImmunoSpot Professional DC Analyzer (v7·0·28·2)). Readouts (one per peptide pool for IFN-γ, one per

peptide pool for IL-5) were expressed as number of $SFU/10^6$ cells and combined as a ratio. The assay's

LLOQ for IFN-γ was 109 $SFU/10^6$ cells and for IL-5 was 43 $SFU/10^6$ cells.

Outcome

The primary outcomes were frequency and intensity of solicited injection site and systemic AEs during 7

days after vaccination; frequency, intensity, and relatedness of clinically significant haematological and

biochemical measurements at 7 days after each vaccination; frequency, intensity, and relatedness of

unsolicited AEs during 28 days after each vaccination; and occurrence of medically-attended AEs, serious

AEs, and AEs of special interest during the interim analysis period of 57 days after-first vaccination. The

secondary immunogenicity outcomes were anti-S IgG and NT_{50} against Wuhan-1 strain SARS-CoV-2

pseudotyped virus assessed on days 29 and 43 and expressed as geometric mean titre (GMT) or

concentration (GMCs, BAU/mL for ELISA, or IU/mL for PNA), geometric mean fold rise (GMFR) from

baseline, and percentage of subjects with ≥4-fold increase and ≥10-fold increase from baseline. The

exploratory immunogenicity outcomes were cell-mediated immunity to SARS-CoV-2 S protein, measured

as the ratio of IFN-γ/IL-5 expressing cells on days 1 and 43 in a random subset of subjects receiving two

vaccine formulations (10 µg or 3 µg+CpG1018) or placebo. We also assessed NT_{50} GMTs and the

percentage of subjects with a NT_{50} titre ≥4-fold higher than the LLOQ (1:10 titre), against vaccine-

heterologous SARS-CoV-2 pseudotyped virus (SARS-CoV-2 variants of concern B.1.351 and P.1) on day

43.

Statistical Analyses

This Phase 1/2 study (ClinicalTrial.gov NCT04764422) has a two-part selection design with group

elimination after the interim analysis. In the first part, 35 subjects per group were randomized across 5

candidate vaccine formulations and a placebo group for a total of 210 subjects. After the interim

analysis, two candidates were selected to advance, at which time 250 additional subjects are to be

randomized 2:2:1 to the two selected candidate groups and the placebo, respectively. The study was

designed to have greater than 90% power to identify the candidate with the highest response as

measured by the NT_{50} by ranked GMCs, assuming the true GMC is at least 1·5-fold larger than the

second highest candidate group and to provide a preliminary safety evaluation of the candidates. An

independent data monitoring committee provided safety oversight.

All statistical tests were two-sided with a significance level of 0·05. All statistical analyses were

performed by an independent statistician using SAS version 9·4. All safety assessments took place in the

treatment-exposed population, according to the treatment received. All subject-level percentages were

supplemented with two-sided 95% confidence intervals computed via the Clopper-Pearson method. The

analysis of immunogenicity was performed in the per protocol population, which excludes subjects with

protocol deviations that would affect the assessment. Immunogenicity data were descriptively analysed.

Geometric mean antibody responses were reported by treatment and time point, accompanied by 95%

CIs. The analysis of geometric means excluded subjects who were seropositive at baseline (defined by

anti-S IgG >LLOQ as measured by ELISA). Geometric mean fold rises (GMFR) were calculated relative to

baseline using the log difference of the paired samples, with corresponding CIs computed via the *t*-

distribution, utilizing the antilog transformation to present the ratio. The proportions of subjects with

GMFRs of NT_{50} ≥4 and ≥10 from baseline were summarized with 95% CIs. The analysis of

immunogenicity relative to baseline included baseline seropositive subjects.

Role of Funding Source

The funders of the study had no role in data collection, data analysis, or writing of the statistical report.

GPO was the clinical trial sponsor and approved the study protocol. GPO employees contributed as

authors by preparing the investigational vaccine, interpreting data, and writing this report.

Results

Between March 22 and April 23, 2021, 210 healthy adults were enrolled and assigned to one of six

treatment groups as shown in Figure 1. All received a first dose of vaccine or placebo; two subjects were

excluded from receipt of a second dose (one became pregnant, one developed mild urticaria within 30

minutes after dose one); three other subjects missed the day 29 visit and got no second dose. The

baseline characteristics are shown by treatment group in table 1; the exposed population was 61%

female, had a median age of 36 years (IQR 28, 43) and a median body mass index of 24·07 (IQR 21·30-

26·72).

All five formulations of NDV-HXP-S were well tolerated with no dose limiting reactogenicity (Table 2).

Most solicited injection site and systemic reactogenicity during 7 days after each vaccination was mild

and transient with no apparent difference between dose 1 and 2. The most common injection site

symptoms (table 2) were pain and tenderness; these were most frequent at the highest dose. The most

common systemic symptoms (table 2) were fatigue, headache, and myalgia, generally in less than one-

third of subjects. Fever was uncommon. Adverse events occurring during 28 days after vaccination (table

3) and judged by the investigator to be treatment-related were infrequent (<15%) and there was no

treatment-related serious adverse event, nor any AE of special interest reported during the 57 day

assessment period. Haematology and serum chemistry laboratory readouts were assessed on day 8

following each vaccination; there was no clinically notable finding relative to baseline assessment. The

independent data monitoring committee expressed no safety concern.

Two doses of NDV-HXP-S were immunogenic in a formulation and dose dependent manner within the

per protocol population. Induction of anti-S IgG was modest following dose one but there was a marked

anamnestic response observed 14 days after vaccine dose two (figure 2A). Seronegative individuals in

the vaccine groups responded 28 days after first vaccination with GMCs of anti-S IgG between 7·79 (1

µg) and 20·93 (10 µg) BAU/mL, with a ≥4-fold increase in 34·3-71·4%. The second dose considerably

increased anti-S-IgG antibody responses after 14 days to GMCs between 151·78 (1 µg) and 479·83 (10

µg) BAU/mL. All individuals in every vaccine group had a ≥ 4-fold increase over baseline after the second

dose; all individuals in the 10 µg and 3 ug+CpG1018 groups had a ≥10-fold increase, as did >90% of

vaccinees in the other three vaccine groups (figure 2C). Notably, the adjuvant effect of CpG was limited

after two vaccine doses (table 4): the 1 µg group had a GMC of 151·78 BAU/mL (95% CI 108·99-211·37)

while the 1 µg+CpG1018 group had a GMC of 199·08 BAU/mL (95% CI 140·25-282·57). Among recipients

of the 3 µg dose, the GMC group difference appeared to be greater: 228·07 BAU/mL (no adjuvant, 95%

CI 154·22-337·27) in contrast to 356·83 BAU/mL (CpG1018, 95% CI (265·89-478·88). GMCs of anti-S IgG

among the vaccine groups on day 43 exceeded the GMC of the HCS panel (N=29, 72·93 95% CI 33.00-

161.14) by 2-6-fold (table 4).

Functional antibody responses were assessed by PNA. Low NT_{50} GMCs were detected in all vaccine

groups after the first vaccination (between 7·49 IU/mL and 12·82 IU/mL) with ≥4-fold rises in 8·8% to

25·7% of the vaccine groups (figure 2B ,2D). The second vaccine dose strongly boosted neutralisation

GMCs to between 122·23 IU/mL (1 µg, 95% CI 86·40-172·91) and 474·35 IU/mL (10 µg, 95% CI 320·90-

701·19), with a ≥4-fold increase over baseline in 93·9% to 100% of vaccine groups and a ≥10-fold rise in

most individuals (100% in the 10 µg group, and between 79·4% and 93·9% in the remaining groups). The

differences in post-second dose GMCs between the unadjuvanted and adjuvanted 1 µg and 3 µg groups

were uncertain: 1 µg, 122·23 IU/mL (95% CI 86·40-172·91) versus 1 µg+CpG1018, 127·92 IU/mL (95% CI

85·08-192·34); 3 µg, 166·54 IU/m: (95% CI 100·19-276·81) versus 3 µg+CpG1018 257·70 IU/mL (95% CI

187·01-355·11).

Based on the vaccine-homologous binding and neutralising antibody responses, there was a clear

ranking of immunogenicity with the 10 µg formulation performing best followed by the 3 µg+CpG1018,

3 µg, 1 µg+CpG1018 and 1 µg formulations. The induction of humoral immunity was strong with post-

boost GMFRs relative to baseline of 48-fold (1 µg) to 152-fold (10 µg) for anti-S IgG and 46-fold (1 µg) to

174-fold (10 µg) for NT_{50} antibodies (figure 3). GMCs of NT_{50} by PNA among the vaccine groups on day 43

exceeded the GMC of the HCS panel (N=32, 36·30 95% CI 19·43-67·79) by 3-13-fold depending on the

vaccine formulation (table 4).

Additionally, neutralisation of variant viruses was assessed by PNA on day 43. The proportion of subjects

attaining a day 43 NT_{50} titre ≥40 increased with higher doses of antigen but the incremental changes in

GMT were small. Reduction in neutralising potency, relative to anti-Wuhan neutralising potency, was

modest for the P.1 variant (2·8 to 5·3-fold) but greater for the B·1·351 variant (7·41 to 20·43-fold) (figure

4, table 5). In groups receiving 3 µg or 10 µg antigen doses, the proportion attaining a NT_{50} titre ≥40

ranged from 80·0% to 94·9% against P.1 and from 43·3% to 58·5% against B.1.351 (table 5). Finally, we

also explored T cell responses to determine if the vaccine induced primarily a type 1 (T_H1) or type 2 (T_H2)

T-helper cell response. In the small subset of subjects evaluated 14 days after a second dose, the IFN-

γ/IL-5 ratio was strongly skewed to a T_H1 response relative to pre-vaccination baseline (figure 5),

suggesting the vaccine induced T cell memory capable of an antiviral response.

Discussion

Current production capacity cannot satisfy the global demand for COVID-19 vaccines[1] and vaccine

distribution is inequitable with most vaccines acquired and used by high income countries while LMICs

have limited access. Local production of COVID-19 vaccines in LMICs would increase global availability

and reduce dependence of countries producing these vaccines on international vaccine supply. Here we

demonstrated for the first time that an engineered inactivated NDV-based vaccine expressing a second-

generation stabilized SARS-CoV-2 spike protein[5], produced in eggs in an existing influenza virus

production facility at GPO in Thailand, shows an acceptable reactogenicity and safety profile in humans

and has immunogenicity that supports its potential clinical benefit. We evaluated a range of vaccine

doses (1 µg, 3 µg, 10 µg) having potency quantified as µg of virus envelope-anchored SARS-CoV-2 spike

protein; the low and medium antigen doses were evaluated in formulations with and without the TLR-9

agonist CpG1018 as a vaccine adjuvant. Over 28 days after each vaccine dose, all formulations were very

well-tolerated with little solicited injection site or systemic reactogenicity aside from mild injection site

pain and tenderness. There was no safety signal issuing from this early interim analysis of the clinical

trial. Moreover, the vaccine was strongly immunogenic in a formulation and dose dependent manner,

inducing levels of vaccine-homologous anti-S IgG and virus neutralising antibodies that exceeded by

several fold the levels measured in 14-day convalescent sera from consecutive cases of health care

workers with mild to moderate COVID-19 illness in 2020. Notably, the adjuvant benefit as measured by

enhanced induction of humoral immunity was uncertain, as the small sample size limited precision. On

the other hand, the vaccine at all dose levels elicited neutralising antibodies against two variants of

concern, B.1.351 and P.1. While neutralising antibody titres decreased modestly against P.1 and more

markedly against B.1.351, this was expected and in the range observed with sera from recipients of the

mRNA vaccines BNT162b2 and mRNA-1273.[14-17] The degree of reduction in neutralisation is dependent

on the assay used and can be especially dramatic with pseudotyped particle inhibition assays as used in

this study.[14,17] Notably, the B.1.351 variant is currently regarded as a worst case example of immune

evasion; accordingly, the NDV-HXP-S would be predicted to have neutralising activity against the now

prevalent delta (B.1.617.2) variant. The T cell response assessed showed a bias towards a T_H1 response

in both evaluated dose groups, alleviating concerns about enhanced disease associated with a T_H2

response (as observed with SARS-CoV-1 in some animal models[18]). These initial data, while sparse,

suggest the vaccine-induced T cell memory capable of an antiviral response.

The study has several limitations. The sample size per treatment group was small, limiting precision, and

assessments were restricted to 43 days for immunogenicity and 57 days for reactogenicity and safety,

narrowing our perspective to acute outcomes only. These are inherent problems of phase 1 trials and

interim analyses in a pandemic response setting. Nevertheless, as clinical trials with similar vaccines are

underway in Vietnam (NCT04830800) and Brazil (NCT04993209), we determined that publication of

early data is a priority. The study had strengths as well. The vaccine construct is a novel platform

expressing a second-generation pre-fusion stabilized S protein in a membrane-bound trimeric

conformation. We hypothesize that these characteristics contribute to the vaccine's notable

immunogenicity, even without the CpG1018 adjuvant. The anti-S ELISA and PNA used to assess vaccine-

homologous NT_{50} potency were validated and results are expressed in International Units[9] for future

comparisons. The induction of anti-S binding and neutralising antibodies was contrasted with mean

levels in human convalescent serum and found to be superior, especially in the mid- and high-dose

groups. However, the neutralisation assay is not a live virus assay; therefore, it is presently uncertain

how our functional antibody readouts can be used to benchmark against live virus neutralising antibody

levels induced by authorized or licensed vaccines. Correlation between neutralising antibody titres and

vaccine efficacy and individual protection has recently been shown; work is ongoing to integrate these

data into this analytic framework.[19-21] On the other hand, mean vaccine anti-S IgG ELISA responses

normalized by the mean in convalescent sera as proposed by Earle and others[19] suggest that the NDV-

HXP-S vaccine will afford important clinical benefit. Furthermore, we plan to conduct a clinical trial in

which the NDV-HXP-S vaccine will be contrasted to an authorized comparator vaccine to generate

relative immunogenicity evidence that may be predictive of clinical benefit.

In summary, we show that the inactivated NDV-HXP-S vaccine candidate has an acceptable safety profile

and is highly immunogenic. This vaccine can be produced at low cost in any facility designed for

production of inactivated influenza virus vaccine; such facilities are present in a number of LMICs.[2]

Based on these results, and acknowledging the imperative to maximize output of vaccine doses from the

manufacturing facility, the 3 µg and 3 µg+CpG1018 formulations were selected for further assessment in

the phase 2 stage of the ongoing clinical trial.

Word count 4,361

Contributors

PPit and SLaw verified the underlying data reported herein. All authors had full access to all the data in the study. Individual author roles are reported using CRediT: Conceptualisation, PPit, SLam, LDM, RSch, AGS, PPal, FK, KP, PW, and BLI; Data curation, PPit and SLaw; Formal analysis, Slaw, LDM, JMC, and JT; Funding acquisition, PN, SSur, RR, NS, AGS, PPal, FK, KP, PW, and BLI; Investigation, PPit, VL, SM, SKam, CS, WS, YL, WP, FA, LG, SKha, SM, IM, BP, SSla, and STra; Methodology, PPit, VL, SM, PN, SSur, SPra, SPuk, SLam, LDM, RR, WS, YL, FA, LG, RK, SKha, ML, IM, BP, RSin, NS, SThe, STra, TV, JAW, FK, and RH, Project administration, PPit, PN, SSur, RSch, YS, NS, NT, TV, and PW; Resources, PPit, SPra, SPuk, WS, CLH, ML, JSM, RSin, SThe, JAW, AGS, PPal, FK, and RH; Supervision, PPit, SPra, RR, RSch, LG, RK, ML, JAW, AGS, PPal, FK, KP, PW, and BLI; Validation, SLaw, SPra, SPuk, LG, RK, SKha, RSin, SThe, and STra; Visualisation, LDM, JMC, JT, and BLI; Writing–original draft, LDM, RR, FK, and BLI; Writing–review & editing, PPit, VL, SLaw, SM, SKam, CS, SSur, SPuk, SLam, WS, JMC, RSch, FA, LG, CLH, RK, SKha, ML, JSM, IM, JT, STra, JAW, PW, and RH.

Declaration of Interest

PN, SSur, SPra, SPuk, RK, RSin, NS, SThe, TV, KP, and PW are salaried employees of the Government Pharmaceutical Organization. The vaccine administered in this study was developed by faculty members at the Icahn School of Medicine at Mount Sinai including FK, AGS, PP, and WS. Mount Sinai is seeking to commercialize this vaccine; therefore, the institution and its faculty inventors could benefit financially. JSM and CLH are inventors on U.S. patent applications concerning the stabilized SARS-CoV-2 S construct.

Data sharing

Under review, to be decided by the sponsor prior to publication.

Acknowledgements

Activities at PATH and Mount Sinai were supported, in part, by the Bill & Melinda Gates Foundation

[INV-021239]. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0

Generic License has already been assigned to the Author Accepted Manuscript version that might arise

from this submission. The findings and conclusions contained within this manuscript are those of the

authors and do not necessarily reflect positions or policies of the Bill & Melinda Gates Foundation. The

salary of PP was partially funded by NIH (Centers of Excellence for influenza Research and Response,

CEIRR,75N93021C00014), NIAID grant (P01 AI097092-07), NIAID grant (R01 AI145870-03), by the NIH

Collaborative Influenza Vaccine Innovation Centers (CIVICs) contract 75N93019C00051 and a grant from

an anonymous philanthropist to Mount Sinai. Design and generation of reagents used in this project in

the Krammer laboratory were supported by CEIRR (75N93021C00014) and CIVIC (75N93019C00051).

Research and development activities in Thailand were funded by the Government Pharmaceutical

Organization Thailand), the National Vaccine Institute (Thailand), and the National Research Council

(Thailand). Testing of clinical trial specimens at Nexelis was supported by the Global Health Vaccine

Accelerator Platform of the Bill & Melinda Gates Foundation. The authors thank Dr. Randy Albrecht for

management of import/export of Newcastle disease virus vectors at the Icahn School of Medicine at

Mount Sinai. We are grateful to Dr. Nina Bardwaj for programmatic and scientific oversight of the Virus

and Cell Therapy Laboratory at the Icahn School of Medicine at Mount Sinai, which produced the NDV-

HXP-S master virus seed.

References

1.	(WHO) WHO. https://www.who.int/director-general/speeches/detail/director-general-s-opening-remarks-at-the-media-briefing-on-covid-19-9-april-2021. 2021.

2.	Sparrow E, Wood JG, Chadwick C, et al. Global production capacity of seasonal and pandemic influenza vaccines in 2019. *Vaccine* 2021; **39**(3): 512-20.

3.	Sun W, McCroskery S, Liu WC, et al. A Newcastle Disease Virus (NDV) Expressing a Membrane-Anchored Spike as a Cost-Effective Inactivated SARS-CoV-2 Vaccine. *Vaccines (Basel)* 2020; **8**(4).

4.	Hsieh CL, Goldsmith JA, Schaub JM, et al. Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science* 2020; **369**(6510): 1501-5.

5.	Sun W, Liu Y, Amanat F, et al. A Newcastle disease virus-vector expressing a prefusion-stabilized spike protein of SARS-CoV-2 induces protective immune responses against prototype virus and variants of concern in mice and hamsters. *bioRxiv* 2021: 2021.07.06.451301.

6.	Campbell JD. Development of the CpG Adjuvant 1018: A Case Study. *Methods Mol Biol* 2017; **1494**: 15-27.

7.	ter Meulen J, van den Brink EN, Poon LL, et al. Human monoclonal antibody combination against SARS coronavirus: synergy and coverage of escape mutants. *PLoS Med* 2006; **3**(7): e237.

8.	Hsieh CL, Goldsmith JA, Schaub JM, et al. Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science* 2020.

9.	Kristiansen PA, Page M, Bernasconi V, et al. WHO International Standard for anti-SARS-CoV-2 immunoglobulin. *Lancet* 2021; **397**(10282): 1347-8.

10.	Bewley KR, Coombes NS, Gagnon L, et al. Quantification of SARS-CoV-2 . neutralising antibody by wild-type plaque reduction neutralisation, microneutralisation and pseudotyped virus neutralisation assays. *Nat Protoc* 2021; **16**(6): 3114-40.

11.	Havranek KE, Jimenez AR, Acciani MD, et al. SARS-CoV-2 Spike Alterations Enhance Pseudoparticle Titers and Replication-Competent VSV-SARS-CoV-2 Virus. *Viruses* 2020; **12**(12).

12.	Tegally H, Wilkinson E, Giovanetti M, et al. Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature* 2021; **592**(7854): 438-43.

13.	Faria NR, Mellan TA, Whittaker C, et al. Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil. *Science* 2021; **372**(6544): 815-21.

14.	Garcia-Beltran WF, Lam EC, St Denis K, et al. Multiple SARS-CoV-2 variants escape neutralisation by vaccine-induced humoral immunity. *Cell* 2021; **184**(9): 2523.

15.	Carreño JM, Alshammary H, Singh G, et al. Reduced . neutralising activity of post-SARS-CoV-2 vaccination serum against variants B.1.617.2, B.1.351, B.1.1.7+E484K and a sub-variant of C.37. *medRxiv* 2021: 2021.07.21.21260961.

16.   Edara VV, Norwood C, Floyd K, et al. Infection- and vaccine-induced antibody binding and
neutralisation of the B.1.351 SARS-CoV-2 variant. *Cell Host Microbe* 2021; **29**(4): 516-21.e3.

17.   Shen X, Tang H, Pajon R, et al. Neutralisation of SARS-CoV-2 Variants B.1.429 and B.1.351. *N Engl*
J Med 2021; **384**(24): 2352-4.

18.   Krammer F. SARS-CoV-2 vaccines in development. *Nature* 2020; **586**(7830): 516-27.

19.   Earle KA, Ambrosino DM, Fiore-Gartland A, et al. Evidence for antibody as a protective correlate
for COVID-19 vaccines. *Vaccine* 2021.

20.   Khoury DS, Cromer D, Reynaldi A, et al. . neutralising antibody levels are highly predictive of
immune protection from symptomatic SARS-CoV-2 infection. *Nat Med* 2021.

21.   Gilbert PB, Montefiori DC, McDermott A, et al. Immune Correlates Analysis of the mRNA-1273
COVID-19 Vaccine Efficacy Trial. *medRxiv* 2021: 2021.08.09.21261290.

Table 1. Baseline characteristics of the exposed population

	1 µg S (N = 35)	1 µg S+CpG (N = 35)	3 µg S (N = 35)	3 µg S+CpG (N = 35)	10 µg S (N = 35)	Placebo (N = 35)
Age, years	33·0 (26·0-39·0)	39·0 (32·0-45·0)	37·0 (29·0-49·0)	34·0 (25·0-44·0)	37·0 (31·0-42·0)	32·0 (27·0-42·0)
Sex						
Male	14 (40·0%)	14 (40·0%)	7 (20·0%)	15 (42·9%)	18 (51·4%)	14 (40·0%)
Female	21 (60·0%)	21 (60·0%)	28 (80·0%)	20 (57·1%)	17 (48·6%)	21 (60·0%)
Ethnicity						
Asian	35 (100%)	35 (100%)	35 (100%)	35 (100%)	35 (100%)	35 (100%)
Body mass index	24·59 (20·76-27·85)	24·85 (21·42-26·33)	23·95 (21·23-27·96)	23·95 (21·70-25·92)	24·52 (21·26-27·68)	23·12 (21·72-27·22)

478   Data are median (q1-q3) or n (%)

480   Table 2. Solicited AEs during 7 days after vaccination

		1 μg S (N = 35)	1 μg S+CpG (N = 35)	3 μg S (N = 35)	3 μg S+CpG (N = 35)	10 μg S (N = 35)	Placebo (N = 35)
		n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)
Any injection site AE	Dose 1	8 (22·9%) (10·4-40·1)	13 (37·1%) (21·5-55·1)	14 (40·0%) (23·9-57·9)	20 (57·1%) (39·4-73·7)	23 (65·7%) (47·8-80·9)	4 (11·4%) (3·2-26·7%)
	Dose 2	10 (28·6%) (14·6-46·3)	14 (42·4%) (25·5-60·8)	16 (47·1%) (29·8-64·9)	20 (58·8%) (40·7-75·4)	24 (68·6%) (50·7-83·1)	10 (29·4%) (15·1-47·5)
Pain	Dose 1	2 (5·7%) (0·7-19·2)	8 (22·9%) (10·4-40·1)	10 (28·6%) (14·6-46·3)	16 (45·7%) (28·8-63·4)	16 (45·7%) (28·8-63·4)	3 (8·6%) (1·8-23·1)
	Dose 2	10 (28·6%) (14·6-46·3)	10 (28·6%) (14·6-46·3)	15 (42·9%) (26·3-60·6)	17 (48·6%) (31·4-66·0)	22 (62·9%) (44·9-78·5)	8 (22·9%) (10·4-40·1)
Tenderness	Dose 1	6 (17·1%) (6·6-33·6)	4 (11·4%) (3·2-26·7)	4 (11·4%) (3·2-26·7)	4 (11·4%) (3·2-26·7)	7 (20·0%) (8·4-36·9)	1 (2·9%) (0·1-14·9)
	Dose 2	0 (0·0%) (0·0-10·0)	4 (11·4%) (3·2-26·7)	1 (2·9%) (0·1-14·9)	3 (8·6%) (1·8-23·1)	2 (5·7%) (0·7-19·2)	2 (5·7%) (0·7-19·2)
Swelling	Dose 1	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)
	Dose 2*	masked	masked	masked	masked	masked	masked
Induration	Dose 1	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)
	Dose 2	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)
Erythema	Dose 1	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)
	Dose 2	0 (0·0%)	0 (0·0%)	0 (0·0%)	0 (0·0%)	0 (0·0%)	0 (0·0%)

		(0·0-10·0)	(0·0-10·0)	(0·0-10·0)	(0·0-10·0)	(0·0-10·0)	(0·0-10·0)
Any systemic AE	Dose 1	12 (34·3%) (19·1-52·2)	8 (22·9%) (10·4-40·1)	19 (54·3%) (36·6-71·2)	14 (4·.0%) (23·9-57·9)	17 (48·6%) (31·4-66·0)	7 (20·0%) (8·4-36·9)
	Dose 2	9 (25·7%) (12·5-43·3)	11 (33·3%) (18·0-51·8)	9 (26·5%) (12·9-44·4)	17 (50·0%) (32·4-67·6)	15 (42·9%) (26·3-60·6)	4 (11·8%) (3·3-2·.5)
Fever >38°C	Dose 1	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	1 (2·9%) (0·1-14·9)	0 (0·0%) (0·0-10·0)	3 (8·6%) (1·8-23·1)	0 (0·0%) (0·0-10·0)
	Dose 2	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	1 (2·9%) (0·1-14·9)	2 (5·7%) (0·7-19·2)	2 (5·7%) (0·7-19·2)	0 (0·0%) (0·0-10·0)
Headache	Dose 1	5 (14·3%) (4·8-30·3)	5 (14·3%) (4·8-30·3)	9 (25·7%) (12·5-43·3)	6 (17·1%) (6·6-33·6)	11 (31·4%) (16·9-49·3)	1 (2·9%) (0·1-14·9)
	Dose 2	4 (11·4%) (3·2-26·7)	5 (14·3%) (4·8-30·3)	6 (17·1%) (6·6-33·6)	8 (22·9%) (10·4-40·1)	4 (11·4%) (3·2-26·7)	1 (2·9%) (0·1-14·9)
Fatigue	Dose 1	8 (22·9%) (10·4-40·1)	4 (11·4%) (3·2-26·7)	12 (34·3%) (19·1-52·2)	6 (17·1%) (6·6-33·6)	6 (17·1%) (6·6-33·6)	7 (20·0%) (8·4-36·9)
	Dose 2	3 (8·6%) (1·8-23·1)	6 (17·1%) (6·6-33·6)	7 (20·0%) (8·4-36·9)	8 (22·9%) (10·4-40·1)	7 (20·0%) (8·4-36·9)	4 (11·4%) (3·2-26·7)
Malaise	Dose 1	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	3 (8·6%) (1·8-23·1)	4 (11·4%) (3·2-26·7)	0 (0·0%) (0·0-10·0)
	Dose 2	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	4 (11·4%) (3·2-26·7)	4 (11·4%) (3·2-26·7)	0 (0·0%) (0·0-10·0)
Myalgia	Dose 1	4 (11·4%) (3·2-26·7)	4 (11·4%) (3·2-26·7)	8 (22·9%) (10·4-40·1)	6 (17·1%) (6·6-33·6)	9 (25·7%) (12·5-43·3)	1 (2·9%) (0·1-14·9)
	Dose 2	6 (17·1%) (6·6-33·6)	2 (5·7%) (0·7-19·2)	4 (11·4%) (3·2-26·7)	11 (31·4%) (16·9-49·3)	11 (31·4%) (16·9-49·3)	1 (2·9%) (0·1-14·9)
Arthralgia	Dose 1	2 (5·7%) (0·7-19·2)	1 (2·9%) (0·1-14·9)	5 (14·3%) (4·8-30·3)	0 (0·0%) (0·0-10·0)	3 (8·6%) (1·8-23·1)	0 (0·0%) (0·0-10·0)

	Dose 2	2 (5·7%) (0·7-19·2)	1 (2·9%) (0·1-14·9)	2 (5·7%) (0·7-19·2)	4 (11·4%) (3·2-26·7)	4 (11·4%) (3·2-26·7)	0 (0·0%) (0·0-10·0)
Nausea	Dose 1	2 (5·7%) (0·7-19·2)	1 (2·9%) (0·1-14·9)	3 (8·6%) (1·8-23·1)	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	0 (0·0%) (0·0-10·0)
	Dose 2	3 (8·6%) (1·8-23·1)	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	2 (5·7%) (0·7-19·2)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)
Vomiting	Dose 1	1 (2·9%) (0·1-14·9)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)	0 (0·0%) (0·0-10·0)
	Dose 2**	masked	masked	masked	masked	masked	masked

481   * Swelling after dose 2 was reported by 1 subject only who remains masked d (0·5%, 95% CI 0·0-2·6)

482   **Vomiting after dose 2 was reported by 1 subject only who remains masked (0·5%, 95% CI 0·0-2·6)

484   Table 3 AEs with onset during 28 days after vaccination

		1 µg S (N = 35)	1 µg S+CpG (N = 35)	3 µg S (N = 35)	3 µg S+CpG (N = 35)	10 µg S (N = 35)	Placebo (N = 35)
		n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)	n (%) (95% CI)
Any	Dose 1	8 (22·9%) (10·4-40·1)	7 (20·0%) (8·4-36·9)	15 (42·9%) (26·3-60·6)	13 (37·1%) (21·5-55·1)	9 (25·7%) (12·5-43·3)	6 (17·1%) (6·6-33·6)
	Dose 2	7 (20·0%) (8·4-36·9)	3 (8·6%) (1·8-23·1)	11 (31·4%) (16·9-49·3)	10 (28·6%) (14·6-46·3)	6 (17·1%) (6·6-33·6)	9 (25·7%) (12·5-43·3)
Vaccine-related	Dose 1	2 (5·7%) (0·7-19·2)	3 (8·6%) (1·8-23·1)	3 (8·6%) (1·8-23·1)	5 (14·3%) (4·8-30·3)	3 (8·6%) (1·8-23·1)	0 (0·0%) (0·0-10·0)
	Dose 2	0 (0·0%) (0·0-10·0)	1 (2·9%) (0·1-14·9)	2 (5·7%) (0·7-19·2)	2 (5·7%) (0·7-19·2)	2 (5·7%) (0·7-19·2)	1 (2·9%) (0·1-14·9)
Serious	Dose 1*	masked	masked	masked	masked	masked	masked
	Dose 2	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	1 (2·9%) (0·1-14·9)	0 (0·0%) (0·0-10·0)	1 (2·9%) (0·1-14·9)	1 (2·9%) (0·1-14·9)
Serious vaccine-related	Dose 1	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)
	Dose 1	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)	0 (0·0%) (0·0-10·0)

485   * Serious AE was reported by one subject only who remains masked (0·5%, 95% CI 0·0-2·6)

487   Table 4. GMCs of anti-S IgG (BAU/mL) and NT_{50} by PNA (IU/mL) on day 43 and GMC ratios, vaccine to
488   HCS panel

		1 µg S	1 µg S+CpG	3 µg S	3 µg S+CpG	10 µg S
Anti-S IgG BAU/mL,	GMC	151·78	199·08	228.07	356.83	479.83
	95% CI	(108·99-211·37)	(140·25-282·57)	(154·22-337·27)	(265·89-478·88)	(360·19-639·20)
	GMC ratio, vaccine to HCS panel	2·08	2·73	3·13	4·89	6·58
	95% CI	(0·89-4·87)	(1·16-6·43)	(1.31-7·48)	(2·12-11·31)	(2·85-15·18)
NT50 by PNA	GMC	122·23	127·92	166·54	257·70	474·35
	95% CI	(86·40-172·91)	(85·08-192·34)	(100·19-276·81)	(187·01-355·11)	(320·90-701·19)
	GMC ratio, vaccine to HCS panel	3·37	3·52	4·59	7·10	13·07
	95% CI	(1·67-6·81)	(1·69-7·34)	(2·08-10·10)	(3·55-14·20)	(6·33-26·99)

491 Table 5. GMT and percentage of subjects with a ≥4-fold rise from baseline (day 43) for NT_{50} by PNA
492 against Wuhan strain and variants of concern

		1 µg S	1 µg S+CpG	3 µg S	3 µg S+CpG	10 µg S	Placebo
Wuhan	GMT	n = 32 228·81 (161·74-323·69)	n = 31 239·47 (159·27-360·06)	n = 30 311·76 (187·56-518·18)	n = 33 482·42 (350·08-664·77)	n = 34 887·99 (600·72-1,312·62)	n = 33 5·89 (4·64-7·48)
	≥4-fold rise	N = 34 34 (100%) (89·7-100)	N = 33 31 (93·9%) (79·8-99·3)	N = 33 32 (97·0%) (84·2-99·9)	N = 33 33 (100%) (89·4-100)	N = 35 35 (100%) (90·0-100)	N = 34 2 (5·9%) (0·7-19·7)
P·1	GMT	n = 31 45·33 (32·18-63·85)	n = 31 74·06 (48·99-111·97)	n = 30 111·28 (72·06-171·83)	n = 33 150·59 (111·15-204·03)	n = 34 167·14 (120·63-231·58)	n = 33 5·57 (4·47-6·94)
	≥4-fold rise	N = 31 15 (48·4%) (30·2 – 66·9)	N = 31 29 (74·2%) (55/4 – 88·1)	N = 30 24 (80·0%) (61·4 – 92·3)	N = 33 29 (87·9%) (71·8 – 96·6)	N = 34 (32 (94·9%) (80·3 – 99·3)	N = 33 1 (3·0%) (0·1 – 5·8)
B·1·351	GMT	n = 32 21·00 (15·12-29·18)	n = 30 32·34 (22·13-47·24)	n = 30 37·43 (24·77-56·56)	n = 33 40·07 (28·09-57·17)	n = 34 43·47 (31·55-59·89)	n = 33 6·12 (4·90-7·64)
	≥4-fold rise	N = 32 5 (15·6%) (5·3 – 32·8)	N = 40 15 (50·0%) (31·3 -68·7)	N = 30 (13 (43·3%) (30·8 – 66·5)	N = 33 16 (48·5%) (30·8 – 66·5)	N = 34 20 (58·8%) (40·7 – 75·4)	N = 33 1 (3·0%) (0·1 – 15·8)

Figure 1. Trial profile

Figure 2. Distribution and GMC of anti-S IgG (BAU/mL) in placebo, vaccine groups and HCS controls (A),

distribution and GMC of NT_{50} by PNA (IU/mL) in placebo, vaccine groups, and HCS controls (B),

percentage of subjects with ≥4-10-fold increase in anti-S IgG (C), and percentage of subjects with ≥4-10-

fold increase in NT_{50} by PNA (D); numbers above data denote number of per-protocol subjects

contributing data

Figure 3. Distribution and GMFR of fold rise in anti-S IgG from baseline (A), distribution and GMFR of fold

rise in NT_{50} by PNA from baseline (B); numbers above data denote number of per-protocol subjects

contributing data

Figure 4. Distribution and GMT of NT_{50} by PNA against variants of concern (day 43); numbers above data

denote number of per-protocol subjects contributing data

Figure 5. Box plot of IFN-γ /IL-5 ratios (ELISpot 2-colour assay); numbers above data denote number of

per-protocol subjects contributing data

514   Figure 1

Figure 3

A

B

Figure 4

Figure 5