**Can we distinguish the roles of demographic and temporal changes in the incidence and prevalence of musculoskeletal disorders? A systematic search and review.**

*Hanifa Bouziri[1*], Alexis Descatha[2], Yves Roquelaure[2], William Dab[1], Kévin Jean[1,3]*

1. Laboratoire MESuRS, Conservatoire national des Arts et Métiers (Cnam), Paris, France.
2. Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) – UMR_S 1085, Univ Angers, Univ Rennes, 28 Roger Amsler, CS 74521, 49045 Anger, Cedex 01, France.
3. MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London, United Kingdom

*Correspondence: hanifa.bouziri@lecnam.net*

**KEYWORDS:**
Musculoskeletal disorders, Temporal trends, Occupational health, Systematic review, Epidemiology, Chronic diseases.

**ACADEMIC DEGREES**
Hanifa Bouziri: PhD student in Public Health
Alexis Descatha: University professor - hospital practitioner
Yves Roquelaure: University Professor and Hospital Practitioner and team director of the occupational health and ergonomics epidemiology team (ESTER)
William Dab: Professor of Health and Safety Chair at Cnam
Kévin Jean: Assistant professor at Cnam

**ABSTRACT:**

Background: Musculoskeletal disorders (MSDs) represent a major public health issue, affecting more than 40 million European workers in 2017. The overall ageing of the working population is expected to increase the burden of disease, but temporal changes in exposures or diagnosis may also drive global trends in MSDs. We, therefore, sought to review and summarize evidence describing the role of demographic and temporal changes in the occurrence of MSDs.

Methods: We conducted a systematic search and review of articles reporting temporal trends in MSDs in the general working-age population. Only articles controlling for age in the analysis were included. The risk of bias was assessed. The main indicators extracted were age-controlled time trends in MSD incidence or prevalence.

Results: Among 2,680 articles, 16 fulfilled the inclusion criteria, representing 23 results according to the indicators extracted. No study was found with a high risk of bias. Twelve

results reported time trends in prevalence and 11 in incidence. After controlling for age, the
reported temporal trends varied, mostly between non-monotonic changes (n=12/23) and
increases (n=10/23); one article also highlighted an increase among women and non-
monotonic changes among men (n = 1/23). Several factors other than ageing were
suggested to explain temporal trends in MSDs, main trends in obesity, changing
occupational exposures, and cultural factors regarding pain tolerance.

<u>Conclusion</u>: This review shows that different kinds of factors in addition to ageing may
contribute to varying or increasing trends in MSDs. This review also highlighted the scarcity
of evidence regarding time trends in the burden of MSDs and their underlying causes.

**INTRODUCTION:**

Musculoskeletal disorders (MSDs) affected more than 40 million workers in Europe in 2017
and were the leading contributors to disability worldwide in 2019 (1, 2). These conditions
refer to a group of painful disorders of muscles, tendons, and nerves. The causes of MSDs
are multifactorial and notably can be induced by occupational, biomechanical and
psychosocial risk factors (3). Since the 1970s, there have been many changes in working
conditions, catalyzed most notably by increased digitization across a range of professions,
and widespread reinforcement of preventive actions has redistributed the risk factors of
MSDs (4-6). There has also been an increase in employment in the service sector, which
contributes to changes in the patterns of exposure to hazards at work (1). The combined
effects of these occupational changes on the temporal evolution of MSDs are thus
challenging to assess.  In addition, the ageing of the workforce could have implications for
the increasing risk of chronic diseases like MSDs (6, 7). In particular, the rising average age
of workers in many high-income countries may increase the risk of MSDs in the absence of
preventive action. This owes in part to degenerative phenomena linked to the ageing
process itself, which induces a reduction in biomechanical tolerance to repetitive and/or
prolonged loading, and in part to prolonged exposure to residual biomechanical stresses and
psychosocial risks accumulated during increasingly long careers (3).

Exposures to leading occupational risk factors of MSDs, such as biomechanical,
organizational, and psychosocial factors, have evolved heterogeneously over time (8-11).
Consequently, successive cohorts of workers have not been exposed to MSD risk factors
with the same intensity and frequency throughout their lives. Moreover, risk factors of MSDs
and their changes over time have mainly been studied individually or by family (e.g.
biomechanical, organizational, psychosocial), but a global view of their simultaneous change
over time is still lacking. Disentangling the respective roles of age and temporal evolution in
exposure to risk factors on the occurrence of MSDs is thus needed to understand current
trends and design adapted prevention policies (3-14). Furthermore, understanding the
evolution of exposures over time while accounting for age would allow for a more accurate
prediction of future trends in MSDs and help to prevent and control their occurrence (15, 16).
87

The objective of this study is to collate and review the existing evidence on the respective
roles of demographic and temporal changes in the occurrence of MSDs. We used the
systematic search and review methodology as previously described by Grant and Booth
(17). This type of review consists of combining a systematic search method with a critical
review analysis and is used to answer broad questions while often incorporating multiple
study designs.

**METHODS:**

*Search strategy*

The study protocol was registered in PROSPERO (CRD42020221499) (18). This protocol is
consistent with the Preferred Reporting Items for Systematic Review and Meta-analysis
(PRISMA) guidelines (19, 21). A detailed PRISMA 2020 checklist is provided in
supplementary material 1. Any modification of the methods stated in the original protocol
was registered in PROSPERO (see the reference mentioned before).

*Literature search*

We searched four different electronic bibliographic databases for studies published between
1990 and 2020: Medline, ScienceDirect, Wiley, and Web of Science. The last source
searched or consulted was checked in November 2020. Details of the search strategy used
for each database are provided in supplementary material S2, including the algorithms of
keywords used, the number of results and the articles preselected for screening.

*Inclusion and exclusion criteria*

For the article identification step, when the databases allowed, we automatically excluded
results related to topics not relevant for our search, such as studies involving animals,
molecular biology, immunology studies or clinical case reports.
114
115  Then, the first round of selective screening was carried out based on titles and abstracts
116  (step 1). Only original articles were included; conference reports, literature reviews, and
117  editorials were excluded. At this stage, only articles that reported MSD or MSD proxy
118  outcomes while mentioning the notion of temporal trends were included.
119
In the full-text assessment (step 2), articles defining MSD as a group or set of diseases
localized at or around the joints (wrists, elbows, shoulders, spine, or knees) were selected.
The pathologies considered here concerned the muscles, tendons and tendon sheaths,
nerves, bursae, joints, ligaments, at the periphery of the joints of the upper limbs, the spine,
and the lower limbs. We excluded MSDs defined as a joint manifestation of organic diseases
(e.g. psoriasis, lupus, gout, etc.) or as the joint location of systemic inflammatory origins (e.g.
secondary osteoarthritis). At this step, only articles reporting temporal trends in incidence
and/or prevalence in MSD while controlling for age were selected. We included studies
conducted among the working-age population. Studies of people under 18 and unpaid
domestic workers were excluded. The prevalence or incidence of MSDs over time that only
addresses the average over a single period were also excluded.
131

### Screening

133  The Covidence Systematic Review software allowed the selection of studies, their download,
134  and the removal of duplicates (22). Both steps 1 and 2 were performed by two independent
135  authors to assess the eligibility of studies identified in the databases. Any conflict in article
136  screening or full-text assessment was resolved by a third senior researcher.
137

### Data extraction

139  All articles included were read for the identification and extraction of the following
140  characteristics: geographic location, population studied, study design and recruitment
141  criteria, start and end date of follow-up, MSD sites (superior limbs, inferior limbs & back, or
142  not specified), criteria used for MSD definition (either based on pain or disability), and the
143  method used for MSD diagnosis.
144

### Assessing risk of bias and quality of evidence

To assess the risk of bias across included studies, we used the RoB-SPEO (23) and the
Navigator guide tool, which we adapted for our study (see supplementary material 4 for
methodological details) (24, 25). The biases we assessed were selection bias, potential
biases linked to misclassification of MSDs, biases due to incorrectly taking confounding
factors into account, and bias due to potential conflict of interest. Each article has been
classified according to its level of bias (low, probably low, probably high, high). We also
assessed the quality of the statistical trends tested by using the following classification:
satisfactory quality, probably satisfactory quality, probably unsatisfactory quality,
unsatisfactory quality. Further details on the criteria and classifications used for the risk of
bias and quality of evidence assessment are available in supplementary material S4.
156

***Analysis of the temporal trends of the occurrence of MSDs***
The following data were extracted from each article: the raw temporal trends of MSD
prevalence and/or incidence (if reported), information concerning methods used to control for
age, and MSD prevalence and/or incidence over time after controlling for age. For each
article, the temporal trends in MSD prevalence and/or incidence were analyzed according to
the location and severity of MSDs. If an article investigated multiple types of MSDs and/or
addressed both temporal trends in the prevalence and incidence of MSDs, we considered
these results independently; therefore, the total number of results could possibly be higher
than the number of studies included. We also distinguished two groups of results based on
the MSD sites and criteria used for MSD definition, either based on pain or repercussion on
work and/or social life (hereafter called disability). The precise definition of MSD used in
each article is provided in supplementary material S3. Temporal changes in MSD
prevalence/incidence were summarized according to whether they decreased, varied non-
monotonically, or increased.
171

***Synthesized evidence***
For articles reporting both raw and age-adjusted MSD time trends, we compared findings to
discern differences between them, and therefore to assess whether it would be possible to
dissociate age from time in the occurrence of MSDs over time. When mentioned, we
summarized the interpretations and hypotheses proposed to explain observed temporal
trends in MSD prevalence/incidence.
178

179 **RESULTS:**
180 ***Studies selected***

A total of 2,680 study records were identified through our systematic search, of which 1,977 were excluded, 335 were duplicates, and 1,642 were deemed irrelevant as per automatic categorization tools provided by some databases (**Figure 1**). A further 658 records were excluded after the title and abstract screening because they did not present original results and/or did not report results based on MSDs and/or did not address temporal trends in their occurrence. Of the 45 full-text articles assessed for eligibility, 29 were classified as non-eligible, 13 of which because they did not control for age is reported indicators. A total of 16 studies fulfilled all eligibility criteria and were thus included in the review.

### General study characteristics

The 16 articles included in the present review were published between 2003 and 2020 (**Table 1**), among which 12 were published after 2010. Overall, the studies were conducted in 3 geographic areas: 12 in Europe (among which 5 were in Scandinavian countries), 3 in the USA, and 1 in Australia. The duration of the study period ranged from 10 to 55 years across studies. Eight studies relied on sampled populations (3 cohort designs, 5 repeated cross-sectional studies) representing a total of 1,387,930 individual working-age adults. Among these, 2 studies focused on the male population only. The other 8 studies relied on a time-series design based on surveillance data collected within 5 countries and 1 subnational administrative area.

The recruitment of individuals for repeated cross-sectional studies was carried out from household-based sampling designs. Among cohort studies, 1 article relied on the recruitment of hospital-based participants, 1 recruited participants from occupational health records, and 1 recruited participants from previous surveys completed at home. Most of the time-series studies relied on hospital-based surveillance systems (5 out of 8 articles).

Among the articles included, 5 defined MSDs based on pain, and 11 defined them according to a disability. These 11 articles relying on a disability-related MSD definition were conducted in the Scandinavian countries, the United Kingdom, and Australia. Among those, the site of MSDs was not specified for 8 articles, 2 articles considered MSD affecting the inferior limbs and the back, and 1 considered MSD affecting the superior limbs.

### Risk of bias and quality of the studies

The studies selected were mainly carried out on the general working population, and the risk of selection bias was considered low or probably low for all studies. Overall, study

participants were carefully selected based on a well-defined sampling strategy based on
random selection from a national longitudinal or cross-sectional survey.

We considered MSDs based on medical diagnosis to be reliable. We classified both MSDs
based on the medical diagnosis and/or disability at low risk of bias. MSDs defined based on
pain were classified as a probably low risk of bias. For the studies which administered a
questionnaire, we considered that they probably had low bias since it is a good method for
detecting chronic pain and disability in the individuals recruited. Studies dealing with
temporal trends in MSDs by controlling for age and then for other factors were considered at
low risk of bias for the confounding factors. Studies not taking other potential confounders
were considered likely to be at low risk of bias since here we are only looking at temporal
trends in MSDs. In the included studies, most of the study authors did not declare a conflict
of interest, nor did they receive any support from a company suggesting that there could be
a financial interest in the results. Therefore, we assessed these studies as having a low risk
of bias in this area. For the studies not clearly mentioning it in the paper, we verified that all
the authors were affiliated with public (research) agencies or scientific institutions and, when
this was the case, we considered that the studies had a low probability of bias. We did not
identify any other biases and therefore assessed all studies as having a probable low risk of
other biases.

Among 16 articles, 9 studies included either tests for temporal trends or confidence intervals
for each value of MSD incidence or prevalence over time. These studies were considered to
be of satisfactory quality (trends tests, or Chi-squared), or of probable satisfactory quality
(95% CI for MSDs incidence and/or prevalence). The studies without statistical tests were
considered to be of a probable unsatisfactory quality. In general, we did not identify studies
where there was a high risk of bias, or where the quality was too low to justify an exclusion
from the review (**Table 2**). Additional results regarding classification of bias and evaluation of
statistical methods are provided in Supplementary material S4.

***Temporal trends of the incidence and prevalence of MSDs***
Some studies simultaneously reported results for several MSDs and/or indicators
(prevalence, incidence), such that the 16 included studies included a total of 23 results
extracted for this review. Among these results, 9 used a definition of MSD based on pain and
14 based on impacts on work or social life. Among all results, 12 presented temporal trends
in prevalence and 11 in incidence. The variability of the definitions of MSDs and body sites
studied in the articles of our sample precludes meta-analysis and calculation of pooled
estimates of temporal trends. Five studies controlled for age in time trends in MSDs by

stratification, 4 articles by adjustment, 3 articles by standardizations, 3 articles by direct
standardization, and 3 by age-period-cohort (see **Table 1**).

Among the 3 articles defining MSD prevalence based on disability (26), 1 showed that
absences due to MSDs decreased over time after adjusting for age. Two articles reported an
increase in MSDs over time, 1 of which reported increases in knee osteoarthritis (27), while
the other reported increases in osteoarthritis (28) (**Figure 2.A**). Among the 9 results based
on pain-related MSD definition, 7 showed non-monotonic change over time, and 2 reported
increasing trends (Großschädl et al (28)  for lower back pain, and Guido et al (29) for pain in
all locations) (**Figure 2.B**). These results demonstrate heterogeneity in MSD time trends,
including both increases and non-monotonic changes (**Table 1**).

The temporal evolution of the incidence of MSDs causing disability also tended to increase
or vary according to their site. Among the 6 articles not specifying MSD location, 3 showed
variable trends, two reported increases over time and 1 (30) reported sex-specific results,
with an increase in women and a bell curve for men. In addition, two articles reported an
increase over time in MSDs located in superior limbs. For the inferior limbs and the back,
two articles showed an increase in MSDs, and 1 article reported a variable evolution of
MSDs (**Figure 2.C** and **Table 1**).

***Synthesized evidence***
We note that few articles analyzed temporal variations in MSDs while controlling for age.

Controlling or not controlling for age may produce divergent pictures of temporal trends in
specific MSDs. In the article by Yu et al (31), the raw data suggested non-monotonic
changes in osteoarthritis incidence over time, whereas increased incidence was observed
when standardizing for age. In the article by Dick et al (32), the unadjusted prevalence of
back and hand pain decreased over time. After adjusting for age, decreased prevalence was
still observed among people < 55 years, but variable time trends were observed for those
aged 55-64 years, and an increase was observed for those ≥ 65 years. Thus, these results
highlight potentially distinct impacts of age and time in the occurrence of MSDs, at least for
older age categories.

Fifteen articles suggested that factors other than ageing could explain temporal MSD trends
(**Table 1**). Regardless of the trends observed, most articles hypothesized a link between
cultural changes around the perception of pain (in both caregivers and patients), and a better

knowledge of pathologies with improvements in detection and treatment techniques (29-33).
We also noticed that 5 articles related the trends they reported in MSDs to similar temporal
trends in obesity and body mass index. Additionally, Dick et al (32) suggested that changes
in psychosocial and organizational factors at work could explain the non-monotonic trends
that they observed between 2002 and 2014. Söderberg et al (34) suggested that the non-
monotonic trends they reported reflected changes in disability eligibility criteria rather than in
the underlying exposures. Finally, the only article which reported a decrease in the instances
of sick leave due to MSD (26) explained it by a probable reduction in the physical demands
of work and better health and safety at work.
298

**DISCUSSION:**
This literature review identified a limited number of articles reporting temporal trends in
MSDs while controlling for age. Study duration ranged from 10 to 55 years, which allowed for
longitudinal analysis of MSD occurrence. Temporal trends in MSDs varied according to the
site of the MSDs, the criteria used to define MSDs (either associated with pain and/or a
disability), and the indicator used (prevalence or incidence). We observed temporal
heterogeneity in the occurrence of MSDs considered, with mainly non-monotonic or
increasing trends reported. Of note, based on studies reporting both crude and age-
controlled indicators, we observed that accounting or not accounting for age could lead to
diverging temporal trends, at least among the highest age categories.
309

This literature review identified some important gaps and residual uncertainty in the evidence
currently available. First, although our inclusion criteria were broad, the systematic review
only identified studies conducted in Western, high-income countries: USA, Europe
(especially Scandinavian countries), and Australia. This lack of evidence considering the
burden of MSDs and their socio-economic implications does not allow us to provide an
interpretation of the evolution of MSDs among the global working population (35).
316

Occurrence of the different groups of MSDs considered in this review (pain vs. disability)
varied over time depending on the indicator considered (prevalence vs. incidence) . To
diagnose the occurrence of MSDs, several scales allow for quickly and easily assessing pain
intensity (visual analog pain scale, simple numeric scale, simple verbal scale) (36, 37). It is
important to note, however, that although the validity of these diagnostic tests is comparable
in educated patients, those who are less or uneducated may be led to answer differently.
These scales do not allow for a complete assessment of the pain component, but they can
allow for repeated self-assessments since they are very quick to complete.  It is also

possible that the pain reported by patients responds more quickly to changes in working

conditions or other factors (such as cultural changes) than do longer disabling pathologies.

Therefore, we must remain cautious about our interpretations of temporal changes in the

occurrence of MSDs, depending on whether the observed outcome relates to self-reported

pain or more disabling pathologies diagnosed by doctors (38). Pain classification

measurements must therefore include aspects such as the severity, frequency, and intensity

of pain as well as measurements of changes in working conditions (39).

We hypothesize that observed heterogeneity in temporal trends of MSD occurrence results

from temporal heterogeneity in the evolution of MSD risk factors in different populations. In

most of the countries covered by this review, a fundamental change in the tertiarization of

work has been observed, resulting in an overall reduction in occupational physical

constraints (40, 41). However, a reduction in MSD is not systematically expected from

decreased exposure to biomechanical factors. The analyses from the ESTEV survey show in

particular that the viscoelastic nature of periarticular soft tissues can also play a role in the

occurrence of low back pain (42). Thus, prolonged exposure from carrying heavy loads can

potentially cause an irreversible deformation of these tissues ("memory of the exposure" or

"creep phenomenon"), which may explain the fact that, in older age groups, some MSDs

have not decreased despite decreased biomechanical exposures. Moreover, a decrease in

occupational physical constraints may have arisen concomitantly with an increase in work-

related mental load, which can also play a significant role in the occurrence of MSDs (8, 43).

The main limitation of this review results from the fact that we exclusively searched

electronic bibliometric databases of scientific literature.  This means that we did not consult

the gray literature or governmental reports on MSDs that were not peer-reviewed by external

readers. Another limitation is that, since we used the generic term musculoskeletal

disorders/disease as a keyword, it is possible that we missed articles on specific MSDs that

did not mention the term MSD in the abstract or key terms. Lastly, variability in MSD

definitions and body sites among our study sample prevented us from conducting a meta-

analysis and computing pooled estimates of time trends.

Finally, we do not have studies capturing MSD data during the health crisis linked to the

COVID-19 pandemic. This sanitary situation could possibly be at the origin of the evolution

and emergence of certain professions which can potentially be at the origin of changes in the

occurrence of MSDs (telework and bad postures, sedentary habits, intensification of work,

work on task linked to a digital platform, increased deliveries carrying heavy loads at

reduced times, stress, etc.) (44-46). In the future, longitudinal data that can capture this

information could be an interesting addition to the interpretation and understanding of the

occurrence of MSDs over time (47).

**CONCLUSION:**

To our knowledge, this is the first systematic search and review of studies reporting on MSD

occurrence while accounting for the key confounding impact of age. Our findings suggest

disparity in the literature regarding the temporal evolution of MSD occurrence, depending on

the site of the MSD and whether accounting for MSDs defined by scales of self-reported pain

or disability. Overall, studies controlling for age reported either non-monotonic changes or

increases in MSD occurrence over time. Factors other than ageing that have also been

suggested to underlie temporal changes in MSD occurrence include changes in obesity,

occupational and cultural exposures, and pain tolerance. The current body of evidence,

however, highlights residual uncertainties, especially given the limited number of articles on

this subject and the fact that we only found articles in wealthy countries. Notably, this review

demonstrates the type of research and data that are lacking to anticipate the temporal trends

in the MSDs occurrence, which is an important question in terms of prevention. We also

showed that in the included articles, the temporal trends of MSDs varied mainly between

increase and non-monotonic changes depending on their site, severity, and age.

## FIGURES AND TABLES

**Figure 1:** Flow chart diagram of study selection using PRISMA Flow Diagram recommendations.

* Automatically excluded when the classification tools of the database allowed

**Figure 2:** Temporal trends of the incidence and prevalence of MSDs according to their

location and severity.

**A.** Temporal trend of the prevalence of MSDs inducing repercussions on work or social life with age

considerations. **B.** Temporal trend of the prevalence of pain with age considerations. **C.** Temporal

trend of the incidence of MSDs inducing repercussions on work or social life with age considerations

Unspecified location refers to MSDs that were not associated with a specified body site.

**Table 1:** Summary of articles reporting temporal trends in MSD occurrence.
457

| References | Country | Recruitment | Period | Population | Sample size | MSD type | Diagnosis | Raw time trends | Time trends tacking into account the age | Dealing with age | Interpretation for age specific time trends |
|---|---|---|---|---|---|---|---|---|---|---|---|
| **3 cohort studies** | | | | | | | | | | | |
| Guido et al, 2020 (29) | Europe 17 studies | Variable | 1991–2015 (24 years) | General population | 660,028 individuals | All pain including MSDs | Self-reported | Untreated | Prevalence: increase | Age-period-cohort (ACP) | Evolving perception of pain that can be explained by cultural or biological changes (in patients and practitioners). |
| Söderberg et al, 2020 (34) | Sweden | National occupational health service | 1977-2010 (33 years) | Construction workers (male 20-60) | 389,132 individuals | Disability pension caused by MSDs | Medical exam | Incidence: variable | Incidence: variable | Stratified ages | Changes in welfare legislation (pension eligibility criteria) rather than underlying exposures. |
| Solomon et al, 2007 (48) | UK | Household | 1949-2004 (55 years) | Men from rural areas | 34,486 individuals | MSD-related job loss | Questionnaire | Untreated | Incidence: increased | Adjusted | Acceptance of evolving occupational diseases that can be explained by cultural changes (in patients and practitioners). |
| **5 repeat cross-sectional studies** | | | | | | | | | | | |
| Dick et al, 2020 (32) | US | Household | 2002-2014 (12 years) | General population | 5135 individuals | Back pain | Questionnaire | Prevalence: Decrease | Prevalence: variable. Decrease until 54 years, 55 – 64: variable >65: Increase | Adjusted | Associated with psychosocial and organizational factors at work |
| | | | | | | Pain in arms | | Prevalence: Decrease | Prevalence: variable. Decrease until 54 years, 55 – 64: variable >65: Increase | | |
| Großschädl et al, 2014 (28) | Austria | Household | 1973–2007 (34 years) | General population | 64,052 individuals | Back pain | Self-reported | Untreated | Prevalence: increase | Standardization | Linked to workload, sedentary activities, BMI and obesity, evolving perception of pain, and cultural changes |
| Martin | US | Household | 1997-2010 | General | 78,328 | Back pain | Question | Untreated | Prevalence: variable | Adjusted | Linked with BMI & obesity |

| Author | Country | Data source | Period | Population | N | Disorder | Method | Treatment | Result | Standardization | Hypothesis |
|---|---|---|---|---|---|---|---|---|---|---|---|
| et al, 2014 (49) | | | (13 years) | population >= 40 years old | Individuals | Neck pain | naire | Untreated | Prevalence: variable | | |
| | | | | | | Other MSDs | | Untreated | Prevalence: variable | | |
| Jimenez-Sanchez et al, 2010 (50) | Spain | Household | 1993–2006 (13 years) | General population | 92,893 individuals | Invalidating MS pain | Self-reported | Prevalence: variable | Prevalence: Bell curve (peaked in2001) | Stratified | Absence of hypothesis |
| Leijon et al, 2009 (51) | Sweden | Household | 1990-2006 (16 years) | General population | 63,876 individuals | Low back pain | Self-reported | Untreated | Prevalence: variable | Direct standardization | Linked to increased professional or economic pressure and/or resulting from cultural changes (in media) |

**8 time-series studies**

| Author | Country | Data source | Period | Population | N | Disorder | Method | Treatment | Result | Standardization | Hypothesis |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Ackerman et al, 2019 (52) | Australia | Medical records | 2003–2013 (10 years) | General population | Not mentioned due to time-series studies | Hip Arthroplasties | Medical exam | Untreated | Incidence: increase | Stratified ages | Linked with BMI & obesity |
| | | | | | | Knee Replacements | | Untreated | Incidence: increase | | |
| Gelfman et al, 2009 (53) | US | Medical record | 1981-2005 (24 years) | General population (Olmsted County, Minnesota) | | Carpal tunnel syndrome (CTS) | Medical records | Untreated | Incidence: increase | Direct standardization | Greater awareness of CTS among the general population and increasing proportion of at-risk occupations |
| Holte et al, 2003 (30) | Norway | Administrative records (pensions) | 1968–1997 (29 years) | General population | | Disability pension: RA, OA, soft tissue rheumatism | Medical exam | Untreated | Incidence: increase among women, bell-shaped among men (peaked in the 80s) | Stratified ages | Linked to an increased general demand for fitness or changes in pain perception explained by cultural changes |
| Paloneva et al, 2015 (33) | Finland | Hospital record & surgery | 1998-2011 (13 years) | General population | | Open and arthroscopic rotator cuff repair | Medical exam & surgery | Untreated | Incidence: increase | Stratified ages | Medical and technical advances leading to improved access to diagnosis and surgery |
| Pekkala et al, 2017 | Finland | Administrative records (sickness | 2005–2014 (9 years) | General population (25-64 yrs.) | | Sickness absence due to MSDs | Medical exam | Untreated | Prevalence: decrease | Adjusted | Probably linked to the alleviation of the physical demands of the work and better occupational |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| (26) | | insurance) | | | | | | | | | health safety |
| Spitael s et al, 2020 (27) | Belgiu m | General practitioner s (primary aid) from a network of registers. | 1992-2013 (21 years) | General population | | Knee osteoarthritis | Medical exam | Incidence: U shape Prevalence: increase | Incidence: U shape Prevalence: increase | Standardiz ed and stratified | Linked to BMI & obesity, better access to diagnosis, surgery and preventive medicine, and cultural changes (in patients and practitioners) |
| Swain et al, 2020 (54) | UK | GP hospitals medical record | 1997-2017 (20 years) | General population | | Osteoarthriti s | Medical exam | Incidence: bell curve Prevalence: augmentatio n | Incidence: bell curve (peaked in 2004-2005) Prevalence: increase | Direct standardiz ation and ACP | Cultural changes in practitioners; cohort effect among people born after the 1960s, who may be less exposed to very physically demanding occupations |
| Yu et al, 2017 (31) | UK | Medical records (primary care) | 1992-2013 (21 years) | General population | | Clinical osteoarthritis | Medical exam | Variable | Incidence: increase | Standardiz ed and ACP | Similar trends in obesity, a risk factor for OA, and the increased reporting of painful symptoms |

**Table 2:** Summary of risk of bias and quality across studies on temporal trends of MSDs

| | Ackerman et al,2019 (52) | Dick et al, 2020 (32) | Gelfman et al, 2009 (53) | Großschädl et al, 2014 (28) | Guido et al, 2020 (29) | Holte et al, 2003 (30) | Jimenez-Sanchez et al, 2010 (50) | Leijon et al, 2009 (51) | Martin et al, 2014 (49) | Paloneva et al, 2015 (33) | Pekkala et al, 2017 (26) | Söderberg et al, 2020 (34) | Solomon et al, 2007 (48) | Spitaels et al, 2020 (27) | Swain et al, 2020 (54) | Yu et al, 2017 (31) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| **Bias** | | | | | | | | | | | | | | | | |
| a. Bias in the selection of study participants | L | L | L | L | L | L | L | L | L | L | L | L | PL | L | L | L |
| b. Bias due to misclassification of MSDs | L | PL | L | PL | PL | L | PL | PL | PL | L | L | PL | PL | L | L | L |
| c. Bias due to poor consideration of confounding factors | L | L | L | L | L | L | L | L | L | L | L | PL | PL | L | PL | L |
| d. Bias due to conflict of interest | L | L | PL | L | L | PL | PL | L | PL | L | L | L | L | L | L | L |
| e. Other biases | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL | PL |
| **Quality of the statistical tests** | | | | | | | | | | | | | | | | |
| Quality of the statistical trend tests | PUS | PSQ | PSQ | PUS | SQ | PUS | SQ | PSQ | SQ | PUS | SQ | PUS | PUS | SQ | PUS | PSQ |

475
476    Legend for the bias: low (L), probably low (PL), probably high (PH), high (H).

477      Legend for the quality of the statistical tests: satisfactory quality (SQ), probable satisfactory quality (PSQ), and probable unsatisfactory quality (PUS).

**FINANCIAL SUPPORT**

All authors are salaried staff members of their respective institutions. The authors acknowledge funding from the Cnam-Malakoff Humanis "Corporations and Health" Chair. The funder had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**DECLARATION OF COMPETING INTEREST**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**ACKNOWLEDGMENTS**

The authors are responsible for the opinions expressed in this article, and they do not necessarily represent the views, decisions, or policies of the institutions with which they are affiliated. We also would like to thank David R M Smith for his precious help and careful proofreading of our work.

**REFERENCES:**

1.  Annual Activity Report 2020 - Safety and health at work - EU-OSHA [Internet]. [cited 2021 Jul 9]. Available from: https://osha.europa.eu/en/publications/annual-activity-report-2020-0/view

2.  Musculoskeletal conditions [Internet]. [cited 2020 Oct 27]. Available from: https://www.who.int/news-room/fact-sheets/detail/musculoskeletal-conditions

3.  Roquelaure Y. Musculoskeletal disorders and psychosocial factors at work [Internet]. [Internet]. Brussels: European Trade Union Institute (ETUI); 2018 p. 82. Report No.: 142. Available from: https://www.etui.org/sites/default/files/EN-Report-142-MSD-Roquelaure-WEB.pdf

4.  Carayon P, Smith MJ, Haims MC. Work organization, job stress, and work-related musculoskeletal disorders. Hum Factors. 1999 Dec;41(4):644–63.

5.  The Impact of Musculoskeletal Disorders on Americans – Opportunities for Action | Scoliosis Research Society [Internet]. [cited 2020 Oct 27]. Available from: https://www.srs.org/about-srs/news-and-announcements/the-impact-of-musculoskeletal-disorders-on-americans--opportunities-for-action

6.  Stock SR, Nicolakakis N, Vézina N, Vézina M, Gilbert L, Turcot A, et al. Are work organization interventions effective in preventing or reducing work-related musculoskeletal disorders? A systematic review of the literature. Scand J Work Environ Health. 2018 Mar 1;44(2):113–33.

7.  Linaker C, Harris EC, Cooper C, Coggon D, Palmer KT. The burden of sickness absence from musculoskeletal causes in Great Britan. Occup Med Oxf Engl. 2011 Oct;61(7):458–64.

8.    Niedhammer I, Milner A, LaMontagne AD, Chastang J-F. Study of the validity of a job-
exposure matrix for the job strain model factors: an update and a study of changes over
time. Int Arch Occup Environ Health. 2018 Jul;91(5):523–36.

9.    Bao SS, Kapellusch JM, Merryweather AS, Thiese MS, Garg A, Hegmann KT, et al.
Relationships between job organisational factors, biomechanical and psychosocial
exposures. Ergonomics. 2016;59(2):179–94.

10.   Roquelaure Y, Garlantézec R, Evanoff BA, Descatha A, Fassier J-B, Bodin J. Personal,
biomechanical, psychosocial, and organizational risk factors for carpal tunnel
syndrome: a structural equation modeling approach. PAIN. 2020 Apr;161(4):749–57.

11.   Harris-Adamson C, Eisen EA, Neophytou A, Kapellusch J, Garg A, Hegmann KT, et al.
Biomechanical and psychosocial exposures are independent risk factors for carpal
tunnel syndrome: assessment of confounding using causal diagrams. Occup Environ
Med. 2016 Nov;73(11):727–34.

12.   Storheim K, Zwart J-A. Musculoskeletal disorders and the Global Burden of Disease
study. Ann Rheum Dis. 2014 Jun 1;73(6):949–50.

13.   Briggs AM, Shiffman J, Shawar YR, Åkesson K, Ali N, Woolf AD. Global health policy in
the 21st century: Challenges and opportunities to arrest the global disability burden
from musculoskeletal health conditions. Best Pract Res Clin Rheumatol [Internet]. 2020
Jul 23 [cited 2020 Oct 29]; Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7377715/

14.   Brennan-Olsen SL, Cook S, Leech MT, Bowe SJ, Kowal P, Naidoo N, et al. Prevalence
of arthritis according to age, sex and socioeconomic status in six low and middle
income countries: analysis of data from the World Health Organization study on global
AGEing and adult health (SAGE) Wave 1. BMC Musculoskelet Disord. 2017 Jun
21;18(1):271.

15.   Notenbomer A, van Rhenen W, Groothoff JW, Roelen CAM. Predicting long-term
sickness absence among employees with frequent sickness absence. Int Arch Occup
Environ Health. 2019;92(4):501–11.

16.   WHO | Integrated chronic disease prevention and control [Internet]. WHO. World Health
Organization; [cited 2020 Oct 27]. Available from:
https://www.who.int/chp/about/integrated_cd/en/

17.   Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and
associated methodologies. Health Inf Libr J. 2009;26(2):91–108.

18.   Bouziri H, Dab W, Jean K, Descatha A. Protocol (CRD42020221499). 2020 [Internet].
https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020221499. 2020
[cited 2022 Feb 7].

19.   Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred
reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015:
elaboration and explanation. BMJ. 2015 Jan 2;350:g7647.

20.   Beller EM, Glasziou PP, Altman DG, Hopewell S, Bastian H, Chalmers I, et al. PRISMA
for Abstracts: Reporting Systematic Reviews in Journal and Conference Abstracts.
PLOS Med. 2013 avr;10(4):e1001419.

21.   Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The
PRISMA statement for reporting systematic reviews and meta-analyses of studies that
evaluate health care interventions: explanation and elaboration. PLoS Med. 2009 Jul
21;6(7):e1000100.

22.   Covidence - Better systematic review management [Internet]. Covidence. [cited 2021
Apr 22]. Available from: https://www.covidence.org/

23.   Pega F, Norris SL, Backes C, Bero LA, Descatha A, Gagliardi D, et al. RoB-SPEO: A
tool for assessing risk of bias in studies estimating the prevalence of exposure to
occupational risk factors from the WHO/ILO Joint Estimates of the Work-related Burden
of Disease and Injury. Environ Int [Internet]. 2020 Feb [cited 2021 Apr 16];135.
Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7479507/

24.   Descatha A, Sembajwe G, Baer M, Boccuni F, Di Tecco C, Duret C, et al. WHO/ILO
work-related burden of disease and injury: Protocol for systematic reviews of exposure
to long working hours and of the effect of exposure to long working hours on stroke.
Environ Int. 2018 Oct 1;119:366–78.

25.   Woodruff TJ, Sutton P. The Navigation Guide systematic review methodology: a
rigorous and transparent method for translating environmental health science into
better health outcomes. Environ Health Perspect. 2014 Oct;122(10):1007–14.

26.   Pekkala J, Blomgren J, Pietiläinen O, Lahelma E, Rahkonen O. Occupational class
differences in diagnostic-specific sickness absence: a register-based study in the
Finnish population, 2005–2014. BMC Public Health [Internet]. 2017 Aug 22 [cited 2020
Oct 27];17. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5568169/

27.   Spitaels D, Mamouris P, Vaes B, Smeets M, Luyten F, Hermens R, et al. Epidemiology
of knee osteoarthritis in general practice: a registry-based study. BMJ Open [Internet].
2020 Jan 20 [cited 2020 Oct 29];10(1). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7044813/

28.   Großschädl F, Freidl W, Rásky É, Burkert N, Muckenhuber J, Stronegger WJ. A 35-
Year Trend Analysis for Back Pain in Austria: The Role of Obesity. PLoS ONE
[Internet]. 2014 Sep 10 [cited 2020 Oct 29];9(9). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4160246/

29.   Guido D, Leonardi M, Mellor-Marsá B, Moneta MV, Sanchez-Niubo A, Tyrovolas S, et
al. Pain rates in general population for the period 1991–2015 and 10-years prediction:
results from a multi-continent age-period-cohort analysis. J Headache Pain [Internet].
2020 May 13 [cited 2020 Oct 29];21(1). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7218619/

30.   Holte HH, Tambs K, Bjerkedal T. Time trends in disability pensioning for rheumatoid
arthritis, osteoarthritis and soft tissue rheumatism in Norway 1968-97. Scand J Public
Health. 2003;31(1):17–23.

31.   Yu D, Jordan KP, Bedson J, Englund M, Blyth F, Turkiewicz A, et al. Population trends
in the incidence and initial management of osteoarthritis: age-period-cohort analysis of
the Clinical Practice Research Datalink, 1992–2013. Rheumatol Oxf Engl. 2017
Nov;56(11):1902–17.

32.   Dick RB, Lowe BD, Lu M-L, Krieg EF. Trends in Work-Related Musculoskeletal
Disorders From the 2002 to 2014 General Social Survey, Quality of Work Life
Supplement. J Occup Environ Med. 2020 Aug;62(8):595–610.

33.   Paloneva J, Lepola V, Äärimaa V, Joukainen A, Ylinen J, Mattila VM. Increasing
incidence of rotator cuff repairs—A nationwide registry study in Finland. BMC
Musculoskelet Disord [Internet]. 2015 Aug 12 [cited 2020 Oct 29];16. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4531533/

34.   Söderberg M, Mannelqvist R, Järvholm B, Schiöler L, Stattin M. Impact of changes in
welfare legislation on the incidence of disability pension. A cohort study of construction
workers. Scand J Public Health. 2020 Jun;48(4):405–11.

35.   GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global,
regional, and national incidence, prevalence, and years lived with disability for 328
diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global
Burden of Disease Study 2016. Lancet Lond Engl. 2017 Sep 16;390(10100):1211–59.

36.   Bailly F, Fautrel B, Gossec L. Évaluer la douleur en rhumatologie – comment faire
mieux□? Revue de la littérature. Rev Rhum. 2016 Mar;83(2):105–9.

37.   Wlodyka-Demaille S, Poiraudeau S, Catanzariti J-F, Rannou F, Fermanian J, Revel M.
The ability to change of three questionnaires for neck pain. Joint Bone Spine. 2004 Jul
1;71(4):317–26.

38.   Descatha A, Dale AM, Franzblau A, Evanoff B. Natural history and predictors of long-
term pain and function among workers with hand symptoms. Arch Phys Med Rehabil.
2013 Jul;94(7):1293–9.

39.   Hagberg M, Violante FS, Bonfiglioli R, Descatha A, Gold J, Evanoff B, et al. Prevention
of musculoskeletal disorders in workers: classification and health surveillance –
statements of the Scientific Committee on Musculoskeletal Disorders of the
International Commission on Occupational Health. BMC Musculoskelet Disord. 2012
Dec;13(1):109.

40.   Memmi S. How have the exposures of private sector employees to occupational risks
evolved over the last twenty years? First results of the SUMER survey [Comment ont
évolué les expositions des salariés aux risques professionnels sur les vingt dernières
années ?]. INRS - Occup Health Ref Fr [Internet]. 2019 Sep; N° 159. Available from:
https://www.inrs.fr/dms/inrs/CataloguePapier/DMT/TI-TF-273/tf273.pdf

41.   Havet N, Fournier J, Stefanelli J, Plantier M, Penot A. Disparate exposure to physically
demanding working conditions in France. Rev DÉpidémiologie Santé Publique. 2020
Nov 1;68(6):327–36.

42.   Derrienic F, Touranchet A, Volkoff S. Enquête E.S.T.E.V. un instrument d'étude des
relations entre âge, santé et travail. Enq ESTEV Un Instrum Détude Relat Entre Âge
Santé Trav. 1992;

43.   Niedhammer I, Lesuffleur T, Memmi S, Chastang J-F. Working conditions in the
explanation of occupational inequalities in sickness absence in the French SUMER
study. Eur J Public Health. 2017 Dec 1;27(6):1061–8.

44.   Bouziri H, Smith DRM, Descatha A, Dab W, Jean K. Working from home in the time of
covid-19: how to best preserve occupational health? Occup Environ Med. 2020 Apr 30;

45.  Lord P. Changing world, changing work. Contemp Soc Sci. 2020 Oct 1;15(4):407–15.

46.  Schulte PA, Streit JMK, Sheriff F, Delclos G, Felknor SA, Tamers SL, et al. Potential
Scenarios and Hazards in the Work of the Future: A Systematic Review of the Peer-
Reviewed and Gray Literatures. Ann Work Expo Health. 2020 Oct 8;64(8):786–816.

47.  ACUNA T. Impact of the Covid-19 confinement measures on telework in France - A
qualitative survey [Internet]. EU Science Hub - European Commission. 2020 [cited
2021 May 6]. Available from: https://ec.europa.eu/jrc/en/publication/eur-scientific-and-
technical-research-reports/impact-covid-19-confinement-measures-telework-france-
qualitative-survey.

48.  Solomon C, Poole J, Palmer KT, Coggon D. Health□related job loss: findings from a
community□based survey. Occup Environ Med. 2007 Mar;64(3):144–9.

49.  Martin LG, Schoeni RF. Trends in Disability and Related Chronic Conditions Among the
Forty-and-Over Population: 1997-2010. Disabil Health J. 2014 Jan;7(1 0):S4-14.

50.  Jimenez-Sanchez S, Jimenez-Garcia R, Hernandez-Barrera V, Villanueva-Martinez M,
Rios-Luna A, Fernandez-de-las-Penas C. Has the Prevalence of Invalidating
Musculoskeletal Pain Changed Over the Last 15 Years (1993-2006)? A Spanish
Population-Based Survey. J Pain. 2010 Jul;11(7):612–20.

51.  Leijon O, Mulder M. Prevalence of low back pain and concurrent psychological distress
over a 16-year period. Occup Environ Med. 2009 Feb;66(2):137–9.

52.  Ackerman IN, Bohensky MA, Zomer E, Tacey M, Gorelik A, Brand CA, et al. The
projected burden of primary total knee and hip replacement for osteoarthritis in
Australia to the year 2030. BMC Musculoskelet Disord [Internet]. 2019 Feb 23 [cited
2020 Nov 20];20. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6387488/

53.  Gelfman R, Melton LJ, Yawn BP, Wollan PC, Amadio PC, Stevens JC. Long-term
trends in carpal tunnel syndrome. Neurology. 2009 Jan 6;72(1):33–41.

54.  Swain S, Sarmanova A, Mallen C, Kuo CF, Coupland C, Doherty M, et al. Trends in
incidence and prevalence of osteoarthritis in the United Kingdom: findings from the
Clinical Practice Research Datalink (CPRD). Osteoarthritis Cartilage. 2020 Jun
1;28(6):792–801.

695