

1 **Genomic Surveillance in Japan of AY.29—A New Sub-lineage of SARS-CoV-2**

2 **Delta Variant with C5239T and T5514C Mutations**

3

4 **Takashi Abe^{1,2*}, Masanori Arita²**

5 ¹Faculty of Engineering, Niigata University, 8050, Ikarashi 2-no-cho, Nishi-ku, Niigata-
6 shi, Niigata-ken 950-2181, Japan

7 ²Bioinformation and DDBJ Center, National Institute of Genetics, 1111, Yata, Mishima-
8 shi, Shizuoka-ken 411-8540, Japan

9

10 ***Corresponding author:** Takashi Abe, PhD

11 Faculty of Engineering, Niigata University, 8050, Ikarashi 2-no-cho, Nishi-ku, Niigata-
12 shi, Niigata-ken, 950-2181, Japan.

13 E-mail: takaabe@ie.niigata-u.ac.jp

14

15 **Running title:** AY.29: new sub-lineage of SARS-CoV-2 Delta variant

16

17 **Keywords:** SARS-CoV-2; Delta variant (B.1.617.2); AY.29 sub-lineage

18

19 **ABSTRACT**

20 In the present study, we report a new sub-lineage of the SARS-CoV-2 Delta variant
21 called AY.29, which has C5239T and T5514C mutations. We investigated the monthly
22 trend of AY.29 in Japan within 37,737 Delta variants downloaded on October 2, 2021.
23 Among the total Japanese Delta variants, the AY.29 sub-lineage accounted for 95.1%. In
24 terms of monthly trends, the sequences became predominant in June, and accounted for
25 95.4%, 97.6% and 90.5% of the reported sequences in July, August and September,
26 respectively. Furthermore, the number of Delta variants imported from abroad during
27 the Tokyo 2020 Olympics and Paralympics (held in August 2021) was extremely low
28 during the fifth wave in Japan. Therefore, the epidemic of the new Delta variant is
29 attributable to a newly occurring mutation in Japan.

30

31 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has spread
32 rapidly across the globe since it was first reported in December 2019, and the
33 momentum of its spread has not diminished (1). To address the trend of SARS-CoV-2
34 pandemic, genome sequencing has been performed on a global scale, and the results
35 have been publicized at the global initiative on sharing avian influenza data (GISAID)
36 (2). Currently, Delta variants (Phylogenetic Assignment of Named Global Outbreak
37 (PANGO) (3) lineage: B.1.617.2) are spread worldwide and predominant in many
38 countries including Japan after its first report in India in December 2020 (4, 5).

39 In the present study, genomic surveillance was performed on the genomic
40 sequence data registered at GISAID to investigate the characteristics of the Delta
41 variants isolated in Japan. First, to investigate the diversity, haplotype networks from
42 genomic Delta single nucleotide variants (SNVs) were constructed using median joining

43 network analysis (6) with PopART version 1.7 (7) against 1,348 strains downloaded on
44 August 12, 2021 (Figure 1). The sequences were aligned using MAFFT version 7.475
45 (8), and sequences with gaps were extracted using trimAl version 1.4.rev22 (9). SNVs
46 were detected using snp-sites version 2.5.1 (10), and clustered using CD-HIT version
47 4.8.1 (11). The resulting haplotype network was divided into five groups, described as
48 N1–N5 in Figure 1. An examination of the hub nodes revealed that N1, N2, N3, and N5
49 originated from airport quarantine strains in Japan, all of which were obtained from
50 returning travelers from India. The remaining N4 was of unknown origin. Most strains
51 collected after June belonged to the nodes C1 to C3, generated from N5 containing
52 Japan/IC-1040/2021. There were two mutations (C5239T and T5514C; ORF1ab:
53 V1750A) from N5 to C1, one mutation (C28170T; ORF8: P93S) from C1 to C2, and
54 one mutation (C5365T) from C2 to C3. Therefore, strains in C1 and C3 differed by two
55 mutations only—i.e., C5239T and T5514C (ORF1ab: V1750A). The relationships
56 between protein functions and mutations T5514C (ORF1ab: V1750A) and C28170T
57 (ORF8: P93S)—which are non-synonymous substitutions—are unknown. No other
58 mutations, except for those characterizing the Delta variant, were detected.

59 Next, to determine whether mutations C5239T and T5514C (ORF1ab:
60 V1750A) are unique to Japanese strains, we searched all Delta variants downloaded on
61 September 1, 2021. Of the 908 sequences detected with these two mutations, the
62 countries and numbers of strains were as follows: Japan (883 strains), Hong Kong (2),
63 Belgium (1), Italy (1), United Kingdom (13), USA (2), and New Zealand (6). Japanese
64 strains accounted for 97.2% of the total population. The earliest strains were detected on
65 April 2021 in Japan, and the most recent strains were detected on August 23, 2021 in
66 the UK and Belgium. When the relationship between these strains on the phylogenetic

67 tree was investigated, they were assigned to the new sub-lineage into the Ultrafast
68 Sample placement on Existing tRee (USHER) global phylogenetic tree (12) (Figure 2).
69 Therefore, mutations C5239T and T5514C (ORF1ab: V1750A) are unique to Japan.

70 Based on these results, on September 2, 2021 we proposed to the PANGO
71 nomenclature team—which manages the genetic lineage information about SARS-CoV-
72 2—that the strains with these mutations constitute a new sub-lineage of the Delta
73 variant. After confirmation by PANGO, AY.29 was assigned as a new sub-lineage for
74 Delta variants (13).

75 Finally, we investigated the monthly trends of AY.29 in Japan within 37,737
76 Japanese Delta variants downloaded on October 2, 2021 (Figure 3). Sequences derived
77 from AY.29 accounted for 95.1% (35,904/37,737 strains). The sequences became
78 predominant in June, and accounted for 95.4% (11,419/11,967 strains), 97.6%
79 (23,110/23,669 strains), and 90.5% (735/812 strains) of the detected sequences in July,
80 August and September, respectively. In addition to these two mutations, strains with
81 C28170T (ORF8: P93S) and C5365T mutations also accounted for approximately half
82 of the AY.29 cases.

83 In conclusion, the predominant strain of the Japanese Delta variants currently
84 prevalent in Japan is AY.29, especially between July and September, 2021. In the
85 GISAID data, the number of Delta variants imported from abroad during the Tokyo
86 2020 Olympics and Paralympics (held in August 2021) was extremely low during the
87 fifth wave in Japan. C5239T and T5514C (ORF1ab: V1750A) are mutation markers that
88 can be used to confirm whether a Delta variant is of Japanese origin. Using these
89 markers, it will be possible to quickly confirm whether new Delta variants originate
90 from AY.29 or from other possibly new variants.

91

92 **ACKNOWLEDGMENTS**

93 We gratefully acknowledge the authors that have submitted sequences to the GISAID
94 database. We are grateful to Dr. So Nakagawa (Tokai University) for helpful
95 discussions.

96

97 **FUNDING INFORMATION**

98 This research was supported by JST and CREST (grant number JPMJCR20H1).

99

100 **CONFLICTS OF INTEREST**

101 The authors have no conflicts of interest to declare.

102

103 **REFERENCES**

- 104 1. World Health Organization (WHO). 2021. Coronavirus Disease (COVID-2019),
105 Situation Reports. Available at <[https://www.who.int/emergencies/diseases/novel-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports)
106 [coronavirus-2019/situation-reports](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports)> . Accessed September 8, 2021.
- 107 2. Shu, Y, McCauley, J. GISAID: Global initiative on sharing all influenza data - from
108 vision to reality. Euro Surveill. 2017;22.
- 109 3. Rambaut A, Holmes EC, O'Toole Á et al. A dynamic nomenclature proposal for
110 SARS-CoV-2 lineages to assist genomic epidemiology. Nat Microbiol.
111 2020;5(11):1403-1407.
- 112 4. World Health Organization (WHO). 2021. Tracking SARS-CoV-2 variants.
113 Available at <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>.
114 Accessed September 8, 2021.

- 115 5. Outbreak.info. 2021. COVID-19 Cases & Deaths. Available at
116 <<https://outbreak.info/>>. Accessed September 8, 2021.
- 117 6. Bandelt, HJ, Forster, P, Röhl, A. Median-joining networks for inferring intraspecific
118 phylogenies. *Mol. Biol. Evol.* 1999;16:37–48.
- 119 7. Leigh, JW, Bryant, D. popart: Full-feature software for haplotype network
120 construction. *Methods Eco. Evol.* 2015;6:1110–1116.
- 121 8. Katoh, K, Standley, DM. MAFFT multiple sequence alignment software version 7:
122 Improvements in performance and usability. *Mol. Biol. Evol.* 2013;30:772–780.
- 123 9. Capella-Gutiérrez, S, Silla-Martínez, JM, Gabaldón, T. trimAl: A tool for automated
124 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*
125 2009;25:1972–1973.
- 126 10. Page AJ, Taylor B, Delaney AJ et al. SNP-sites: rapid efficient extraction of SNPs
127 from multi-FASTA alignments. *Microb Genom.* 2016;29:2(4):e000056.
- 128 11. Fu L, Niu B, Zhu Z et al. CD-HIT: accelerated for clustering the next-generation
129 sequencing data. *Bioinformatics.* 2012;1:28(23):3150-3152.
- 130 12. Turakhia Y, Thornlow B, Hinrichs AS et al. Ultrafast Sample Placement on Existing
131 Trees (UShER) Empowers Real-Time Phylogenetics for the SARS-CoV-2
132 Pandemic. *Nat Genet.* 2021; 53(6):809-816.
- 133 13. Proposal for new sub-lineage of AY with C5239T and T5514C (ORF1ab:V1750A)
134 in Japan. Available at < [https://github.com/cov-lineages/pango-](https://github.com/cov-lineages/pango-designation/issues/208)
135 [designation/issues/208](https://github.com/cov-lineages/pango-designation/issues/208)>. Accessed September 8, 2021.
- 136

137 **FIGURE LEGENDS**

138 Figure 1. Haplotype networks. Representative groups are shown in N1 to N5. Nodes in
139 which the strains isolated after June were mainly classified are indicated in C1 to C3.

140 The colors indicate the months in which the strains were isolated at the bottom right of
141 the figure. The sizes of the circles indicate the number of strains.

142

143 Figure 2. Phylogenetic tree derived from candidate sequences assigned by UShER. The
144 colors indicate the clades. The candidate strains are shown in red, and the other strains
145 are shown in the upper right of the figure.

146

147 Figure 3. Monthly trends of (A) the AY.29 sub-lineage and (B) subgroups of AY.29 in
148 Japan. The subgroups C1 to C3 are the group IDs defined in Figure 1. The numbers in
149 parentheses are the total numbers of strains isolated in a particular month. The number
150 of strains belonging to each group is shown at the bottom of the figure.




Figure 1




Figure 2

(A)

Percentage (%)




	2021-04 (73)	2021-05 (287)	2021-06 (927)	2021-07 (11967)	2021-08 (23669)	2021-09 (812)
■ Airport quarantine strains	53	93	76	221	219	66
■ Other Delta variants	10	174	241	327	340	11
■ AY.29	10	20	610	11419	23110	735

■ AY.29 ■ Other Delta variants ■ Airport quarantine strains

(B)

Percentage (%)



	2021-04 (73)	2021-05 (287)	2021-06 (927)	2021-07 (11967)	2021-08 (23669)	2021-09 (812)
■ Airport quarantine strains	53	93	76	221	219	66
■ Other Delta variants	10	174	241	327	340	11
■ C3	7	1	386	6511	11715	393
■ C2	0	4	38	15	6	0
■ C1	3	15	186	4893	11389	342

■ C1 ■ C2 ■ C3 ■ Other Delta variants ■ Airport quarantine strains

Figure 3