Nationally Representative Social Contact Patterns in the United States, August 2020-April 2021

Kristin N. Nelson
Aaron J Siegler
Patrick S Sullivan
Heather Bradley
Eric Hall
Nicole Luisi
Palmer Hipp-Ramsey
Travis Sanchez
Kayoko Shioda
Benjamin A Lopman

Affiliations
Department of Epidemiology, Rollins School of Public Health, Emory University (Nelson, Siegler, Sullivan, Sanchez, Luisi, Lopman)
Department of Population Health Sciences, Georgia State University School of Public Health (Bradley)
Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University (Shioda)
School of Public Health, Oregon Health & Science University (Hall)

Contact information for corresponding author
Kristin N. Nelson, MPH, PhD
1518 Clifton Road NE
Room CNR 4007
Atlanta GA, 30322
Email: knbratt@emory.edu

Declarations of interest: none

Author statement of roles:
KNN conceptualized the study, conducted data analysis, and write the original draft, EH and NL curated and analyzed data, PH was responsible for project administration, AJS, PSS, HB, TS, and BAL designed the larger COVIDVu study and acquired funding, all authors reviewed and edited the manuscript.

Funding source: All were funded by the COVIDVu study, NIAID 3R01AI143875-02S1, the Woodruff Foundation, and the California Department of Public Health. KNN and BAL were funded by GlobalMix NIH/NICHD R01 HD097175 and CorporateMix CDC U01CK000572.

52
53
Abstract
The response to the COVID-19 pandemic in the U.S prompted abrupt and dramatic changes to social contact patterns. Monitoring changing social behavior is essential to provide reliable input data for mechanistic models of infectious disease, which have been increasingly used to support public health policy to mitigate the impacts of the pandemic. While some studies have reported on changing contact patterns throughout the pandemic., few have reported on differences in contact patterns among key demographic groups and none have reported nationally representative estimates. We conducted a national probability survey of US households and collected information on social contact patterns during two time periods: August-December 2020 (before widespread vaccine availability) and March-April 2021 (during national vaccine rollout). Overall, contact rates in Spring 2021 were similar to those in Fall 2020, with most contacts reported at work. Persons identifying as non-White, non-Black, non-Asian, and non-Hispanic reported high numbers of contacts relative to other racial and ethnic groups. Contact rates were highest in those reporting occupations in retail, hospitality and food service, and transportation. Those testing positive for SARS-CoV-2 antibodies reported a higher number of daily contacts than those who were seronegative. Our findings provide evidence for differences in social behavior among demographic groups, highlighting the profound disparities that have become the hallmark of the COVID-19 pandemic.

Keywords

103
Introduction

The response to the COVID-19 pandemic in the U.S., including the closing of schools, workplaces, and
businesses, prompted abrupt and dramatic changes to social contact patterns. In the first months of 2020,
reducing social contact was the only measure available to 'flatten the curve' and blunt the severity of the
early pandemic. A synchronous, near-universal decline in contact rates occurred across countries in North
America, Western Europe, and Asia in the Spring and Summer of 2020, with mean daily contacts
dropping from 7-26 contacts pre-pandemic to 2-5 contacts per person in the early 'lockdown' period. [1]
Through Summer and Fall, as restrictions began to ease, contact patterns slowly rebounded. A key
inflection point occurred in November 2020, when the first COVID-19 vaccines became available,
signaling the start of a massive national vaccination campaign. Widespread vaccination reduced
individuals' risks of infection and led to declining case rates and hospitalizations, contributing to
perceptions of reduced pandemic severity and leading to further relaxation of social distancing policies.
[2] Characterizing changing social contact patterns across this time period is critical to better understand
behavioral drivers of the trajectory of the pandemic and inform ongoing efforts to estimate the impact of
interventions. Systematic data collected in other countries has helped to explain the interplay between
contact patterns and transmission dynamics [3, 4], but studies of contact patterns in the U.S. during this
period draw primarily from convenience samples which survey unrepresentative segments of the
population. [5, 6]

While some studies have reported on changing contact patterns throughout the COVID-19
pandemic in the U.S. [1], few have reported on differences in contact patterns among key demographic
groups. This is particularly important since the burden of the COVID-19 pandemic has disproportionately
fallen on low-income and minority populations, with a heavier burden of COVID-19 cases and deaths in
low-income and minority populations. [7-10] To date, research has suggested that such disparities reflect
limited capacity for these groups to markedly change their contact patterns to protect themselves and their
communities from SARS-CoV-2 infection due to social and structural factors. [11-13] Indeed, studies on
neighborhood-level mobility support the notion that certain demographic groups were more likely to self-
isolate than others during the early pandemic. [14, 15] Disparities in the impact of COVID-19 persisted as
vaccines became available. Inequities in vaccine access and uptake have resulted in suboptimal
vaccination rates in many U.S. communities. [16-18] A person's vaccination status, as well as the
vaccination coverage in their broader community, influences the real and perceived risk of infection,
shaping differences in social contact patterns across demographic groups as the pandemic continues.

Data are needed on contact patterns over the period of the COVID-19 pandemic in the U.S.;
ideally these data should be both nationally representative and investigate differences by demographic
groups. Such data are critical to illustrate a national picture of contact patterns driving the changing
epidemiology of COVID-19 and provide robust data on the behavioral patterns that may be perpetuating
inequities in the impact of the pandemic. Specifically, unbiased estimates of contact will be useful in
parameterizing mechanistic models of infectious disease transmission, informing underlying assumptions
about the rate at which individuals come into contact with one another which are necessary to accurately
simulate spread of infection and the potential impact of interventions. In particular, it is important to
capture contact patterns in the period during which restrictions began to be substantially relaxed but
before the availability of vaccination in the general population and after in order to guide models aiming
to understand the role of vaccines in altering population-level transmission dynamics. With infectious
disease models increasingly informing both domestic and global policy, it is essential that the data
underlying models maintains fidelity to rapidly changing contact patterns, which represents one of the key
sources of uncertainty in COVID-19 transmission models.

We conducted a national probability survey of US households (The COVIDVu Study) with the
primary aim to generate nationally representative estimates of the cumulative incidence of SARS-CoV-2
infection. [19-21] As a part of this study, we collected information on social contact patterns during two
time periods: August-December 2020 (before widespread vaccine availability) and March-April 2021
(during national vaccine rollout). We calculate nationally representative contact rates for the U.S. by
demographic group and assess the relationship between contact rates and SARS-CoV-2 serostatus in both
periods.
160
161
Methods
Sampling
Sampling methods for the COVIDVu study have been previously described. [19] Briefly, we used a
national address-based household sample derived from the USPS Computerized Delivery Sequence File,
which covers all residential delivery points in the US. To achieve a total sample of at least 4,000
responding households, a total of 39,500 addresses were sampled. We oversampled households in census
tracts with >50% Black residents and households with surnames likely to represent Hispanic ethnicity.
[22] One household member aged ≥18 years was randomly selected and offered participation in the study.
Baseline surveys were conducted from August-December 2020 and a follow-up survey was conducted in
March – April 2021.
172
173 Sample weights were used to estimate key parameters that represent the non-institutionalized, housed
174 adult population in the U.S. The COVIDVu study was approved by the Emory University Institutional
175 Review Board (STUDY00000695).
176
Survey and specimen collection
Consenting participants completed an online survey, which collected the age, sex, race, ethnicity, income,
household size, and occupation of participants. We also included questions about person-to-person
contacts on the previous day, which we adapted from previous, similar surveys. [6, 23, 24] We defined a
physical contact as any contact involving physical touch, such as a handshake, hug, or kiss and a non-
physical contact by an interaction in which the participant was within 6 feet of the other person and
exchanged three or more words. For each contact reported in the survey, participants were asked to record
the age of the contact and the location where contact occurred (home, work, school, or other). Occupation
categories were adapted from those used in the 2018 American Community Survey and based on the
North American Industry Classification System. [25]
187
188 Participants self-collected and returned a dried blood spot (DBS) specimen to a central laboratory via
189 prepaid mailer. DBS specimens were tested using the BioRad Platelia Total Antibody test that targets the
190 nucleocapsid protein (i.e., IgA, IgM, IgG; BioRad, Hercules, California). [26, 27]
191
192 Statistical Analysis
193
We restricted our analysis to participants who completed the baseline and follow-up surveys and returned
DBS specimens from each period with valid total Ig result. Where missing, we imputed responses for all
variables involved in calculation of weights, including age, gender, race/ethnicity, and income. [19, 28]
Using the survey weights, we calculated the mean and median daily number of contacts reported in each
time period, overall and by age, sex, location of contact, race/ethnicity, income, occupation, and SARS-
CoV-2 serostatus. We performed two-sample t-tests and Kolmogorov-Smirnov tests for the difference in
contacts reported at baseline and follow-up, which assess differences in the mean and overall distribution,
respectively, of highly skewed data. For comparability with analyses of social contact from other
countries, we truncated data at 50 contacts per day per contact age group to report total contacts [4],
though we explored other cutoffs to assess how the impact on our findings. All analysis was conducted in

R version 3.5.3 and all code is available on GitHub at https://github.com/kbratnelson/covidvu-
socialcontact.
206
Results
39,500 registration materials and kits were mailed to sampled US households. 4,654 (12.6% of sampled
households) returned a baseline survey and a DBS specimen with a valid total Ig result during the period
August 9-December 8, 2020. Of these, 3,112 (66.9% of participants in the baseline survey) returned a
follow-up survey and a DBS specimen with a valid total Ig result during the period March 2-April 18,
2021.
213
Overall, the mean number of daily contacts was 13.9 (IQR: 2, 10) at baseline and 14.5 (IQR: 2, 11) at
follow-up (p of two-sample t-test = 0.552). (Table 1) Physical contacts increased slightly from 5.9 (IQR:
1, 4) to 7.8 (IQR: 0, 5) contacts (p=0.322), and non-physical contacts stayed similar (10.2 (IQR: 0, 5) at
baseline and 9.6 (IQR: 0, 6) at follow-up, p=0.722). Most contacts in both survey periods were reported at
work (65% at baseline and 54% at follow-up), with fewer contacts reported at home (17% at baseline and
21% at follow-up) and other locations (17% at baseline and 22% at follow-up); the fewest contacts were
reported at school (1% at baseline and 3% at follow-up). (Figure 1) (Of note, nearly all participants were
adults, so contacts at school were not expected to be common.) While the distribution of contacts in all
locations was skewed, this skew was particularly apparent among workplace contacts (90[th] percentile of
the distribution: 16 and 15 for baseline and follow-up, respectively).

255
256 Figure 1. Frequency of reported contacts per day by location of contact for a probability sample of 3,112
257 households representing the population aged ≥18 years at study baseline, United States, August-
258 December 2020

259
260 Contacts reported in each location are truncated at 100 to display the lower end of the distribution for
261 each plot. Vertical dotted line shows the mean for each location (as reported in Table 1).
262
263
264

Trends in contact varied by age group (Figure 2). The number of contacts did not substantially change
among 18-24 year-olds (p=0.461), 35-44 year-olds (p=0.097), and 45-54 year-olds (p=0.796) from
baseline to follow-up; comparatively, contact increased among 25-34 year-olds (mean of 16 contacts at
baseline and 23.3 contacts at follow-up, p=0.030). Contact rates among the elderly (65+) were the lowest
of all age groups and remained similar from baseline to follow-up (mean of 5.5 and 6.0 contacts at
baseline and follow-up, respectively, p=0.592). The number of contacts among men were generally higher
than among women and were similar from baseline to follow-up (15.0 to 17.2, p=0.251), contacts
reported among women were also similar between the two surveys (mean of 12.5 and 11.3 at baseline and
follow-up, respectively, p=0.158).

Contact patterns were distinct across socioeconomic and racial groups in both surveys (Figure 2), but
showed little change from baseline to follow-up. Those identifying as Asian reported the lowest contact
rates (6.8 and 11.3 mean contacts at baseline and follow-up, respectively, p=0.019), and those who
identified as non-White, non-Black, non-Asian, and non-Hispanic ('Other') reported the highest contact
rates (baseline mean: 25.5, follow-up mean 28.2, p=0.838). Number of reported contacts was highest
among those reporting annual incomes from $10,000-$49,999. The number of contacts reported at work
varied substantially by industry, with jobs in retail (mean 92 contacts), accommodation and food service
(mean 29 contacts at baseline), transportation (mean 24 contacts at baseline), healthcare (mean 16
contacts at baseline), manufacturing (mean 12.3 contacts at follow-up), and education (mean 12 contacts
at follow-up) accounting for the highest levels of contact.

Participants with a positive serostatus had slightly higher number of contacts than those with a negative
serostatus in both periods (16.6 compared to 13.3 contacts at baseline and 19.6 compared to 13.7 contacts
at follow-up). (Figure 2) Those who were seronegative at both surveys reported a lower mean number of
contacts than those who were seropositive at either survey (13.4 and 16.8 contacts at baseline and follow-
up, respectively). Those who seroconverted from baseline to follow-up reported a higher number of
contacts at both surveys (17.0 and 18.2 contacts) than those who did not (13.6 and 14.1 contacts).

316 Figure 2. Mean reported contacts per day by demographic group and serostatus for a probability sample
317 of 3,112 households representing the population aged ≥18 years at study baseline, United States, August-
318 December 2020

319
320 Violin plots show the distribution of the non-zero numbers of contacts reported by group and grey dots
321 indicate each individual data point. Black dots show the mean for each group (as reported in Table 1).
322
323

We chose to truncate contacts at 50 per day per contact age group (total of 450 contacts) to enable
comparison with other pandemic-focused studies of contact. Moreover, we reasoned that the nature of
some occupations could reasonably lead to several hundred contacts a day. Very few participants reported
above the 50 per day per group limit of contacts (5, or 0.2% of participants, at baseline and 10, or 0.3% of
participants, at follow-up). When we used a truncation limit of 29 total contacts per day, consistent with
other previous studies of contact [23], we found qualitatively similar results to those reported (mean 7.7
contacts (IQR: 2, 10) at baseline to 8.1 contacts (IQR: 2, 11) at follow-up. A moderate number of
participants reported more than 29 contacts (220 participants (7%) at baseline and 249 participants (8%)
at follow-up). In general, those reporting a very high number of contacts reported them at work or at
'other' locations.
334
335   **Discussion**
336
Overall, national contact rates in Spring 2021 were similar to those in Fall 2020, with most contacts in
both surveys reported at work. The number of contacts reported was not uniform across groups, with
those identifying as non-White, non-Black, non-Asian, and non-Hispanic reporting high rates of contact
relative to other racial and ethnic groups. Contact rates were highest among those with lower incomes and
in specific occupational categories, including retail, hospitality and food service, and transportation.
While the number of contacts reported were mostly similar from baseline to follow-up, younger adults
(aged 25-34) reported higher numbers of contacts at follow-up as compared to baseline. Finally, we found
that those testing positive for SARS-CoV-2 antibodies reported a higher number of daily contacts than
those who were seronegative. Collectively, these findings provide robust empirical evidence for
differences in social behavior among demographic groups, highlighting the profound disparities that have
become the hallmark of the COVID-19 pandemic.
348
349   The 'opening up' of much of the U.S. in mid-2020 saw the lifting of restrictive social distancing
measures put into place across much of the country at the beginning of the epidemic. Social contact
reached a nadir in Spring 2020, began to rebound by Fall 2020, and according to our findings stayed
largely consistent through Spring 2021. Although pre-pandemic data on social contact is limited,
available data suggest that contact rates have rebounded to pre-pandemic levels, estimated at 12-14
contacts per day for the average adult. [29, 30] The consistency of overall contact rates across these two
periods is ostensibly the result of a complex interplay of factors. The expiration of many local mitigation
policies in late Summer 2020 [31] likely prompted increases in contact, before the severe wintertime
'wave' arrived in November caused contact rates to decline once again. Optimism about the introduction
of vaccines in early 2021 gradually began to counterbalance the caution prompted by rising case counts in
winter [2]. Together, these factors likely led to largely consistent social contact rates across the two
survey periods.
361
362   Our finding that social contact rates varied by demographic group in the U.S. has been shown in
smaller, less broadly representative studies [5] We found high contact rates among the youngest age
group, 18-24 year-olds, consistent with other data showing that this group typically reported lowest
prevalence of mitigation behaviors. [32-34] We also found that adults 24-35 increased their contact rates
the most from baseline to follow-up, which may reflect the opening of schools and workplaces over this
period, and later, increased contacts after becoming partially or fully vaccinated. Minority populations
and those with lower incomes reported high mean rates of contact. Together with the fact that the majority
of contacts at both baseline and follow-up were reported in the workplace, this is consistent with reports
that lower-income and minority groups were more likely to be classified as 'essential' workers, a
designation which may have required them to continue working in-person or to return to work shortly
after the most strict social distancing restrictions were lifted and prior to our baseline survey. [13] At least
some of this disparity in contact rates is likely due to inflexible and unstable employment situations that
are less likely to allow for remote work. Differential contact rates by job type were expected but striking

in magnitude. We found contact rates up to ten times the population average in occupations which involve
working environments that place employees in close contact with many other people, such as retail,
transportation, and food processing. Notably, workplaces in general are an important source of infection
and transmission risk for respiratory infections [35-37] and the industries associated with high contact
rates in our survey were the settings of several highly publicized outbreaks in 2020. [38, 39] Ensuring that
effective prevention measures are enacted in workplaces, where exposure rates are high, may
disproportionately reduce infection risk.
382
Finally, we found that seropositive individuals were more likely to report higher contact rates at
both survey rounds. Many studies have reported on risk factors for infection related to social
environment, which have included poverty and crowding. [40-42] We coupled behavioral information
with serological status in nationally representative sample to show that higher rates of contact are indeed
linked with previous infection status. Linking information on mitigation behaviors and serostatus will be
important in understanding epidemic dynamics as population-level immunity increases, which are
predicted to be characterized by highly localized outbreaks in space and time driven by 'pockets' of
susceptible persons. [43, 44]
391
The availability of nationally representative data on contact patterns by demographic group fills
an important gap in efforts to accurately forecast and explain the trajectory of the COVID-19 pandemic.
Traditionally, infectious disease models have rarely incorporated social factors, though they are
instrumental in determining disease risk. This is partially because of limited availability of data to
properly formulate models that track many population groups, but also because average population
models are often favored for their simplicity and greater ease of computation. However, accounting for
exposure differences between groups is a critical component of accurately modeling disease dynamics by
population segment. To date, most models that have aimed to explain differences in the burden of cases
and deaths by demographic group have used proxies of contact, such as mobility measured by mobile
phone location data, to represent the frequency of person-person interactions rather than data on direct
interactions. However, recent work has shown that during 'lockdowns', mobility was likely to be a poor
indicator of contact patterns, calling into question the validity of this assumption. [45] Data on contact
patterns by race, ethnicity, socioeconomic group, and occupation provides the necessary granularity to
investigate disease dynamics driving sociodemographic inequities in disease burden. Indeed, modeling
studies that have investigated these questions have noted the lack of longitudinal social contact surveys
that would provide data to allow proper parameterization of models. [46] Systematically collected data
that is finely stratified by demographic group has to date been a major barrier to COVID-19 research
efforts. [47] Collecting and making this data available to the modeling community can support the
development of more accurate models to inform public health policymaking on continued use of vaccines
as well as non-pharmaceutical interventions. As vaccine rollout continues and transmission becomes
increasingly concentrated among unvaccinated and undervaccinated populations, understanding between
and within-group interactions will be particularly critical to track trends in transmission over time and to
understand the potential utility of targeted interventions to mitigate population-level risk.
415
There were several limitations to this study. First, our response rates were low (<15%) but
standard for address-based surveys. For certain groups at risk of being underrepresented by our initial
survey responses, we intentionally oversampled to ensure that such groups were adequately represented,
and survey weights were used to help ensure the sample represented the underlying population. Second,
contact patterns may have changed over the period spanned by our survey (four months in the baseline
survey, one month in the follow-up). In addition, our surveys were conducted before and after the winter
holidays, which likely saw transient increases in travel and contact rates; we could not capture these
changes with our surveys. Third, differences in contact patterns on the state and local levels are likely to
be significant due to variation in the nature of and adherence to distancing and other mitigation policies as
well as differences in the speed of vaccine rollout and eligibility criteria. The national focus of our survey

limits the inferences we can make about contact patterns at a finer geographic scale. With respect to the
contact survey, we asked participants to record contacts from one day (the day prior to completing the
survey) and we do not know to what extent this day might represent general patterns of contact.
Moreover, this prevents an analysis of whether reported contacts were unique on that day, or repeated
across days. Generally, a higher number of unique contacts would represent higher infection and
transmission risk. Fourth, we do not have information on mitigation measures relevant for each contact,
which impact the likelihood of transmitting infection during an interaction. Lastly, the age range of the
surveyed population limits the generalizability of these results to only adults, as there were not aa
sufficient number of participants under 18 to make conclusions about contact rates among younger
people. This also limits our view into school-based contact, which is potentially an important setting of
transmission.
437
As the pandemic continues, continued collection of curation of data on behavioural indicators by
key demographic groups can support the development of detailed COVID-19 models that accurately
represent disease dynamics in varied sociodemographic contexts. Continued, systematic collection and
analysis of social contact data is critical to understand the past and future trajectory of the epidemic as
well as explain differences in disease burden by population segment and through time. Longitudinal
social contact data is particularly powerful alongside seropositivity estimates that are finely stratified by
demographic group, and can further support efforts to properly fit dynamic models aiming assess
interventions to mitigate the continuing impacts of the pandemic.

References

1. Liu CY, Berlin J, Kiti MC, et al. Rapid review of social contact patterns during the COVID-19 pandemic. medRxiv **2021**:2021.03.12.21253410.

2. Trends in Number of COVID-19 Cases and Deaths in the US Reported to CDC, by State/Territory. Accessed July 29 2021.

3. Zhang J, Litvinova M, Liang Y, et al. Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. Science **2020**; 368:1481-6.

4. Jarvis CI, Van Zandvoort K, Gimma A, et al. Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. BMC Medicine **2020**; 18:124.

5. Feehan DM, Mahmud AS. Quantifying population contact patterns in the United States during the COVID-19 pandemic. Nature Communications **2021**; 12:893.

6. Kiti MC, Aguolu OG, Liu CY, et al. Social contact patterns among employees in 3 U.S. companies during early phases of the COVID-19 pandemic, April to June 2020. Epidemics **2021**; 36:100481.

7. Escobar GJ, Adams AS, Liu VX, et al. Racial Disparities in COVID-19 Testing and Outcomes : Retrospective Cohort Study in an Integrated Health System. Ann Intern Med **2021**; 174:786-93.

8. Macias Gil R, Marcelin JR, Zuniga-Blanco B, Marquez C, Mathew T, Piggott DA. COVID-19 Pandemic: Disparate Health Impact on the Hispanic/Latinx Population in the United States. The Journal of Infectious Diseases **2020**; 222:1592-5.

9. Yehia BR, Winegar A, Fogel R, et al. Association of Race With Mortality Among Patients Hospitalized With Coronavirus Disease 2019 (COVID-19) at 92 US Hospitals. JAMA Network Open **2020**; 3:e2018039-e.

10. Jung J, Manley J, Shrestha V. Coronavirus infections and deaths by poverty status: The effects of social distancing. J Econ Behav Organ **2021**; 182:311-30.

11. Chang S, Pierson E, Koh PW, et al. Mobility network models of COVID-19 explain inequities and inform reopening. Nature **2021**; 589:82-7.

12. Selden TM, Berdahl TA. Risk of Severe COVID-19 Among Workers and Their Household Members. JAMA Internal Medicine **2021**; 181:120-2.

13. Roberts JD, Dickinson KL, Koebele E, et al. Clinicians, cooks, and cashiers: Examining health equity and the COVID-19 risks to essential workers. Toxicology and Industrial Health **2020**; 36:689-702.

14. Kissler SM, Kishore N, Prabhu M, et al. Reductions in commuting mobility correlate with geographic differences in SARS-CoV-2 prevalence in New York City. Nature Communications **2020**; 11:4674.

15. Kishore N, Kahn R, Martinez PP, De Salazar PM, Mahmud AS, Buckee CO. Lockdown related travel behavior undermines the containment of SARS-CoV-2. medRxiv **2020**:2020.10.22.20217752.

16. COVID-19 Vaccination Equity. Accessed July 29 2021.

17. M Reitsma SA, J Goldhaber-Fiebert , N Joseph , J Kates, L Levitt , A Rouw,, Salomon J. Disparities in Reaching COVID-19 Vaccination Benchmarks: Projected Vaccination Rates by Race/Ethnicity as of July 4: Kaiser Family Foundaation, **2021**.

18. Disparities in COVID-19 Vaccination Rates across Racial and Ethnic Minority Groups in the United States. Washington, D.C.: U.s. Department of Health and Human Services, Office of the Assistant Secretary for Planning and Evaluation, Office of Science and Data Policy, **2021**.

19. Siegler AJ, Sullivan PS, Sanchez T, et al. Protocol for a national probability survey using home specimen collection methods to assess prevalence and incidence of SARS-CoV-2 infection and antibody response. Ann Epidemiol **2020**; 49:50-60.

20. Lamba K, Bradley H, Shioda K, et al. SARS-CoV-2 Cumulative Incidence and Period Seroprevalence: Results From a Statewide Population-Based Serosurvey in California. Open Forum Infect Dis. Vol. 8, **2021**:ofab379.

21. Sullivan PS, Siegler AJ, Shioda K, et al. SARS-CoV-2 cumulative incidence, United States, August-December 2020. Clin Infect Dis **2021**.

22. Lavange LM, Kalsbeek WD, Sorlie PD, et al. Sample design and cohort selection in the Hispanic Community Health Study/Study of Latinos. Ann Epidemiol **2010**; 20:642-9.

23. Mossong J, Hens N, Jit M, et al. Social Contacts and Mixing Patterns Relevant to the Spread of
Infectious Diseases. PLOS Medicine **2008**; 5:e74.
24. Gimma A, Munday JD, Wong KL, et al. CoMix: Changes in social contacts as measured by the
contact survey during the COVID-19 pandemic in England between March 2020 and March 2021.
medRxiv **2021**:2021.05.28.21257973.
25. Industry and Occupation Code Lists and Crosswalks. Available at:
https://www.census.gov/topics/employment/industry-occupation/guidance/code-lists.html. 2021.
26. Sullivan PS, Sailey C, Guest JL, et al. Detection of SARS-CoV-2 RNA and Antibodies in Diverse
Samples: Protocol to Validate the Sufficiency of Provider-Observed, Home-Collected Blood, Saliva, and
Oropharyngeal Samples. JMIR Public Health Surveill **2020**; 6:e19054.
27. Guest JL, Sullivan PS, Valentine-Graves M, et al. Suitability and Sufficiency of Telehealth Clinician-
Observed, Participant-Collected Samples for SARS-CoV-2 Testing: The iCollect Cohort Pilot Study.
JMIR public health and surveillance **2020**; 6:e19731-e.
28. Andridge RR, Little RJA. A Review of Hot Deck Imputation for Survey Non-response. Int Stat Rev
2010; 78:40-64.
29. Prem K, Cook AR, Jit M. Projecting social contact matrices in 152 countries using contact surveys
and demographic data. PLOS Computational Biology **2017**; 13:e1005697.
30. Feehan DM, Cobb C. Using an Online Sample to Estimate the Size of an Offline Population.
Demography **2019**; 56:2377-92.
31. J Raifman KN, D Jones, J Bor, S Lipson, J Jay, M Cole, N Krawczyk, E Benfer, P Chan, S Galea
COVID-19 US State Policy Database. Ann Arbor, MI: Inter-university Consortium for Political and
Social Research.
32. Coroiu A, Moran C, Campbell T, Geller AC. Barriers and facilitators of adherence to social distancing
recommendations during COVID-19 among a large international sample of adults. PLOS ONE **2020**;
15:e0239795.
33. Ingram M, Zahabian A, Hur C. Prediction of COVID-19 Social Distancing Adherence (SoDA) on the
United States county-level. Humanities and Social Sciences Communications **2021**; 8:87.
34. HJ Hutchins BW, R Leeb, JY Ko, E Odom, J Willey, A Friedman, R Bitsko. COVID-19 Mitigation
Behaviors by Age Group — United States, April–June 2020. MMWR Morb Mortal Wkly Rep. Vol. 69.
Atlanta, GA: Centers for Disease Control and Prevention, **2020**:1584-90.
35. Edwards CH, Tomba GS, de Blasio BF. Influenza in workplaces: transmission, workers' adherence to
sick leave advice and European sick leave recommendations. European Journal of Public Health **2016**;
26:478-85.
36. Lan F-Y, Wei C-F, Hsu Y-T, Christiani DC, Kales SN. Work-related COVID-19 transmission in six
Asian countries/areas: A follow-up study. PLOS ONE **2020**; 15:e0233588.
37. Murti M, Achonu C, Smith BT, et al. COVID-19 Workplace Outbreaks by Industry Sector and Their
Associated Household Transmission, Ontario, Canada, January to June, 2020. Journal of Occupational
and Environmental Medicine **2021**; 63:574-80.
38. MA Waltenburg TV, CE Rose, et al. Update: COVID-19 Among Workers in Meat and Poultry
Processing Facilities ― United States, April–May 2020. In: Prevention CfDCa, ed. MMWR Morb Mortal
Wkly Rep. Vol. 69. Atlanta, GA, **2020**:887-92.
39. Contreras Z, Ngo V, Pulido M, et al. Industry Sectors Highly Affected by Worksite Outbreaks of
Coronavirus Disease, Los Angeles County, California, USA, March 19-September 30, 2020. Emerg
Infect Dis **2021**; 27:1769-75.
40. Saloner B, Parish K, Ward JA, DiLaura G, Dolovich S. COVID-19 Cases and Deaths in Federal and
State Prisons. JAMA **2020**; 324:602-3.
41. Ahmad K, Erqou S, Shah N, et al. Association of poor housing conditions with COVID-19 incidence
and mortality across US counties. PLOS ONE **2020**; 15:e0241327.
42. Emeruwa UN, Ona S, Shaman JL, et al. Associations Between Built Environment, Neighborhood
Socioeconomic Status, and SARS-CoV-2 Infection Among Pregnant Women in New York City. JAMA
2020; 324:390-2.

43. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission dynamics of
SARS-CoV-2 through the postpandemic period. Science **2020**; 368:860-8.
44. Baker RE, Park SW, Yang W, Vecchi GA, Metcalf CJE, Grenfell BT. The impact of COVID-19
nonpharmaceutical interventions on the future dynamics of endemic infections. Proceedings of the
National Academy of Sciences **2020**; 117:30547-53.
45. Buckee C, Noor A, Sattenspiel L. Thinking clearly about social aspects of infectious disease
transmission. Nature **2021**; 595:205-13.
46. Ma KC, Menkir TF, Kissler S, Grad YH, Lipsitch M. Modeling the impact of racial and ethnic
disparities on COVID-19 epidemic dynamics. eLife **2021**; 10:e66601.
47. Krieger N, Testa C, Hanage WP, Chen JT. US racial and ethnic data for COVID-19 cases: still
missing in action. The Lancet **2020**; 396:e81.

Table 1. Mean, median, and quantiles of reported contacts per day by contact location, type, demographic group, and serostatus for a probability sample of 3,112 households representing the population aged ≥18 years, United States, 2020-2021

	Baseline			Follow-up			Sample		Weighted Sample		Statistical tests**	
Characteristic	Mean contacts	Median contacts	(25th, 75th, 90th percentiles)	Mean contacts	Median contacts	(25th, 75th, 90th percentiles)	N	% (column)	N	% (column)	T-test p-value***	K-S test p-value***
Overall	13.9	4.0	2.0, 10.0, 27.0	14.5	5.0	2.0, 11.0, 26.0	3112	100	252117112	100	0.552	0.180
Nature of contact												
Physical	5.9	4.0	1.0, 4.0, 10.0	7.8	2.0	0.0, 5.0, 10.0	-	-	-	-	0.322	0.654
Non-physical	10.2	2.0	0.0, 5.0, 16.0	9.6	2.0	0.0, 6.0, 15.0	-	-	-	-	0.722	0.685
Contact Location												
Home	2.7	1.0	0.0, 3.0, 5.0	3.7	2.0	1.0, 3.0, 5.0	-	-	-	-	0.410	0.596
Work	10.4	0.0	0.0, 1.0, 16.0	9.4	0.0	0.0, 2.0, 15.0	-	-	-	-	0.568	0.993
School	0.2	0.0	0.0, 0.0, 0.0	0.6	0.0	0.0, 0.0, 0.0	-	-	-	-	0.243	1.000
Other	2.7	0.0	0.0, 2.0, 6.0	3.8	0.0	0.0, 2.8, 7.0	-	-	-	-	0.674	1.000
Age category												
18 - 24	20.9	5.0	3, 20, 60	17.7	7.0	3, 16, 62	169	5.4	24823133	9.8	0.461	0.768
25 - 34	16	6.0	3, 13, 35	23.3	7.0	3, 14, 40	448	14.4	42037821	16.7	0.030	1.000
35 - 44	18.3	5.0	3, 13, 35	14.5	6.0	3, 14, 23	503	16.2	43010914	17.1	0.097	1.000
45 - 54	17.7	5.0	2, 11, 32	16.8	5.0	2, 12, 29	509	16.4	41649386	16.5	0.796	1.000
55 - 64	10.7	3.0	1, 7, 26	12.7	4.0	1, 10, 20	649	20.9	44192201	17.5	0.283	0.999
65+	5.5	2.0	1, 5, 10	6.0	2.0	1, 6, 12	834	26.8	56403657	22.4	0.592	1.000
Sex												
Female	12.5	4.0	2, 9, 26	11.3	5.0	2, 10, 20	1907	61.3	115127131	45.7	0.158	0.576
	15	4.0	2, 10, 32	17.2	5.0	2, 12, 32	1205	38.7	136989980	54.3	0.251	1.000

Race/Ethnicity

Hispanic	11.8	4.0	2, 12, 34	14.8	6.0	2, 13, 31	388	12.5	41498755	16.5	0.242	0.931
Non-Hispanic White	14.3	4.0	2, 9, 26	14.0	5.0	2, 11, 23	2118	68.1	161730015	64.1	0.796	1.000
Non-Hispanic Black	16.1	3.0	1, 10, 40	15.5	3.0	1, 9, 42	409	13.1	26371146	10.5	0.846	1.000
Non-Hispanic Asian	6.8	3.0	2, 7, 19	11.3	5.0	2, 9, 26	145	4.7	16297586	6.5	0.019	0.626
Non-Hispanic Other	25.5	6.0	1, 13, 40	28.2	6.0	2, 21, 208	52	1.7	6219608	2.5	0.838	0.999

Income

$0 to $9,999	9.9	4.0	2, 6, 23	14.1	5.0	1, 10, 54	156	5.0	9175348	3.6	0.235	0.624
$10,000 to $24,999	28.9	4.0	1, 9, 71	19.2	4.0	1, 10, 31	282	9.1	21014175	8.3	0.102	1.000
$25,000 to $49,999	16.7	4.0	1, 12, 39	24.5	5.0	2, 14, 43	594	19.1	42517322	16.9	0.020	0.998
$50,000 to $74,999	9.8	4.0	2, 9, 21	12.1	5.0	2, 11, 20	540	17.4	40975725	16.3	0.214	1.000
$75,000 to $99,999	14	4.0	2, 10, 35	12.9	4.0	2, 9, 24	454	14.6	34753959	13.8	0.614	1.000
$100,000 to $149,999	12.9	4.0	2, 10, 32	9.2	4.0	2, 9, 17	534	17.2	44554817	17.7	0.009	1.000
$150,000 to $199,999	12.2	4.0	2, 9, 26	11.8	4.0	2, 11, 30	246	7.9	25854439	10.3	0.871	1.000
$200,000 or higher	9.5	4.0	2, 9, 22	12.4	6.0	2, 12, 22	306	9.8	33271326	13.2	0.150	0.943

Occupation*

Utilities	1.3	0.0	0, 0, 4	10.1	0.0	0, 0, 75	16	0.5	1624138	0.6	0.205	1.000
Construction	8.5	5.0	0, 16, 26	5.9	3.0	1, 10, 15	50	1.6	7468477	3.0	0.166	0.905
Manufacturing	4.4	2.0	0, 8, 15	12.3	0.0	0, 6, 30	50	1.6	8095010	3.2	0.085	0.982
Transportation and Warehousing	92	12.0	0, 65, 171	72.1	4.0	0, 30, 142	84	2.7	12398395	4.9	0.638	0.690
	24.3	6.0	0, 26, 85	3.2	0.0	0, 4, 10	57	1.8	4028061	1.6	0.000	0.446
Information	0.8	0.0	0, 0, 2	0.8	0.0	0, 0, 4	55	1.8	4702720	1.9	0.926	1.000
Finance and Insurance	1.7	0.0	0, 0, 10	2.2	0.0	0, 0, 6	105	3.4	9429022	3.7	0.520	1.000
Real Estate and Rental and Leasing	1.5	0.0	0, 3, 8	1.9	0.0	0, 5, 6	29	0.9	2407083	1.0	0.635	1.000
Professional, Scientific, and Technical Services	1.5	0.0	0, 0, 5	2.1	0.0	0, 0, 6	254	8.2	17627427	7.0	0.327	1.000

Management of Companies and Enterprises	4	0.0	0, 0, 26	8.0	0.0	0, 0, 30	29	0.9	2279303	0.9	0.672	1.000
Administrative Services	4	0.0	0, 3, 15	2.1	0.0	0, 2, 7	25	0.8	1804659	0.7	0.380	1.000
Educational Services	8.5	0.0	0, 10, 20	12.2	0.0	0, 11, 30	227	7.3	11316238	4.5	0.160	1.000
Health Care and Social Assistance	16.4	2.0	0, 17, 46	10.0	2.0	0, 12, 29	313	10.1	24401663	9.7	0.003	1.000
Arts, Entertainment, and Recreation	4.7	0.0	0, 0, 10	3.6	0.0	0, 0, 5	50	1.6	3116292	1.2	0.717	1.000
Accommodation and Food Services	29.3	0.0	0, 10, 49	18.1	6.0	0, 26, 90	55	1.8	7428078	2.9	0.943	1.000
Other Services (except Public Administration)	10.4	0.0	0, 5, 10	4.2	0.0	0, 6, 13	75	2.4	6013537	2.4	0.292	1.000
Public Administration	3.7	0.0	0, 7, 12	2.3	0.0	0, 2, 9	37	1.2	2938101	1.2	0.326	1.000
Other	12.7	0.0	0, 3, 10	14.9	0.0	0, 5, 17	250	8.0	19646552	7.8	0.731	1.000
Household size												
1	6.8	2.0	0, 5, 12	8.7	2.0	0, 8, 19	850	27.3	40909934	16.2	0.104	0.956
2	11.5	3.0	1, 8, 23	12.0	3.0	1, 8, 21	1228	39.5	104424631	41.4	0.722	1.000
3	14.9	5.0	2, 12, 32	14.4	7.0	3, 14, 28	448	14.4	42787020	17.0	0.817	0.958
4	20.3	6.0	3, 14, 42	18.2	7.0	3, 14, 28	383	12.3	39212704	15.6	0.466	0.999
5+	23.6	7.0	4, 21, 40	28.7	7.0	3, 14, 39	203	6.5	24782822	9.8	0.464	1.000
Serostatus at baseline (August-December 2020)												
Positive	16.6	6.0	3, 36, 44	-			-	138	0.0	12316038	4.9	-	-
Negative	13.3	4.0	2, 9, 26	-			-	2974	1.0	239801073	95.1	-	-
Serostatus at follow-up (March-April 2021)												
Positive	-		-	19.6	8.0	3, 14, 31	350	0.1	31360651	12.4	-	-
Negative	-		-	13.7	4.0	2, 10, 25	2762	0.9	220756460	87.6	-	-

* work, not total, contacts

**Statistical tests performed on

weighted data

***The t-test is sensitive to differences in means of the distributions; the K-S test is sensitive to shape of the distribution.

N, total participants. Weighted N: sum of the weights of participants. K-S: Kolmogorov-Smirnov.