

AnthropoAge, a novel approach to integrate body composition into the estimation of biological age

Carlos A. Fermín-Martínez^{1,2*} & Alejandro Márquez-Salinas^{1,2*}, Enrique C. Guerra^{1,2}, Lilian Zavala-Romero⁴, Neftali Eduardo Antonio-Villa^{1,2}, Luisa Fernández-Chirino^{1,3}, Eduardo Sandoval-Colin², Daphne Abigail Barquera-Guevara⁴, Alejandro Campos Muñoz⁴, Arsenio Vargas-Vázquez², César Daniel Paz-Cabrera⁵, Daniel Ramírez-García⁴, Luis Miguel Gutiérrez-Robledo¹, Omar Yaxmehen Bello-Chavolla¹

¹Research Division, Instituto Nacional de Geriátría, ²MD/PhD (PECEM) Program, Facultad de Medicina, Universidad Nacional Autónoma de México, ³Facultad de Química, Universidad Nacional Autónoma de México, ⁴Facultad de Medicina, Universidad Nacional Autónoma de México, ⁵Facultad de Medicina, Universidad Veracruzana Campus Minatitlán

***These authors contributed equally to the drafting of this manuscript**

Correspondence: Omar Yaxmehen Bello-Chavolla. Dirección de Investigación. Instituto Nacional Geriátría. Anillo Perif. 2767, San Jerónimo Lídice, La Magdalena Contreras, 10200, Mexico City, Mexico. Phone: +52 (55) 5548486885. E-mail: oyaxbell@yahoo.com.mx

CONFLICT OF INTERESTS: The authors declare that they have no conflict of interests.

5,375 text words; 57 references; 6 figures, 1 table.


Running headline: AnthropoAge, a novel estimator of biological aging

ABSTRACT (250 WORDS)

Aging is believed to occur across multiple domains, one of which is body composition; however, attempts to integrate it into biological age (BA) have been limited. Here, we consider the sex-dependent role of anthropometry for prediction of 10-year all-cause mortality using data from 18,794 NHANES participants to generate and validate a new BA metric. Our data-driven approach pointed to sex-specific contributors for BA estimation: WHtR, arm and thigh circumferences for men; weight, WHtR, thigh circumference, subscapular and triceps skinfolds for women. We used these measurements to generate AnthroAge, which predicted all-cause mortality (AUROC 0.876, 95%CI 0.864-0.887) and cause-specific mortality independently of race, sex, and comorbidities; AnthroAge was a better predictor than PhenoAge for cerebrovascular, Alzheimer and COPD mortality. A metric of age acceleration was also derived and used to assess sexual dimorphisms linked to accelerated aging, where women had an increase in overall body mass plus an important subcutaneous to visceral fat redistribution, and men displayed a marked decrease in fat and muscle mass. Finally, we showed that consideration of multiple BA metrics may identify unique aging trajectories with increased mortality (HR for multidomain acceleration 2.43, 95%CI 2.25-2.62) and comorbidity profiles. A simplified version of AnthroAge (S-AnthroAge) was generated using only BMI and WHtR, all results were preserved using this metric. In conclusion, AnthroAge is a useful proxy of BA that captures cause-specific mortality and sex dimorphisms in body composition, and it could be used for future multidomain assessments of aging to better characterize the heterogeneity of this phenomenon.

Keywords: AnthroAge, PhenoAge, biological age, chronological age, aging process, anthropometry, body composition

GRAPHICAL ABSTRACT



INTRODUCTION

Aging is a complex phenomenon that researchers have been attempting to characterize for decades, and it is well known that chronological age (CA) does not fully capture its heterogeneity (Kennedy et al., 2014). Methods developed to quantify aging rates need to address the most common physiological changes that occur with aging and capture the variability of individual aging trajectories (López-Otín et al., 2013; Sebastiani et al., 2017). The concept of biological age (BA) goes beyond CA and refers to the underlying processes that modify the susceptibility for development of age-related diseases, disability, and functional impairment, ultimately increasing mortality risk (Belsky et al., 2015; Levine, 2013). Besides calculations derived from epigenetic and omics-based markers, previous efforts have used easily-accessible clinical tools to develop BA estimations, such as Phenotypic Age (PhenoAge), which uses CA and nine blood biomarkers (albumin, creatinine, glucose, C-reactive protein [CRP], mean corpuscular volume [MCV], red blood cell distribution width [RDW], alkaline phosphatase [ALP], white blood cell count [WBC] and lymphocyte percentage) for prediction of 10-year mortality risk (Levine et al., 2018; Liu et al., 2018). Aging is known to be a systemic process, but it is also believed to take place across domains that act in synergy to produce functional changes (Margolick & Ferrucci, 2015; P. -L. Kuo et al., 2020). In line with this idea, it has been shown that some BA estimations tend to be poorly correlated with each other (Li et al., 2020; Earls et al., 2019), while other studies have described BA metrics that capture distinct aspects of aging with unique genetic determinants (C. Kuo et al., 2021). We expect that a broader assessment of aging may provide a more comprehensive overview of its mechanisms, including additional aging domains not previously accounted for in BA estimations (Rivero-Segura et al., 2020). Changes in anthropometry and body composition have been one of the most extensively documented aspects of aging. Overall, aging is characterized by a linear decline in body length, while

body mass initially increases in early and middle adulthood but starts decreasing in older individuals (Fernihough & McGovern, 2015; P. -L. Kuo et al., 2020), a reduction in muscle mass that accelerates at older ages also occurs, as well as a gradual increase in fat mass that is accompanied by a redistribution of adipose tissue from subcutaneous to visceral depots (Wilkinson et al., 2018; Tchernof & Després, 2013); in general, this shift is reflected by a shortening and broadening of the upper body in relation to the lower body (Frenzel et al., 2020). Interestingly, these changes take place at different rates and magnitudes for men and women; to be specific, men appear to have a steeper decline in muscle mass, and the redistribution of adipose tissue is more prominent in women (P. -L. Kuo et al., 2020; Frenzel et al., 2020). These modifications may reflect changes in lifestyle, but they can also be attributed to physiological alterations that occur with aging. Particularly, decline in lean mass can be explained by impaired responses to anabolic stimuli due to systemic inflammation, intramuscular lipid accumulation, stem cell exhaustion, or mitochondrial dysfunction, resulting in a poor regenerative potential of skeletal muscle (López-Otín et al., 2013; Wilkinson et al., 2018). On the other hand, neuroendocrine disturbances such as decrease in sexual hormones (specially in post-menopausal women), insulin resistance and systemic inflammation can hinder the capacity of the subcutaneous adipose tissue to store lipids, resulting in a disproportionate accumulation of visceral and ectopic fat (Tchernof & Després, 2013). Thus, changes in body composition can interact with each other, as well as with other genetic and environmental factors to produce unique age-related risk phenotypes that likely capture alterations in body homeostasis, including frailty, sarcopenic obesity, osteosarcopenia and metabolically unhealthy obesity (April-Sanders & Rodriguez, 2021; Atkins & Wannamethee, 2020; Fried et al., 2021). With all this, it is unsurprising that body composition is often recognized as one of the main domains in which aging takes place (P. -L. Kuo et al., 2020). However, despite the overwhelming evidence supporting the

contribution of body composition and anthropometry to predict age and age-related outcomes, their proper implementation in the development of BA estimations has been limited (Kang et al., 2012; Negasheva et al., 2017). Here, we hypothesize that the integration of sex-based differences in multiple anthropometric measurements would result in the development of a reliable and easily implementable BA estimation that captures the body composition domain of aging. Moreover, we use a combination of PhenoAge and our new metric to investigate whether the simultaneous assessment of more than one domain of aging would identify unique trajectories with differences in mortality risk and body composition patterns. To test these hypotheses, we use anthropometric, laboratory and mortality data from two independent cycles of the National Health Examination Survey, with NHANES-III (1988-1994) used as the training cohort and NHANES-IV (1999-2008) as the validation cohort.

RESULTS

Anthropometric measurements included

For the purposes of our study, we divided anthropometric measurements available in all NHANES cycles into five groups that represent the property of body composition that they likely capture. 1) Body length: height, upper arm length and upper leg length. 2) Body mass: weight and body mass index (BMI). 3) Visceral adiposity: waist circumference and waist-to-height ratio (WHtR). 4) Subcutaneous adiposity: subscapular and triceps skinfolds. 5) Primarily lean mass: mid-upper arm circumference and mid-thigh circumference. (See **Supplementary Table 1**).

Study population

We included only participants ≥ 20 years old with complete anthropometric and mortality data, resulting in a total of 18,794 subjects (NHANES-III: 11,774, NHANES-IV: 7,020). A flowchart diagram of participant selection for each analysis is shown in **Figure 1**, and a

comprehensive overview of our study population is available in **Supplementary Table 2**. Briefly, among participants from both NHANES-III and NHANES-IV, 9,505 were female (50.6%) with a median CA of 45 years (IQR 31-64 years) and a median PhenoAge of 56.8 years (IQR 42.6-76.2 years), most participants self-identified as non-Hispanic White (8,761, 46.6%), followed by Mexican American (4,760, 25.3%), and non-Hispanic Black (4,246, 22.6%). Regarding medical history, 9,268 participants (49.3%) had at least one comorbidity, with hypertension (26.6%) being the most frequent. When assessing the distribution of anthropometric measurements, we found that men had greater values than women for all measurements except for WHtR, subscapular skinfold and triceps skinfold (**Figure 2**). We obtained all-cause mortality follow-up data using information from the National Death Index through March 2020, and a total of 7,853 deaths (31.1%) were recorded, with a mean follow-up of 191 months (IQR 131-288 months). Finally, we compared data from NHANES-III vs NHANES-IV and observed no significant differences in CA; however, median PhenoAge values were higher in NHANES-III (57.9 years [IQR 43.8-78.1] vs. 55.1 years [IQR 40.6-72.6], $p < 0.001$). We observed that all anthropometric variables significantly differed between individuals from both cohorts except for subscapular skinfold for men and BMI for women.

Anthropometric measurements are associated with sex-specific mortality patterns

We first sought to comprehensively characterize the non-linear relationship between anthropometric measurements and mortality risk independently of CA, comorbidities and race/ethnicity using Gompertz proportional hazards models (see **Methods**). Body lengths, particularly height and leg length, displayed an inverse linear relationship with mortality (**Figure 2a-c**). Weight and BMI had similar U-shaped curves as previously reported in numerous studies (Donini et al., 2020; Lavie et al., 2009), where mortality risk was the highest at lower values and virtually invariant thereafter, except for a small increase at extremely high values (particularly in women) (**Figure 2d-e**). Visceral adiposity measured

by WHtR was associated with a progressive risk increase for both genders (**Figure 2f**), while higher reserves of subcutaneous adiposity were related to a marked decline in mortality risk in women, and to a lesser extent in men (**Figure 2g-h**). Interestingly, mortality showed a similar behavior with arm and thigh circumferences as the one seen for BMI (**Figure 2i-j**); we initially established that these variables would capture primarily lean mass, but this has been shown to be influenced by age, gender, and total adiposity (Cavedon et al., 2020). Thus, lower values of BMI, arm and thigh circumferences confer a dramatic increase in mortality likely due to reduced muscle mass, while the slight risk increase observed at greater values could be attributed to higher fat-mass. In summary, highest mortality was observed for decreased total and lean mass, moreover, in most cases men had a higher mortality risk than women except for instances of very low subcutaneous adiposity and extreme obesity.

Estimation of Anthropometric Age

We then developed our own BA estimator which we termed Anthropometric Age (AnthropoAge) by adapting the methods previously described to develop PhenoAge using Gompertz models (Levine et al., 2018). As the relationship between anthropometry and mortality risk is heavily influenced by sex, we decided to perform variable selection separately for men and women. In women, AnthropoAge includes weight, WHtR, thigh circumference, subscapular skinfold, and triceps skinfold; while in men it only includes WHtR, thigh circumference, and arm circumference (**Supplementary Table 3**). Our data driven approach confirms that anthropometric variables contribute to BA and mortality risk in a sex-dependent fashion. Roughly, the best predictors are variables that capture visceral adiposity and lean mass for men, and variables that capture visceral and subcutaneous adiposity for women. Anthropometry is an unexpensive and easily implementable tool for clinical practice (Padilla et al., 2021); however, some of these measurements are not as

commonly used. To promote a wider application of the concept of anthropometric aging into large-scale epidemiological studies, we sought to develop a simpler version of AnthroAge using exclusively BMI and WHtR given the robust association of changes in body fat distribution with aging (Bello-Chavolla et al., 2020; Tchernof & Després, 2013) and the relationships that these metrics displayed with mortality risk; the resulting metric was termed simplified AnthroAge (S-AnthroAge) (**Supplementary Table 4**). AnthroAge and S-AnthroAge act as a proxy of BA and represent the predicted 10-year mortality risk based on an individual's anthropometry. Notably, we found no significant discrepancies or systematic bias between AnthroAge and S-AnthroAge, with overall low bias (0.239 years, 95% CI: 0.208 to 0.270) and a high intra-class correlation coefficient (ICC 0.9940, 95%CI 0.9937-0.9943, $p < 0.001$) (**Supplementary Figure 3**).

Estimation of Anthropometric Age Acceleration

Next, we calculated Phenotypic Age Acceleration (PhenoAgeAccel) as described elsewhere (Liu et al., 2018) and followed this method to obtain a new metric of age acceleration that we called Anthropometric Age Acceleration (AnthroAgeAccel and S-AnthroAgeAccel for the simplified version). These metrics represent the divergence of BA from CA, where values >0 indicate a 10-year mortality risk greater than that predicted by CA (accelerated aging) and values ≤ 0 represent an equal or lower risk (physiological aging). All of our newly developed metrics (AnthroAge, S-AnthroAge, AnthroAgeAccel and S-AnthroAgeAccel) have been deployed within a ShinyApp along with the estimation of PhenoAge to facilitate their use for clinical and research purposes, available at <https://bellolab.shinyapps.io/anthropoage/>.

BA estimates across sex and race/ethnicities

There were no significant differences between men and women in AnthroAge (45.3 years [IQR: 31.7 – 62.6] vs. 44.4 years [IQR: 31.5 – 62.8], $p = 0.671$) nor AnthroAgeAccel (-0.18

years [IQR: -1.88, 1.57] vs -0.21 years [IQR: -2.31, 2.04], $p=0.4406$) in the validation cohort. In contrast, both PhenoAge (56.3 years [IQR: 41.7 – 74.1] vs. 53.7 years [IQR: 39.4 – 71.2], $p<0.001$) and PhenoAgeAccel (-0.08 years [IQR: -2.92, 3.13] vs -2.01 years [IQR: -5.19, 1.79], $p<0.001$) were significantly increased in men (**Figure 3**). On the other hand, we compared age acceleration metrics across race/ethnicities, and observed higher PhenoAgeAccel values in non-Hispanic Black participants, while AnthroAgeAccel and S-AnthroAgeAccel were higher among Mexican Americans (**Supplementary Figure 4**).

AnthroAge and S-AnthroAge predict 10-year all-cause mortality

We evaluated the performance of AnthroAge for prediction of all-cause mortality compared to PhenoAge, CA and individual anthropometric measurements. AnthroAge had a significantly higher AUROC compared to CA, BMI, WHtR, thigh circumference and triceps skinfold. In the training cohort, AnthroAge had a significantly lower AUROC compared to PhenoAge ($p<0.001$), however, they had a similar performance in the validation cohort ($p=0.307$) (**Figure 4a-b**). When assessing AUROC stratified by sex and by number of comorbidities, we observed that AnthroAge performed similarly between men and women and had a better performance for subjects with 1 comorbidity. PhenoAge was slightly superior to AnthroAge in subjects with ≥ 2 comorbidities ($p=0.041$), but there were no significant differences in any other subgroup (**Figure 4c-d, Supplementary Table 5**).

Cause-specific mortality is heterogeneously predicted by different aging metrics

We hypothesized that AnthroAge and PhenoAge would assess different domains of aging. For this purpose, we evaluated whether AnthroAge and PhenoAge were better at predicting distinct mortality causes. All metrics were overall strong to predict cause-specific mortality after adjustment by sex, ethnicity, and number of chronic comorbidities. However, AnthroAge and S-AnthroAge had better predictive performance for cerebrovascular, Alzheimer's disease and chronic lower respiratory disease related mortality, while

PhenoAge was superior for diabetes and nephritis/nephrosis related mortality. All metrics had a roughly comparable performance for cardiovascular, cancer and influenza/pneumonia related mortality (**Table 1, Supplementary Table 6**). Next, we evaluated a possible interaction effect between BA and sex on cause-specific mortality. We found that AnthroAge significantly interacted with female sex to predict cardiovascular disease (HR for interaction 1.014, 95%CI: 1.007-1.020), diabetes mellitus (HR for interaction 1.028, 95%CI: 1.014-1.042), stroke (HR for interaction 1.022, 95%CI: 1.010- 1.034), and cancer (HR for interaction 0.989, 95%CI: 0.984-0.994) related mortality. These results a) suggest that AnthroAge and PhenoAge could be assessing different aspects of aging and b) strengthen the notion that sex dimorphisms link changes in body composition to poor age-related outcomes.

Multidomain age acceleration increases mortality risk and comorbidity burden

We also explored mortality trajectories of subjects with and without accelerated aging using Kaplan-Meier curves. Notably, subjects with accelerated aging displayed higher risk of overall mortality compared to those with physiological aging; this result was replicated for all age acceleration metrics (**Figure 5a-c**). Furthermore, we built an indicator of multidomain age acceleration with the following categories: physiological aging (both AnthroAgeAccel and PhenoAgeAccel ≤ 0), accelerated AnthroAge (AnthroAgeAccel > 0 but PhenoAgeAccel ≤ 0), accelerated PhenoAge (AnthroAgeAccel ≤ 0 but PhenoAgeAccel > 0) and multidomain acceleration (both AnthroAgeAccel and PhenoAgeAccel > 0) and found that individuals with multidomain acceleration had the highest risk of mortality (**Figure 5d**). Cases with accelerated AnthroAge (HR 1.46, 95%CI: 1.35-1.59), accelerated PhenoAge (HR 1.74, 95%CI: 1.60-1.88) and multidomain acceleration (HR 2.43, 95%CI: 2.25-2.62) had higher risk for all-cause mortality compared to individuals with physiological aging after adjustment for age, sex, number of comorbidities, and ethnicity. A similar pattern was

observed with S-AnthropoAge (HR for multidomain acceleration: 2.22, 95%CI: 2.06-2.40). Aging metrics have been previously shown to be associated with comorbidity profiles (C. Kuo et al., 2021; Liu et al., 2018; Sayed et al., 2021), on that basis, we investigated whether specific comorbidity profiles were captured by anthropometric aging. Subjects with an increasing number of comorbidities had higher AnthroAgeAccel and S-AnthroAgeAccel independently of CA categories (**Supplementary Figures 5a, 6a**). Similarly, when using the multidomain aging indicator, subjects had a higher comorbidity burden in the multidomain acceleration category, indicating that simultaneous consideration of these metrics increases the likelihood of identifying accumulation of comorbidities (**Supplementary Figures 5b, 6b**).

Accelerated aging determines specific body composition and biochemical phenotypes

Next, we characterized body composition and biochemical profiles in 6,284 cases with complete DXA information (**Figure 1**). We stratified individuals by sex and compared those with AnthroAgeAccel >0 (accelerated anthropometric aging) to those with AnthroAgeAccel ≤0. These variables are presented as spider plots to facilitate interpretation and pattern-recognition (**Figure 6**), the following description summarizes the most relevant changes found in subjects with accelerated aging. In women, accelerated anthropometric aging was related to a larger overall body mass (comprising fat and lean mass), and an important accumulation of visceral adiposity (greater fat mass and fat-to-lean ratio in trunk, and greater WHtR) disproportionate to their subcutaneous fat reserves (smaller skinfolds, greater trunk-to-appendicular fat mass ratio); notably, these women also presented larger amounts of total and appendicular lean mass. Men with accelerated aging had markedly lesser amounts of fat and lean mass in all body compartments (the most dramatic difference was seen in appendicular lean mass). Although they had a lower subcutaneous adiposity, evidence of visceral fat accumulation was negligible; they also presented reduced bone mineral density. On the other hand, laboratory data showed that

women with accelerated anthropometric aging had a distinctive proinflammatory profile with greater CRP, WBC and ALP and lower albumin and lymphocyte percentage; they also had higher PhenoAge values. Meanwhile, in men with accelerated anthropometric aging the aforementioned changes were not present or were modest (ALP, WBC), however, these subjects had a significantly lower creatinine, which could be attributed to less muscle mass. Using S-AnthropoAgeAccel led to similar body composition patterns linked to accelerated aging, with the only major changes being 1) a more prominent accumulation of abdominal fat and reduction of bone mineral density in both sexes, and 2) no increase in lean mass in women with accelerated aging (**Supplementary Figure 7**). Finally, we sought to evaluate how multidomain acceleration would impact on body composition and biochemical profiles and found that most of the changes described above seem to be greatly augmented when both AnthropoAge and PhenoAge are accelerated (**Supplementary Figure 8**). Our findings denote sharp sexual dimorphisms in body composition and inflammatory biomarkers in individuals at higher risk of mortality, which requires further evaluation.

DISCUSSION

In this work, we developed a novel aging metric based on anthropometric parameters, aiming to predict 10-year mortality risk as a proxy of BA. We showed that AnthropoAge and S-AnthropoAge predict all-cause mortality independently of sex, CA, ethnicity, and number of chronic comorbidities. Notably, AnthropoAge and S-AnthropoAge were better predictors for Alzheimer's, stroke, and chronic lower respiratory related mortality than PhenoAge. We also found that subjects with accelerated anthropometric aging have sharp sexual dimorphisms in body composition and in cause-specific mortality risk. Finally, we demonstrate that PhenoAgeAccel and AnthropoAgeAccel, when considered simultaneously, may identify aging phenotypes with unique comorbidity profiles and

differential risk of all-cause mortality, which may indicate domain or tissue-specific aging rates (Ahadi et al., 2020; C. Kuo et al., 2021).

Body composition has been previously explored for the evaluation of BA, identifying sex-based differences of anthropometric parameters that are linearly associated with CA (Kang et al., 2012; Negasheva et al., 2017). In contrast to these previous studies, AnthroAge uses a non-linear approach to link body composition patterns to mortality risk, which has been speculated to be a better surrogate of BA; this in turn may clarify the relationship of body composition with aging and age-related outcomes beyond individual metrics such as BMI, given that these alone do not completely capture the complexity of this phenomenon (Donini et al., 2020; Lavie et al., 2009). Thus, inclusion of a richer diversity of variables paired with refined modeling methods may lead to more precise assessments of BA (Ahadi et al., 2020; Elliott et al., 2021; Sayed et al., 2021; Xia et al., 2020).

In addition to aging mechanisms that may lead to changes in body composition, numerous studies have also drawn connections between body features and poor age-related outcomes such as disability, age-related diseases, and mortality (Kim & Won, 2022; Santanasto et al., 2016; Schorr et al., 2018). Evidence suggests that the bidirectional relationship between aging and body composition can eventually turn into a vicious cycle that results in an aggravation of age acceleration (Salvestrini et al., 2019), and that this phenomenon is heavily influenced by sex dimorphisms (Goossens et al., 2021; MAGIC et al., 2010; Pomatto et al., 2018; Sampathkumar et al., 2020; Wells, 2007). A relevant addition of our study is that we characterized sexual dimorphisms in the context of accelerated anthropometric aging. Particularly, we observed that females with accelerated aging display increased body mass and a disproportionately larger collection of visceral fat in relation to subcutaneous fat, while males present a phenotype of deeply decreased lean and fat mass with a mild accumulation of visceral adiposity. Furthermore, anthropometric aging significantly

increased cardiovascular, diabetes and stroke-related mortality for women, and cancer-related mortality for men. A plausible explanation for these findings could be postmenopausal estrogens loss, which have been shown to elicit beneficial effects on several processes such as glucose and lipid metabolism in skeletal muscle, adipose tissue, and liver (Goossens et al., 2021), bone homeostasis (Khosla & Monroe, 2018), and cognitive function (Gurvich et al., 2018), among others. Peripheral estrogen production in subcutaneous adipose tissue can offer some of these beneficial effects (Sampathkumar et al., 2020), as a result, a marked shift from subcutaneous to visceral adiposity could precipitate cardiometabolic risk (Goossens et al., 2021; Schorr et al., 2018), as seen in women with accelerated aging from our study. On the other hand, men have been reported to have less muscle mitochondrial content and activity (Rosa-Caldwell & Greene, 2019) and greater systemic proinflammatory responses (Márquez et al., 2020), which may increase their susceptibility for inflammation-induced muscle loss as seen in several age-related diseases including chronic obstructive pulmonary disease and cancer (Machado et al., 2021; Montalvo et al., 2018). Thus, AnthropoAge, S-AnthropoAge and their age acceleration metrics widen our understanding of disparities in aging rates and longevity attributable to sex. Further studies are required to determine the extent to which sex differences in fat distribution and muscle functionality may aid in the estimation of BA and its relative performance in comparison to blood biomarkers.

A previous analysis using data from the Baltimore Longitudinal Study of Aging proposed four domains to integrate functional, phenotypic, and biological aging rates including: body composition, energy regulation, homeostatic mechanisms, and neurodegeneration/neuroplasticity (P. -L. Kuo et al., 2020). Similarly, the work by Kuo et al. demonstrated that PhenoAge and BioAge evaluate different domains of aging, with genetic data suggesting unique aging pathways for each BA indicator (C. Kuo et al., 2021; Levine, 2013). In line with

these findings, we observed that PhenoAge and AnthroAge have distinct abilities to predict cause-specific mortality; furthermore, simultaneous consideration of both accelerated metrics establishes unique mortality risk trajectories and comorbidity profiles, which may better reflect the heterogeneity of aging rates. Consideration of multiple markers of aging may increase the likelihood of modeling complex interactions between different biological domains and it may prove useful to tailor specific strategies to reduce the burden associated with unhealthy aging (Scott et al., 2021). Our newly developed metric could represent a step further in this direction by providing a precise and easily implementable estimation of the body composition domain of biological aging.

This study has numerous strengths that support the validity and clinical relevance of our findings. By using large population-based data, we were able to capture a diverse population, with a wide-range of body phenotypes and varying comorbidity burden. Furthermore, we aimed to shed light on the multidimensionality of the aging process by contrasting and combining multiple aging metrics. Lastly, AnthroAge and S-AnthroAge offer unique opportunities to translate the concept of BA onto clinical practice or routine use, as they can be implemented with simpler measurements compared to those using omics technologies or genomic data (Jansen et al., 2021; Li et al., 2020; Rivero-Segura et al., 2020). Nonetheless, our results are subject to some limitations. First, despite taking races/ethnicities into account to estimate AnthroAge and S-AnthroAge, we found substantial differences in aging rates across racial and ethnic categories, this implies that specific validation studies may be required to adapt these metrics to different populations due to the marked racial and ethnic diversity in aging rates and body composition (Caleyachetty et al., 2021). Second, we were not able to assess how longitudinal changes in AnthroAge, S-AnthroAge and their accelerated metrics modify mortality risk; future longitudinal studies may allow to translate these metrics onto specific strategies to intervene

on aging rates by targeting body composition. Third, despite being readily accessible, the reliability of anthropometric parameters depends upon adequate techniques and reproducibility for their measurement, which may hinder their adequate usage (Wang et al., 2000).

In conclusion, we characterized the CA-independent contribution of body composition to mortality risk for men and women and implemented anthropometric measurements into the estimation of biological age, resulting in the development of complete and simplified versions of AnthroAge and its age acceleration indicator AnthroAgeAccel. These BA estimators capture mortality risk, comorbidity profiles and identify body composition phenotypes with marked sexual dimorphisms linked to accelerated aging, and they can be easily implemented into clinical practice to monitor and promote healthier aging. Finally, we showed that consideration of multiple BA metrics may address the aging process from different perspectives, thus improving the identification of aging trajectories and shedding light on its complexity and heterogeneity.

METHODS

NHANES

NHANES is a nationally representative cross-sectional survey designed to assess health and nutritional status of US population conducted in waves by the Centers for Disease Control and Prevention (CDC). NHANES-III refers to the third wave conducted from 1988 to 1994, and NHANES-IV (also known as continuous NHANES) corresponds to the ongoing wave, which comprises 2-year cycles starting in 1999 (in this study we used data up to 2008). NHANES underwent National Center for Health Statistics (NCHS) Research Ethics Review Board approval, and all participants provided informed consent. Complete methods for recruitment, procedures, and study design for NHANES are described in detail elsewhere (*NHANES - National Health and Nutrition Examination Survey Homepage*, 2022). This

project was registered and approved by the Ethics and Research Committee at Instituto Nacional de Geriatria, project number DI-PI-006/2020.

Body composition data

In addition to demographic, socioeconomic and health-related questionnaires, a subset of NHANES underwent further biochemical and anthropometric evaluations. We only included anthropometric measurements available in both NHANES-III and IV (since calf and hip circumference were not available in all cycles, these parameters were not considered for this study). Normal distribution was assessed with the Anderson-Darling test and variable transformations were performed as specified in **Supplementary Figure 1**; an overview of measuring techniques is provided in **Supplementary Table 1**. For a subset of NHANES-IV, whole-body Dual X-ray Absorptiometry (DXA) assessments were acquired using Hologic QDR 4500A fan-beam bone densitometers.

All-cause and cause-specific mortality data

We obtained all-cause mortality follow-up data using information from the National Death Index for NHANES-III and IV through March 2020. Cause-specific mortality information was evaluated for 8 out of 10 underlying causes of death: cardiovascular, chronic lower respiratory disease, cerebrovascular disease, malignant neoplasms, Alzheimer's disease, diabetes mellitus, influenza/pneumonia, and nephritis/nephrosis; excluding accidents and non-specified causes of death (*NCHS Data Linkage - Mortality Data*, 2022). Follow-up time was estimated from date of initial interview to last follow-up in person-month time.

Estimation of biological age

PhenoAge: PhenoAge was previously developed in NHANES-III and validated in NHANES-IV by Levine et al. (Levine et al., 2018; Liu et al., 2018) using a Gompertz proportional hazards model to predict 10-year mortality risk based on CA and the nine blood biomarkers

previously mentioned. We obtained PhenoAge values by using the equation described elsewhere (Liu et al., 2019).

AnthropoAge: We used a similar approach to develop AnthropoAge using proportional hazards regression models with the parametric Gompertz distribution. First, we fitted two Gompertz models to predict all-cause 10-year mortality risk: the first one had only CA as predictor, and the second had anthropometric measurements as predictors. We equated the risk from both models and solved for age to convert the mortality risk into units of years, thus obtaining AnthropoAge (for a detailed description see **Supplementary Methods**). Importantly, we used orthogonal polynomials to model the non-linear relationship between anthropometry and mortality risk, adjusted for CA and with ethnicity being included in the shape parameter of the Gompertz distribution. To select which anthropometric variables would be included and the optimal number of degrees for orthogonal polynomials, we systematically tested all possible combinations and chose the best model according to minimization of the Bayesian Information Criterion (BIC). We controlled for variable correlation (**Supplementary Figure 2**) and multicollinearity by dropping variables with a Variance Inflation Factor (VIF) ≥ 10 . All Gompertz models were carried out with the *flexsurv* R package, which allows modeling of time-to-event data using any parametric distribution; in particular, the Gompertz distribution assumes an exponential rise in mortality rate as the population ages (Jackson, 2016; Wilson, 1994). To assess average bias and limits of agreement between AnthropoAge and S-AnthropoAge we carried out Bland-Altman analyses with the *blandr* R package.

Estimation of age acceleration

Following the methods by Liu et al. used to calculate PhenoAgeAccel (Liu et al., 2018), we developed a metric to estimate aging acceleration. We regressed AnthropoAge onto CA in a least-squares linear regression and extracted residuals from this model, which represent

the deviation of anthropometric age from CA; a noteworthy difference from method originally described is that AnthroAgeAccel and S-AnthroAgeAccel were calculated separately for men and women.

Assessment of AnthroAge performance

We validated the utility of AnthroAge for prediction of 10-year all-cause mortality in NHANES-IV using Gompertz proportional hazard regression models and the area under receiver operating characteristic curves (AUROC) and compared its performance with PhenoAge, CA, and individual anthropometric measurements using non-parametric ROC tests with bootstrapping (B=1,000) in the *pROC* R package (Robin et al., 2011).

Cause-specific mortality analysis

We evaluated cause-specific mortality using Fine & Gray semiparametric competitive risk regression models with the *survival* R package (Therneau et al., 2022) to contrast predictive performance between AnthroAge, S-AnthroAge and PhenoAge in subjects with complete mortality data. All models were adjusted by sex, ethnicity, and number of chronic comorbidities. Predictive performance was assessed by the C-statistic and BIC differences ($\Delta\text{BIC} = \text{BIC}_{\text{Model-1}} - \text{BIC}_{\text{Model-2}}$), where a $\Delta\text{BIC} < -2$ indicates that the first model is better, a $\Delta\text{BIC} > 2$ indicates that the second model is better, and a ΔBIC in between indicates that both models are comparable for the prediction of a specific outcome.

Assessment of mortality trajectories

We used Kaplan-Meier curves and the log-rank test with the *survival* R package to compare mortality trajectories between subjects with physiological and accelerated aging for all metrics (AnthroAgeAccel, S-AnthroAgeAccel and PhenoAgeAccel). We also plotted Kaplan-Meier curves to compare mortality between categories from the multidomain aging indicator.

Body composition and biochemical phenotypes

We evaluated sex-specific differences in body composition and biochemical profile between subjects with accelerated and non-accelerated aging using laboratory data, anthropometry and DXA measurements from NHANES-IV cycles. Regarding DXA, we divided fat mass and lean mass variables by squared height, resulting in standardized fat mass and lean mass indexes (FMI and LMI, respectively); this was done for whole-body, trunk and appendicular compartments. We also calculated fat-to-lean ratios (fat mass divided by lean mass) in whole-body and trunk and a trunk-to-appendicular fat ratio (trunk fat mass divided by appendicular fat mass) to assess disproportionate fat accumulations. All variables were Z-transformed to facilitate visualization using spider plots with the *fmsb* R package.

Comparisons between categorical and continuous variables were performed with the chi-squared and the Mann-Whitney U test, respectively. All statistical analyses were conducted using R version 4.1.1 and p-values thresholds are estimated for a two-sided significance level of $\alpha=0.05$.

ACKNOWLEDGMENTS

This project was registered and approved by the Research Committee at Instituto Nacional de Geriatria, project number DI-PI-006/2020. NEAV, CAFM, AMS, ECG, and ESC are enrolled at the PECEM Program of the Faculty of Medicine at UNAM. NEAV and ESC supported by CONACyT. The graphical abstract and figure 1 were designed using resources created by *Victoruler*, *Smashicons*, *Freepik* and *Surang* from www.flaticon.com

AUTHOR CONTRIBUTIONS

Research idea and study design: CAFM, AMS, ECG, NEAV, LMGR, OYBC; data acquisition: CAFM, AMS, ECG, OYBC; analysis/interpretation: CAFM, AMS, OYBC; statistical analysis: CAFM, AMS, OYBC; manuscript drafting: CAFM, AMS, ECG, LZR, LFC, DABG, NEAV, ESC, ACM, AVV, CDPC, DRG, LMGR, OYBC; supervision or mentorship: OYBC. Each author contributed important intellectual content during manuscript drafting or

revision and accepts accountability for the overall work by ensuring that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved.

DATA AVAILABILITY: All code, datasets and materials are available for reproducibility of results at <https://github.com/oyaxbell/anthropoage/>

CONFLICT OF INTEREST/FINANCIAL DISCLOSURE: Nothing to disclose.

FUNDING: This study was supported by Instituto Nacional de Geriátría.

REFERENCES

- Ahadi, S., Zhou, W., Schüssler-Fiorenza Rose, S. M., Sailani, M. R., Contrepois, K., Avina, M., Ashland, M., Brunet, A., & Snyder, M. (2020). Personal aging markers and ageotypes revealed by deep longitudinal profiling. *Nature Medicine*, 26(1), 83–90. <https://doi.org/10.1038/s41591-019-0719-5>
- April-Sanders, A. K., & Rodriguez, C. J. (2021). Metabolically Healthy Obesity Redefined. *JAMA Network Open*, 4(5), e218860. <https://doi.org/10.1001/jamanetworkopen.2021.8860>
- Atkins, J. L., & Wannamethee, S. G. (2020). Sarcopenic obesity in ageing: Cardiovascular outcomes and mortality. *British Journal of Nutrition*, 124(10), 1102–1113. <https://doi.org/10.1017/S0007114520002172>
- Bello-Chavolla, O. Y., Antonio-Villa, N. E., Vargas-Vázquez, A., Viveros-Ruiz, T. L., Almeda-Valdes, P., Gomez-Velasco, D., Mehta, R., Elias-López, D., Cruz-Bautista, I., Roldán-Valadez, E., Martagón, A. J., & Aguilar-Salinas, C. A. (2020). Metabolic Score for Visceral Fat (METS-VF), a novel estimator of intra-abdominal fat content and cardio-metabolic health. *Clinical Nutrition*, 39(5), 1613–1621. <https://doi.org/10.1016/j.clnu.2019.07.012>

- Belsky, D. W., Caspi, A., Houts, R., Cohen, H. J., Corcoran, D. L., Danese, A., Harrington, H., Israel, S., Levine, M. E., Schaefer, J. D., Sugden, K., Williams, B., Yashin, A. I., Poulton, R., & Moffitt, T. E. (2015). Quantification of biological aging in young adults. *Proceedings of the National Academy of Sciences*, 112(30). <https://doi.org/10.1073/pnas.1506264112>
- Caleyachetty, R., Barber, T. M., Mohammed, N. I., Cappuccio, F. P., Hardy, R., Mathur, R., Banerjee, A., & Gill, P. (2021). Ethnicity-specific BMI cutoffs for obesity based on type 2 diabetes risk in England: A population-based cohort study. *The Lancet Diabetes & Endocrinology*, 9(7), 419–426. [https://doi.org/10.1016/S2213-8587\(21\)00088-7](https://doi.org/10.1016/S2213-8587(21)00088-7)
- Cavedon, V., Milanese, C., & Zancanaro, C. (2020). Are body circumferences able to predict strength, muscle mass and bone characteristics in obesity? A preliminary study in women. *International Journal of Medical Sciences*, 17(7), 881–891. <https://doi.org/10.7150/ijms.41713>
- Donini, L. M., Pinto, A., Giusti, A. M., Lenzi, A., & Poggiogalle, E. (2020). Obesity or BMI Paradox? Beneath the Tip of the Iceberg. *Frontiers in Nutrition*, 7, 53. <https://doi.org/10.3389/fnut.2020.00053>
- Earls, J. C., Rappaport, N., Heath, L., Wilmanski, T., Magis, A. T., Schork, N. J., Omenn, G. S., Lovejoy, J., Hood, L., & Price, N. D. (2019). Multi-Omic Biological Age Estimation and Its Correlation With Wellness and Disease Phenotypes: A Longitudinal Study of 3,558 Individuals. *The Journals of Gerontology: Series A*, 74(Supplement_1), S52–S60. <https://doi.org/10.1093/gerona/glz220>
- Elliott, M. L., Caspi, A., Houts, R. M., Ambler, A., Broadbent, J. M., Hancox, R. J., Harrington, H., Hogan, S., Keenan, R., Knodt, A., Leung, J. H., Melzer, T. R., Purdy, S. C., Ramrakha, S., Richmond-Rakerd, L. S., Righarts, A., Sugden, K., Thomson, W. M.,

- Thorne, P. R., ... Moffitt, T. E. (2021). Disparities in the pace of biological aging among midlife adults of the same chronological age have implications for future frailty risk and policy. *Nature Aging*, 1(3), 295–308. <https://doi.org/10.1038/s43587-021-00044-4>
- Fernihough, A., & McGovern, M. E. (2015). Physical stature decline and the health status of the elderly population in England. *Economics & Human Biology*, 16, 30–44. <https://doi.org/10.1016/j.ehb.2013.12.010>
- Frenzel, A., Binder, H., Walter, N., Wirkner, K., Loeffler, M., & Loeffler-Wirth, H. (2020). The aging human body shape. *Npj Aging and Mechanisms of Disease*, 6(1), 5. <https://doi.org/10.1038/s41514-020-0043-9>
- Fried, L. P., Cohen, A. A., Xue, Q.-L., Walston, J., Bandeen-Roche, K., & Varadhan, R. (2021). The physical frailty syndrome as a transition from homeostatic symphony to cacophony. *Nature Aging*, 1(1), 36–46. <https://doi.org/10.1038/s43587-020-00017-z>
- Goossens, G. H., Jocken, J. W. E., & Blaak, E. E. (2021). Sexual dimorphism in cardiometabolic health: The role of adipose tissue, muscle and liver. *Nature Reviews Endocrinology*, 17(1), 47–66. <https://doi.org/10.1038/s41574-020-00431-8>
- Gurvich, C., Hoy, K., Thomas, N., & Kulkarni, J. (2018). Sex Differences and the Influence of Sex Hormones on Cognition through Adulthood and the Aging Process. *Brain Sciences*, 8(9), 163. <https://doi.org/10.3390/brainsci8090163>
- Jackson, C. (2016). **flexsurv**: A Platform for Parametric Survival Modeling in R. *Journal of Statistical Software*, 70(8). <https://doi.org/10.18637/jss.v070.i08>
- Jansen, R., Han, L. K., Verhoeven, J. E., Aberg, K. A., van den Oord, E. C., Milaneschi, Y., & Penninx, B. W. (2021). An integrative study of five biological clocks in somatic and mental health. *ELife*, 10, e59479. <https://doi.org/10.7554/eLife.59479>

- Kang, Y. G., Chul-Young Bae, Young-Sung Suh, Jee Hye Han, Sung-Soo Kim, & Kyung Won Shim. (2012). A model for estimating body shape biological age based on clinical parameters associated with body composition. *Clinical Interventions in Aging*, 11. <https://doi.org/10.2147/CIA.S38220>
- Kennedy, B. K., Berger, S. L., Brunet, A., Campisi, J., Cuervo, A. M., Epel, E. S., Franceschi, C., Lithgow, G. J., Morimoto, R. I., Pessin, J. E., Rando, T. A., Richardson, A., Schadt, E. E., Wyss-Coray, T., & Sierra, F. (2014). Geroscience: Linking Aging to Chronic Disease. *Cell*, 159(4), 709–713. <https://doi.org/10.1016/j.cell.2014.10.039>
- Khosla, S., & Monroe, D. G. (2018). Regulation of Bone Metabolism by Sex Steroids. *Cold Spring Harbor Perspectives in Medicine*, 8(1), a031211. <https://doi.org/10.1101/cshperspect.a031211>
- Kim, S., & Won, C. W. (2022). Sex-different changes of body composition in aging: A systemic review. *Archives of Gerontology and Geriatrics*, 102, 104711. <https://doi.org/10.1016/j.archger.2022.104711>
- Kuo, C., Pilling, L. C., Liu, Z., Atkins, J. L., & Levine, M. E. (2021). Genetic associations for two biological age measures point to distinct aging phenotypes. *Aging Cell*, 20(6). <https://doi.org/10.1111/accel.13376>
- Kuo, P. -L., Schrack, J. A., Shardell, M. D., Levine, M., Moore, A. Z., An, Y., Elango, P., Karikkineth, A., Tanaka, T., Cabo, R., Zukley, L. M., AlGhatrif, M., Chia, C. W., Simonsick, E. M., Egan, J. M., Resnick, S. M., & Ferrucci, L. (2020). A roadmap to build a phenotypic metric of ageing: Insights from the Baltimore Longitudinal Study of Aging. *Journal of Internal Medicine*, 287(4), 373–394. <https://doi.org/10.1111/joim.13024>

- Lavie, C. J., Milani, R. V., & Ventura, H. O. (2009). Obesity and Cardiovascular Disease. *Journal of the American College of Cardiology*, *53*(21), 1925–1932. <https://doi.org/10.1016/j.jacc.2008.12.068>
- Levine, M. E. (2013). Modeling the Rate of Senescence: Can Estimated Biological Age Predict Mortality More Accurately Than Chronological Age? *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *68*(6), 667–674. <https://doi.org/10.1093/gerona/gls233>
- Levine, M. E., Lu, A. T., Quach, A., Chen, B. H., Assimes, T. L., Bandinelli, S., Hou, L., Baccarelli, A. A., Stewart, J. D., Li, Y., Whitsel, E. A., Wilson, J. G., Reiner, A. P., Aviv, A., Lohman, K., Liu, Y., Ferrucci, L., & Horvath, S. (2018). An epigenetic biomarker of aging for lifespan and healthspan. *Aging*, *10*(4), 573–591. <https://doi.org/10.18632/aging.101414>
- Li, X., Ploner, A., Wang, Y., Magnusson, P. K., Reynolds, C., Finkel, D., Pedersen, N. L., Jylhävä, J., & Hägg, S. (2020). Longitudinal trajectories, correlations and mortality associations of nine biological ages across 20-years follow-up. *ELife*, *9*, e51507. <https://doi.org/10.7554/eLife.51507>
- Liu, Z., Kuo, P.-L., Horvath, S., Crimmins, E., Ferrucci, L., & Levine, M. (2018). A new aging measure captures morbidity and mortality risk across diverse subpopulations from NHANES IV: A cohort study. *PLOS Medicine*, *15*(12), e1002718. <https://doi.org/10.1371/journal.pmed.1002718>
- Liu, Z., Kuo, P.-L., Horvath, S., Crimmins, E., Ferrucci, L., & Levine, M. (2019). Correction: A new aging measure captures morbidity and mortality risk across diverse subpopulations from NHANES IV: A cohort study. *PLOS Medicine*, *16*(2), e1002760. <https://doi.org/10.1371/journal.pmed.1002760>

- López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2013). The Hallmarks of Aging. *Cell*, 153(6), 1194–1217. <https://doi.org/10.1016/j.cell.2013.05.039>
- Machado, F. V. C., Spruit, M. A., Coenjaerds, M., Pitta, F., Reynaert, N. L., & Franssen, F. M. E. (2021). Longitudinal changes in total and regional body composition in patients with chronic obstructive pulmonary disease. *Respirology*, 26(9), 851–860. <https://doi.org/10.1111/resp.14100>
- MAGIC, Heid, I. M., Jackson, A. U., Randall, J. C., Winkler, T. W., Qi, L., Steinthorsdottir, V., Thorleifsson, G., Zillikens, M. C., Speliotes, E. K., Mägi, R., Workalemahu, T., White, C. C., Bouatia-Naji, N., Harris, T. B., Berndt, S. I., Ingelsson, E., Willer, C. J., Weedon, M. N., ... Lindgren, C. M. (2010). Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nature Genetics*, 42(11), 949–960. <https://doi.org/10.1038/ng.685>
- Margolick, J. B., & Ferrucci, L. (2015). Accelerating aging research: How can we measure the rate of biologic aging? *Experimental Gerontology*, 64, 78–80. <https://doi.org/10.1016/j.exger.2015.02.009>
- Márquez, E. J., Chung, C., Marches, R., Rossi, R. J., Nehar-Belaid, D., Eroglu, A., Mellert, D. J., Kuchel, G. A., Banchereau, J., & Ucar, D. (2020). Sexual-dimorphism in human immune system aging. *Nature Communications*, 11(1), 751. <https://doi.org/10.1038/s41467-020-14396-9>
- Montalvo, R. N., Counts, B. R., & Carson, J. A. (2018). Understanding sex differences in the regulation of cancer-induced muscle wasting. *Current Opinion in Supportive & Palliative Care*, 12(4), 394–403. <https://doi.org/10.1097/SPC.0000000000000380>
- NCHS Data Linkage—Mortality Data. (2022, July 28). <https://www.cdc.gov/nchs/data-linkage/mortality.htm>

- Negasheva, M. A., Zimina, S. N., Lapshina, N. E., & Sineva, I. M. (2017). Express Estimation of the Biological Age by the Parameters of Body Composition in Men and Women over 50 Years. *Bulletin of Experimental Biology and Medicine*, 163(3), 405–408. <https://doi.org/10.1007/s10517-017-3814-y>
- NHANES - National Health and Nutrition Examination Survey Homepage. (2022, September 29). <https://www.cdc.gov/nchs/nhanes/index.htm>
- Padilla, C. J., Ferreyro, F. A., & Arnold, W. D. (2021). Anthropometry as a readily accessible health assessment of older adults. *Experimental Gerontology*, 153, 111464. <https://doi.org/10.1016/j.exger.2021.111464>
- Pomatto, L. C. D., Tower, J., & Davies, K. J. A. (2018). Sexual Dimorphism and Aging Differentially Regulate Adaptive Homeostasis. *The Journals of Gerontology: Series A*, 73(2), 141–149. <https://doi.org/10.1093/gerona/glx083>
- Rivero-Segura, N. A., Bello-Chavolla, O. Y., Barrera-Vázquez, O. S., Gutierrez-Robledo, L. M., & Gomez-Verjan, J. C. (2020). Promising biomarkers of human aging: In search of a multi-omics panel to understand the aging process from a multidimensional perspective. *Ageing Research Reviews*, 64, 101164. <https://doi.org/10.1016/j.arr.2020.101164>
- Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C., & Müller, M. (2011). pROC: An open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics*, 12(1), 77. <https://doi.org/10.1186/1471-2105-12-77>
- Rosa-Caldwell, M. E., & Greene, N. P. (2019). Muscle metabolism and atrophy: Let's talk about sex. *Biology of Sex Differences*, 10(1), 43. <https://doi.org/10.1186/s13293-019-0257-3>
- Salvestrini, V., Sell, C., & Lorenzini, A. (2019). Obesity May Accelerate the Aging Process. *Frontiers in Endocrinology*, 10, 266. <https://doi.org/10.3389/fendo.2019.00266>

- Sampathkumar, N. K., Bravo, J. I., Chen, Y., Danthi, P. S., Donahue, E. K., Lai, R. W., Lu, R., Randall, L. T., Vinson, N., & Benayoun, B. A. (2020). Widespread sex dimorphism in aging and age-related diseases. *Human Genetics*, *139*(3), 333–356. <https://doi.org/10.1007/s00439-019-02082-w>
- Santanasto, A. J., Goodpaster, B. H., Kritchevsky, S. B., Miljkovic, I., Satterfield, S., Schwartz, A. V., Cummings, S. R., Boudreau, R. M., Harris, T. B., & Newman, A. B. (2016). Body Composition Remodeling and Mortality: The Health Aging and Body Composition Study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, glw163. <https://doi.org/10.1093/gerona/glw163>
- Sayed, N., Huang, Y., Nguyen, K., Krejciova-Rajaniemi, Z., Grawe, A. P., Gao, T., Tibshirani, R., Hastie, T., Alpert, A., Cui, L., Kuznetsova, T., Rosenberg-Hasson, Y., Ostan, R., Monti, D., Lehallier, B., Shen-Orr, S. S., Maecker, H. T., Dekker, C. L., Wyss-Coray, T., ... Furman, D. (2021). An inflammatory aging clock (iAge) based on deep learning tracks multimorbidity, immunosenescence, frailty and cardiovascular aging. *Nature Aging*, *1*(7), 598–615. <https://doi.org/10.1038/s43587-021-00082-y>
- Schorr, M., Dichtel, L. E., Gerweck, A. V., Valera, R. D., Torriani, M., Miller, K. K., & Bredella, M. A. (2018). Sex differences in body composition and association with cardiometabolic risk. *Biology of Sex Differences*, *9*(1), 28. <https://doi.org/10.1186/s13293-018-0189-3>
- Scott, A. J., Ellison, M., & Sinclair, D. A. (2021). The economic value of targeting aging. *Nature Aging*, *1*(7), 616–623. <https://doi.org/10.1038/s43587-021-00080-0>
- Sebastiani, P., Thyagarajan, B., Sun, F., Schupf, N., Newman, A. B., Montano, M., & Perls, T. T. (2017). Biomarker signatures of aging. *Aging Cell*, *16*(2), 329–338. <https://doi.org/10.1111/acel.12557>

- Tchernof, A., & Després, J.-P. (2013). Pathophysiology of Human Visceral Obesity: An Update. *Physiological Reviews*, 93(1), 359–404. <https://doi.org/10.1152/physrev.00033.2011>
- Therneau, T. M., until 2009), T. L. (original S.->R port and R. maintainer, Elizabeth, A., & Cynthia, C. (2022). *survival: Survival Analysis* (3.4-0). <https://CRAN.R-project.org/package=survival>
- Wang, J., Thornton, J. C., Kolesnik, S., & Pierson, R. N. (2000). Anthropometry in body composition. An overview. *Annals of the New York Academy of Sciences*, 904, 317–326. <https://doi.org/10.1111/j.1749-6632.2000.tb06474.x>
- Wells, J. C. K. (2007). Sexual dimorphism of body composition. *Best Practice & Research Clinical Endocrinology & Metabolism*, 21(3), 415–430. <https://doi.org/10.1016/j.beem.2007.04.007>
- Wilkinson, D. J., Piasecki, M., & Atherton, P. J. (2018). The age-related loss of skeletal muscle mass and function: Measurement and physiology of muscle fibre atrophy and muscle fibre loss in humans. *Ageing Research Reviews*, 47, 123–132. <https://doi.org/10.1016/j.arr.2018.07.005>
- Wilson, D. L. (1994). The analysis of survival (mortality) data: Fitting Gompertz, Weibull, and logistic functions. *Mechanisms of Ageing and Development*, 74(1–2), 15–33. [https://doi.org/10.1016/0047-6374\(94\)90095-7](https://doi.org/10.1016/0047-6374(94)90095-7)
- Xia, X., Chen, X., Wu, G., Li, F., Wang, Y., Chen, Y., Chen, M., Wang, X., Chen, W., Xian, B., Chen, W., Cao, Y., Xu, C., Gong, W., Chen, G., Cai, D., Wei, W., Yan, Y., Liu, K., ... Han, J.-D. J. (2020). Three-dimensional facial-image analysis to predict heterogeneity of the human ageing rate and the impact of lifestyle. *Nature Metabolism*, 2(9), 946–957. <https://doi.org/10.1038/s42255-020-00270-x>

TABLES

TABLE 1. Performance of Fine & Gray semiparametric competitive risk regression models to evaluate cause-specific mortality using AnthroAge, S-AnthroAge and PhenoAge. We show the C-statistic (95% confidence interval) and the difference in the Bayesian Information Criterion ($\Delta\text{BIC} = \text{BIC}_{\text{Model-1}} - \text{BIC}_{\text{Model-2}}$) to contrast model predictions. $\Delta\text{BIC} < -2$ indicates that model 1 is better. $\Delta\text{BIC} > 2$ indicates that model 2 is better. All models are adjusted for age, sex, race/ethnicity, and number of comorbidities. Hazard ratios with 95% confidence intervals and standard errors for all models are available within supplementary material.

Cause-specific mortality	Events (n)	AnthropoAge C-statistic	S-AnthropoAge C-statistic	PhenoAge C-statistic	Δ BIC (AnthropoAge vs. S-AnthropoAge)	Δ BIC (AnthropoAge vs. PhenoAge)	Δ BIC (S-AnthropoAge vs. PhenoAge)
Cardiovascular	1197	0.847 (0.839-0.855)	0.845 (0.837-0.854)	0.851 (0.842-0.859)	-0.490	2.976	3.466
Diabetes Mellitus	167	0.845 (0.819-0.87)	0.843 (0.818-0.869)	0.876 (0.854-0.899)	0.580	77.499	76.918
Stroke	343	0.828 (0.808-0.847)	0.827 (0.808-0.847)	0.821 (0.801-0.841)	1.124	-51.242	-52.365
Cancer	1190	0.764 (0.752-0.775)	0.763 (0.751-0.775)	0.766 (0.754-0.778)	1.571	-0.704	-2.275
Influenza or Pneumonia	151	0.859 (0.833-0.885)	0.857 (0.83-0.883)	0.855 (0.827-0.883)	-4.712	-6.196	-1.485
Nephritis or Nephrosis	76	0.842 (0.805-0.878)	0.842 (0.805-0.879)	0.871 (0.838-0.904)	0.561	26.787	26.225
Alzheimer	144	0.878 (0.862-0.895)	0.878 (0.862-0.895)	0.859 (0.84-0.877)	2.270	-50.546	-52.815
Chronic lower respiratory	241	0.843 (0.826-0.861)	0.844 (0.826-0.862)	0.835 (0.816-0.853)	3.749	-34.446	-38.195

FIGURES

FIGURE 1. Flowchart diagram of participant selection from the National Health and Nutrition Examination Survey (NHANES) along with the key points from each analysis in our study. A brief conceptual framework is also provided at the top, illustrating how changes in body composition can modify aging rates, ultimately leading to accelerated aging and poor age-related outcomes, and this relationship is strongly influenced by sex and by changes in other domains of aging.

FIGURE 2. Anthropometric variables predict 10-year all-cause mortality in a non-linear and sex-dependent fashion. Variables are grouped according to the property of body composition that they capture: body length (a-c), body mass (d-e), visceral adiposity (f), subcutaneous adiposity (g-h), primarily lean mass (i-j). We used Gompertz models stratified by sex and adjusted by CA and number of comorbidities, with race/ethnicity included in the shape parameter.

FIGURE 3. AnthroAge and PhenoAge scatter plots show that these BA metrics have different dispersion patterns with CA in both the training (a-b) and validation cohorts (c-d). The horizontal distances between each point and regression lines represent divergence from CA and were used to estimate age acceleration metrics, which are presented as density plots stratified by sex (e-h). Only AnthroAgeAccel was calculated separately for each sex, as reflected by a more homogeneous distribution between men and women.

FIGURE 4. Areas under the receiving operating characteristic curves (AUROC) for prediction of 10-year all-cause mortality demonstrate that AnthroAge and S-AnthroAge (not shown) have a comparable performance to PhenoAge, and that it outperforms CA and

individual anthropometric measurements (WHtR, thigh circumference, triceps skinfold and BMI) (a-b). This prediction is equally good between men and women (c), but superior in subjects with 1 comorbidity (b). There were no significant differences between the three BA metrics in any category except for subjects with ≥ 2 comorbidities.


FIGURE 5. AnthroAgeAccel (a), S-AnthroAgeAccel (b) and PhenoAgeAccel (c) identify subjects with impaired survival linked to accelerated aging (values >0) visualized with Kaplan-Meier curves. Simultaneous consideration of AnthroAgeAccel and PhenoAgeAccel shows that acceleration in multiple domains of aging produces a higher mortality risk (d).

FIGURE 6. Spider plots stratified by sex show that subjects with accelerated AnthroAge (AnthroAgeAccel >0) display unique patterns of anthropometry, DXA-derived body composition and biochemical profiles (PhenoAge components). DXA variables were divided by squared height to obtain whole-body, trunk and appendicular fat mass and lean mass index (FMI and LMI, respectively), we also calculated fat-to-lean mass ratios and trunk-to-appendicular fat ratio. Variables were scaled and medians were compared using Mann-Whitney's U test (p-value <0.05 : *, <0.01 : **, <0.001 : ***). Spider plots assessing S-AnthroAgeAccel and multidomain aging are shown in supplementary figures 7 and 8.

Can we use anthropometric data to estimate biological age?

NHANES-III (n=23,907)
NHANES-IV (n=30,501)
n=54,408

≥20 years + complete mortality and anthropometric data



ANALYSIS #1

Training: NHANES-III (n=11,774)
Validation: NHANES-IV (n=7,020)
n= 18,794

Complete laboratory data to calculate PhenoAge

Development and validation of AnthroAge

- ✓ Non-linear association with 10-year all-cause mortality.
- ✓ Ethnic and sex-based differences are taken into account.
- ✓ Complete and simplified versions (only BMI and WHtR).

ANALYSIS #2

n= 17,540 (III: 10,823, IV: 6,627)

Complete body composition assessment by DXA

Multidomain aging (PhenoAge and AnthroAge)

- ✓ Comparison of predictions for cause-specific mortality.
- ✓ Risk stratification based on multidomain aging acceleration.

ANALYSIS #3

n= 6,284 (Only NHANES-IV)

Patterns linked to accelerated aging

- ✓ Subjects with accelerated vs non-accelerated aging.
- ✓ Body composition, laboratory data and comorbidity burden.



AnthroAge and S-AnthroAge can be estimated at: <https://bellolab.shinyapps.io/anthroage/>

