

## **Diagnostic Yield of Screening for SARS-CoV-2 among Patients Admitted for Alternate Diagnoses**

Phil Davis<sup>1</sup>, Rhonda J. Rosychuk<sup>2</sup>, Jeffrey P Hau<sup>3,14</sup>, Ivy Cheng<sup>4,5</sup>, Andrew D. McRae<sup>6</sup>, Raoul Daoust<sup>7</sup>,  
Eddy Lang<sup>8</sup>, Joel Turner<sup>9</sup>, Jaspreet Khangura<sup>10</sup>, Patrick T. Fok<sup>11</sup>, Maja Stachura<sup>12</sup>, Baljeet Brar<sup>13</sup>, and  
Corinne Hohl<sup>14</sup> **on behalf of the CCEDRRN investigators, and for the Network of Canadian  
Emergency Researchers and the Canadian Critical Care Trials Group**

Correspondence to: Dr. Philip J.B. Davis; 103 Hospital Drive, Saskatoon, SK, S7N 0W8;  
[Phil.davis@usask.ca](mailto:Phil.davis@usask.ca)

### Author affiliations:

<sup>1</sup> Department of Emergency Medicine, University of Saskatchewan, Saskatoon, SK, Canada

<sup>2</sup> Department of Pediatrics, University of Alberta, Edmonton, AB, Canada

<sup>3</sup> Centre for Clinical Epidemiology and Evaluation, Vancouver Coastal Health Research Institute, 828 W 10th Ave, Vancouver BC V5Z1M9

<sup>4</sup> Department of Emergency Medicine, Sunnybrook Health Sciences Center

<sup>5</sup> Division of Emergency Medicine, Dept of Medicine, University of Toronto

<sup>6</sup> Department of Emergency Medicine, University of Calgary

<sup>7</sup> Département Médecine de Famille et Médecine d'Urgence, Faculté de Médecine, Université de Montréal, Department of Emergency Medicine, Research Centre, CIUSSS-Nord-de-l'Île de-Montréal Montréal, Québec, Canada

<sup>8</sup> Department of Emergency Medicine, University of Calgary

<sup>9</sup> Department of Emergency Medicine, McGill University, Montreal, Qc

<sup>10</sup> Department of Emergency Medicine, University of British Columbia

<sup>11</sup> Division of EMS, Department of Emergency Medicine, Dalhousie University

<sup>12</sup> Department of Emergency Medicine, University of British Columbia

<sup>13</sup> Department of Emergency Medicine, University of British Columbia

<sup>14</sup> Department of Emergency Medicine, Faculty of Medicine, University of British Columbia, 2775 Laurel St., Vancouver BC V5Z 1M9.

Word count (excluding title page, abstract, references, figures, and tables): 2282

**Keywords:** screening, COVID-19, coronavirus disease, health services utilization, diagnostic testing, pandemic

**Abstract:**

*Objectives:* To determine the diagnostic yield of screening patients for SARS-CoV-2 who were admitted with a diagnosis unrelated to COVID-19, and identify risk factors for positive tests.

*Design:* Cohort from the Canadian COVID-19 Emergency Department Rapid Response Network (CCEDRRN) registry

*Setting:* 30 acute care hospitals across Canada

*Participants:* Patients hospitalized for non-COVID-19 related diagnoses who were tested for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) between March 1, and December 29, 2020

*Main outcome:* Positive nucleic acid amplification test (NAAT) for SARS-CoV-2

*Outcome measure:* Diagnostic yield

*Results:* We enrolled 15,690 consecutive eligible adults who were admitted to hospital without clinically suspected COVID-19. Among these patients, 122 tested positive for COVID-19, resulting in a diagnostic yield of 0.8% (95% CI 0.64% – 0.92%). Factors associated with a positive test included presence of a fever, being a healthcare worker, having a positive household contact or institutional exposure, and living in an area with higher 7-day average incident COVID-19 cases.

*Conclusions:* Universal screening of hospitalized patients for COVID-19 across two pandemic waves had a low diagnostic yield and should be informed by individual-level risk assessment in addition to regional COVID-19 prevalence.

Trial registration: NCT04702945

## **SUMMARY BOXES**

*Section 1:* Universal screening of admitted patients for SARS-CoV-2 was implemented in many hospitals at the beginning of the pandemic. The Infectious Diseases Society of America (IDSA) recommended avoiding universal screening of asymptomatic hospitalized patients in areas and times of low-COVID prevalence (defined as <2% prevalence) with very low certainty of evidence, based on studies of COVID-19 prevalence among asymptomatic individuals in the community.

*Section 2:* This study supports IDSA recommendations to avoid universal screening for COVID-19 in times and areas of low COVID prevalence and identifies patient-level risk factors strongly associated with positive testing that should be considered for screening.

## INTRODUCTION

Healthcare institutions initiated widespread testing of admitted patients for coronavirus disease 2019 (COVID-19) in the spring of 2020 (1). Patients without any reported symptoms of COVID-19 were routinely tested even in jurisdictions where COVID-19 rates were low in order to identify asymptomatic carriers and prevent hospital outbreaks (2). Some jurisdictions have continued this practice without robust evidence to support this practice. An Alberta study of 3,375 patients admitted to hospital for alternate diagnoses during the first wave of the pandemic when COVID-19 prevalence was very low found that none of the patients tested positive (3). In contrast, other studies from times and regions with higher COVID-19 prevalence reported positive tests in between 2.6 and 15.5% of otherwise asymptomatic patients (4–8).

Universal testing has several potential downsides if diagnostic yield is low. First, it may worsen Emergency Department (ED) crowding, as admitted patients with pending COVID-19 tests are boarded in EDs until their test results are reported. While ED volumes were lower than usual in the early pandemic such that EDs could absorb this delay, high patient volumes have since returned, exacerbating the impact of this practice on hospital crowding (9). This in turn increases patient morbidity and mortality (10). In addition, diagnostic workups and therapeutic interventions may be delayed until COVID-19 test results are back, as it takes longer to move patients on isolation precautions through the system. This can further exacerbate patient outcomes and hospital crowding. Thirdly, diagnostic testing capacity may be limited, potentially delaying processing of tests for symptomatic patients. In addition, the use of Personal Protective Equipment (PPE) may be increased as institutions have placed patients with pending COVID-19 tests under isolation precautions. The use of PPE during resuscitation has been associated with worse patient outcomes (11). Lastly, there is also an opportunity cost for hospitals as money spent on universal testing could be allocated to other areas. These unintended consequences of liberal testing policies need to be weighed carefully against the anticipated diagnostic yield and potential benefits.

As for any diagnostic test, rational COVID-19 testing guidelines should be informed by the level of risk of the patient, such that testing is reduced in situations when risk is low, and more widespread when risk is high. Based on expert opinion, the Infectious Diseases Society of America (IDSA) recommended a testing strategy based on the prevalence of the disease in the community (12). They recommended universal testing of asymptomatic hospitalized patients in times and places of high disease prevalence defined as  $\geq 10\%$  or  $\geq 10,000$  active cases per 100,000 population and did not recommend universal testing in times and places of low prevalence, defined as under 2% prevalence of disease, or less than 2,000 active cases per 100,000. Most jurisdictions never met the proposed screening threshold as public health measures were enacted to reduce disease prevalence to avoid overwhelming hospital capacity. The IDSA was unable to provide further guidance due to lack of available evidence. Our aim was to determine the diagnostic yield of screening patients for SARS-CoV-2 who had been admitted with a diagnosis unrelated to COVID-19 and identify risk factors for positive tests.

## **METHODS**

### *Study Design and Setting*

The Canadian COVID ED Rapid Response Network (CCEDRRN, pronounced ‘sedrin’) is a pan Canadian population-based registry that has enrolled consecutive eligible patients presenting with suspected or confirmed COVID-19 from EDs across Canada starting on March 1, 2020. The study population, data collection, data quality assurance, management and governance structure are described in the network’s methods paper (13). The research ethics boards of all participating institutions approved this study with a waiver of informed consent for data collection and linkage. Thirty CCEDRRN sites in 7 provinces contributed data to this study (Appendix A).

### *Patient and Public Involvement*

CCEDRRN has an active patient engagement committee with patient partners who have lived experience with COVID-19 from geographically representative areas of Canada. Patient partners provided input into the development of this research question and study protocol and the final manuscript.

### *Study Patients*

Participating sites needed to demonstrate  $\geq 99\%$  compliance in enrolling consecutive eligible patients for their data to be included in this study. Data from sites and periods that did not meet this quality threshold were excluded. We included consecutive eligible patients who were admitted to hospital and swabbed for SARS-CoV-2 using a nucleic acid amplification test (NAAT) within 24 hours of ED arrival. We enrolled patients between March 1, 2020 and December 29, 2020. To identify a population of admitted patients in whom COVID-19 disease was not suspected, we excluded patients with ED diagnoses that would have been clinically suspicious for COVID-19. These included all patients with ED diagnoses of suspected or confirmed COVID-19, influenza-like-illness (ILI), upper respiratory infections, and pneumonia or viral pneumonia for which testing would have been indicated based on clinical suspicion. We excluded patients who were discharged directly from the ED, diagnosed with COVID-19 before ED arrival (based on a NAAT done in the community), those whose first swab occurred more than 24 hours after their arrival, and repeat admissions. We also excluded patients in whom initial SARS-CoV-2 testing was negative and repeat testing became positive more than 5 days after arrival, as these patients could have contracted nosocomial COVID.

### *Data Collection*

Trained research assistants collected data retrospectively from electronic and/or paper-based medical records into a central, web-based REDCap database (Vanderbilt University; Nashville, TN, USA). Research assistants captured demographics, infection risk, ED vital signs, presenting symptoms, comorbid conditions and the results of COVID-19 tests. The coordinating centre implemented regular data quality checks, including logic checks in REDCap as well as site-level record verifications for nonsensical or outlying values.

In addition to these clinical variables, we calculated the seven-day moving average incident COVID-19 case count for the health region of each participating site using publicly available epidemiological data (14). For each calendar day within each health region represented in the study, we calculated the average daily incident rate of new infections per 100,000 population over the preceding seven days. This seven-day moving average incidence was assigned to each patient based on the date of their index emergency department encounter and the health region of their postal code of residence. We allocated patients with no fixed address to the health region of the hospital in which they were tested. We imputed values for the first five weeks of the pandemic by modeling the reported COVID-19 cases that had accumulated in every health region over time using linear interpolation (0.1% missing), COVID-19 case data were not publicly available for the early pandemic. The seven-day moving average incident COVID-19 case count was categorized as 0 – 1.99 per 100,000 population, 2 – 7.99 per 100,000 population, and  $\geq 8$  per 100,000 population based on the relationship between incidence and COVID-19 positive results in a previous analysis (15).

*Outcome:*

The primary outcome was a positive NAAT for SARS-CoV-2 in patients admitted with non-COVID-related diagnoses.

*Data Analysis:*

We divided the cohort into two groups, those without symptoms of COVID-19 and those with symptoms compatible with COVID-19 that were attributed to an alternate diagnosis (i.e., CHF, COPD, Asthma, etc.). We considered cough, dyspnea, fever, general weakness, chest pain, diarrhea, nausea and vomiting, headache, chills, myalgia, sore throat, altered level of consciousness, and dysgeusia/anosmia to be COVID-19-compatible symptoms. We used descriptive statistics to describe the population. We calculated the diagnostic yield by dividing the number of positive NAATs over the total number of NAATs performed. We calculated the exact binomial proportion 95% confidence intervals (95% CI) for

all proportions and used the modified Clopper-Pearson interval for small samples. We completed a planned subgroup analyses for patients presenting with and without COVID compatible symptoms to determine associated factors for a positive test. The initial multivariable logistic regression model to identify factors associated with a positive NAAT considered candidate variables with a p-value cut-off point of 0.20 based on the Wald test from univariable analyses. From the full model, a step-down procedure reduced the model to key predictors based on Akaike's information criterion (AIC) scores (e.g., chose the model with the smallest AIC score). Candidate variables included seven-day moving average incident COVID-19 case count category, patient age, gender, infection risk, and presenting symptoms. We limited the number of predictor variables in the model to one variable for every 10 outcomes in our data to avoid overfitting. Statistical analysis was performed using Stata (Version 16.1, StataCorp, College Station, Texas).

## RESULTS

We identified 19,791 patients admitted to hospital who presented to a participating ED between March 1, 2020, and December 29, 2020 (Figure 1). We excluded 4,101 patients, of which 2,769 had ED diagnoses that were clinically suspicious for COVID-19 and warranted SARS-CoV-2 testing on clinical grounds. The final cohort contained 15,690 patients. During the study period Canada experienced two pandemic waves with the local 7-day average incident case count ranging from between 0 and 42.6 cases per 100,000 population across sites.

We divided the cohort into two groups, those without any COVID-19 compatible symptoms, and those with COVID-19 compatible symptoms that were attributed to an alternate diagnosis in the ED (Table 1). Most patients arrived from home and were full code. The most common comorbidities were hypertension, diabetes and mental health illness. Of 3,113 patients admitted without COVID-19 compatible symptoms, 13 (0.4%, 95% CI 0.19% – 0.64%) tested positive for COVID-19. Of the 12,570 with COVID-19-

compatible symptoms, 109 patients (0.9%, 95% CI 0.70% – 1.03%) tested positive for COVID-19.

Among the 122 individuals who tested positive for COVID-19, 33 (27.0%, 95% CI 19.0% - 35.0%) were from a geographic region that had a moving average daily incident rate of  $\geq 8$  infections per 100,000 population. The diagnostic yield of testing among patients with COVID-19 compatible symptoms admitted for alternative diagnoses did not vary substantially by presenting symptom (Figure 2) or ED diagnosis (Figure 3).

When examining the association between patient factors and screening positive self-reported fever, being a healthcare worker, having a positive household contact or institutional exposure, and being from an area where the seven-day moving average incident COVID-19 case count was  $\geq 8$  per 100,000 population were associated with a greater risk of testing positive. The most important risk factor was reporting a household contact or being the caregiver of a known COVID-19 case.

## **DISCUSSION**

Our aim was to evaluate the diagnostic yield of screening non-COVID-19 admissions for SARS-CoV-2 across Canada in 2020 and identify patient-level risk factors for positive tests. The diagnostic yield of screening patients with non-COVID-19 related ED diagnoses who were admitted to hospital was low overall, and extremely low in patients without COVID compatible symptoms. The most important patient factors associated with a positive test were having a positive household contact, being a healthcare worker, or having had an institutional exposure to COVID-19. Those factors were more important than a high ( $\geq 8$  daily cases per 100,000 population) 7-day moving average incident COVID-19 case count.

Our study has several strengths. We used data from a large pan-Canadian registry that enrolls from large geographically and culturally diverse areas and is one of the largest registries in the world. CCEDRRN's patient enrolment and data verification protocols are rigorous, ensuring consecutive eligible patients and high-quality clinical data (13). We have previously demonstrated the inter-rater reliability for our data collection methods, including for symptoms (13).

Prior studies have examined the diagnostic yield of universal screening in single centers with varied diagnostic yield estimates between 0 and 15.5% (3–8). Many of these were case series with limited methods from the early pandemic. There is one known multi-center study which examines the benefit of universal screening for elective and emergent surgical admissions at 14 centers in the Netherlands (1). Like our study, the authors found that the overall COVID-19 NAAT positivity varied with community prevalence. Our finding that positive SARS-CoV-2 tests were associated with self-reported or measured fever is in keeping with a prior Cochrane systematic review that noted considerable variability in COVID-19 associated symptoms (17).

Our study is interesting in the context of the IDSA recommendations which were based on expert opinion and of “very low certainty” (12). The IDSA panel recommended avoiding universal screening for COVID-19 in times and areas of low COVID prevalence, defined as a disease prevalence of under 2% , or fewer than 2,000 active cases per 100,000 population, a threshold so high that it was never met at any of our study sites, even though multiple sites were in COVID-19 hotspots in 2020 (12). This threshold would have equated to over 6 million cases of active COVID-19 infection in the United States at any given time, which would have vastly overwhelmed hospital capacity, and thus represents an untenable threshold for hospitals. It is therefore not surprising that the prevalence of COVID-19 during the study period was far below the IDSA recommended threshold for initiating screening. While the number needed to screen to identify one positive case among admitted patients in our study was between 110 and 250 among unvaccinated patients, we propose that the IDSA screening threshold likely needs to be adopted.

A limitation of our study is that we only considered NAATs and did not consider the diagnostic yield of antigen-based COVID-19 tests, as they were not widespread in Canada in 2020 (16). We were unable to examine the sensitivity and specificity of the SARS-CoV-2 NAATs as we were unable to define false positive tests, so it is possible that some of the positive test results we encountered are false positives, leading to an overestimation of diagnostic yield. While our study is based on a Canadian population without international sites, we believe our findings are generalizable given their wide geographic spread,

and the cultural and racial diversity of our patient population. Finally, as data becomes available on the fourth wave of the pandemic, a future study should examine the impact of widespread vaccination on the yield of screening. As a larger proportion of the population is protected from severe disease and death through vaccination, decision makers should carefully consider the low diagnostic yield of a universal testing strategy going forward.

## **ACKNOWLEDGEMENTS**

We gratefully acknowledge the assistance of Mr. Rajan Bola in the preparation of this manuscript. And we thank the UBC clinical coordinating center staff, the UBC legal, ethics, privacy and contract staff and the research staff at each of the participating institutions in the network outlined in the attached Supplement (Supplement Tables 1-4). The network would not exist today without the dedication of these professionals.

Thank you to all of our patient partners who shared their lived experiences and perspectives to ensure that the knowledge we co-create addresses the concerns of patients and the public. Creating the largest network of collaboration across Canadian Emergency Departments would not have been feasible without the tireless efforts of Emergency Department Chiefs, and research coordinators and research assistants at participating sites. Finally, our most humble and sincere gratitude to our colleagues in medicine, nursing, and the allied health professions who have been on the front lines of this pandemic from day one staffing our ambulances, Emergency Departments, hospitals and ICUs bravely facing the risks of COVID-19 to look after our fellow citizens and after one another. We dedicate this network to you.

## **COMPETING INTERESTS**

None identified.

## **FUNDING**

The Canadian Institutes of Health Research (447679), Ontario Ministry of Colleges and Universities (C-655-2129), Saskatchewan Health Research Foundation (5357), Genome BC (COV024) Foundation du CHU de Québec (Octroi No. 4007) and the Public Health Agency of Canada provided peer-reviewed funding. The BC Academic Health Science Network and BioTalent Canada provided non-peer reviewed funding. These organizations are not-for-profit, and had no role in study conduct, analysis, or manuscript preparation.

## REFERENCES

1. Puylaert CAJ, Scheijmans JCG, Borgstein ABJ, Andeweg CS, Bartels-Rutten A, Beets GL, et al. Yield of Screening for COVID-19 in Asymptomatic Patients Before Elective or Emergency Surgery Using Chest CT and RT-PCR (SCOUT): Multicenter Study. *Ann Surg*. 2020 Dec;272(6):919–24.
2. Sutton D, Fuchs K, D’Alton M, Goffman D. Universal Screening for SARS-CoV-2 in Women Admitted for Delivery. *N Engl J Med*. 2020 May 28;382(22):2163–4.
3. Ravani P, Saxinger L, Chandran U, Fonseca K, Murphy S, Lang E, et al. COVID-19 screening of asymptomatic patients admitted through emergency departments in Alberta: a prospective quality-improvement study. *CMAJ Open*. 2020 Oct;8(4):E887–94.
4. Goldfarb IT, Diouf K, Barth WH, Robinson JN, Katz D, Gregory KE, et al. Universal SARS-CoV-2 testing on admission to the labor and delivery unit: Low prevalence among asymptomatic obstetric patients. *Infect Control Hosp Epidemiol*. 2020 Sep;41(9):1095–6.
5. Ossami Saily RR, Globke B, Pratschke J, Schoening W, Eurich D. Successful implementation of preventive measures leads to low relevance of SARS-CoV-2 in liver transplant patients: Observations from a German outpatient department. *Transpl Infect Dis [Internet]*. 2020 Dec [cited 2021 Aug 9];22(6). Available from: <https://onlinelibrary.wiley.com/doi/10.1111/tid.13363>
6. Gruskay JA, Dvorzhinskiy A, Konnaris MA, LeBrun DG, Ghahramani GC, Premkumar A, et al. Universal Testing for COVID-19 in Essential Orthopaedic Surgery Reveals a High Percentage of Asymptomatic Infections. *J Bone Jt Surg*. 2020 Aug 19;102(16):1379–88.
7. London V, McLaren R, Atallah F, Cepeda C, McCalla S, Fisher N, et al. The Relationship between Status at Presentation and Outcomes among Pregnant Women with COVID-19. *Am J Perinatol*. 2020 Aug;37(10):991–4.
8. Bianco A, Buckley AB, Overbey J, Smilen S, Wagner B, Dinglas C, et al. Testing of Patients and Support Persons for Coronavirus Disease 2019 (COVID-19) Infection Before Scheduled Deliveries. *Obstet Gynecol*. 2020 Aug;136(2):283–7.
9. Lee DD, Jung H, Lou W, Rauchwerger D, Chartier L, Masood S, et al. The Impact of COVID-19 on a Large, Canadian Community Emergency Department. *West J Emerg Med [Internet]*. 2021 Jun 9 [cited 2021 Jul 26];22(3). Available from: <https://escholarship.org/uc/item/0xk8j1fj>
10. Sun BC, Hsia RY, Weiss RE, Zingmond D, Liang L-J, Han W, et al. Effect of Emergency Department Crowding on Outcomes of Admitted Patients. *Ann Emerg Med*. 2013 Jun;61(6):605–611.e6.
11. Shao F, Xu S, Ma X, Xu Z, Lyu J, Ng M, et al. In-hospital cardiac arrest outcomes among patients with COVID-19 pneumonia in Wuhan, China. *Resuscitation*. 2020 Jun;151:18–23.
12. Hanson KE, Caliendo AM, Arias CA, Hayden MK, Englund JA, Lee MJ, et al. The Infectious Diseases Society of America Guidelines on the Diagnosis of COVID-19: Molecular Diagnostic Testing. *Clin Infect Dis*. 2021 Jan 22;ciab048.
13. Hohl CM, Rosychuk RJ, McRae AD, Brooks SC, Archambault P, Fok PT, et al. Development of the Canadian COVID-19 Emergency Department Rapid Response Network population-based registry: a methodology study. *CMAJ Open*. 2021 Jan;9(1):E261–70.

14. Health Regional Archive (Public View). [Internet]. [cited 2021 Jun 4]. Available from: <https://resourcescovid19canada.hub.arcgis.com/datasets/3aa9f7b1428642998fa399c57dad8045/data?layer=1>
15. McRae AD, Hohl CM, Rosychuk RJ, Vatanpour S, Ghaderi G, Archambault PM, et al. Development and validation of a clinical risk score to predict SARS-CoV-2 infection in emergency department patients: The CCEDRRN COVID-19 Infection Score (CCIS) [Internet]. *Emergency Medicine*; 2021 Jul [cited 2021 Aug 9]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2021.07.15.21260590>
16. Prince-Guerra JL, Almendares O, Nolen LD, Gunn JKL, Dale AP, Buono SA, et al. Evaluation of Abbott BinaxNOW Rapid Antigen Test for SARS-CoV-2 Infection at Two Community-Based Testing Sites — Pima County, Arizona, November 3–17, 2020. *2021*;70(3):6.
17. Struyf T, Deeks JJ, Dinnes J, Takwoingi Y, Davenport C, Leeflang MM, et al. Signs and symptoms to determine if a patient presenting in primary care or hospital outpatient settings has COVID-19 disease. Cochrane Infectious Diseases Group, editor. *Cochrane Database Syst Rev* [Internet]. 2020 Jul 7 [cited 2021 Jul 26]; Available from: <https://onlinelibrary.wiley.com/doi/10.1002/14651858.CD013665>

## SUPPLEMENT

**Supplement Table 1.** Network coordinating center staff at the University of British Columbia

<b>Name</b>	<b>Roles</b>	<b>Contributions</b>
Jeffrey Hau	Data manager	REDCap, data processing and analysis for manuscripts.
Vi Ho	National coordinator	Coordinate with provincial coordinators and training/onboarding of research assistants.
Serena Small	Research coordinator	Ethics & privacy reviews, data management plan, privacy impact assessment, and qualitative analyses
Amber Cragg	Research manager	Data and manuscript management
Wei Zhao	Data analyst	Data processing and analysis for manuscripts.
Vicky Wu	Data analyst	Data processing and analysis for manuscripts.
Elnaz Bodaghkhani	Research associate	Data and manuscript management

**Supplement Table 2.** Provincial Coordinators

<b>Name</b>	<b>Province</b>	<b>Institutional affiliation</b>	<b>Contributions to CCEDRRN</b>
Corinne DeMone	NS	Dalhousie University, Halifax, Nova Scotia	Research ethics board submission, manages research assistants, data cleaning and quality.
Jacqueline Fraser	NB	Dalhousie University, St. John New Brunswick	Site coordinator as well as research assistant.
Martyne Audet	QC	Centre intégré de santé et de services sociaux de Chaudière-Appalaches (Hôtel-Dieu de Lévis site), Lévis	Provincial research coordinator, translation of research material to French, ethics management.
Connie Taylor	ON	Queen's University, Kingston	Coordination of research assistants in Ontario, maintenance of REB applications for the province
Kate Mackenzie	MB	Health Sciences Centre, Winnipeg	Lead RA for the province
Aimee Goss	SK	University of Saskatchewan, Saskatoon	Screens records in Saskatoon, data/extraction and entry, coordinates research assistants.
Hina Walia	AB	University of Calgary, Calgary	Provincial coordinator lead for Alberta, oversight of all Alberta sites.
Josie Kanu	BC	University of British Columbia, Vancouver	Provincial coordinator lead for BC, oversight of all BC sites.

**Supplement Table 3.** Institutional research assistant (RA) leads


<b>Name</b>	<b>Province</b>	<b>Institutional affiliation(s)</b>
Corinne DeMone	NS	Dartmouth General Hospital, Cobequid Community Health Centre, Hants Community Hospital Secondary Assessment Centers of the Dartmouth General Hospital, and Halifax Infirmary, Halifax
Jacqueline Fraser	NB	Saint John Regional Hospital, Saint John
Alexandra Nadeau	QC	CHU de Québec Université Laval, Quebec City
Audrey Nolet	QC	Centre intégré de santé et de services sociaux de Chaudière-Appalaches (Hôtel-Dieu de Lévis site), Lévis
Xiaoqing Xue	QC	Jewish General Hospital, Montréal
David Iannuzzi	QC	McGill University Health Center, Montréal
Chantal Lanthier	QC	Hôpital du Sacré-Cœur de Montréal, Montréal
Konika Nirmalanathan	ON	University Health Network, Toronto
Vlad Latiu	ON	Kingston General Hospital, Hotel Dieu Hospital, Kingston
Joanna Yeung	ON	Sunnybrook Health Sciences Center, Toronto
Natasha Clayton	ON	Hamilton General Hospital, Juravinski Hospital, Hamilton
Tom Chen	ON	London Health Sciences Centre, London
Jenna Nichols	ON	Health Sciences North, Sudbury
Kate Mackenzie	MB	Health Sciences Centre, Winnipeg
Aimee Goss	SK	St. Paul's Hospital, Royal University Hospital, Saskatoon City Hospital, Saskatoon
Stacy Ruddell	AB	Foothills Medical Centre, Peter Lougheed Centre, Rockyview General Hospital, South Health Campus, Calgary
Natalie Runham	AB	University of Alberta Hospital, Edmonton
Karlin Su	AB	Royal Alexandra Hospital/Northeast Community Health Center, Edmonton
	BC	St. Paul's Hospital, Mount Saint Joseph, Vancouver
Bernice Huynh	BC	Abbotsford Regional Hospital and Cancer Center, Abbotsford
Amanda Swirhun	BC	Royal Columbian Hospital, New Westminster
Tracy Taylor	BC	Eagle Ridge Hospital and Health Care Centre, Port Moody
Mai Hayashi	BC	Royal Inland Hospital, Kamloops
Mackenzie Cheyne	BC	Kelowna General Hospital, Kelowna
Sarim Asim	BC	Surrey Memorial Hospital, Surrey
Katherine Lam	BC	Vancouver General Hospital, Vancouver
Kelsey Compagna	BC	Lions Gate Hospital, Vancouver

**Supplement Table 4.** Contributing Study Sites and Investigators

<b>Lead Investigator</b>	<b>Contributing Site / Code</b>	<b>Member Investigators</b>
<b>Maritime</b>		
Patrick Fok		
<b>Nova Scotia</b>		
Hana Wiemer	Halifax Infirmary/ 902	Patrick Fok
	Dartmouth General Hospital/ 903	Hana Wiemer
	Hants Community Hospital/ 904	Samuel Campbell
	Cobequid Community Health Centre/ 905	Kory Arsenaault
	Secondary Assessment Centers of Dartmouth General and Halifax Infirmary/ 908	Tara Dahn
<b>New Brunswick</b>		
Kavish Chandra	Saint John Regional Hospital/ 901	Kavish Chandra
<b>Quebec</b>		
Patrick Archambault	Hotel-Dieu de Lévis/ 701	Patrick Archambault
	Jewish General Hospital/ 702	Joel Turner
	Centre Hospitalier de l'Université Laval (CHU de Québec)/ 703	Éric Mercier
	L'hôpital Royal Victoria - Royal Victoria Hospital/ 705	Greg Clark
	Hôpital de l'Enfant-Jésus, CHU de Québec/ 706	Éric Mercier
	Hôpital du Saint-Sacrement, CHU de Québec/ 707	Éric Mercier
	Hôpital Saint-François d'Assise, CHU de Québec/ 708	Éric Mercier
	Hôtel-Dieu de Québec, CHU de Québec/ 709	Éric Mercier
	IUCPQ: Institut universitaire de cardiologie et de pneumologie de Québec/ 710	Sébastien Robert
	Hôpital du Sacré-Coeur de Montreal/ 711	Raoul Daoust
<b>Ontario</b>		
Laurie Morrison & Steven Brooks	Sunnybrook/ 401	Ivy Cheng
	The Ottawa Hospital - Civic Campus/ 403	Jeffrey Perry
	The Ottawa Hospital - General Campus/ 404	Jeffrey Perry
	Kingston/Queens/ 406	Steven Brooks
	Hamilton General Hospital/ 407	Michelle Welsford
	Health Science North, Sudbury Ontario/ 408	Rob Ohle
	University Hospital – LHSC/ 409	Justin Yan
	North York General Hospital, Toronto/ 410	Rohit Mohindra
	Victoria Hospital – LHSC/ 412	Justin Yan
Toronto Western Hospital/ 414	Megan Landes	
<b>Manitoba</b>		
Tomislav Jelic	Health Sciences Centre/ 307	Tomislav Jelic
<b>Saskatchewan</b>		
Phil Davis	St Paul's Hospital, Saskatoon/ 303	Phil Davis
	Royal University, Saskatoon/ 304	Phil Davis

	Saskatoon City Hospital, Saskatoon/ 305	Phil Davis
<b>Alberta</b>		
Andrew McRae	University of Alberta Hospital, Edmonton/ 201	Brian Rowe
	Foothills, Calgary/ 202	Katie Lin
	Rockyview, Calgary/ 203	Andrew McRae
	Peter Lougheed Centre/ 204	Andrew McRae
	South Campus, Calgary/ 205	Stephanie VandenBerg
	Northeast Community Health Centre, Edmonton/ 206	Jake Hayward, Jaspreet Khangura
	Royal Alexandra Hospital, Edmonton/ 306	Jake Hayward, Jaspreet Khangura
<b>British Columbia</b>		
Corinne Hohl	Vancouver General Hospital/ 101	Daniel Ting
	Lions Gate Hospital/ 102	Maja Stachura
	Saint Paul's Hospital/ 103	Frank Scheuermeyer
	Mount St Joseph's/ 104	Frank Scheuermeyer
	Surrey Memorial Hospital/ 105	Balijeet Braar
	Royal Columbian Hospital/ 106	John Taylor
	Abbotsford Regional Hospital/ 107	Ian Martin
	Eagle Ridge Hospital/ 108	Sean Wormsbecker
	Royal Inland Hospital/ 112	Ian Martin
Kelowna General / Hospital/ 115	Lee Graham	

**Figure 1: Patient Flow Diagram**




**Table 1:** Baseline Characteristics of admitted patients without clinical suspicion of COVID-19 (N=15,690)

	Patients without COVID-19 compatible symptoms  (N=3,113)	Patients with COVID-19 compatible symptoms attributed to an alternate diagnosis  (N=12,570)
<b>Demographics</b>		
Age (mean, SD)	57.6 (22.6)	64.6 (20.4)
Female (%)	1,418 (45.6)	5,924 (47.1)
<b>Arrival From (%)</b>		
Home	2,552 (82.0)	10,943 (87.0)
Long-term care or rehab facility	217 (7.0)	832 (6.6)
Unstable housing*	190 (6.1)	414 (3.3)
Corrections	7 (0.2)	14 (0.1)
Interfacility transfer	121 (3.9)	262 (2.1)
<b>Risk for Infection (%)</b>		
Travel	32 (1.0)	134 (1.1)
Institutional (LTC/prison)	231 (7.4)	721 (5.7)
Household contact	28 (0.9)	144 (1.1)
Occupational	10 (0.3)	38 (0.3)
Unknown	1,502 (48.2)	5,377 (42.8)
<b>Pre-ED Goals of Care (%)</b>		
Full code	2,946 (94.6)	11,259 (89.5)
Intermediate GOC	18 (0.6)	173 (1.4)
Do not resuscitate	149 (4.8)	1,142 (9.1)
<b>Acuity</b>		
Arrival by Ambulance (%)	1,724 (55.4)	7,189 (57.2)
CTAS 1 (Resuscitation)	241 (7.7)	1,053 (8.4)
CTAS 2 (Emergent)	1,000 (32.1)	5,786 (46.0)
CTAS 3 (Urgent)	1,527 (49.1)	5,086 (40.4)
CTAS 4 (Less Urgent)	295 (9.5)	572 (4.6)
CTAS 5 (Non Urgent)	40 (1.3)	59 (0.5)
<b>Arrival Vital Signs, Mean (SD)</b>		
Heart Rate, beats per min	91.2 (21.2)	95.5 (23.9)
Systolic BP, mm Hg	134.7 (25.1)	133.6 (27.9)
Oxygen saturation, (%)	96.6 (3.4)	95.7 (4.1)
Respiratory Rate, beats per min	18.6 (4.4)	21.2 (6.3)
Temperature, degrees Celsius	36.6 (0.6)	36.8 (0.9)
<b>Comorbidities (%)</b>		
Hypertension	951 (30.6)	5,321 (42.3)
Psychiatric Condition	728 (23.4)	2,134 (17.0)
Dyslipidemia	425 (13.6)	2,434 (19.4)
Diabetes	427 (13.7)	2,577 (20.5)
Chronic Neuro Disorder	322 (10.3)	1,406 (11.2)
Coronary Artery Disease	284 (9.1)	1,796 (14.3)
Rheumatologic Disorder	229 (7.4)	1,249 (9.9)


Dementia	199 (6.4)	696 (5.5)
Active Cancer	231 (7.4)	1,647 (12.9)
Chronic Kidney Disease	195 (6.3)	1,319 (10.5)
Chronic Lung Disease (not asthma)	199 (6.4)	1,691 (13.5)
Congestive Heart Failure	159 (5.1)	1,392 (11.1)
Asthma	125 (4.0)	712 (5.7)
Obesity	57 (1.8)	344 (2.7)
<b>Symptoms (%)</b>		
Cough	-	2,763 (22.0)
Dyspnea	-	4,757 (37.8)
Fever	-	2,531 (20.1)
General Weakness	-	3,183 (25.3)
Chest Pain	-	2,714 (21.6)
Diarrhea	-	1,339 (10.7)
Nausea/Vomiting	-	3,345 (26.6)
Headache	-	784 (6.2)
Chills	-	957 (7.6)
Myalgia	-	466 (3.7)
Sore Throat	-	374 (3.0)
Altered Consciousness	-	2,502 (19.9)
Dysgusea/Anosmia	-	41 (0.3)
<b>Social Factors (%)</b>		
Pregnant (%)	18 (0.6)	45 (0.4)
Tobacco use (%)	491 (15.8)	1,656 (13.2)
Illicit substance use (%)	421 (13.5)	967 (7.7)
<b>ED Diagnosis (%)</b>		
Respiratory Disease, not specified	8 (0.3)	118 (0.9)
COPD Exacerbation	11 (0.4)	648 (5.2)
Asthma Exacerbation	<5	97 (0.8)
Congestive Heart Failure	44 (1.4)	1,003 (8.0)
Shortness of Breath, NYD*	-	466 (3.6)
Cough, NYD*	-	63 (0.5)
Fever, NYD*	-	482 (3.8)
<b>Outcome (%)</b>		
Positive SARS-CoV-2 NAAT	13 (0.4)	109 (0.9)

\*NYD denotes "not yet determined"

**Figure 2:** Diagnostic Yield by Presenting Symptoms with 95% confidence intervals



**Figure 3:** Diagnostic Yield by ED Diagnosis with 95% confidence intervals.



\*NYD denotes "not yet determined"

Table 2: Multivariable analysis of factors associated with positive SARS-CoV-2 NAATs (N=15,690)

	Univariate analysis odds ratio (95% CI)	Final model with fully adjusted odds ratio (95% CI) <sup>1</sup>	P-value
<b>Sex</b>			
Male	<i>Reference</i>	<i>Reference</i>	0.18
Female	0.84 (0.59 – 1.21)	0.78 (0.54 – 1.12)	
<b>Age</b>			
	1.00 (1.00 – 1.02)	1.00 (0.99 – 1.01)	0.27
<b>7-day average incident COVID-19 cases</b>			
0 – 1.99 daily cases per 100,000 population	<i>Reference</i>	<i>Reference</i>	< 0.001
2 to 7.99 daily cases per 100,000 population	1.42 (0.91 – 2.22)	1.47 (0.94 – 2.31)	
≥8 daily cases per 100,000 population	2.99 (1.95 – 4.59)	3.17 (2.05 – 4.89)	
<b>COVID-19 compatible symptoms present</b>			
No	<i>Reference</i>	<i>Reference</i>	0.08
Yes	2.08 (1.71 – 3.71)	1.65 (0.90 – 3.00)	
<b>Self-reported fever, or temperature ≥ 37.5 °C</b>			
No	<i>Reference</i>	<i>Reference</i>	< 0.001
Yes	2.72 (1.89 – 3.90)	2.53 (1.74 – 3.67)	
<b>Diarrhea present</b>			
No	<i>Reference</i>	<i>Reference</i>	0.11
Yes	1.74 (1.04 – 2.92)	1.57 (0.93 – 2.67)	
<b>Healthcare worker</b>			
No	<i>Reference</i>	<i>Reference</i>	< 0.001
Yes	5.62 (1.35 – 23.43)	4.67 (1.05 – 20.54)	
<b>Household contact or caregiver</b>			
No	<i>Reference</i>	<i>Reference</i>	< 0.001
Yes	9.48 (5.01 – 17.96)	7.74 (3.98 – 15.04)	
<b>Institutional exposure</b>			
No	<i>Reference</i>	<i>Reference</i>	< 0.001
Yes	3.46 (2.17 – 5.52)	3.39 (2.10 – 5.47)	
<b>Dysgeusia or anosmia present</b>			
No	<i>Reference</i>	-	
Yes	3.21 (0.43 – 23.52)	-	
<b>Dyspnea present</b>			
No	<i>Reference</i>	-	
Yes	1.16 (0.80 – 1.70)	-	
<b>Nausea or vomiting present</b>			
No	<i>Reference</i>	-	

Yes

0.81 (0.51 – 1.29) -

---

<sup>1</sup> Final model determined by including variables with a p-value of  $p < 0.20$  during the sex and age adjusted analysis and using the Akaike Information Criterion (AIC) to determine additional variables to exclude from the final model. Variables adjusted for all other variables present in the final model

Appendix A: List of hospital sites, with an inclusion start and end date for this study.

Site Name	Province	Start Date	End Date
Vancouver General Hospital	British Columbia	1-Mar-2020	31-Aug-2020
Lions Gate Hospital	British Columbia	1-Mar-2020	29-Apr-2020
Saint Paul's Hospital	British Columbia	1-Mar-2020	23-May-2020
Mount Saint Joseph Hospital	British Columbia	1-Mar-2020	24-Mar-2020
Surrey Memorial Hospital	British Columbia	19-Mar-2020	30-Apr-2020
Royal Columbian Hospital	British Columbia	1-Mar-2020	31-May-2020
Abbotsford Regional Hospital	British Columbia	20-Apr-2020	15-Jul-2020
University of Alberta Hospital	Alberta	8-Apr-2020	7-May-2020
Foothills Medical Centre	Alberta	1-Mar-2020	7-Apr-2020
Rockyview General Hospital	Alberta	1-Mar-2020	7-Apr-2020
Peter Lougheed Centre	Alberta	1-Mar-2020	12-Dec-2020
South Health Campus	Alberta	1-Mar-2020	12-Dec-2020
St Paul's Hospital	Saskatchewan	17-Mar-2020	30-Apr-2020
Royal University Hospital	Saskatchewan	17-Mar-2020	31-Oct-2020
Saskatoon City Hospital	Saskatchewan	17-Mar-2020	30-Apr-2020
Sunnybrook Health Sciences Centre	Ontario	14-May-2020	31-Oct-2020
The Ottawa Hospital - Civic Campus	Ontario	14-May-2020	31-May-2020
Health Science North	Ontario	14-May-2020	29-Dec-2020
Toronto Western Hospital	Ontario	1-Sep-2020	31-Sep-2020
Hotel-Dieu de Lévis	Quebec	4-May-2020	18-May-2020
Jewish General Hospital	Quebec	1-Mar-2020	4-Jun-2020
Hôpital de l'Enfant-Jésus, CHU de Québec	Quebec	4-May-2020	23-Jul-2020
IUCPQ: Institut universitaire de cardiologie et de pneumologie de Québec	Quebec	4-May-2020	13-May-2020
Hôpital du Sacré-Coeur de Montreal	Quebec	4-May-2020	18-May-2020
Saint John Regional Hospital	New Brunswick	12-Mar-2020	12-Apr-2020
Halifax Infirmary	Nova Scotia	5-Apr-2020	15-Apr-2020
Dartmouth General Hospital	Nova Scotia	5-Apr-2020	15-Apr-2020
Hants Community Hospital	Nova Scotia	5-Apr-2020	15-Apr-2020
Cobequid Community Health Centre	Nova Scotia	5-Apr-2020	15-Apr-2020
Secondary Assessment Centers of Dartmouth General and Halifax Infirmary	Nova Scotia	26-Mar-2020	15-May-2020