

1 The impact of SARS-CoV-2 vaccination on Alpha & Delta variant 2 transmission

3

4 David W Eyre^{1,2}, Donald Taylor³, Mark Purver³, David Chapman⁴, Tom Fowler^{3,5}, Koen B
5 Pouwels^{2,6}, A Sarah Walker^{2,7}, Tim EA Peto^{2,7}

6

7 ¹Big Data Institute, Nuffield Department of Population Health, University of Oxford, Oxford,
8 UK

9 ²NIHR Health Protection Research Unit in in Healthcare Associated Infections and
10 Antimicrobial Resistance, University of Oxford, Oxford, UK

11 ³Department of Health and Social Care, UK Government, London, UK

12 ⁴Deloitte MCS Ltd, London, UK

13 ⁵William Harvey Research Institute, Queen Mary University of London, London UK

14 ⁶Health Economics Research Centre, Nuffield Department of Population Health, University
15 of Oxford, Oxford, UK

16 ⁷Nuffield Department of Medicine, University of Oxford

17

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

18 Abstract

19

20 Background

21 Pre-Delta, vaccination reduced transmission of SARS-CoV-2 from individuals infected despite
22 vaccination, potentially via reducing viral loads. While vaccination still lowers the risk of
23 infection, similar viral loads in vaccinated and unvaccinated individuals infected with Delta
24 question how much vaccination prevents onward transmission.

25

26 Methods

27 We performed a retrospective observational cohort study of contacts of SARS-CoV-2-
28 infected index cases using contact testing data from England. We used multivariable logistic
29 regression to investigate the impact of index case and contact vaccination on transmission,
30 and how this varies with Alpha and Delta variants (classified using S-gene
31 detection/calendar trends) and time since second vaccination.

32

33 Results

34 51,798/139,164(37.2%) contacts tested were PCR-positive. Two doses of BNT162b2 or
35 ChAdOx1 vaccines in Alpha variant index cases independently reduced PCR-positivity in
36 contacts (aOR, adjusted odds ratio vs. unvaccinated=0.18[95%CI 0.12-0.29] and 0.37[0.22-
37 0.63] respectively). The Delta variant attenuated vaccine-associated reductions in
38 transmission: two BNT162b2 doses reduced Delta transmission (aOR=0.35[0.26-0.48]), more
39 than ChAdOx1 (aOR=0.64[0.57-0.72]; heterogeneity $p < 0.001$). Variation in viral load (Ct
40 values) explained only a modest proportion of vaccine-associated transmission reductions.

41 Transmission reductions declined over time since second vaccination, for Delta reaching
42 similar levels to unvaccinated individuals by 12 weeks for ChAdOx1 and attenuating
43 substantially for BNT162b2. Protection from vaccination in contacts also declined in the 3
44 months after second vaccination.

45

46 **Conclusions**

47 Vaccination reduces transmission of Delta, but by less than the Alpha variant. The impact of
48 vaccination decreased over time. Factors other than PCR-measured viral load are important
49 in vaccine-associated transmission reductions. Booster vaccinations may help control
50 transmission together with preventing infections.

51

52 Introduction

53 SARS-CoV-2 vaccines have been shown in randomised controlled trials¹⁻³ and real-world
54 population studies^{4,5} to prevent infection and adverse outcomes from several SARS-CoV-2
55 variants including Alpha (B.1.1.7) and Delta (B.1.617.2).⁶⁻⁸

56

57 Vaccination also potentially prevents onward transmission by at least two mechanisms.

58 Firstly, by reducing symptomatic and asymptomatic infections and therefore the number of
59 infectious individuals, and secondly via reduced onward spread from those who become
60 infected despite vaccination. Household studies have demonstrated vaccination reduces
61 onward transmission of the Alpha variant from those infected despite vaccination in the
62 UK,⁹ Israel^{10,11} and Finland.¹² One hypothesised mechanism is lower viral loads observed in
63 post-vaccination Alpha infections^{7,13} compared in unvaccinated individuals, as viral load is
64 associated with the likelihood of infection in contacts.^{14,15}

65

66 However, viral loads in Delta variant infections occurring after vaccination are similar in
67 vaccinated and unvaccinated individuals,^{8,16} although the duration of viral shedding may be
68 reduced.^{17,18} This questions whether vaccination can control Delta spread as effectively as
69 Alpha, and whether, with increased transmissibility,¹⁹ this explains the rapid global
70 dissemination of Delta despite rising vaccination coverage.

71

72 We use national contact testing data from England to investigate the impact of vaccination
73 on onward transmission of SARS-CoV-2, and how this varies with Alpha and Delta variants

74 and time since second vaccination. We also investigate how much reductions in
75 transmission after vaccination are explained by variation in PCR cycle threshold (Ct) values.

76

77 [Methods](#)

78 [Setting and variants](#)

79 We performed a retrospective observational cohort study of contacts of symptomatic and
80 asymptomatic SARS-CoV-2-infected index cases. Data were obtained from the national
81 contact tracing and testing service in England, NHS Test and Trace. Contacts were eligible for
82 inclusion if they accessed NHS Test and Trace PCR testing 1-10 days after the index case's
83 PCR test (typically following symptoms, but also after positive asymptomatic antigen
84 screening tests), i.e. including contact pairs where the index case was the most likely source
85 for any infection in the contact.¹⁵ Only index cases with PCR tests performed by three
86 national "lighthouse" laboratories (Milton Keynes, Alderley Park, Glasgow) were included, as
87 these tests used the same standardised workflow and PCR assay (Thermo Fisher TaqPath,
88 assessing for S gene, N gene and ORF1ab targets). Contacts could be tested by any
89 community/hospital laboratory reporting results to NHS Test and Trace. Vaccination status
90 in cases and contacts was obtained from National Immunisation Management Service (see
91 Supplement).

92

93 Contacts of index cases tested between 01-January-2021 and 31-July-2021 were included as
94 follows. Index cases were classified as the Alpha (B.1.1.7) variant based on S-gene target
95 failure (SGTF), while this was considered a reliable proxy for Alpha, namely to 06-June-2021
96 (after which <5% cases had SGTF). From 10-May-2021 national spread of Delta meant that

97 >98% of sequenced cases were either due to the Alpha or Delta variants,¹⁹ such that we
98 used detection of S-gene on or after 10-May-2021 as a proxy for Delta (see Supplement).

99

100 We restricted our analysis to contacts undergoing testing, excluding untested contacts, to
101 control as much as possible for biases related to health-seeking behaviour (including
102 differences before and after vaccination), access to testing, and case ascertainment.²⁰

103

104 [Statistical analysis](#)

105 We used multivariable logistic regression to investigate how onward transmission, i.e.,
106 SARS-CoV-2 PCR-positive tests in contacts, varied with index case vaccination status. Index
107 case vaccination status was defined using administrative classifications as: unvaccinated,
108 partially vaccinated (from day of first vaccine to 13 days after second vaccine), or fully
109 vaccinated (≥ 14 days after second vaccine), further considering whether vaccination was
110 AstraZeneca ChAdOx1 or Pfizer-BioNTech BNT162b2. We also investigated how onward
111 transmission varied with Alpha vs. Delta index cases and whether any effects varied by
112 vaccine by including pre-specified interaction terms. We additionally included a model term
113 for time since 14 days after second BNT162b2 or ChAdOx1 vaccine to estimate the effect of
114 time since second vaccine.

115

116 We adjusted for the following additional covariates: contact event type; index case factors -
117 age, sex, and symptom status; contact factors - age, sex, vaccination status and time since
118 vaccination (as above); local deprivation, local SARS-CoV-2 incidence, and calendar time (to
119 capture changes in behaviour/social distancing, the likelihood of acquisition from a third
120 party, population-wide vaccine uptake, and the percentage of unvaccinated people

121 previously infected) (Table S1). We used natural cubic splines and log transformation to
122 account for non-linearity and tested for interactions (see Supplement).

123

124 We refitted models including index case Ct values to investigate whether the effect of index
125 case vaccination status was explained by viral load (approximated Ct value²¹).

126

127 Ethics

128 The study was performed as public health surveillance and NHS Test and Trace program
129 quality assurance, under Section 251 of the NHS Act 2006 with approvals from Public Health
130 England (PHE), the Department of Health and Social Care and NHS Test and Trace. PHE's
131 Research Ethics and Governance Group (PHE's Research Ethics Committee) reviewed the
132 study protocol and confirmed compliance with all regulatory requirements. As no regulatory
133 or ethical issues were identified, it was agreed that full ethical review was not needed, and
134 the protocol was approved.

135

136 Results

137 151,821 contacts of 99,597 index cases underwent PCR testing between 02-January-2021
138 and 02-August-2021. 12,657 contacts were excluded with incomplete data (8.3%, see
139 Supplement). Of the remaining 139,164 contacts (95,716 index cases), 51,798(37.2%) tested
140 PCR-positive. The median(IQR)[range] index case and contact ages were 38(26-50)[0-102]
141 and 38(23-50)[0-104] years respectively. 50,356(53%) index cases and 77,277(56%) contacts
142 were female (see Table S1-S2 for details by case and contact vaccine status). Contact events

143 were predominantly within households (97,387;70%), but also in household visitors
144 (14,066;10%), at events and activities (14,270;10%) and at work/education (13,441;10%).

145

146 [Index case vaccination and onward transmission](#)

147 27,666/55,977(49%) contacts of unvaccinated index cases tested PCR-positive, as did
148 5,256/14,398(37%) and 9,623/36,085(27%) contacts of partially ChAdOx1 and BNT162b2
149 vaccinated cases, and 7,559/25,422(30%) and 1,694/7,282(23%) contacts of fully ChAdOx1
150 and BNT162b2 vaccinated cases. For index cases vaccinated twice with BNT162b2 or
151 ChAdOx1 the median(IQR) days from second vaccine to an Alpha variant PCR-positive test
152 was 41(26-62) or 27(18-43) respectively and 89(69-110) and 50(34-69) for Delta.

153

154 In a multivariable model (Tables 1, S4), BNT162b2 vaccination in Alpha variant index cases
155 independently reduced PCR-positive results in contacts, with two doses (aOR, adjusted odds
156 ratio at 14 days post-second vaccine vs. unvaccinated=0.18[95%CI 0.12-0.29]) reducing
157 onward transmission more than one (aOR=0.74[0.70-0.80]). There was weak evidence that
158 ChAdOx1 was less effective than BNT162b2 at preventing transmission after one (aOR vs.
159 unvaccinated=0.82[0.76-0.88], p=0.11 vs BNT162b2) and two doses (aOR=0.37[0.22-0.63],
160 p=0.085 vs BNT162b2).

161

162 The Delta variant was associated with increased onward transmission vs. Alpha for
163 symptomatic index cases (aOR=1.30[1.10-1.54]) and to a greater extent for asymptomatic
164 index cases (aOR=2.14[1.75-2.60]). Post-second dose vaccine-associated reductions on
165 onward transmission were also attenuated with Delta, for both BNT162b2 by 1.9-fold
166 (aOR=1.93[1.25-2.99]) and ChAdOx1 by 1.7-fold (aOR=1.71[1.01-2.90]). This resulted in two

167 BNT162b2 doses reducing onward transmission of Delta by a greater extent than ChAdOx1
168 (aOR=0.35[0.26-0.48] vs aOR=0.64[0.57-0.72], respectively, heterogeneity $p<0.001$).

169

170 [Vaccination in contacts](#)

171 The estimated effect of contact vaccination status does not reflect overall vaccine
172 effectiveness, as study inclusion was conditional on the contact being tested (typically after
173 developing symptoms). However, as expected, PCR-positivity was highest in unvaccinated
174 contacts (34,137/70,370(49%)), followed by those partially vaccinated with ChAdOx1,
175 (3,401/10,325(33%)) and BNT162b2 (5,809/18,017(32%)); those fully vaccinated with
176 ChAdOx1 (6,163/27,343(23%)) and BNT162b2 (2,288/13,109(17%)) had the lowest rates. In
177 a multivariable model (Tables 1, S4), with Alpha contacts fully vaccinated with BNT162b2
178 had lower rates of PCR-positive tests than ChAdOx1 (aOR at 14 days post second vaccine vs.
179 unvaccinated=0.06[0.04-0.10] vs. aOR=0.29[0.17-0.49] respectively, heterogeneity $p<0.001$).
180 With Delta, more BNT162b2 vaccinated contacts tested PCR-positive than with Alpha, but
181 there was no evidence that PCR-positivity changed in fully ChAdOx1 vaccinated contacts.
182 Nevertheless, two doses of BNT162b2 remained more effective against Delta than ChAdOx1
183 (aOR vs. unvaccinated=0.10[0.08-0.13] vs aOR=0.28[0.25-0.32], respectively, heterogeneity
184 $p<0.001$).

185

186 [Duration of protection and transmission reductions](#)

187 Vaccine-associated reductions in onward transmission declined over time since second
188 vaccination in index cases (Figure 1A, Table S4). Independently of contact vaccination status,
189 for each doubling of weeks since 14 days after second vaccination in index cases, the odds
190 of a contact testing PCR-positive increased 1.13-fold (95%CI 1.09-1.17) for ChAdOx1 and

191 1.20-fold (1.10-1.31) for BNT162b2 with no evidence of a difference between vaccines
192 ($p=0.19$). There was no evidence that fitting different rates by variant improved model fit.
193 However, higher probabilities of PCR-positive results in contacts 14 days after second
194 vaccination for Delta vs. Alpha meant that by 12 weeks post second ChAdOx1 dose there
195 was no evidence that onward Delta transmission rates differed between those not
196 vaccinated and those having received two ChAdOx1 doses and the impact of BNT162b2 had
197 also attenuated substantially.

198

199 Although 14 days post second vaccination in contacts, those receiving BNT162b2 vs.
200 ChAdOx1 were at lower risk of testing positive, the protective effect of vaccination in
201 contacts waned faster for BNT162b2 than ChAdOx1 (Figure 1B, aOR per doubling of weeks
202 since 14 days after second vaccination=1.37[95%CI 1.27-1.48] vs. 1.16[1.11-1.21] for
203 ChAdOx1; heterogeneity $p=0.002$).

204

205 [Other transmission risk factors](#)

206 Multiple other factors were associated with contacts testing positive (Figure 2, Table S4,
207 Figures S2-S4), including contact event type and index case age, with the highest rates of
208 PCR-positivity after household contact with index cases aged ≥ 30 years and the lowest rates
209 following contact with index cases < 20 years at work or education (Figure 2A). Contacts in
210 their 30s, 40s and 70s had the highest rates of positive tests after household contact, while
211 contacts in their 20s had the highest rates after contact events outside their own home
212 (Figure 2B). Contacts of index cases of the opposite sex were more likely to test positive,
213 except for children where contacts of girls were more likely to test positive (Figure 2C). Male
214 contacts were more likely than female contacts to be infected outside the home (Figure S2).

215 Case-contact pairs of similar ages were most likely to test positive, particularly with
216 increasing age (Figure 2D). Contacts of asymptomatic index cases were less likely to test
217 positive (aOR for Alpha=0.28[95%CI 0.26-0.30], Delta=0.61[0.50-0.73]) likely related to both
218 lower viral loads (Figure 3) and symptoms. Contacts living in more deprived areas and areas
219 with higher SARS-CoV-2 incidence (Figure S3) were more likely to test positive. Positivity
220 also varied by calendar time (Figure S4).

221

222 [Extent of vaccine impact on transmission explained by viral load](#)

223 Consistent with previous reports, vaccination with BNT162b2 or ChAdOx1 was associated
224 with lower viral loads in Alpha index cases at the time of their positive test, e.g. after two
225 vaccinations in the presence of symptoms, median Ct values (IQR) were 27.9(19.7-32.8) and
226 25.5(18.1-33.0) vs. 18.4(15.8-22.8) if unvaccinated. However, Delta variant infections had
227 similar viral loads independent of vaccination status (Figure 3), and higher viral loads than
228 Alpha in both symptomatic and asymptomatic infections.

229

230 We investigated if adjusting for index case Ct values could explain the attenuated reductions
231 in onward transmission after vaccination seen with the Delta variant. Estimates, adjusted for
232 Ct value, of the impact of index case vaccination status and variants showed variation in
233 measured Ct value accounted for only a modest proportion of the impact of vaccination
234 overall (Figure 4A). This potentially explains why vaccination still reduces onward
235 transmission of Delta despite Ct values at the index positive test being similar regardless of
236 vaccination status. Higher viral loads (lower Ct values) were independently associated with
237 increased transmission for both Alpha and Delta, but with a greater reduction in
238 transmission of Alpha vs. Delta at lower viral loads (Figure 4B).

239

240 Discussion

241 Using large-scale contact tracing data, we show that BNT162b2 and ChAdOx1 vaccination
242 both reduce onward transmission of SARS-CoV-2 from individuals infected despite
243 vaccination. However, reductions in transmission are lower for both vaccines for the Delta
244 variant compared to Alpha. Vaccines continue to provide protection against infection with
245 Delta, but to a lesser degree than with Alpha, particularly considering symptomatic
246 infections or infections with moderate/high viral loads.⁸ Therefore, Delta erodes vaccine-
247 associated protection against transmission by both making infection more common and
248 increasing the likelihood of transmission from vaccinated individuals who become infected.

249

250 It has been hypothesised that vaccines reduce onward transmission from infected
251 vaccinated individuals by reducing viral loads, as higher viral loads are associated with
252 transmission.^{14,15} Therefore, it is perhaps surprising we found that most of the effect of
253 vaccines persisted after adjusting for Ct values, i.e., factors other than PCR-measured viral
254 load at diagnosis are important in vaccine-associated transmission reductions. The single
255 measured Ct value only approximates viral load at the time of transmission, as viral loads
256 are dynamic over time.²² Hence, observed viral loads may not be representative of viral
257 loads at transmission, however, the strong relationship between measured Ct values and
258 risk of onward transmission argues against this¹⁵ (replicated here, Figure 4B). Therefore, it is
259 possible vaccination acts by facilitating faster clearance of viable infectious virions,^{17,18} but
260 leaving damaged ineffective virions behind that still contain PCR-detectable RNA. This may

261 mean antigen assays have advantages in predicting the risk of onward transmission in those
262 vaccinated, but this needs further study.

263

264 We found that index cases infected with the Delta variant and vaccinated with BNT162b2
265 had lower odds of having PCR-positive contacts compared to index cases receiving
266 ChAdOx1, with potentially insufficient power to resolve differences for Alpha. Contacts
267 vaccinated twice with BNT162b2 also had lower rates of Alpha and Delta infections than
268 those vaccinated with ChAdOx1.

269

270 Protection against onward transmission waned within 3 months post second vaccination.

271 For Alpha this still left good levels of protection against transmission, but for Delta this

272 eroded much of the protection against onward transmission, particularly for ChAdOx1,

273 which by 3 months post second vaccine had no evidence of difference in transmission

274 compared to that seen in unvaccinated individuals. “Waning” of protective behaviour over

275 time may also underlie some of the differences seen, with vaccination facilitating reduced

276 social distancing and mask wearing. However, reductions in antibody levels²³ and vaccine

277 effectiveness⁸ over time suggest biological explanations for increasing transmission over

278 time are likely important. Additionally, some of the observed decline may be attributable to

279 the fact that those clinically vulnerable with weaker immune systems were vaccinated

280 longer ago. We also find that the probability of a contact testing positive increased with

281 time since their second vaccination. Although BNT162b2 provided higher levels of

282 protection for contacts throughout the 3 months post-second vaccine, protection against

283 infection waned faster for BNT162b2 than ChAdOx1, as also seen in a representative UK

284 survey.⁸

285

286 This study has several limitations. We considered only contacts who underwent PCR testing,
287 to minimise bias introduced by differences in testing behaviour that may occur for multiple
288 reasons including vaccination of contacts. This means we cannot estimate secondary attack
289 rates by case and contact vaccination status, and that absolute protective effects of
290 vaccination on transmission may be under-estimated as vaccine-protected uninfected
291 contacts may not have sought testing. Our approach is also not likely to eliminate bias,
292 particularly if test-seeking behaviour is related to perceived vaccine efficacy, given non-
293 specificity of many symptoms.²⁴ We did not have sufficient data to consider the impact of
294 previous infection status, which is also imperfectly ascertained in national testing programs.
295 It is likely that part of the explanation for the declines over time in the adjusted probability
296 of contacts testing positive (Figure S4), is increasing prevalence of prior infection in the
297 unvaccinated group, along with changes in test seeking behaviour and the incidence of
298 other infections causing similar symptoms.²⁵ We also had to use SGTF and time as a proxy
299 for Alpha vs. Delta infection rather than sequencing, which means some low viral load Delta
300 infections with SGTF may have been misclassified as Alpha, however we restricted the time
301 period of our dataset to minimise this. As we considered all PCR results in contacts, not just
302 those tested with assays including an S-gene target, we could not assess SGTF concordance
303 as supporting evidence for transmission between case-contact pairs. Finally, we did not
304 have data to adjust for comorbidities; with clinically vulnerable individuals and healthcare
305 workers vaccinated earlier, this may have partly impacted some of our findings, particularly
306 on waning over time and differences by vaccine type.

307

308 The Delta variant has spread globally and caused resurgences of infection even in the setting
309 of high vaccination coverage. Increased onward transmission from individuals who become
310 infected despite vaccination is an important reason for its spread. Booster vaccination
311 campaigns being considered and implemented²⁶ are likely to help control transmission as
312 well as preventing infections.

313

314 Data availability

315 Applications to use the data in this study can be made to NHS Digital's Data Access Request
316 Service, please see <https://digital.nhs.uk/services/data-access-request-service-dars> for more
317 details.

318

319 Declarations

320 DWE declares lecture fees from Gilead outside the submitted work. No other author has a
321 conflict of interest to declare.

322

323 Funding

324 This study was funded by the UK Government's Department of Health and Social Care. This
325 work was supported by the National Institute for Health Research Health Protection
326 Research Unit (NIHR HPRU) in Healthcare Associated Infections and Antimicrobial Resistance
327 at Oxford University in partnership with Public Health England (PHE) (NIHR200915), and the
328 NIHR Biomedical Research Centre, Oxford. The views expressed in this publication are those
329 of the authors and not necessarily those of the NHS, the National Institute for Health

330 Research, the Department of Health or Public Health England. DWE is a Robertson

331 Foundation Fellow and an NIHR Oxford BRC Senior Fellow. ASW is an NIHR Senior

332 Investigator.

333

334 References

- 335 1 Polack FP, Thomas SJ, Kitchin N, *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19
336 Vaccine. *New Engl J Med* 2020; 383: 2603–15.
- 337 2 Voysey M, Clemens SAC, Madhi SA, *et al.* Safety and efficacy of the ChAdOx1 nCoV-19
338 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled
339 trials in Brazil, South Africa, and the UK. *Lancet* 2021; 397: 99–111.
- 340 3 Baden LR, Sahly HME, Essink B, *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2
341 Vaccine. *New Engl J Med* 2020; 384: 403–16.
- 342 4 Bernal JL, Andrews N, Gower C, *et al.* Effectiveness of the Pfizer-BioNTech and Oxford-
343 AstraZeneca vaccines on covid-19 related symptoms, hospital admissions, and mortality in
344 older adults in England: test negative case-control study. *Bmj* 2021; 373: n1088.
- 345 5 Dagan N, Barda N, Kepten E, *et al.* BNT162b2 mRNA Covid-19 Vaccine in a Nationwide
346 Mass Vaccination Setting. *New Engl J Med* 2021. DOI:10.1056/nejmoa2101765.
- 347 6 Bernal JL, Andrews N, Gower C, *et al.* Effectiveness of Covid-19 Vaccines against the
348 B.1.617.2 (Delta) Variant. *New Engl J Med* 2021; 385: 585–94.
- 349 7 Pritchard E, Matthews PC, Stoesser N, *et al.* Impact of vaccination on new SARS-CoV-2
350 infections in the United Kingdom. *Nat Med* 2021; 27: 1370–8.
- 351 8 Pouwels KB, Pritchard E, Matthews PC, *et al.* Impact of Delta on viral burden and vaccine
352 effectiveness against new SARS-CoV-2 infections in the UK. *Medrxiv* 2021; :
353 2021.08.18.21262237.

- 354 9 Harris RJ, Hall JA, Zaidi A, Andrews NJ, Dunbar JK, Dabrera G. Effect of Vaccination on
355 Household Transmission of SARS-CoV-2 in England. *New Engl J Med* 2021; 385: 759–60.
- 356 10 Layan M, Gilboa M, Gonen T, *et al.* Impact of BNT162b2 vaccination and isolation on
357 SARS-CoV-2 transmission in Israeli households: an observational study. *Medrxiv* 2021; :
358 2021.07.12.21260377.
- 359 11 Prunas O, Warren JL, Crawford FW, *et al.* Vaccination with BNT162b2 reduces
360 transmission of SARS-CoV-2 to household contacts in Israel. *Medrxiv* 2021; :
361 2021.07.13.21260393.
- 362 12 Salo J, Hägg M, Kortelainen M, *et al.* The indirect effect of mRNA-based Covid-19
363 vaccination on unvaccinated household members. *Medrxiv* 2021; : 2021.05.27.21257896.
- 364 13 Levine-Tiefenbrun M, Yelin I, Katz R, *et al.* Initial report of decreased SARS-CoV-2 viral
365 load after inoculation with the BNT162b2 vaccine. *Nat Med* 2021; 27: 790–2.
- 366 14 Marks M, Millat-Martinez P, Ouchi D, *et al.* Transmission of COVID-19 in 282 clusters in
367 Catalonia, Spain: a cohort study. *Lancet Infect Dis* 2021; 21: 629–36.
- 368 15 Lee LYW, Rozmanowski S, Pang M, *et al.* SARS-CoV-2 infectivity by viral load, S gene
369 variants and demographic factors and the utility of lateral flow devices to prevent
370 transmission. *Clin Infect Dis* 2021; : ciab421-.
- 371 16 Brown CM, Vostok J, Johnson H, *et al.* Outbreak of SARS-CoV-2 Infections, Including
372 COVID-19 Vaccine Breakthrough Infections, Associated with Large Public Gatherings —

- 373 Barnstable County, Massachusetts, July 2021. *Morbidity Mortal Wkly Rep* 2021; 70: 1059–
374 62.
- 375 17 Singanayagam A, Hakki S, Dunning J, *et al.* Community Transmission and Viral Load
376 Kinetics of SARS-CoV-2 Delta (B.1.617.2) Variant in Vaccinated and Unvaccinated Individuals.
377 *Ssrn Electron J* 2021. DOI:10.2139/ssrn.3918287.
- 378 18 Chia PY, Ong SWX, Chiew CJ, *et al.* Virological and serological kinetics of SARS-CoV-2
379 Delta variant vaccine-breakthrough infections: a multi-center cohort study. *Medrxiv* 2021; :
380 2021.07.28.21261295.
- 381 19 Public Health England. SARS-CoV-2 variants of concern and variants under investigation
382 in England: Technical briefing 21. 2021; published online Aug 20.
383 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1012644/Technical_Briefing_21.pdf)
384 [data/file/1012644/Technical_Briefing_21.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1012644/Technical_Briefing_21.pdf) (accessed Sept 2, 2021).
- 385 20 Fukushima W, Hirota Y. Basic principles of test-negative design in evaluating influenza
386 vaccine effectiveness. *Vaccine* 2017; 35: 4796–800.
- 387 21 Public Health England. Understanding cycle threshold (Ct) in SARS-CoV-2 RT-PCR. 2020;
388 published online Oct 1.
389 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/926410/Understanding_Cycle_Threshold__Ct__in_SARS-CoV-2_RT-PCR_.pdf)
390 [data/file/926410/Understanding_Cycle_Threshold__Ct__in_SARS-CoV-2_RT-PCR_.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/926410/Understanding_Cycle_Threshold__Ct__in_SARS-CoV-2_RT-PCR_.pdf)
391 (accessed Sept 27, 2021).
- 392 22 He X, Lau EHY, Wu P, *et al.* Temporal dynamics in viral shedding and transmissibility of
393 COVID-19. *Nat Med* 2020; 26: 672–5.

- 394 23 Wei J, Stoesser N, Matthews PC, *et al.* Antibody responses to SARS-CoV-2 vaccines in
395 45,965 adults from the general population of the United Kingdom. *Nat Microbiol* 2021; 6:
396 1140–9.
- 397 24 Lewnard JA, Patel MM, Jewell NP, *et al.* Theoretical Framework for Retrospective Studies
398 of the Effectiveness of SARS-CoV-2 Vaccines. *Epidemiology* 2021; 32: 508–17.
- 399 25 Vihta K-D, Pouwels KB, Peto T, *et al.* Symptoms and SARS-CoV-2 positivity in the general
400 population in the UK. *Medrxiv* 2021; : 2021.08.19.21262231.
- 401 26 Bar-On YM, Goldberg Y, Mandel M, *et al.* Protection of BNT162b2 Vaccine Booster
402 against Covid-19 in Israel. *New Engl J Med* 2021. DOI:10.1056/nejmoa2114255.
- 403

404 Table

Characteristic	Alpha			Delta			Alpha vs Delta		
	aOR	95% CI	p-value	aOR	95% CI	p-value	Interaction OR	95% CI	Interaction p value
Impact on onward transmission: Case vaccination status									
Unvaccinated	—	—	—	—	—	—	—	—	—
Partial ChAdOx1	0.82	0.76, 0.88	<0.001	0.98	0.90, 1.06	0.99	1.20	1.07, 1.34	0.003
Partial BNT162b2	0.74	0.70, 0.80	<0.001	0.87	0.81, 0.94	0.004	1.17	1.06, 1.30	0.009
Full ChAdOx1	0.37	0.22, 0.63	0.002	0.64	0.57, 0.72	<0.001	1.71	1.01, 2.90	0.045
Full BNT162b2	0.18	0.12, 0.29	<0.001	0.35	0.26, 0.48	<0.001	1.93	1.25, 2.99	0.003
Contact vaccination status									
Unvaccinated	—	—	—	—	—	—	—	—	—
Partial ChAdOx1	0.89	0.82, 0.97	0.076	0.54	0.50, 0.58	<0.001	0.60	0.54, 0.67	<0.001
Partial BNT162b2	0.69	0.64, 0.75	<0.001	0.58	0.55, 0.61	<0.001	0.83	0.76, 0.92	<0.001
Full ChAdOx1	0.29	0.17, 0.49	<0.001	0.28	0.25, 0.32	<0.001	0.97	0.57, 1.65	0.91
Full BNT162b2	0.06	0.04, 0.10	<0.001	0.10	0.08, 0.13	<0.001	1.64	1.07, 2.51	0.024

405


406 **Table 1. Relationship between PCR-positive results in contacts, and index case and contact vaccination status according to Alpha/Delta**

407 **variant in the index case.** Results for those with two vaccine doses are estimated at day 14 post second vaccine, see Figure 3 for trends with

408 time post-second vaccine. aOR, adjusted odds ratio, CI confidence interval. Adjustment made for contact event type; index case factors - age,

409 sex, and symptom status; contact factors - age, sex; local deprivation, local SARS-CoV-2 incidence and calendar time (see Table S4 and Figures
410 S2-S8 for details).

411 **Figures**




412
413

414 **Figure 1. Estimated probability of a positive PCR test in contacts by time since second**
415 **vaccination in index cases (panel A) and in contacts (panel B), variant, and vaccine type.**

416 For panel A estimates are displayed for an unvaccinated contact and for panel B for an
417 unvaccinated index case. The relationships shown also apply to vaccinated contacts and
418 cases, as the underlying odds ratios are independent of contact and case vaccine status
419 respectively, however absolute probabilities differ. The dashed horizontal lines indicate the

420 probability of a positive-PCR result in an unvaccinated contact of an unvaccinated index
421 case. The shaded area indicates the 95% confidence interval. Adjustment made for
422 covariates, set to reference values: contact event type (set to Household or
423 accommodation); index case factors – age (median), sex (female), and symptom status
424 (symptomatic); contact factors – age (median), sex (female); local deprivation (median),
425 local SARS-CoV-2 incidence (median) and calendar time (median).
426



427

428

429 **Figure 2. Estimated probability of a positive PCR test in contacts by contact event type and**


430 **index case age (panel A) or contact age (panel B), contact age and case and contact sex**

431 **(panel C) and case and contact age (panel D).** For each panel all other covariates are set to

432 reference values for categorical values and median values for continuous variables, i.e.

433 contact event type set to Household or accommodation; index case factors – age (median),

434 sex (female), vaccination status (unvaccinated) and symptom status (symptomatic); contact
435 factors – age (median), sex (female), vaccination status (unvaccinated); local deprivation
436 (median), local SARS-CoV-2 incidence (median) and calendar time (median). Shaded ribbons
437 and error bars indicate 95% confidence intervals.
438




439

440 **Figure 3. Distribution of viral loads by index case vaccination status, variant, and**

441 **symptoms.** The solid line in each violin plot indicates the median. See Lee *et al* for details of

442 equivalent viral loads in copies per ml ($\log_{10}(\text{VL}) = 12.0 - 0.328 \cdot \text{Ct}$).¹⁵

443



444

445

446 **Figure 4. Extent of vaccine-associated transmission reductions explained by change in**

447 **viral load (Ct values) in index case (panel A) and relationship between Ct values and**

448 **transmission by lineage (panel B). Adjusted odds ratios are shown in panel A and error bars**

449 **(panel A) or the shaded ribbon (panel B) indicate the 95% confidence interval.**