SARS-CoV-2 RNA and antibody dynamics in a Dutch household study with dense sampling frame

Wanda G.H. Han[1]*
Arno Swart[1]*
Axel Bonacic Marinovic[1]
Dirk Eggink[1]
Johan Reimerink[1]
Lisa A. Wijsman[1]
Bas van der Veer[1]
Sharon van den Brink[1]
Anne-Marie van den Brandt[1]
Fion Brouwer[1]
Marieke Hoogerwerf[1]
On behalf of the Dutch FFX-COVID-19 research group
Daphne F.M. Reukers[1]
Nynke Rots[1]
Chantal Reusken[1]
Adam Meijer[1]
18
19 [1]Centre for Infectious Disease Control, National Institute for Public Health and the Environment,
20 Bilthoven, the Netherlands
21 *Equal contribution
22 Membership of the Dutch FFX-COVID-19 research group is provided in the Acknowledgments
23
Corresponding author contact information:
Adam Meijer
Centre for Infectious Diseases Research, Diagnostics and Laboratory Surveilance, Centre for
Infectious Disease Control (CIb), National Institute for Public Health and the Environment (RIVM)
PO Box 1, 3720 BA Bilthoven, the Netherlands
Tel.: +31 302743595
E-mail: Adam.Meijer@rivm.nl
31

Running title: SARS-CoV-2 diagnostics and symptom onset

Abstract

This study investigated the dynamics of SARS-CoV-2 infection and diagnostics in household members

of different ages and with different symptom severity after SARS-CoV-2 exposure during the early

phase of the pandemic. Households with a SARS-CoV-2 confirmed positive case and at least one child

in the Netherlands were followed for 6 weeks. Naso (NP)- and oropharyngeal (OP) swabs, oral fluid

and feces specimens were analyzed for SARS-CoV-2 RNA and serum for SARS-CoV-2-specific

antibodies. The dynamics of the presence of viral RNA and the serological response was modeled to

determine the sampling time-frame and sample type with the highest sensitivity to confirm or reject

a SARS-CoV-2 diagnosis. Transmission of SARS-CoV-2 between adults and children within a household

was correlated with symptom severity of index cases. In children higher viral loads compared to adults

were detected at symptom onset. Early in infection, higher viral loads were detected in NP and OP

specimens, while RNA in especially feces were longer detectable. SARS-CoV-2-specific antibodies have

a 90% probability of detection from 7 days (total Ig) and 18 days (IgG) since symptom onset. In

conclusion this study has shown that on average, children carry higher loads of virus as compared to

adults early after infection. For highest probability of detection in SARS-CoV-2 diagnostics early in

infection, RT-PCR on NP and OP specimens are more sensitive than on oral fluid and feces. For SARS-

CoV-2 diagnostics late after infection, RT-PCR on feces specimens and serology are more valuable.

Keywords: SARS-CoV-2, Household study, molecular diagnostics, serological diagnostics, infection

dynamics

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has spread rapidly across the world since January 2020 [1]. In the Netherlands, the first COVID-19 (the syndrome caused by SARS-CoV-2) case was detected on 27 February 2020. From March until May 2020, the Dutch government mandated a partial lockdown. This included social distancing, self-quarantine and self-isolation orders, closing of schools, bars and restaurants, and urging people to work from home [2]. Yet, households are close-contact settings with high probability of (pre/a-symptomatic) transmission of SARS-CoV-2 after introduction of the virus. In this period, a prospective cohort study was performed including 55 complete households with a RT-PCR-confirmed SARS-CoV-2 positive case (index case) and at least one child below 18 years of age. All household contacts were tested as soon as possible after a SARS-CoV-2 infection in the household was identified. At multiple timepoints, various clinical samples were collected for molecular and serological diagnostics. Using a dense sampling strategy, SARS-CoV-2 transmission and kinetics of diagnostic parameters could be closely monitored within the households. Earlier we described that the estimated Secondary Attack Rate (SAR) in this cohort that was high (35% in children, 51% in adults), with reduced susceptibility of children compared to adolescents and adults (0.67; 95%CI: 0.40-1.1) [3]. Here we looked further into the kinetics of infection.

In the present study, we use the results of the dense sampling and various molecular and serological assays to identify participants with an acute or recent SARS-CoV-2 infection to analyse household transmission patterns in relation to disease severity. Secondly, we describe the dynamics of the infection per individual based on viral RNA and antibody presence. Lastly, we compared the dynamics of the different diagnostic methods (test and sample type), by modeling the outcomes per assay in relation to the days post symptom onset (dps), disease severity and age.

Methods

Study protocol

A prospective cohort study was performed following households where one symptomatic household

member was tested RT-PCR positive for SARS-CoV-2 in the period 24 March – 6 April 2020 [3]. In brief,

persons 18 years and older testing positive for SARS-CoV-2 RT-PCR (i.e. the index case) who had at

least one child in their household below the age of 18 could be included in this study (METC nr:

NL13529.041.06). Table 1 describes the sampling scheme (See Reukers et al [3] for more details). We

defined adults as individuals of 18 years of age or older and individuals as SARS-CoV-2 infection

positive when they tested positive in at least one RT-PCR or serological assay.

COVID-19 severity

The day of onset of possible COVID-19 associated symptoms, i.e. respiratory symptoms (including sore

throat, cough, dyspnea or other respiratory difficulties, rhinorrhoea), fever, chills, headache, anosmia

or ageusia, muscle pain, joint ache, diarrhoea, nausea, vomiting, loss of appetite or fatigue, as

reported by the participant was defined 1 dps. Laboratory confirmed SARS-CoV-2 infected cases with

any clinical symptoms other than pneumonia and/or requiring medical consultation were defined as

mild cases. Moderate cases showed clinical signs of pneumonia, including dyspnea. Severe cases

reported dyspnea and consulted a health care professional for their symptoms, or reported having

been admitted to the hospital for COVID-19 [3, 4].

Transmission categories

We categorized the household transmission patterns in three groups. In the 'no transmission'

category, SARS-CoV-2 infection was only detected in the index case. In the 'adult transmission'

category, SARS-CoV-2 infection was detected in adults other than the index case only. In the 'family

transmission' category, SARS-CoV-2 infection was detected in children and possibly also adults other

than the index case. In order to assess the significance of differences in severities over transmission

categories a generalized linear model (GLM) for count data with Poisson family and exponential link

was set up with interactions between the transmission category and severity classes.

Molecular diagnostics

Nasopharyngeal swabs (NP) and oropharyngeal swabs (OP) were collected in gelatin-lactalbumin-

yeast (GLY) viral transport medium (Mediaproducts BV, Groningen, The Netherlands), transported to

the laboratory in a cooling box and stored at maximum a few days at 4° C until being processed for

RT-PCR. Feces specimens were self-collected by the patient and send to the laboratory by regular mail,

stored frozen at -20° C until being processed for RT-PCR. Oral fluid specimens were collected with a

Oracol sponge (Malvern Medical Developments Ltd, U.K.), transported to the laboratory in a cooling

box, processed for storage according to the manufacturer's instructions, and aliquots stored frozen at

-80° C until being used for RT-PCR. Total nucleic acid was extracted from NP-, OP swab, oral fluid or

feces using MagNApure 96 (MP96) with total nucleic acid kit small volume (Roche). Of the feces

specimens a 5% suspension was made in MEM with Hanks' salts and penicillin and streptomycin,

vortex for 15 seconds and 1 minute centrifuged at 16,000 Relative Centrifugal Force. Two-hundred µl

supernatant was mixed with 275 µl MP96 lysis buffer including equine arteritis virus (EAV) internal

control and yeast tRNA stabilizer. Total nucleic acid was eluted in 50 µl Tris EDTA buffer. RT-qPCR was

performed on 5 µl total nucleic acid using TaqMan® Fast Virus 1-Step Master Mix (Thermo Fisher) on

Roche LC480 II thermal cycler with SARS-like beta coronavirus (Sarbeco) specific E-gene primers and

probe and EAV as described previously [5, 6]. As no other Sarbeco viruses are currently detected in

humans, a positive Sarbeco E-gene RT-qPCR is validly taken as positive for SARS-CoV-2. For modeling

purposes no detection of SARS-CoV-2 RNA was given an artificial cycle threshold (Ct) value of 40.

Serological diagnostics

The Wantai SARS-CoV-2 total antibody ELISA (Beijing Wantai Biological Pharmacy Enterprise, Beijing,

China; catalogue number WS1096) was performed according to the manufacturer's instructions [7].

This assay is a double-antigen sandwich ELISA using the recombinant receptor-binding domain of

SARS-CoV-2 as antigen. Optical density (OD) is measured at 450 nm and the antibody OD ratio for each

sample is calculated as the ratio of the OD of that sample to the reading of a calibrator (included in

the kit).

Sera were tested for the presence of IgG antibodies reactive with the SARS-CoV-2 S1 and SARS-

CoV-2 N antigens in a protein microarray, in duplicate 2-fold serial dilutions starting at 1:20, essentially

as described previously [8]. For each antigen, a 4-parameter loglogistic calibration curve was

generated. Antibody titers (EC50 value) were defined as the interpolated serum dilution that gave a

fluorescence intensity of 50% of the corresponding calibration curve. Raw data were processed with

the R 4.04 statistical software as described previously [9].

Modeling RT-PCR

All available RT-PCR outcomes (Table 1) were modelled by a Bayesian hierarchical model of the form

$$Ct_i \sim N(\mu_i, \sigma)$$

$$\mu_i = \alpha_{\text{test}[i]} + \alpha_{\text{test}[i], \text{id}[i]} + \alpha_{\text{test}[i],\text{agecat}[i]} + \alpha_{\text{test}[i],\text{severity}[i]}$$
$$+ +\big(\beta_{\text{test}[i]} + \beta_{\text{test}[i],\text{id}[i]} + \beta_{\text{test}[i],\text{agecat}[i]} + \beta_{\text{test}[i],\text{severity}[i]} +\big)d_i$$

Here, Ct_i is the measured Ct value for sample i, and σ the overall variation. There is a part dependent

on the days since onset of symptoms (d_i) with coefficients β, and a constant part with coefficients α.

Both the α and β parameters include several contributions stratified by categorical variables: id[i] is

the person specific identifier for sample i enabling longitudinal modelling, agecat[i] is either 'child'

or 'adult', severity[i] is either 'asymptomatic', 'mild', 'moderate' or 'severe', and test[i] is either 'pcr

NP', 'pcr OP', 'pcr oral fluid', 'pcr feces', 'wantai', 'microarray S1' or 'microarray N'.

For the id's a hierarchical model is built (i.e. a random effect),

$$\alpha_{\text{test}[i], \text{id}[i]} \sim N(0, \sigma_{\text{test}[i]}), \quad \beta_{\text{test}[i], \text{id}[i]} \sim N(0, \tau_{\text{test}[i]}).$$

Whenever a Ct-value of 40 is encountered, we apply censoring in the model by changing the

probability density function $f(x; \mu_i, \sigma)$ into its cumulative counterpart $1 - F(40; \mu_i, \sigma)$, thereby

encoding that we have an unknown Ct value which would either indicate the absence of amplifiable

RNA or presence of RNA but well below the detection limit of the used RT-PCR. All parameters are

given weakly informative priors, and the posterior distributions are obtained using the JAGS software

[10], interfaced from R [11]. Bayesian credible intervals were obtained from the samples of the

posterior as calculated by JAGS. Prediction intervals were calculated by drawing randomly from

$N(\hat{\mu}_j, \hat{\sigma}_j)$, where the indicated means and standard deviations are samples from the posterior

distributions. The posterior probability of being positive is modelled by $F(40; \hat{\mu}_j, \hat{\sigma}_j)$. The modeling

did not include the Ct values of the inclusion RT-PCR NP+OP which was performed in index cases just

before start of the study, since these values are not known to us.

Modeling serology

The dynamics of serology cannot be assumed to be linear as is the case for Ct-values. Rather,

seronegative individuals have a titer (OD ratio for Wantai or EC50 for protein microarray) varying

around a low value, and seropositive individuals have a titer varying around a high value. In the case

of the ELISA-test and microarray-based assays used in the current study, we find that a cut-off value

to distinguish seropositives and seronegatives works well, since the two components are well

separated (Figure S1). Using the cut-off values 1 for Wantai (according to manufacturer's instructions)

and 10 for microarray [8], we classify each measurement X_i as either positive or negative. Using a

Bernouilli distribution and logit link for the probability we model the outcomes as

$$X_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \alpha_{\text{test}[i]} + \alpha_{\text{test}[i],\text{id}[i]} + \alpha_{\text{test}[i],\text{agecat}[i]} + \alpha_{\text{test}[i],\text{severity}[i]}$$
$$+ \left(\beta_{\text{test}[i]} + \beta_{\text{test}[i],\text{id}[i]} + \beta_{\text{test}[i],\text{agecat}[i]} + \beta_{\text{test}[i],\text{severity}[i]}\right)d_i$$

Parameter estimation proceeds analogous to the RT-PCR model.

Assessing differences between factors

We assessed the difference between posterior estimates of parameters using the Region of Practical

Equivalence (ROPE) [12-14]. The ROPE is an interval chosen based on domain knowledge that indicates

values that are practically indistinguishable. For Ct-values our ROPE interval is [-1,1], which means that

we consider differences between Ct-values of less than one as not meaningful. For changes in Ct-value

per day (the slope) we choose [-1/7, 1/7], which means that we consider differences between Ct-

values of less than one per week as not meaningful. For serology detection probability (dps) our ROPE

interval is [-2,2], which means that we consider differences between days of less than 2 as not

meaningful. The ROPE is compared to the 89% highest posterior density interval (HDI). When the ROPE

contains the HDI, no meaningful difference exists, when the ROPE is completely outside of the HDI,

there is a difference, when the ROPE and HDI overlap we withhold a decision because of too high

uncertainty.

Results

Household transmission SARS-CoV-2

A total of 242 participants from 55 complete households were included in this study. The number of

analyses performed per assay and specimen type at the various timepoints with the day of the first

home visit (so the start of the study within the particular household) defined as day 1 are described in

Table 1 and Table S1. To identify different transmission patterns, we visualized SARS-CoV-2 infection

detection by the different assays and specimen types per participant and household in heatmaps. We

identified the transmission pattern 'no transmission' in 16 households (Figure 1A), 'adult transmission'

in 11 households (Figure 1B) and 'family transmission' in 28 households (Figure 1C). Eight of the 28

households in the 'family transmission' category did not show transmission to adults.

Symptom severity of COVID-19 index cases correlated with transmission of SARS-CoV-2

between adults and children within a household as reflected by the overrepresentation of index cases

with severe symptoms in the family transmission group (p=0.03, 54% of indexes with severe COVID-

19) compared to the other groups (Figure 2). In the no transmission category, more than half (56.3%)

of the index cases had mild symptoms, whilst only 19% had severe symptoms. In the adult

transmission category, there were 4 severe index cases out of 10 (36%).

SARS-CoV-2 infection dynamics

We investigated the SARS-CoV-2 infection dynamics in the participants of the study. Using an 'upset

plot' [15], Figure 3 shows patterns of positive and negative results in the various molecular and

serological assays. Of the 242 participants, 136 individuals were positive for SARS-CoV-2 infection by

either molecular and/or serological diagnostics. Of these 136 SARS-CoV-2 infection positive

individuals, 125 (91.9%) were symptomatic and severe symptomatic individuals were mainly (29 out

of 32) RT-PCR- and serology-positive (Figure 3B). Most individuals were found SARS-CoV-2 positive by

multiple diagnostic assays and/or materials, but 19 individuals tested positive with only one assay type

and/or material (only Wantai n=5, RT-PCR NP n=3, RT-PCR oral fluid n=3, RT-PCR OP n=2, RT-PCR feces

n=2, Inclusion PCR NP+OP n=2, MA-S n=1, MA-N n=1) during the study period (Figure 3A). In most of

the infected cases (81.6%, 111 of the 136) both SARS-CoV-2 RNA and SARS-CoV-2-specific antibodies

were detected (Figure 3B). For 12 individuals only at one timepoint during the study one positive test

was found (Figure 1, red rectangles).

Next, for the individuals with at least one RT-PCR and one serological result at visit 1, 2 and 3

(n=198), we could analyze rough dynamics of the infection process (Figure 4). The median dps relative

to visit 1 is indicated in Figure 4. Six common patterns in 173 individuals, ranked A – F based on

frequency, could be identified. Within laboratory confirmed SARS-CoV-2 infected individuals, the

common patterns B (n=28) and C (n=27) included individuals with a positive PCR and serological assay

at visit 1. As can be expected, SARS-CoV-2 viral RNA was not detected anymore in these cases at the

end of the study (4-6 weeks after inclusion), while SARS-CoV-2 antibodies remained present.

Individuals with pattern E (n=12) did not have detectable SARS-CoV-2 RNA at any visit, but did have

detectable SARS-CoV-2 antibodies at visit 1, 2 and 3. This pattern is in line with an earlier onset of

symptoms compared to the individuals with pattern B and C, thus these individuals were included in

the study later in their infection process resulting in already diminished viral RNA and present

antibodies at visit 1. Pattern D (n=18) and F (n=9) included individuals with a positive RT-PCR at visit 1

and developed antibodies after visit 1. Compared to pattern B and C, these individuals reported their

onset of symptoms 2-4 days later, thus at study inclusion (visit 1) they were earlier in their infection

process. Pattern A included individuals with negative RT-PCR and serology results at all visits. These

rough patterns underline that there are optimal time windows in which detection of SARS-CoV-2 viral

RNA or SARS-CoV-2-specific antibodies are most appropriate in diagnostics. To further investigate this

we used a modelling approach.

Dynamics SARS-CoV-2 diagnostics

Bayesian modeling on all available RT-PCR data demonstrated a difference in Ct-values at symptoms

onset between adults and children (Figure 5A, B and S2-G1) and in Ct-values at symptom onset and in

Ct-value increase per day between different specimen types (Figure 5C and S2-A to F). The predicted

Ct value (inversely correlated with viral load), was on average 2.6 Ct lower in children (Ct 27.5; all

specimens) compared to adults (Ct 30.1; all specimens) at the day of symptom onset (intercept)

(Figure 5A, B and S2-G1). The decay in time in viral load (slope) was comparable between adults and

children (Figure 5B and S2-G2). In line with this, there is a longer probability of SARS-CoV-2 RNA

detection with increasing dps in children (99% detection until 13 dps), compared to adults (99%

detection until 7.6 dps) (Figure S3A). When analyzing all ages, the predicted viral load seems slightly

higher in NP and OP swabs (Ct 28.8 and 28.7) compared to feces (Ct 30.1) and oral fluid (Ct 30.7) at

the day of symptom onset (intercept) (Figure 5C) indicating higher sensitivity of SARS-CoV-2 RNA

detection in NP and OP specimens compared to oral fluid (Figure S2-B1 and D1) and possibly feces

(Figure S2-C1 and E1). In contrast, there seems te be a slower decay (slope) in viral load in oral fluid

and feces specimens (0.25 and 0.22 Ct per day) compared to NP and OP specimens (0.35 and 0.36 Ct

per day) (Figure 5C). The relevance of these findings is uncertain as there is partial overlap between

the ROPE and HDI (Figure S2-B1 to E1 and B2 to E2). Overall, as a most likely estimate, the estimated

viral load is higher in NP and OP swabs compared to oral fuid until 21.1 dps and 19.0 dps, respectively

(Figure S4-B and D), and compared to feces until 10.2 dps and 9.7 dps, respectively (Figure S4-C and

E). Furthermore, there is a longer probability of SARS-CoV-2 RNA detection with increasing dps in feces

263 (90% detection until 27 dps), compared to NP, OP and oral fluid specimens (90% detection until 19.4,

264 20.3 and 22.7 dps, respectively) (Figure 5D and Figure S5-C, E and F). Similar trends are shown for 50%

265 and 10% detection probability (Figure S5 and S6). We could not find a clear correlation in severity of

266 symptoms and the dynamics of the SARS-CoV-2 RNA detection (data not shown).

267  Furthermore, we investigated the dynamics of SARS-CoV-2 antibody detection (Figure 6). The

268 Wantai assay (total Ig) demonstrated a higher sensitivity for detection of anti-S1 antibodies than the

269 micro-array (IgG) as the probability for detection was earlier using Wantai upon onset of illness (Figure

270 6A and S7-A). The dps at which 90% detection probability was reached for Wantai was 7.1 compared

271 to 16.9 and 18 for Nucleoprotein (N)- and S1-protein microarray respectively (Figure 6B). The protein

272 microarray for S1 and N had comparable sensitivity, in line with a previous study [8]. The probability

273 of detecting N-specific IgG antibodies in children was delayed by 3.0 days (at 90% probability

274 detection) versus adults (Figure 6C and D), while this was not the case for detection of S1-specific

275 antibodies (Figure 6C). The relevance of this finding is uncertain as there is much overlap between the

276 distributions in children and adults (Figure S8). The N-specific IgG antibody titers (at visit 3;

277 convalescent phase) were not significantly lower in children compared to adults (Figure S9).

278 Furthermore, we could not find a correlation in severity of symptoms and the dynamics of the SARS-

279 CoV-2 infection detection by the serological assays (data not shown).

281 **Discussion**

282 We studied SARS-CoV-2 RNA and antibody kinetics in a household cohort during the early phase of

283 the pandemic using an unusual dense sampling schedule allowing for high resolution analysis. For

284 highest probability of detection of SARS-CoV-2 RNA by RT-PCR early in infection NP and OP are more

285 suitable than oral fluid and feces. SARS-CoV-2-specific antibodies have a 90% probability of detection

286 from 7 dps with the Wantai assay and 18 dps with the microarray S1 and N assay. This study has been

287 performed in a naïve population during the early phase of the pandemic. Although SARS-CoV-2

288 vaccination, previous infection and the circulation of other SARS-CoV-2 variants may influence the

dynamics of SARS-CoV-2 infection and thereby diagnostics, our study provides valuable reference

insights into this subject.

Households present close-contact settings with high risk of SARS-CoV-2 transmission after

introduction of SARS-CoV-2 in the household [3, 16]. We observed a positive correlation between

transmission of SARS-CoV-2 between adults and children and the severity of disease in the household

indexes. This is in line with studies that report that the severity of SARS-CoV-2 infection of the index

case was associated with higher infectiousness [16, 17]. It should be noted that at the time of the

study, SARS-CoV-2 testing in the Netherlands was limited to symptomatic healthcare workers and

symptomatic vulnerable individuals. The index cases were thus mainly symptomatic healthcare

workers [3]. As schools and daycare centres were closed during the study period, transmission outside

the households among children was minimalized. Due to limited sample size, we could only

categorized children as those of 17 years of age or younger for the Bayesian hierarchical modelling. It

would be of interest to stratify the children in more groups, e.g. primary school age and adolescent

age.

Seroconversion rates in mild to severe symptomatic SARS-CoV-2 RT-PCR positive cases have

been reported in the range of 93-100% after 3-4 week [18-20]. During the course of this study, a vast

majority of RT-PCR confirmed SARS-CoV-2 infected participants (91.0%, 111/122) developed SARS-

CoV-2 specific antibodies (Figure 3B). In 6 RT-PCR positive cases, serology data was missing. In 11 of

122 (9.0%) RT-PCR confirmed SARS-CoV-2 infected individuals there was no seroconversion (Figure

3B). These individuals may have experienced a relative mild infection, or in the cases with only one

positive RT-PCR test (including the available results of the extra sampling between visit 1 and 2, n=8)

and no evidence of antibody response, the SARS-CoV-2 diagnosis of these individuals is disputable.

Four of the 8 RT-PCR-negative cases that did have SARS-CoV-2-specific antibodies during the study

period, showed only one positive serological test, suggesting also a disputable SARS-CoV-2 diagnosis.

In the other 4 individuals, SARS-CoV-2-specific antibodies were detected at multiple visits including at

visit 1, therefore likely having experienced infection before inclusion. Nevertheless, technically there

was no evidence of false positivity as all negative controls had correct results. Therefore, the disputable results remain unexplained. Transient exposure without established infection and rapid waning of immune response might be one explanantion.The individuals with negative RT-PCR and serology results at all visits were probably not infected with SARS-CoV-2 (Figure 3B and Figure 4, pattern A). Those reporting symptoms were not tested for alternative diagnoses (i.e. other respiratory viruses). From national surveillance reports we known that also other viruses causing COVID-19-like acute respiratory infection symptoms circulated (https://www.rivm.nl/virologische-weekstaten). Especially in March 2020, the first study month, before the COVID-19 measures were put in place. Later on rhinoviruses continued circulating during the measures.

In the current study we used Bayesian hierarchical models to determine the sampling time-frame and sample type with the highest sensitivity to confirm or reject a SARS-CoV-2 diagnosis. Due to the close-contact settings with high probability of (pre/a-symptomatic) transmission of SARS-CoV-2 after introduction of the virus in households and the longitudinal dense sampling performed in this study, the analysis yielded results with relatively narrow credibility intervals which support our findings. For highest probability of a SARS-CoV-2 diagnosis (lower Ct), RT-PCR on NP and OP specimens were more suitable than feces and oral fluid until 10 dps and 20 dps, respectively (Figure S4C, E and B, D). Our study confirms that NP and OP (or combined NP+OP) were the preferred sample type for RT-PCR-based SARS-CoV-2 diagnostics within 1 week upon onset of symptoms [21-23]. A systematic review concluded that of the alternative specimens to NP and OP swabs, oral fluid (saliva) has an estimated sensitivity of 83.9% (95% CI: 77.4-88.8) and specificity of 96.4% (95% CI: 89.5-98.8) compared to reference NP and OP swabs in nucleic acid assays [23]. The sensitivity and specificity of feces specimens seems much lower, although limited data is available [23]. For SARS-CoV-2 diagnostics late in infection or in past infections, RT-PCR on feces and oral fluid specimens are more valuable than NP and OP specimens, since the presence of viral RNA in especially feces remain present over a longer time compared to NP and OP swab specimens. This is in line with findings of other studies that indicate that SARS-CoV-2 RNA can be detected up to 126 days in feces compared to 83 days in

respiratory specimens and that beyond 10 dps, feces sampling may be preferred [22, 24, 25]. Although

SARS-CoV-2 RNA can remain present in respiratory and feces specimens for a long time, the duration

of presence of viable virus is relatively short-lived [26, 27]. Alternatively, from a week after symptom

onset testing for the presence of SARS-CoV-2 Spike-specific total Ig using Wantai ELISA can confirm a

recent or past SARS-CoV-2 infection. SARS-CoV-2 diagnostics using protein microarray detecting SARS-

CoV-2 Spike- and Nucleoprotein-specific IgG antibodies, is useful 2 weeks after infection or symptom

onset.

The infection dynamics of SARS-CoV-2 may be influenced by characteristics of the tested

population, such as age and the severity of COVID-19. We, however, could not find a clear correlation

in severity of symptoms or age and the RNA and SARS-CoV-2-specific antibody kinetics, although our

study may be underpowered to detect these differences (data not shown). Children in general report

milder symptoms compared to adults (Table S1) [3]. It is known that with age, the expression of ACE-

2 increases in nasal epithelium [28]. Since SARS-CoV-2 uses the ACE-2 receptor for host entry, a lower

expression of *ACE-2* in children relative to adults might explain the lower susceptibility and milder

infection course in children. However, our findings and other studies show that viral loads in children

are similar or higher than viral loads in adults [29, 30]. In our study children displayed lower Ct values

(higher viral loads) at the day of symptom onset compared to adults, while the decay in viral load was

comparable (Figure 5) . This suggests that if children become infected with SARS-CoV-2, they can carry

high loads of virus for a longer time compared to adults. Therefore, children are potentially longer

infectious than adults after symptom onset. Whether this observation holds for new Variants of

concern (VOC) e.a. delta, warrants further investigation. No clear differences between adults and

children were found in the dynamics of SARS-CoV-2 serology, yet the detection of N-specific antibodies

seems slightly delayed in children compared to adults (Figure 6C, D and S8). A study showed a reduced

breadth of anti-SARS-CoV-2-specific antibodies, predominantly generating IgG antibodies specific for

the S protein but not the N protein in children compared to adults [31]. Whether this has

consequences for the development of immunity to SARS-CoV-2 is not yet clear.

In summary, our study allowed for a high resolution analysis of the sensitivity of molecular

and serology-based detection of recent SARS-CoV-2 infections due to the unusual dense sampling

strategy in a confined setting. For highest probability of SARS-CoV-2 diagnostics early in infection, PCR

on NP and OP specimens are in favor over oral fluid and feces. For SARS-CoV-2 diagnostics late in

infection or in past infection, RT-PCR on feces specimens and serology are more valuable. Children

seem to carry higher loads of virus for a prolonged time in comparison to adults. The data presented

here strengthen the evidence-basis for SARS-CoV-2 testing strategies.

Acknowledgements

We thank the Public Health Service Utrecht for assistance in the recruitment of households. We thank

Alper Çevirgel, Anneke Westerhof, Anoek Backx, Elma Smeets-Roelofs, Elsa Porter, Elske Bijvank,

Francoise van Heiningen, Gabriel Goderski, Gert-Jan Godeke, Harry van Dijken, Helma Lith, Hinke ten

Hulscher, Ilse Akkerman, Ilse Schinkel, Jeroen Hoeboer, Jolanda Kool, Josine van Beek, Joyce Greeber,

Kim Freriks, Lidian Izeboud, Lisa Beckers, Liza Tymchenko, Maarten Emmelot, Maarten Vos, Margriet

Bisschoff, Marit de Lange, Marit Middeldorp, Marjan Bogaard, Marjan Kuijer, Martien Poelen, Nening

Nanlohy, Olga de Bruin, Rogier Bodewes, Ruben Wiegmans, Sakinie Misiedjan, Saskia de Goede,

Sophie van Tol, Titia Kortbeek, and Yolanda van Weert for assistance in logistics and laboratory

analyses. We thank Bettie Voordouw for critical review of manuscript. The Dutch FFX-COVID-19

research group of the Centre for Infectious Disease Control, National Institute for Public Health and

the Environment, the Netherlands, includes the following members: Arianne B. van Gageldonk-

Lafeber, Wim van der Hoek, Susan van den Hof, Adam Meijer, Daphne F.M. Reukers, Chantal Reusken,

Inge Roof and Nynke Rots.

This study was funded by the Dutch Ministry of Health, Welfare, and Sport (VWS).

None of the authors has any potential conflicts of interest to disclose.

References

[1] World Health Organization (WHO). Timeline of WHO's response to COVID-19. https://www.who.int/news/item/29-06-2020-covidtimeline.

[2] Minister Bruno Bruins MIvE, Minister Arie Slob, Minister Tamara van Ark. Kamerbrief met nieuwe aanvullende maatregelen om de COVID 19 uitbraak te bestrijden. Medische Zorg en Sport OCeW, Basis- en Voortgezet Onderwijs en Media, Sociale Zaken en Werkgelegenheid. Den Haag, 15 March 2020.

[3] Reukers DFM, van Boven M, Meijer A, Rots N, Reusken C, Roof I, et al. High infection secondary attack rates of SARS-CoV-2 in Dutch households revealed by dense sampling. Clin Infect Dis. 2021.

[4] (WHO) WHO. Clinical Management of COVID-19, 2020.

[5] Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill. 2020;25.

[6] Scheltinga SA, Templeton KE, Beersma MF, Claas EC. Diagnosis of human metapneumovirus and rhinovirus in patients with respiratory tract infections by an internally controlled multiplex real-time RNA PCR. J Clin Virol. 2005;33:306-11.

[7] Lassaunière R FA, Harboe ZB, Nielsen AC, Fomsgaard A, Krogfelt KA, Jørgensen CS. Evaluation of nine commercial SARS-CoV-2 immunoassays. medRxiv 2020040920056325. 2020.

[8] van Tol S, Mogling R, Li W, Godeke GJ, Swart A, Bergmans B, et al. Accurate serology for SARS-CoV-2 and common human coronaviruses using a multiplex approach. Emerg Microbes Infect. 2020;9:1965-73.

[9] Koopmans M, de Bruin E, Godeke GJ, Friesema I, van Gageldonk R, Schipper M, et al. Profiling of humoral immune responses to influenza viruses by using protein microarray. Clin Microbiol Infect. 2012;18:797-807.

[10] Martyn Plummer (2003). JAGS: A Program for Analysis of Bayesian Graphical Models Using Gibbs Sampling, Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC 2003), March 20–22, Vienna, Austria. ISSN 1609-395X.

[11] R Core Team (2021). R: A language and environment for statistical computing. Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

[12] Kruschke JK, Liddell TM. The Bayesian New Statistics: Hypothesis testing, estimation, meta-analysis, and power analysis from a Bayesian perspective. Psychon Bull Rev. 2018;25:178-206.

[13] R M. Statistical rethinking: A bayesian course with examples in r and stan2018.

[14] Piironen J VA. Comparison of bayesian predictive methods for model selection. Statistics and Computing. 2017;27:711–35.

[15] Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. UpSet: Visualization of Intersecting Sets. IEEE Trans Vis Comput Graph. 2014;20:1983-92.

[16] Madewell ZJ, Yang Y, Longini IM, Jr., Halloran ME, Dean NE. Household Transmission of SARS-CoV-2: A Systematic Review and Meta-analysis. JAMA Netw Open. 2020;3:e2031756.

[17] Fung HF, Martinez L, Alarid-Escudero F, Salomon JA, Studdert DM, Andrews JR, et al. The household secondary attack rate of SARS-CoV-2: A rapid review. Clin Infect Dis. 2020.

[18] Fu Y, Li Y, Guo E, He L, Liu J, Yang B, et al. Dynamics and Correlation Among Viral Positivity, Seroconversion, and Disease Severity in COVID-19 : A Retrospective Study. Ann Intern Med. 2021;174:453-61.

[19] Long QX, Liu BZ, Deng HJ, Wu GC, Deng K, Chen YK, et al. Antibody responses to SARS-CoV-2 in patients with COVID-19. Nat Med. 2020;26:845-8.

[20] Wajnberg A, Mansour M, Leven E, Bouvier NM, Patel G, Firpo-Betancourt A, et al. Humoral response and PCR positivity in patients with COVID-19 in the New York City region, USA: an observational study. Lancet Microbe. 2020;1:e283-e9.

[21] Moreira VM, Mascarenhas P, Machado V, Botelho J, Mendes JJ, Taveira N, et al. Diagnosis of SARS-Cov-2 Infection by RT-PCR Using Specimens Other Than Naso- and Oropharyngeal Swabs: A Systematic Review and Meta-Analysis. Diagnostics (Basel). 2021;11.

[22] Stanoeva KR, van der Eijk AA, Meijer A, Kortbeek LM, Koopmans MPG, Reusken C. Towards a
sensitive and accurate interpretation of molecular testing for SARS-CoV-2: a rapid review of 264
studies. Euro Surveill. 2021;26.
[23] Lee RA, Herigon JC, Benedetti A, Pollock NR, Denkinger CM. Performance of Saliva,
Oropharyngeal Swabs, and Nasal Swabs for SARS-CoV-2 Molecular Detection: a Systematic Review
and Meta-analysis. J Clin Microbiol. 2021;59.
[24] Mallett S, Allen AJ, Graziadio S, Taylor SA, Sakai NS, Green K, et al. At what times during
infection is SARS-CoV-2 detectable and no longer detectable using RT-PCR-based tests? A systematic
review of individual participant data. BMC Med. 2020;18:346.
[25] Cevik M, Tate M, Lloyd O, Maraolo AE, Schafers J, Ho A. SARS-CoV-2, SARS-CoV, and MERS-CoV
viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-
analysis. Lancet Microbe. 2021;2:e13-e22.
[26] van Kampen JJA, van de Vijver D, Fraaij PLA, Haagmans BL, Lamers MM, Okba N, et al. Duration
and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-
2019 (COVID-19). Nat Commun. 2021;12:267.
[27] Wolfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Muller MA, et al. Virological
assessment of hospitalized patients with COVID-2019. Nature. 2020;581:465-9.
[28] Bunyavanich S, Do A, Vicencio A. Nasal Gene Expression of Angiotensin-Converting Enzyme 2 in
Children and Adults. JAMA. 2020;323:2427-9.
[29] Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related Differences in
Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in Patients
With Mild to Moderate Coronavirus Disease 2019 (COVID-19). JAMA Pediatr. 2020;174:902-3.
[30] L'Huillier AG, Torriani G, Pigny F, Kaiser L, Eckerle I. Culture-Competent SARS-CoV-2 in
Nasopharynx of Symptomatic Neonates, Children, and Adolescents. Emerg Infect Dis. 2020;26:2494-
7.
[31] Weisberg SP, Connors TJ, Zhu Y, Baldwin MR, Lin WH, Wontakal S, et al. Distinct antibody
responses to SARS-CoV-2 in children and adults across the COVID-19 clinical spectrum. Nat Immunol.
2021;22:25-31.

Legends

Figure 1. Various transmission patterns of SARS-CoV-2 infection based on different assays and

specimen types collected at visits 1, 2 and 3 in households visualized in heatmaps. (A) households with

no transmission. (B) household with only transmission in adults. (C) heatmaps of household with

transmission in children and possibly also adults. Symptoms can be unrelated to a SARS-CoV-2

infection. On the left side, age category (A = adult and C = Child) of the participant is indicated. * Index

case. Blanks = not available/tested. Red rectangle: individual with only one test positive on one

timepoint. On the right side the Household ID (number) is indicated.

Figure 2. Symptom severity of COVID-19 index cases in the households. The GLM (generalized linear

model) revealed that there were less individuals in the "Severe" category than in the "Mild" category

(p=0.1), and that households in "Family transmission" category was overrepresented in the "Severe"

category (p=0.03). Maximum severity score of index is used.

Figure 3. (A) Overview of the (combination of) various positive SARS-CoV-2 infection diagnosis in the

242 participants of the study cohort for all various diagnostic assays or specimens. All available RT-

PCR and serology outcomes (Table 1) were incuded in these analyses.The black dots indicate a

positive test at any (or multiple) moment(s) during the study, except for 'Inclusion PCR NP+OP'

which was performed in index cases just before start of the study. The number of individuals with a

particular combination of positive tests are indicated in the top of the figure. The numbers at the

right after each test indicates the overall number of positive tests. (B) The number of individuals

with a positive or negative tests or missing data for serological and molecular (PCR) diagnostics

combined. In brackets the number of symptomatic or severe symptomatic individuals, respectively,

are indicated. Color intensity is related to the frequency.

Figure 4. Common SARS-CoV-2 infection dynamics patterns, based on the presence or absence of a

positive or negative RT-PCR or serological assay at visit 1, 2 and 3. Common patterns were named A

– F based on frequency, with the number of individuals (n) displaying the pattern indicated. The

median timing of onset of symptoms relative to visit 1 is indicated on the left. The black line

indicates whether on average symptoms were reported at visit 1, 2 or 3.

Figure 5. Dynamics SARS-CoV-2 infection diagnosis by RT-PCR in various specimens since symptoms

onset (dps). All available RT-PCR outcomes (Table 1) were incuded in these analyses. A) Predicted viral

load (Ct values RT-PCR) in relation to dps and specimen type. The shadow indicates the 95% Bayesian

confidence interval and the dotted lines indicate the prediction interval (variation over individuals). B

and C) Ct-value distribution at day symptom onset (intercept) and increase of Ct-value per day (slope)

in relation to age category (B) and different specimens (C). D) Average dps until when different

specimens have at least 90% detection probability.

Figure 6. Dynamics SARS-CoV-2 infection diagnosis by different serological assays. All available

serology outcomes (Table 1) were incuded in these analyses. Probability SARS-CoV-2-specific antibody

detection (A) and average dps from when the Wantai (Spike-specific IgM and IgG), microarray S1

(Spike-specific IgG) and microarray N (Nucleoprotein-specific IgG) assays have at least 90% detection

probability (B) for all ages. C) Probability SARS-CoV-2-specific antibody detection by Wantai,

microarray S1 and microarray N in adults and children D) Average dps from when microarray N has at

least 90% detection probability for adults and children. The shadows in (C) and (D) indicate the 95%

Bayesian confidence interval.

Table 1

	Visit 1														Visit 2	Visit 3
Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15 (range 14-21)	35 (range 28-42)
Start questionnaire	x															
Symptoms diary		x	x	x	x	x	x	x	x	x	x	x	x	x		
Final questionnaire																x
Blood (serum) - Wantai	222														210	149
Blood (serum) - Microarray S1 and N	230														221	208
Nasopharyngeal swab	183 [1]		53			46			51			55			221	
Oropharyngeal swab	187 [1]		53			47			51			54			230	
Oral fluid	219														206	189
Feces	222														226	195

Table 1. Schedule of administering questionnaires, symptom diaries and home visits for sampling. The numbers in the table indicate the amount of analyzed specimens in 242 participants. [1] A naso- and oropharyngeal swab was not collected for the index case at the first home visit, as these persons were already swabbed a few days before and tested SARS-CoV-2 positive.

Figure 1

C. Family transmission

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6