

# Information, Involvement, Self-care and Support - The Needs of Caregivers of People with Stroke: A Grounded Theory Approach

Elton H. Lobo<sup>1\*,2</sup>, Anne Frølich<sup>2,3</sup>, Mohamed Abdelrazek<sup>1</sup>, Lene J. Rasmussen<sup>4,5</sup>, John Grundy<sup>6</sup>, Patricia M. Livingston<sup>7</sup>, Sheikh Mohammed Shariful Islam<sup>8</sup> and Finn Kensing<sup>9</sup>

<sup>1</sup>School of Information Technology, Deakin University, Geelong, VIC, Australia;

<sup>2</sup>Department of Public Health, University of Copenhagen, Copenhagen, Denmark;

<sup>3</sup>Innovation and Research Centre for Multimorbidity, Slagelse Hospital, Region Zealand, Denmark;

<sup>4</sup>Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen, Denmark;

<sup>5</sup>Center for Healthy Aging, University of Copenhagen, Copenhagen, Denmark;

<sup>6</sup>Faculty of Information Technology, Monash University, VIC, Australia;

<sup>7</sup>Faculty of Health, Deakin University, Geelong, VIC, Australia;

<sup>8</sup>Institute for Physical Activity and Nutrition (IPAN), Deakin University, Geelong, Australia;

<sup>9</sup>Department of Computer Science, University of Copenhagen, Copenhagen, Denmark;

**\*Corresponding author:**

Email: [elobo@deakin.edu.au](mailto:elobo@deakin.edu.au) (EL)

1 **Abstract**

2 **Background:** Globally, stroke is a leading cause of death and disability, with most care  
3 undertaken by caregivers who are generally family and friends without prior experience of  
4 care. The lack of experience or unpreparedness results in feelings of uncertainty, burnout,  
5 anxiety, burden, etc. Hence, it is necessary to identify the needs of caregivers to support them  
6 in their caregiving journey better and improve the quality of care delivered.

7 **Methods:** The study employed a grounded theory methodology that utilizes information  
8 gathered from literature reviews and social media to represent the needs and create a storyline  
9 visually. The storyline is further refined and evaluated using an online survey of 72  
10 participants recruited through online stroke caregiving communities.

11 **Results:** The study identified four core categories of needs, i.e., information, involvement,  
12 self-care and support that need to be delivered at appropriate times to ensure the caregivers  
13 feel prepared to assume their role of care and are supported throughout the care trajectory.

14 **Conclusions:** There is a need to create a caregiver-centered approach in stroke recovery to  
15 ensure limited obstruction to care and reduced uncertainty in recovery. Moreover, through the  
16 inclusion of caregivers in the recovery process, it may be possible to reduce the burden of  
17 care to the caregiver and ensure the satisfaction of the healthcare system throughout recovery.

18

19 **Keywords:** Needs, Experience, Stroke, Caregiver, Information, Involvement, Self-Care,  
20 Support

21

22

23

24

## 25 Introduction

26 Globally, there are over 13.7 million new incident cases of stroke each year, with more than  
27 116 million years of healthy life is lost due to stroke-related deaths and disabilities [1]. With  
28 the recent advancements in the medical field, the stroke mortality rate has decreased [2],  
29 resulting in more than 80 million people currently living after experiencing a stroke [1]. Of  
30 these survivors, up to 50% of individuals are chronically disabled [3], with family members  
31 and/or friends assuming the role of caregiver to provide care [4].

32 The role of caregiving is arguably the most important aspect of stroke recovery. However,  
33 caregiving is a complex concept and is dependent on the condition of the survivor (i.e., age,  
34 impairments and living situations), individual characteristics (i.e. personal beliefs, coping  
35 styles and social expectations) and the relationship between the survivor and the caregiver [5,  
36 6]. Moreover, since the occurrence of stroke is sudden [5], caregivers often have to adjust to  
37 the diagnosis [7], assume new roles and responsibilities [8] and face the challenge of  
38 becoming a caregiver with limited or no preparation [9]. This leads to a substantial increase  
39 in caregiver strain or burden, leaving the caregiver to feel abandoned and unsupported [10].

40 As a result, several researchers have attempted to explore the needs of stroke caregivers at  
41 different stages of stroke. For example, a systematic review by Luker, Murray [11]  
42 highlighted the need for stroke caregivers during in-patient rehabilitation. At the same time, a  
43 longitudinal study by Tsai, Yip [12] described the changing needs of stroke caregivers at  
44 different stages of the care trajectory.

45 Although numerous reviews are available that understand the caregiver's needs and  
46 experiences during recovery, an improved understanding is necessary to create effective  
47 interventions to support the caregiver during the transition into the caregiver role. Hence, this  
48 study utilizes a grounded theory methodology that combines data acquired from literature and

49 social media sources to identify needs specific to the caregiver and refine and evaluate these  
50 needs using an online-based survey to ensure comprehensiveness.

51

## 52 **Materials and Methods**

### 53 **Study Design**

54 The study was designed and conducted based on the grounded theory methodology. The  
55 grounded theory aims to generate a theory grounded within data [13]. This approach allowed  
56 for investigating a particular phenomenon in diverse environments to develop an explanatory  
57 theory [14].

58

### 59 **Data Collection**

60 The three-phase study was conducted from September 2020 to May 2021. Phase one and two  
61 were completed in English, while phase three was conducted in English and Danish. Data  
62 collected in Danish was translated into English by another author (AF). Five authors (EL,  
63 MA, FK, AF and SI) coded the data collected in English based on three essential coding  
64 schemes; initial, immediate, and advanced coding. To ensure anonymity and confidentiality,  
65 personal information related to the participant was excluded from the study.

66

### 67 **Phase 1: Literature Review**

68 The first phase involves systematically reviewing the literature to determine the needs and  
69 inform the type of caregiver groups to be involved in the study.

- 70 • **Selection Criteria:** Qualitative studies that included the needs, experiences, and  
71 perspectives of caregivers supporting people living with stroke at their home was  
72 eligible for inclusion. Articles were excluded if they (i) were not available in English,

73 (ii) were protocols or abstracts, (iii) solely reported quantitative data, and (iv) reported  
74 needs, experiences, and perspectives within the hospital. Studies were excluded if they  
75 did not acquire data directly from the caregiver. Furthermore, reference lists of included  
76 reviews were added to this study to ensure comprehensiveness.

77

78 **Table 1. Search Terms**

|                    |  |
|--------------------|--|
| <b>Disease</b>     | "cerebrovascular disorders" OR "basal ganglia cerebrovascular disease" OR "brain ischemia" OR "carotid artery diseases" OR "intracranial arterial diseases" OR "intracranial embolism and thrombosis" OR "intracranial haemorrhage*" OR "stroke" OR "brain infarction" OR "cerebrovascular accident" OR vasospasm OR "vertebral artery dissection" |
| <b>Concept</b>     | "transitioning from hospital to home" OR "transition home" OR "discharged home" OR outpatient OR transition  |
| <b>Methodology</b> | "Qualitative Research" OR "Cohort Studies" OR "Observational Study" OR "Focus Groups" OR semi-structured OR semistructured OR unstructured OR informal OR in-depth OR indepth OR face-to-face OR structure OR guide OR interview* OR discussion* OR question?aire*   |
| <b>Study Focus</b> | "Patient centered" OR patient centred OR Patient-centered OR patient-centred OR "patient satisfaction" OR "consumer satisfaction" OR "caregiver focus" OR "carer focused" OR "caregiver centred" OR "caregiver centered" OR "caregiver satisfaction" OR "carer support" OR "carer centered" OR "carer centred" OR "carer satisfaction"             |

79

80 • **Data Sources and Searches:** A systematic search was conducted on five electronic  
81 databases: Medline, Embase, CINAHL, PsycINFO and Web of Science, from inception  
82 to September 2020 for keywords related to stroke caregiving needs, experiences and  
83 perspectives (**Table 1**). After removing duplicates, the primary author (EL) reviewed  
84 titles and abstracts and was supervised by another author (MA). The potentially relevant

85 articles were downloaded in full-text and independently reviewed by two authors (EL  
86 and FK). All authors discussed any discrepancies until a consensus was achieved.

87 • **Data Extraction and Representation:** User needs data was extracted from thirty-one  
88 accepted articles by the primary author and was subsequently reviewed by another  
89 author for accuracy. The data extracted were represented using a concept mapping  
90 approach to visually illustrate different concepts into meaningful connections [15].

91

## 92 **Phase 2: Social Media Analysis**

93 The data extracted from the first phase led to further data gathering. Phase two data collection  
94 was based on user posts on popular information-based social media platforms (i.e., Facebook  
95 and Twitter). As the data analysis proceeded, theoretical sampling was implemented to decide  
96 how to collect data based on emerging theories and categories.

97 • **Selection Criteria:** Social media posts were included if they (i) were available in  
98 English, (ii) were made publically available, (iii) included stroke caregiver discussions,  
99 and (iv) discussed stroke caregiver needs during care and recovery. The posts were  
100 excluded if they (i) included discussions from patients, clinicians, or community  
101 administrators, (ii) were not available publically, (iii) discussed promotions of products  
102 or services, and (iv) were not related to the needs of stroke caregivers.

103 • **Data Sources and Searches:** A systematic search was conducted on popular social  
104 media platforms (i.e., Facebook and Twitter) from inception to January 2021 to identify  
105 relevant stroke communities using keywords from popular internet searches from  
106 December 2010 to January 2021 (**Table 2**).

107 • **Data Extraction and Representation:** The primary author (EL) extracted user posts  
108 from identified stroke communities using a Python-based web-scraping tool. These  
109 posts were filtered by the primary author (EL) based on the selection criteria and were

110 reviewed by two author (AF and SI). The 1,017 accepted posts were analyzed with the  
111 data from the previous phase to develop the initial storyline.

112

113 **Table 2. Keywords used in Social Media Searches**

| Keywords   |  |
|--|--|
| <ul style="list-style-type: none"><li>• Stroke</li><li>• Stroke Care</li><li>• Stroke Recovery</li><li>• Apoplexy</li><li>• Cerebrovascular Accident</li></ul> | <ul style="list-style-type: none"><li>• Stroke Unit</li><li>• Traumatic Brain Injury</li><li>• Lacunar Infarct</li><li>• Stroke Medication</li><li>• Aphasia</li></ul> |

114

115 **Phase 3: Caregiver Survey**

116 In phase three, theoretical sampling is utilized to refine the storyline to facilitate an improved  
117 theory integration process. This process would determine the suitable area for integrating the  
118 generated theories using an anonymous online survey.

- 119
- **Ethics:** Prior to phase three data collection, the study received approval from the Deakin University Human Research Ethics Committee (HREC): 2020-225.
  - **Selection Criteria:** Caregivers of people with stroke receiving active or palliative treatment over the past five years. The caregivers were excluded if they were aged below 18 years and did not understand English or Danish language(s).
  - **Participant Recruitment and Study Setting:** Seventy-two caregivers were recruited (Table 3) through stroke health organizations, caregiver organizations and social media communities from March 2021 to May 2021 using a recruitment flyer. All participants were informed about the research and provided informed consent to participate in the anonymous online survey.
- 128

129 • **Data Representation:** The participants received survey questions based on the concept  
 130 map and the storyline identified in the previous two phases to determine its relevance to  
 131 the stroke caregiving needs. The online survey included questions related to the type of  
 132 resources and support required, and aspects that are challenging to the caregiver in their  
 133 daily activities.

134

135 **Table 3. Participant Demographics**

| <b>Caregiver</b> |                  |    |
|------------------|------------------|----|
| Gender           | Male             | 13 |
|                  | Female           | 59 |
| Age              | 18-27            | 2  |
|                  | 28-37            | 11 |
|                  | 38-47            | 8  |
|                  | 48-57            | 18 |
|                  | ≥58              | 33 |
| Living Location  | Metropolitan     | 36 |
|                  | Smaller City     | 11 |
|                  | Rural Area       | 25 |
| Education        | Primary School   | 1  |
|                  | Secondary School | 10 |
|                  | Diploma          | 5  |
|                  | Bachelor's       | 30 |
|                  | Master's         | 14 |
|                  | Doctoral         | 2  |
|                  | Other            | 10 |
| Relationship     | Grandparent      | 2  |
|                  | Parent           | 15 |
|                  | Partner          | 50 |
|                  | Child            | 5  |
| <b>Survivor</b>  |                  |    |
| Gender           | Male             | 54 |
|                  | Female           | 18 |
| Age              | 18-27            | 1  |
|                  | 28-37            | 3  |
|                  | 38-47            | 2  |
|                  | 48-57            | 15 |
|                  | ≥58              | 51 |

136

137

## 138 **Data Analysis**

139 The data extracted in each phase was analyzed based on a constant comparative analysis  
140 involving three prominent coding processes, i.e., initial, intermediate and advanced coding.  
141 All data collected was organized and managed using NVivo 12. Based on the constant  
142 comparative method, the “core category” of the grounded theory is identified from different  
143 parts of the data, including emerging codes, properties, dimensions and categories as well as  
144 some parts of the data which are compared with the identified themes to identify variations,  
145 differences and similarities [16]. This means that the incoming data or new codes in each  
146 phase were compared with those from either the same or previous phase(s) to generate a  
147 theory.

148 The raw data is broken down into fragments and labeled to form new codes in the initial  
149 coding process. The process of labeling stems from conceptual actions or observable needs.  
150 The data from the initial coding is condensed to create a more focused coding. At this stage,  
151 the codes identified formed significant categories. The categories undergo constant  
152 comparative analysis until the advanced coding phase. In the advanced coding phase, the  
153 high-level codes look to address the theory, i.e., the needs of stroke caregivers during care  
154 and recovery.

155

## 156 **Results**

157 Findings highlight four critical needs identified by analyzing the data of individual  
158 caregivers’ experiences across the stroke disease trajectory. The four needs are information,  
159 involvement, support and self-care. **S1 File** presents the identified needs and describes  
160 individual requirements for individual needs.

161

## 162 **Need 1: Information**

163 Most caregivers emphasized the importance of information to reduce uncertainty and be  
164 better prepared in stroke caregiving. Due to the sudden onset of stroke, most caregivers find it  
165 challenging to understand its occurrence, leaving them shocked and helpless due to this  
166 experience.

167

*“It was very hard to come to terms with what just happened.. there was  
no warning.”*

*“I don’t know how to get through these dark days.. I feel isolated and  
alone”*

168

169 Due to the uncertainty followed by the onset of stroke and the inability to understand the  
170 cause of the disease, many caregivers often searched for an explanation to cope with the  
171 disease.

172

*“Frankly speaking, I’m quite confused now, because it is a sudden  
thing...ya, of course, this is my first priority (to know more about the  
condition)...we have to learn a lot of things? I don’t know” [17]*

173

174 Most caregivers relied on healthcare professionals such as physicians, nurses and therapists  
175 for information to help cope and prepare them to navigate through a ‘completely foreign  
176 world’ [18]. However, medical professionals primarily provided the stroke survivor with  
177 information, with little consideration to the caregiver. Moreover, some caregivers reported  
178 poor physician service attitudes during post-stroke discussions with the clinicians [19].

179

*“I don’t think they told me anything, I was just left out in the cold ... I didn’t have a clue what was going on ...”* [20]

*“His attitude was somewhat arrogant. He won’t answer everything that we asked. I don’t know if it is because of our manner or what... I feel he did not tell us much information. We need to try on our own to understand it.”* [19]

*“I know as a family member or a patient we have the right to know about what the treatment is for and the medication for. You should give me an explanation. But he didn’t. Every time when I asked him, his attitude was always like this way...Then he answered me, “Are you the doctor or am I? You don’t trust me?!”* [19]

180

181 Despite the limited information and communication with the healthcare professionals, a few  
182 caregivers mentioned that the information provided to them by the healthcare professional  
183 was established based on brochures and pamphlets [21, 22], which were not particularly  
184 useful [17]. One caregiver mentioned, *“Honestly, a layperson might not be able to*  
185 *understand those leaflets and they won’t read them! So I think the leaflets are not*  
186 *worthwhile”* [19]. The lack of information provided to the caregiver during recovery left the  
187 caregiver distressed.

188

*“They [medical professionals] weren’t forthcoming with information of things to do...you feel like you don’t know what to do ...it felt pretty*

*helpless” [21]*

*“With the lack of information, I really don’t know what to expect.”*

189

190 Most caregivers discussed the need to have sufficient information provided at the correct  
191 times. For example, they requested information about the disease, cause, effects, signs and  
192 symptoms, risk factors, prevention techniques, healthcare procedures, home care procedures,  
193 treatment and recovery options, locally available services, medications, and aid/tools as  
194 shown in **S1 File**. The caregivers preferred the information to be delivered clearly in ‘layman  
195 terms’.

196

*“I was told by the hospital that they are removing him [stroke survivor]  
from the high dependency ward to the general ward where beds were  
available. I am not sure what this means. I need someone to explain to  
me in layman’s terms what this means.”*

*“My wife and I care for my grandmother, and we have no healthcare  
background, which has resulted in a very tough learning curve with the  
information provided to us.”*

197

198 The caregivers also mentioned the importance of timing of information post-stroke, as the  
199 information needs are said to change over time [23]. This is because caregivers found it easier  
200 to recall information based on their current setting than provided at the time of stroke or  
201 discharge [21]. Further, caregivers also discussed the need to have personalized information  
202 based on individual needs based on personal experiences delivered in several different ways

203 based on their literacy and health literacy levels. Some examples discussed by the caregivers  
204 for the delivery of stroke information include (i) oral discussions and demonstrations by  
205 healthcare professionals and (ii) visual and written methods. Oral and verbal discussions were  
206 particularly valuable methods for information delivery. They allowed the caregivers to  
207 generate skills at the beginning to help support the person affected with stroke.

208

*“My preference would be the first-hand experience. I mean to have the nurses, doctors, and medical staff actually sharing with me: ‘Your father has stroke, so the best thing that you can do now is to...’ Preferably, they can share some advice on things that I can carry out at home to look after my dad...”* [17]

*“these [care practices] are not difficult. They just teach me one time then I can do it”* [19]

*“They played a videotape for us and some papers...such as how to prevent a fall, how to care for a stroke person...but I still don’t understand after reading those papers and watching the videotape. I needed a real situation to increase my understanding”* [19]

209

210 Besides oral and verbal discussions, caregivers preferred to receive visual and written  
211 information if someone explained it more comprehensively. Furthermore, visual and written  
212 information allowed the caregivers to access the information later should the need arise with  
213 no other form of support available [18].

214

*“They do have some brochures and pamphlets but those are not really— I won’t say not up-to-date— but those just explain the stroke condition very basically... What I reckon would be really helpful is to have someone brief us and go through them with us.” [17]*

*“It would have been nice to have somebody sit down with me and say this is what’s happened, this is why it’s happened, this is what you can expect. Okay, so it is there in the leaflets but you just kind of flick through the leaflets” [20]*

*“But the fact that she, instead of just handing me information, she sat down and went through each point even though I was so tired and I’m going to remember it so much more. When I need to, I’ll know in the sheets that she gave me where to go to look for the information.” [23]*

215

## 216 **Need 2: Involvement**

217 Caregivers discussed the need to be actively involved in the care process, as it added to their  
218 sense of preparation and confidence. Moreover, it would allow the caregiver to ask questions  
219 leading to a good learning experience instead of providing the clinicians with progress  
220 updates.

221

*“I think it’s good for the family to be involved and to know what’s going on. I realize that when you are dealing with the public some of them*

*don't know what you're talking about and some of them don't care. But I think that the majority of us want to know what's going on ... and if we don't understand it immediately, then we need to have it explained to us in some terms that we do. We all need to be involved in our own health care, and we need to be involved in whatever needs to happen for people for whom we're taking at least some responsibility for" [24]*

222

223 Some caregivers expressed that they saw the caregiving role for their parents as an  
224 opportunity to give back as retribution for love and care when they were a child. In contrast,  
225 some caregivers did it out of an expression of loyalty, religious beliefs or lack of confidence  
226 in the survivor to take care of themselves. Hence, leading to a significant emotional impact  
227 on the caregiver's decision to provide care.

228

*"I just thought that she had looked after me since I was a baby. When she can't help herself, I have to help her." [25]*

*"I feel that it's my duty. No one is free to take care of her, so I have to do it. If I get a job or become more stable than this, I might hire someone to take care of her. I am 28 years old now. I need to go to work. I feel that it's my duty to look after her so I do it. In my opinion, no one can do it better than me."*

*"I can't stop myself sometimes in letting her know when she makes the error. It is just to help her understand that she still does need help for things."*

229

230 In addition to the emotional factors, caregivers felt that by being more engaged in care they  
231 would be able to get reassurances of their fears and worries through encouragement derived  
232 from feedback from medical professions during recovery [23]. Moreover, it would allow  
233 them to share more information with the medical professional regarding the pre-stroke life of  
234 the survivor required for them to make informed decisions based on the survivor's  
235 preferences for care.

236

*“By being involved in care I feel reassured. I can engage with providers discuss my fears and worries and share with them the needs of my loved one. This allows me to carry on in my daily activities and feel a part of the recovery.”*

237

238 Despite caregivers expressing their needs to be involved in care, most caregivers feel left out.  
239 Caregivers expressed their interactions with medical professionals as uninvolved or  
240 disconnected [9]. When approached by caregivers, medical professionals either did not  
241 provide enough information or expressed that they were “too busy”, which left them  
242 uncertain [9].

243

*“...but we do not get much information [about] his condition. There is no update. You see, I have to go and ask them. They say, ‘The doctor will come but don’t know what time.’ Do you understand?” [17]*

*“The communication between me and the hospital leaves a lot to be desired. Every day I wake up and I’m still in the twilight zone.”*

244

245 The uncertainty led to numerous concerns regarding the rehabilitation received by the

246 survivor and the caregiver's own preparation to assume the full-time role as a caregiver. One

247 caregiver mentioned

248 *"Well one of the things that's uncertain is not knowing when she's coming home and also not*

249 *knowing what level of care she'll need"* [9]. This led to the caregivers developing negative

250 emotions.

251

*"...we're still very shocked... we're trying but we're not coping well...our family is going haywire..."*

*"I am scrambling to get a caregiver...I am so stressed...In fact, we are very worried..."*

*"I'm probably still coping, but I still get stressed up sometimes...because he's never been so sick...all of a sudden..."*

*"...at least he can eat...if he needs tube feeding, then we're in trouble...might need to send him to a nursing home."* [17]

252

253 Furthermore, the lack of communication with healthcare professionals has led to improper

254 treatment for the survivor resulting in the caregiver being frustrated and affecting the

255 survivor's progress during their recovery.

256

*"...the hospital never answered my questions about care package, they didn't tell the home that my mum requires one on one care - they also didn't tell us that she can get up and walk on her own. Her speech is very muddled, but some make sense. I asked the hospital three times*

*over a two week period to provide me with details of the care package and they never got back to me. The home has called emergency doctor out, and she's said that my mum shouldn't have been brought to this home, as clear she needs further physio, speech and ot"*

*"I have found the lack of treatment pretty appalling ... I do feel quite angry ... she needs more treatment ... she could make far better progress if she were getting better treatment." [26]*

257

258 The lack of preparation and communication was not the only reason for the caregivers'  
259 involvement in care. One caregiver mentioned that the survivor would constantly 'push' them  
260 away during recovery. While another caregiver mentioned that her husband was 'refusing  
261 treatment'. The lack of motivation amongst the survivor causes friction with the caregiver  
262 leading to depression and withdrawal from care.

263

*"... he [survivor] has been going to speech therapy off and on ... He is stating that he does not want to go anymore. Typically, when he doesn't want to do something, he is pretty ornery and stubborn about it. Should I push the issue?"*

*"Sometimes getting along and sometimes arguing, arguing just because he doesn't want to do these exercises ...and he doesn't want to do it he just sits there and I say come on let's go let's get up let's do something...and I feel so frustrated sometimes and I get angry and that's what we argue about..." [27]*

264

265 Regardless of these issues, caregivers have provided care to the survivor in planning, caring,  
266 managing, motivating, and goal-setting activities. These activities are critical in ensuring the  
267 survivor feels supported throughout recovery. Caregivers have employed a range of strategies  
268 in their efforts to care for the survivor. For example, some caregivers considered taking one  
269 day at a time and did not plan for the future. In contrast, other caregivers created ‘meaningful  
270 goals’ for the survivor to create ongoing reassurances of their recovery. Furthermore, the  
271 caregivers also set up medical and follow-up appointments and participated in physical  
272 therapy to add to their sense of preparation and confidence.

273

*“I train her cognition and memory. I write some words or numbers and show it to her. Such as I write our four kids’ names on four cards then show her first. Later I pick one and ask her what’s the name on it and which child’s name....things like this. Those strategies pop out of my mind. I use those skills to train her cognition” [19]*

*“I often talked with her, very often.... I played with her the childhood games she taught us. And I sang her favorite songs to her..... One day I asked her: Mom, do you want some Zong Zi (a pyramid-shaped mass of glutinous rice wrapped in bamboo leaves)? She answered, “No”...And on July 4th, she could say my aunt’s name. Now she can sing her favorite songs with us...At first, she only can sing in her mouth, but now she has the voice. She cannot recognize us, but she can sing those songs!” [19]*

*“I faced him and put his arms on my shoulder; then I walked backward to guide him to walk. His left foot could not move, so I used my right foot to hook his left foot and pulled his leg forward. I did this during his way to the bathroom and back to his bed.” [19]*

274

275 Caregivers in stroke also discussed methods to be considered to be more involved in care.

276 The primary consideration is for healthcare professionals to consider the caregiver as the

277 ‘central role’ in the survivors’ rehabilitation [28]. This can be achieved by creating a close

278 collaboration with clear communication between the caregivers, patients and healthcare

279 professionals. The preferred mode of communication amongst caregivers was face-to-face

280 communication to build a trusting relationship, which may be critical during the counseling

281 process [29]. For example, one caregiver mentioned, *“I need someone who gives me the time*

282 *I need and where I can also show emotions” [29].*

283 Other than efficiently communicating with the healthcare professional, the caregiver

284 preferred being a part of the decision-making process for the next step of rehabilitation and

285 receiving training from a healthcare professional to provide improved care. The inclusion in

286 the decision-making process allowed them to share their opinions, such as the type of care,

287 availability (or time) to attend care and survivors' preferences for care. By ensuring tailored

288 or individualized care, caregivers felt more satisfied and confident in their ability to support

289 the survivor. Moreover, it was considered to be ‘very important in maintaining stroke

290 survivor health’.

291 As for care training, most caregivers stated that they would ‘require training to help their

292 family member specifically those skills needed for transfer and at home after discharge. One

293 caregiver explained, *“I felt so hard because I never bumped into this before. I never faced a*

294 *seriously sick person. This is my first time. I had no idea what I have to do for her....but if I*

295 *want to take care of her, I must learn”* [19]. The caregivers believed that hospitalization  
296 would be the best period for them to learn from experts (i.e., healthcare professionals) and  
297 gain hands-on experience. Moreover, it could allow them to gain some feedback during care.

298

*“if my father is referred to the rehab centre, I could probably be there to watch them and learn a lot...on-the-job training.”*

*“...or the rehab centre. I don’t think the hospital will be able to help [because] they are so busy.”*

*“The community hospital will train the maid at the bedside and ask her to do [the care tasks]. She [survivor] will need to stay there for one to two months. That should be enough time to train the maid.”* [17]

*“...we used the chance to learn. We were in learning process, so we had to do hands-on it. If we learned it, we might feel less pressure. We could not start to learn it after going home. So in the hospital, if we are unable to do it, we can ask for help from them [healthcare professionals]”* [19]

299

### 300 **Need 3: Self-care**

301 The physical and psychological impact of caregiving was common amongst most post-stroke  
302 caregivers. Caregivers reported having feelings of anxiety, loneliness, fatigue, frustration,  
303 burnout, fear, social isolation and helplessness during their role as a caregiver due to the  
304 alterations of their lifestyles, life status and routines in a negative way.

305

*“Ok I admit I am defeated ... I can’t keep the happy me anymore or keep*

*up with everything. Maybe just tired or trying to deal with too much, but I do not have a way to fix it or to even try to make a way to make it work. I have bitten off more than I can deal with... I need to be here & go to work & also get chores done & also be able to try & get me time. It's just not possible."*

*"We are exhausted, drained & overwhelmed feeling. It's a lot to take in!!"*

306

307 Most caregivers felt that caregiving's physical and psychological impact was the lack of  
308 independence to engage in other activities that previously ensured the fulfillment of life.  
309 Before the stroke, caregivers participated in activities such as work, social gatherings,  
310 volunteer work, religious activities, vacation and trips, which were halted due to the  
311 dependence of the survivor on the caregiver.

312

*"Since he's [survivor] come home I've not really gone out very much. Normally I would just go out and do whatever, but I haven't been able to do that since he's come home from the hospital." [20]*

*"I've quit doing gym. I have virtually given myself up. I only live for my mother and my husband here at home, and don't do anything else." [30]*

313

314 Caregivers often put their dreams on hold as they were responsible for providing constant  
315 care for the survivor. For example, one caregiver decided not to commit to future  
316 employment opportunities, while another caregiver couldn't go on a vacation with their loved

317 ones. The reason for putting their ‘lives’ or ‘dreams’ on hold was the survivors needs and  
318 uncertainty during recovery. However, giving up their dreams was not always done readily  
319 [31].

320

*“We put a lot of our dreams on hold, because we don’t see how we can go on vacation. We think about who can watch our loved one, or how can I take him/her [survivor] with us.”*

*“I’ve been angry. I resented having to quit the full time job where I was making good money and I loved the job...I didn’t want to have to do that” [31]*

321

322 Work and social participation were two critical themes discussed by caregivers to ensure  
323 fulfillment of life during their care journey. Currently, most caregivers have either given up  
324 their work or have reduced their work hours to accommodate survivor care. Caregivers who  
325 have quit their jobs feel ‘homebound’ or ‘isolated’ [32, 33], while those who have managed  
326 to continue their work report having difficulties in maintaining work and care responsibilities.

327

*“this [work] has taken a big toll on me – I am juggling between my job, my husband and children, my husband’s business and caring for my mother – sometimes I can’t see any light at the end of the tunnel” [33]*

328

329 In terms of social participation, stroke has altered the lives of the caregivers in such a way  
330 that they can no longer participate in the activities that give them ‘joy’ or ‘happiness’. One  
331 caregiver discussed “not having enough time for himself”; while another caregiver mentioned

332 that they felt “*restrained from going out because of the caring activities and housework*”  
333 [34]. The inability to participate in social activities strains the relationship between the  
334 caregiver and the survivors due to the mental and emotional aspects associated with being  
335 isolated at home and the loss of independence.  
336

*“I used to have lots of outing activities eight or nine years ago. Now I  
feel so alone” [34]*

*“The kids used to love camping, but now they have to go with friends,  
my wife can’t do that. They also love going to the movies but my wife is  
sensitive to light so we don’t go anymore. One of us has to stay with my  
wife at all times, we take turns. I miss the spontaneity; the stroke  
restricts our lives...” [22]*

*“My dilemma is finding me time without making them feel left out, which  
is what's happening. It's straining our relationship because of the mental  
and emotional toll... I'm trying to be compassionate, forgiving and  
understanding; to see things from their perspective but I'm almost burnt  
out.”*

337

338 Caregivers also discussed practices currently employed for self-care. The practices involved:  
339 (i) distracting their partners by tempting them to engage in other activities, (ii) participating  
340 in relaxing activities such as cooking, going to church or other household activities, and (iii)  
341 taking time off during the survivors’ rehabilitation appointments. Caregivers believed that  
342 these practices allowed them to ‘take a break’ or ‘get time for themselves’.

343

*“And I do [activities of my own], because my activities also help me get distracted, because it happens that I stress myself with my mother's illness. It is worse, when I get sick, though. So, what I do is do my normal activities as I did before” [35]*

344

345 Despite developing practices to ensure self-care, caregivers often report wanting more time to  
346 engage in leisure and self-care activities, which has contributed to poor health. To further  
347 improve self-care practices, caregivers mentioned potential considerations to improve  
348 physical and psychological well-being. Caregivers discussed the need to take breaks from  
349 physical care, and either ‘going back to work’, ‘travel’ and/or ‘participate in recreational  
350 activities’ as illustrated in **S1 File**. Some described this situation as ‘out of sight, and out of  
351 mind’. However, it was only possible for the caregiver to take a break when they were able to  
352 share responsibilities with other family members and/or friends during recovery.

353

#### 354 **Need 4: Support**

355 Caregivers discussed the importance of receiving support – in terms of resources, services  
356 and finances to sustain their ability to provide quality care (**S1 File**). As caregivers mainly  
357 were family members or friends of people living with stroke without prior experience in  
358 providing care, they expressed the need for help or support, especially from the healthcare  
359 system.

360

*“the system could have provided more psychological and emotional support; also a long-term follow-up of how we are doing would be*

*important” [33]*

361

362 The lack of care support can affect the caregiver mentally, leading to uncertainty, fear,

363 loneliness, anxiety, anger and frustration leading to financial and relationship impact.

364

*“How am I going to cope? ... I don’t know I really don’t know ... when the worry is on my shoulders, that’s all I’m thinking about ... I don’t think I could take the worry. I don’t—maybe I’ll have to.” [26]*

*“... I am trying to stay positive for him [survivor] but I don’t have any answers and I am feeling so alone and anxious. I just want to know it will get better.”*

*“I feel so overwhelmed at times, feel like I’m doing it all wrong, when it comes to important stuff like our finances or dealing with Dr. or the government I feel so inadequate. just scared I’m gonna screw up something important that can’t get fixed. Only been doing this for 8 months. Not getting much help .”*

365

366 Caregivers discussed that uncertainty was due to the lack of understanding of the disease and

367 future life. For example, one caregiver mentioned, *“And what the future will bring nobody*

368 *knows. Sometimes I think it can come back, because they told me, it could happen, another*

369 *stroke. Because most people come back with another stroke” [36]. Further, the uncertainty*

370 *towards the future could result in fear in terms of recurrence of stroke, falls, finance and*

371 relationships. As a result, most caregivers choose to be at home to observe and care for the  
372 survivor, contributing to isolation and loneliness.

373

*“... now I have to hurry even to go to the supermarket because I’m afraid ... to leave him ... He may have a CVA again at any time, as the doctor said. Like the first time it happened. He fell to the floor, urinated on himself, he fell in the middle of the night. He fell off the bed, and I didn’t even notice. So, I’m afraid. So, I hurry to supermarket, but my mind stays here ..” [30]*

*“I am struggling with feeling very alone. My husband had a stroke in May and it's getting to the point where I don't feel like a wife or partner anymore. I'm a caregiver 24/7.”*

374

375 During caregiving, caregivers have also discussed feeling of anger and frustration, which is  
376 contributed by issues related to care, financial, relationship, emotional and social hardships.

377

*“Sometimes I think I may be going crazy! We are going on almost 3 years post-stroke and it's so hard to deal with still. I lose my temper so easily now, full of anxiety and anger which I end up taking out on my husband who has had a stroke. I feel so bad after it happens and it's not any particular thing that sets it off. I'm supposed to be the strong one for us - hes' the survivor - I'm the one angry and emotional all the time now?? It's almost as if I'm experiencing his symptoms if that makes sense. I should probably speak with someone but I have such crappy*

*insurance and can't afford another Dr. bill.....ugh!"*

*"Ever get to the point where you have so much on your plate that you cry and laugh at the same time and then have nothing but oreos for dinner and want to throw your hands up and scream at the universe to back the f\*\*k off?!Aaaand I go back to my wine."*

378

379 The financial impact is a common stress factor amongst caregivers. Most caregivers have  
380 stopped working altogether to support their loved ones, which made them entirely dependent  
381 on financial support from other sources such as family members, insurance, pensions, etc.  
382 Some caregivers described that these sources of support were not sufficient to cover the  
383 survivor's expenses and as a result resumed working. However, this contributed to increasing  
384 workload, caregiver burnout, and guilt due to the inability to provide continuous care.

385

*"This [the financial situation] has most affected me, because now I have to wait for my son to give me money, while before I always solved all of the problems related to the household on my own." [35]*

*"My God, what will my life be like now? I won't work." And then I started thinking: "What will I do if I don't work?"*

386

387 Caregivers also discussed the impact of caregiving on their relationships with the survivors  
388 and other family members. For example, one caregiver mentioned, *"I wasn't able to be there  
389 as much as I really, really wanted to be, because obviously I was seeing to [partner's] needs  
390 ...we would have had a much more hands-on relationship with the four grandchildren ..."*

391 [28]. The caregivers' inability to maintain relationships was due to their failure to balance the  
392 different roles. One caregiver discussed that their partners were upset with them for being the  
393 second priority in their relationships and would often “complain” about the time and effort  
394 taken by the caregiver in caregiving. While another caregiver described that the lack of  
395 support received from their partner in the caregiving role resulted in them working for “two  
396 people”, which resulted in friction or conflict and thereby increasing their burden.  
397

*“With my partner who is far away, that’s really hard because he feels like he’s second most important.... because a lot of my time is spent at home with my parents and helping my dad. So that’s kind of suffering”*  
[31]

*“At the beginning, when my father got sick, ... I argued with my husband and he left. We were separated for almost a year. Then, he came back. Then, he decided to separate again because I was too nervous. He even said that he could understand me, but couldn’t take that situation anymore ... It’s difficult! You see. I had my house, my life there. My husband and I. Then, we left there and moved here, thinking that things would be a certain way, and now that we are here, he has to be everyone’s father ... he’s playing a role that shouldn’t be his. My husband is a very good man, but who can take all this?”* [30]

*“All that being said....we have been raising our 3 grandchildren as well. They have been a handful for me for 6 years, all being neglected in the past. So they are special needs as well, each with different needs. They*

*are state kids, wards of the court, and we fought for them to come to us instead of being separated and adopted out, or placed in-group homes/foster homes. Majority of the work fell on me, counseling, doctors, court meetings...etc.”*

*“There was maybe just a little bit of tension initially with my children because they’re small and don’t understand the seriousness of the situation and I guess they couldn’t understand why I always had to go to the hospital” [31]*

398

399 The impact of caregiving, i.e., the mental, financial, relationships and care described above,  
400 can obstruct recovery. For example, caregivers with financial difficulties explained the  
401 inability to afford rehabilitation or community services to improve survivor recovery, while  
402 caregivers with relationship and mental hardships discussed the issues in engaging in care of  
403 the survivor.

404

*“We hired a person who performs exercises. Rs. 1000 per day. Later, we had to face some financial difficulties. so we couldn’t continue that.”*  
[32]

*“I guess from my perspective probably the friendship that is most lacking is the friendship with him. That’s – (now I’m teary sorry) – it doesn’t exist. No it’s, I have become nothing more than a carer, you know – the secretary that sits at the computer typing, doing his emails, reading everything to him, telling him about everything that happens.*

*But that's all there is to it" [27]*

*"My mom had a stroke almost 1 month ago and is now in rehab. We lived in the same town. My brother lives in 2 provinces over and came when she had a stroke. We live in Canada. We are her power of attorney. Can't agree on anything. And now my mom seems to be agreeing with him and I'm being pushed out! I don't understand....."*

405

406 As a result, caregivers often look for support to ensure better care for their loved ones, and  
407 limit their burden (or burnout). Currently, some caregivers have discussed receiving financial  
408 and care support from relatives. Relatives assisted caregivers, including financial aid,  
409 visitation, food, advice, materials and equipment, engaging in treatment, and occasionally  
410 supporting the caregivers at tasks. Caregivers found that receiving support from relatives  
411 could prevent loneliness and isolation and allow them to engage in social activities, work,  
412 self-training classes, volunteer groups, and travel. One caregiver stated, *"...my daughters and  
413 my sons-in-law have helped me. They are very good to me. Our family is very attached.  
414 Whenever I need it, I call, and we can always manage to arrange schedules, even though they  
415 have a job"* [30]; while another caregiver discussed the opportunities sought by  
416 relative support *"Having more caregivers other than family members [would help]...because  
417 it would give us, the family members, an opportunity to get away from the parent...give us  
418 some time that we can go out and have time on our own... Give me a break!"* [31]. However,  
419 not all caregivers received support from relatives to care for the survivor. These caregivers  
420 found it challenging to adjust to post-stroke conditions. One caregiver stating that *"...our  
421 children have been less supportive than ...expected ...because the ball game has changed for  
422 them too, there is no-one around to support them, so they have been of less practical*

423 *assistance*” [8]. These caregivers often decided to live alone and relied on friends, neighbors,  
424 and healthcare providers to ease their burden. In addition to relatives and friends, caregivers  
425 often drew on the experiences of other caregivers to cope and understand the situation.

426

*“I know from another group I’m in that other members are wonderful about giving advice and I hope I can learn from them, and maybe offer some advice of my own.”*

*“being in a group [support] of individuals with similar experiences provides me with comfort and hope, and abundance of information for which I am grateful to God.”*

*“ya, I think it will definitely help me, because um... some of them may have the experience to be caregivers for stroke patients for many years. For me, I’m just starting out, I’m very new. Definitely there will be lots of mistakes, and my learning curve will be like ups and downs. So definitely if the more experienced caregivers will share with me advice or tips on how to look after a stroke patient. Definitely like I will find it very useful. Like, for me, I need a mentor to become my buddy you know. I’m really very blur, I do not know what to do.” [17]*

427

428 Caregivers often found it challenging to find caregiving peers and support groups, and often  
429 discussed meeting other caregivers on internet-based support groups and outpatient waiting  
430 clinics. One caregiver stated, *“My Dad just had a TIA Wednesday. He had one three weeks  
431 prior also, but never went to the hospital. This time I made him go. All tests came back*

432 *normal, which to me is concerning cause then they don't know why these are happening. It*  
433 *makes me so nervous about another TIA or even worse. So, I'm happy to be on this online*  
434 *support page for him”.*

435 Other than learning from the experiences of others, caregivers often found it crucial to have  
436 coping strategies. Coping strategies allowed caregivers to adjust to their new roles and stay  
437 ‘strong’ and ‘positive’ [36]. Self-encouragement, strong determination and self-strengthening  
438 of the mind were considered critical strategies to cope with the caregiving role [32]. Another  
439 coping mechanism discussed by caregivers was including relaxing activities such as going to  
440 church, cooking or participating in leisure activities that allowed the caregiver to feel  
441 independent.

442

*“[...] I sit down and start to pray and say “please, the life of my  
husband is in your hands. I don't have faith in anything else other than  
in You. You will give me enough courage,” and I pray for half an hour,  
ten minutes, five minutes, the time that I have, then [...] afterward I feel  
better to go on facing the problems I have. This is my way, it is not, as I  
told you, it is not that I pray and then the things get resolved, to resolve  
them, I have to resolve them myself.”*

443

444 Further, the caregiver frequently relied on acceptance and patience, the release of temper, rest  
445 or relaxation, self-treatment, time management, seeking and receiving help, letting go, using  
446 materials and equipment and creativity to manage complex issues with care.

447

*“I think we have all learned to adjust ...that's a big word because our  
lives changed drastically when it happened. We have to learn to adjust*

*and accept.” [8]*

*“Just the fact that [survivor] wasn’t able to do what he had been able to do previously ...acceptance was a big thing in that regard” [8]*

*“You couldn’t return to where you were ...people think that getting better is getting back to as you were. She got better but in a different way. We evolved our life in a different way” [28]*

*“Things are good, getting better all the time, it’s not stopped. We’re not in the situation where we’re going to be like this for the rest of our lives. We’ve still got our lives to live and we will” [28]*

448

449 While caregivers found creative methods to deal with the burden of stroke caregiving, they  
450 often looked for support from the healthcare system. The type of support valued by these  
451 caregivers for themselves included information, emotional, social and financial support.

452

*“We did not get as much help as we needed – I could have used more respite, especially at the beginning, my questions were poorly addressed by the [local community health clinic]” [33]*

*“this is difficult financially because the supplies are very expensive, and the insurance doesn’t cover very much”*

453

454 Caregivers also looked for current healthcare services that were tailored to their needs. The  
455 needs included flexibility in visiting healthcare staff or rehabilitation programs, as this  
456 allowed for them to a break from their caregiving duty for a few hours each day.

457 In addition to receiving support from healthcare systems, caregivers also highlighted the  
458 importance of informal support. For example, one caregiver demonstrated the ability to take  
459 refuge in family members *“It does help the fact that, that I know that what I feel, he feels*  
460 *exactly the same way ...I phone up and moan at him about things and feel better for getting it*  
461 *o□ my chest and he phones me about things, and I know it makes him feel better, but*  
462 *nevertheless for both of us it is still di□cult to cope with things.”* [26]; while another  
463 discussed the importance of supportive and understanding relationships during the care  
464 journey.

465 Caregivers also stressed the importance of financial, social, emotional, and physical support.  
466 Financial support was critical for caregivers to provide efficient care and would allow  
467 caregivers to remain less uncertain about the future as described by one caregiver *“Hardest*  
468 *part is cognition and him [survivor] not wanting to let go of me - he fears/anxious about*  
469 *being alone, but I HAVE to return home to hubby and my job! Can’t keep up the overdrawn*  
470 *bank account”*. On the other hand, social and emotional support is essential to support the  
471 vulnerability of the caregiver as it enables the caregiver to ‘master their fear’ of caregiving  
472 and ensures they ‘feel cared for by others’.

473

*“My daughter helps me – she calls every day to see how I am doing  
because she knows this is very hard for me and she has us over to dinner  
once a week”* [33]

*“I mean, we [siblings] have always been very united,[...] we are one;*

*this situation connected us even more,[...] in life money is not everything, not everything in life is about having something. Life is living in harmony, and this has united us more...” [35]*

*“I have some help from friends to keep up my yard... most people don’t understand what I am going through or how much energy it takes to look after someone who is not well” [33]*

474

475 Some caregivers expressed difficulties in managing the physical issues of caregiving,  
476 especially those that require physical strength, i.e., moving or carrying the survivor. Hence,  
477 caregivers often looked for materials and equipment that can help reduce the stress of  
478 physical activities, especially during emergencies. However, most caregivers expressed a lack  
479 of proper understanding of the tools essential for recovery and local services available to  
480 support the survivor. Hence, requiring additional support from healthcare professionals to  
481 ensure limited physical stress on the caregiver.

482 Caregivers also expected healthcare professionals to assist them in gaining skills to ‘pacify’  
483 and motivate the survivor. One caregiver mentioned, “...is there any possibility that any plan  
484 is drawn up, to really let the patient—instead of us (the family)— know what he may expect  
485 now...if they are still conscious of what’s going on, I think it will be good to tell them (stroke  
486 patients) what they are or what they will be going through” [17]. The ability to gain skills to  
487 support the survivor would allow the caregiver to remain ‘positive’ and ‘hopeful’ about the  
488 future.

489

## 490 **Discussion**

### 491 **Summary of main findings**

492 The study drew on grounded theory to explore the needs of caregivers of the people living  
493 with stroke based on their experiences in care. The core categories identified, i.e.,  
494 information, involvement, self-care, and support, need to be addressed to empower the  
495 caregiver while reducing uncertainty, mental impact, financial stress, and burnout. In  
496 particular, it was clear that there are several barriers in the current healthcare system leading  
497 to unpreparedness and unwanted effects on the caregiver. Hence, requiring the inclusion of a  
498 more caregiver-focused approach in clinical practice and health care planning.

499

## 500 **Comparison with literature**

501 Key findings from this grounded theory study demonstrate that caregivers with limited  
502 support feel unprepared to care for the survivor and are not recognized as key stakeholders in  
503 recovery. Further, the caregiver is overwhelmed with emotions and feels unsupported during  
504 the care trajectory. The needs, however, differ based on the individual and the services  
505 provided to the caregiver and the survivor. For example, some caregivers received a verbal  
506 explanation about the disease and practices leading to increased satisfaction of healthcare  
507 services. In contrast, others did not, hence highlighting differences in care that bring  
508 challenges in understanding care needs. However, it is important to note that the role of a  
509 caregiver is well documented [22, 37-41]. Further, by addressing the individual needs of  
510 stroke caregivers, it is possible to limit the feelings of fear, uncertainty, isolation, burnout,  
511 fatigue, etc.

512 In this study, we have, therefore, highlighted four critical themes based on the individual  
513 needs of caregivers verified through multiple stages of review and evaluation. While these  
514 themes are unique, there may be several synergies to other studies that address stroke  
515 caregiver needs. For example, Tsai, Yip [12] classified caregiver needs into (i) health  
516 information, (ii) instrumental support, (iii) emotional support, (iv) involvement with patient

517 care, (v) professional support, and (vi) community networks; while Bakas, Austin [42]  
518 categorized the needs and concerns of stroke caregivers into (i) information, (ii) emotions and  
519 behaviors, (iii) physical care, (iv) instrumental care and (v) personal responses to caregiving.  
520 The needs in this study and past studies show a change in needs based on the survivor's  
521 disease progression and health situation. The change in needs has been well documented in  
522 the study by Cameron, Naglie [23], which has classified caregivers' needs into five stages:  
523 event/diagnosis, stabilization, preparation, implementation, and adaption. The change in  
524 needs is the highest during the first few months post-discharge, i.e., in preparation and  
525 implementation phases, especially for those caregivers who manage survivors with high  
526 levels of physical, communication, and/or cognitive disability [23], which is consistent with  
527 the findings in this study.

528

## 529 **Strengths and Limitations**

530 The study was conducted in three phases, which include: (i) literature analysis, (ii) social  
531 media analysis, and (iii) caregiver survey to identify and evaluate critical needs in stroke  
532 caregiving. Due to existing restrictions, most of the collected data did not involve face-to-  
533 face interactions with the caregiver. Face-to-face interactions with the caregiver may have  
534 allowed for a more detailed understanding of the caregiver's needs, which may have affected  
535 the richness and quality of data collected. However, all findings presented in this study have  
536 undergone rigorous analysis and documentation to ensure the caregivers' needs are well  
537 represented.

538

## 539 **Implications for future research**

540 The findings in this study have several implications for clinical practice, healthcare planning,  
541 and healthcare policy. In particular, this study sheds light on the needs of stroke caregivers

542 across the care trajectory. Thus, considerations are made to address these needs to bring about  
543 preparedness, reduce uncertainty and promote behavior change. One plausible approach is the  
544 use of technology.

545 Currently, several caregivers in stroke have employed technology connected to the internet to  
546 ensure better care for their loved ones [9, 19, 21, 36]. However, the reliability of these  
547 technologies to support the constantly changing needs of stroke caregivers during recovery  
548 has not been fully explored [43]. Hence there is a need to develop a tool that addresses all the  
549 aspects mentioned above.

550 With the growth of new technological approaches, it is possible to provide the caregiver with  
551 context-aware support, i.e., allow the system to understand the varying needs of the  
552 caregivers and provide support at critical times to address these needs [10]. However, it  
553 would require a more detailed understanding of individual needs to create efficient algorithms  
554 and infrastructure [44] for improved support and care. Techniques such as participatory  
555 design and user-centered design approaches could be implemented to ensure the design of a  
556 sustainable system by creating actionable strategies to test, refine and integrate the solution in  
557 the daily activities of the caregiver [10]. Hence, addressing all the critical needs of stroke  
558 caregiving and potentially creating a commercially viable solution for support.

559

## 560 **Conclusions**

561 The ground theory study provides new insights into the needs and preferences of stroke  
562 caregivers during recovery and exposes several areas that need to be addressed to support the  
563 care process and prevent obstruction in care and reduce uncertainty. Moreover, by  
564 considering these needs, the healthcare system can help navigate the caregiver through the  
565 disease trajectory and focus on designing a more inclusive approach towards caregiver-  
566 centered care.

567

## 568 **Acknowledgements**

569 None

570

## 571 **Funding**

572 This study was supported through doctoral scholarships from the School of Information  
573 Technology, Deakin University, and the Department of Public Health, University of  
574 Copenhagen. Further, Prof John Grundy is supported by ARC Laureate Fellowship  
575 FL190100035 and Dr Islam is supported by NHMRC Emerging Leadership Fellowship.

576

## 577 **Conflicts of Interest**

578 The authors declare that they have no competing interests

579

## 580 **References**

- 581 1. Lindsay MP, Norrving B, Sacco RL, Brainin M, Hacke W, Martins S, et al. Global  
582 Stroke Fact Sheet 2019: World Stroke Organization (WSO); 2019 [cited 2021 7th July].  
583 Available from: [https://www.world-stroke.org/assets/downloads/WSO\\_Fact-](https://www.world-stroke.org/assets/downloads/WSO_Fact-sheet_15.01.2020.pdf)  
584 [sheet\\_15.01.2020.pdf](https://www.world-stroke.org/assets/downloads/WSO_Fact-sheet_15.01.2020.pdf).
- 585 2. Lackland DT, Roccella EJ, Deutsch AF, Fornage M, George MG, Howard G, et al.  
586 Factors influencing the decline in stroke mortality: a statement from the American Heart  
587 Association/American Stroke Association. *Stroke*. 2014;45(1):315-53. Epub 2013/12/05. doi:  
588 10.1161/01.str.0000437068.30550.cf. PubMed PMID: 24309587.
- 589 3. Donkor ES. Stroke in the 21(st) Century: A Snapshot of the Burden, Epidemiology,  
590 and Quality of Life. *Stroke Res Treat*. 2018;2018:3238165-. doi: 10.1155/2018/3238165.  
591 PubMed PMID: 30598741.
- 592 4. Torregosa MB, Sada R, Perez I. Dealing with stroke: Perspectives from stroke  
593 survivors and stroke caregivers from an underserved Hispanic community. *Nursing & health*  
594 *sciences*. 2018;20(3):361-9. Epub 2018/03/08. doi: 10.1111/nhs.12414. PubMed PMID:  
595 29512845.
- 596 5. Camak DJ. Addressing the burden of stroke caregivers: a literature review. *Journal of*  
597 *clinical nursing*. 2015;24(17-18):2376-82. Epub 2015/06/23. doi: 10.1111/jocn.12884.  
598 PubMed PMID: 26095074.

- 599 6. Tosun ZK, Temel MJJoCS. Burden of caregiving for stroke patients and the role of  
600 social support among family members: an assessment through home visits. 2017;10(3):1696.  
601 7. Soufi S, Chabrier S, Bertoletti L, Laporte S, Darteyre S. Lived experience of having a  
602 child with stroke: A qualitative study. *European Journal of Paediatric Neurology*.  
603 2017;21(3):542-8. doi: <https://doi.org/10.1016/j.ejpn.2017.01.007>.  
604 8. Graven C, Sansonetti D, Moloczij N, Cadilhac D, Joubert L. Stroke survivor and carer  
605 perspectives of the concept of recovery: a qualitative study. *Disability and rehabilitation*.  
606 2013;35(7):578-85. Epub 2012/08/15. doi: 10.3109/09638288.2012.703755. PubMed PMID:  
607 22889405.  
608 9. Creasy KR, Lutz BJ, Young ME, Ford A, Martz C. The impact of interactions with  
609 providers on stroke caregivers' needs. *Rehabilitation nursing : the official journal of the*  
610 *Association of Rehabilitation Nurses*. 2013;38(2):88-98. Epub 2013/03/27. doi:  
611 10.1002/rnj.69. PubMed PMID: 23529947; PubMed Central PMCID: PMCPMC3742102.  
612 10. Lobo EH, Frølich A, Rasmussen LJ, Livingston PM, Grundy J, Abdelrazek M, et al.  
613 Understanding the Methodological Issues and Solutions in the Research Design of Stroke  
614 Caregiving Technology. *Frontiers in public health*. 2021;9:647249. Epub 2021/05/04. doi:  
615 10.3389/fpubh.2021.647249. PubMed PMID: 33937175; PubMed Central PMCID:  
616 PMCPMC8085388.  
617 11. Luker J, Murray C, Lynch E, Bernhardsson S, Shannon M, Bernhardt J. Carers'  
618 Experiences, Needs, and Preferences During Inpatient Stroke Rehabilitation: A Systematic  
619 Review of Qualitative Studies. *Archives of physical medicine and rehabilitation*.  
620 2017;98(9):1852-62.e13. Epub 2017/04/02. doi: 10.1016/j.apmr.2017.02.024. PubMed  
621 PMID: 28363703.  
622 12. Tsai PC, Yip PK, Tai JJ, Lou MF. Needs of family caregivers of stroke patients: a  
623 longitudinal study of caregivers' perspectives. *Patient preference and adherence*. 2015;9:449-  
624 57. Epub 2015/04/04. doi: 10.2147/ppa.S77713. PubMed PMID: 25834409; PubMed Central  
625 PMCID: PMCPMC4370911.  
626 13. Chun Tie Y, Birks M, Francis K. Grounded theory research: A design framework for  
627 novice researchers. *SAGE Open Medicine*. 2019;7:2050312118822927. doi:  
628 10.1177/2050312118822927.  
629 14. Starks H, Brown Trinidad S. Choose Your Method: A Comparison of  
630 Phenomenology, Discourse Analysis, and Grounded Theory. *Qualitative Health Research*.  
631 2007;17(10):1372-80. doi: 10.1177/1049732307307031.  
632 15. Ligita T, Nurjannah I, Wicking K, Harvey N, Francis K. From textual to visual: the  
633 use of concept mapping as an analytical tool in a grounded theory study. *Qualitative*  
634 *Research*. 2020:1468794120965362. doi: 10.1177/1468794120965362.  
635 16. Hallberg LRM. The "core category" of grounded theory: Making constant  
636 comparisons. *International Journal of Qualitative Studies on Health and Well-being*.  
637 2006;1(3):141-8. doi: 10.1080/17482620600858399.  
638 17. Ang SY, Tin AS, Pavitar G, Ng WM, Lee KE, Lim LH, et al. A Qualitative Study into  
639 Stroke Caregivers' Educational Needs — Perspectives of Caregivers and Healthcare  
640 Professionals. *Proceedings of Singapore Healthcare*. 2013;22(3):166-74. doi:  
641 10.1177/201010581302200303.  
642 18. Howe T, Davidson B, Worrall L, Hersh D, Ferguson A, Sherratt S, et al. 'You needed  
643 to rehab ... families as well': family members' own goals for aphasia rehabilitation.  
644 *International journal of language & communication disorders*. 2012;47(5):511-21. Epub  
645 2012/09/04. doi: 10.1111/j.1460-6984.2012.00159.x. PubMed PMID: 22938062.  
646 19. Wu C-m. Learning to be a family caregiver for severely debilitated stroke survivors  
647 during the first year in Taiwan: The University of Iowa; 2009.

- 648 20. Cobley CS, Fisher RJ, Chouliara N, Kerr M, Walker MF. A qualitative study  
649 exploring patients' and carers' experiences of Early Supported Discharge services after stroke.  
650 *Clinical rehabilitation*. 2013;27(8):750-7. Epub 2013/03/05. doi:  
651 10.1177/0269215512474030. PubMed PMID: 23455948.
- 652 21. Danzl MM, Harrison A, Hunter EG, Kuperstein J, Sylvia V, Maddy K, et al. "A Lot  
653 of Things Passed Me by": Rural Stroke Survivors' and Caregivers' Experience of Receiving  
654 Education From Health Care Providers. *The Journal of rural health : official journal of the*  
655 *American Rural Health Association and the National Rural Health Care Association*.  
656 2016;32(1):13-24. Epub 2015/06/24. doi: 10.1111/jrh.12124. PubMed PMID: 26100171.
- 657 22. Thomas M, Greenop K. Caregiver experiences and perceptions of stroke %J *Health*  
658 *SA Gesondheid (Online)*. 2008;13:29-40.
- 659 23. Cameron JI, Naglie G, Silver FL, Gignac MA. Stroke family caregivers' support  
660 needs change across the care continuum: a qualitative study using the timing it right  
661 framework. *Disability and rehabilitation*. 2013;35(4):315-24. Epub 2012/06/13. doi:  
662 10.3109/09638288.2012.691937. PubMed PMID: 22686259.
- 663 24. Cameron JI, Bastawrous M, Marsella A, Forde S, Smale L, Friedland J, et al. Stroke  
664 survivors', caregivers', and health care professionals' perspectives on the weekend pass to  
665 facilitate transition home. *Journal of rehabilitation medicine*. 2014;46(9):858-63. Epub  
666 2014/08/29. doi: 10.2340/16501977-1854. PubMed PMID: 25166922.
- 667 25. Subgranon R, Lund DA. Maintaining caregiving at home: a culturally sensitive  
668 grounded theory of providing care in Thailand. *Journal of transcultural nursing : official*  
669 *journal of the Transcultural Nursing Society*. 2000;11(3):166-73. Epub 2002/05/02. doi:  
670 10.1177/104365960001100302. PubMed PMID: 11982104.
- 671 26. Hunt D, Smith JA. The personal experience of carers of stroke survivors: an  
672 interpretative phenomenological analysis. *Disability and rehabilitation*. 2004;26(16):1000-11.  
673 Epub 2004/09/17. doi: 10.1080/09638280410001702423. PubMed PMID: 15371048.
- 674 27. El Masry Y, Mullan B, Hackett M. Psychosocial experiences and needs of Australian  
675 caregivers of people with stroke: prognosis messages, caregiver resilience, and relationships.  
676 *Topics in stroke rehabilitation*. 2013;20(4):356-68. Epub 2013/07/31. doi: 10.1310/tsr2004-  
677 356. PubMed PMID: 23893835.
- 678 28. Buschenfeld K, Morris R, Lockwood S. The experience of partners of young stroke  
679 survivors. *Disability and rehabilitation*. 2009;31(20):1643-51. Epub 2009/05/30. doi:  
680 10.1080/09638280902736338. PubMed PMID: 19479557.
- 681 29. Krieger T, Feron F, Dorant E. Developing a complex intervention programme for  
682 informal caregivers of stroke survivors: The Caregivers' Guide. *Scandinavian journal of*  
683 *caring sciences*. 2017;31(1):146-56. Epub 2016/07/22. doi: 10.1111/scs.12344. PubMed  
684 PMID: 27440738.
- 685 30. Bocchi SC, Angelo M. Between freedom and reclusion: social support as a quality-of-  
686 life component in the family caregiver-dependent person binomial. *Revista latino-americana*  
687 *de enfermagem*. 2008;16(1):15-23. Epub 2008/04/09. doi: 10.1590/s0104-  
688 11692008000100003. PubMed PMID: 18392525.
- 689 31. Bastawrous M, Gignac MA, Kapral MK, Cameron JI. Adult daughters providing post-  
690 stroke care to a parent: a qualitative study of the impact that role overload has on lifestyle,  
691 participation and family relationships. *Clinical rehabilitation*. 2015;29(6):592-600. Epub  
692 2014/09/27. doi: 10.1177/0269215514552035. PubMed PMID: 25258424.
- 693 32. Wagachchige Muthucumarana M, Samarasinghe K, Elgán C. Caring for stroke  
694 survivors: experiences of family caregivers in Sri Lanka - a qualitative study. *Topics in stroke*  
695 *rehabilitation*. 2018;25(6):397-402. Epub 2018/07/22. doi: 10.1080/10749357.2018.1481353.  
696 PubMed PMID: 30028654.

- 697 33. Barbic SP, Mayo NE, White CL, Bartlett SJ. Emotional vitality in family caregivers:  
698 content validation of a theoretical framework. *Quality of life research : an international*  
699 *journal of quality of life aspects of treatment, care and rehabilitation*. 2014;23(10):2865-72.  
700 Epub 2014/05/24. doi: 10.1007/s11136-014-0718-4. PubMed PMID: 24853533.
- 701 34. Chow C, Tiwari A. Experience of family caregivers of community-dwelling stroke  
702 survivors and risk of elder abuse: a qualitative study. *J Adult Pro*. 2014;16(5):276-93. doi:  
703 10.1108/JAP-03-2014-0007.
- 704 35. Pesantes MA, Brandt LR, Ipince A, Miranda JJ, Diez-Canseco F. An exploration into  
705 caring for a stroke-survivor in Lima, Peru: Emotional impact, stress factors, coping  
706 mechanisms and unmet needs of informal caregivers. *eNeurologicalSci*. 2017;6:33-50. Epub  
707 2017/10/11. doi: 10.1016/j.ensci.2016.11.004. PubMed PMID: 28989982; PubMed Central  
708 PMCID: PMC5627498.
- 709 36. Smith SD, Gignac MA, Richardson D, Cameron JI. Differences in the experiences  
710 and support needs of family caregivers to stroke survivors: does age matter? *Topics in stroke*  
711 *rehabilitation*. 2008;15(6):593-601. Epub 2009/01/23. doi: 10.1310/tsr1506-593. PubMed  
712 PMID: 19158067.
- 713 37. Jullamate P, de Azeredo Z, Pául C, Subgranon R. Thai stroke patient caregivers: who  
714 they are and what they need. *Cerebrovascular diseases (Basel, Switzerland)*. 2006;21(1-  
715 2):128-33. Epub 2005/12/13. doi: 10.1159/000090211. PubMed PMID: 16340188.
- 716 38. Steiner V, Pierce L, Drahuschak S, Nofziger E, Buchman D, Szirony T. Emotional  
717 support, physical help, and health of caregivers of stroke survivors. *The Journal of*  
718 *neuroscience nursing : journal of the American Association of Neuroscience Nurses*.  
719 2008;40(1):48-54. Epub 2008/03/12. doi: 10.1097/01376517-200802000-00008. PubMed  
720 PMID: 18330410; PubMed Central PMCID: PMC2442227.
- 721 39. Chow SK, Wong FK, Poon CY. Coping and caring: support for family caregivers of  
722 stroke survivors. *Journal of clinical nursing*. 2007;16(7b):133-43. Epub 2007/06/23. doi:  
723 10.1111/j.1365-2702.2006.01711.x. PubMed PMID: 17584422.
- 724 40. Mei YX, Lin B, Zhang W, Yang DB, Wang SS, Zhang ZX, et al. Benefits finding  
725 among Chinese family caregivers of stroke survivors: a qualitative descriptive study. *BMJ*  
726 *open*. 2020;10(10):e038344. Epub 2020/10/21. doi: 10.1136/bmjopen-2020-038344. PubMed  
727 PMID: 33077565; PubMed Central PMCID: PMC5757495.
- 728 41. Araújo O, Lage I, Cabrita J, Teixeira L. Training informal caregivers to care for older  
729 people after stroke: A quasi-experimental study. *Journal of advanced nursing*. 2018. Epub  
730 2018/05/13. doi: 10.1111/jan.13714. PubMed PMID: 29752812.
- 731 42. Bakas T, Austin JK, Okonkwo KF, Lewis RR, Chadwick L. Needs, concerns,  
732 strategies, and advice of stroke caregivers the first 6 months after discharge. *The Journal of*  
733 *neuroscience nursing : journal of the American Association of Neuroscience Nurses*.  
734 2002;34(5):242-51. Epub 2002/10/24. doi: 10.1097/01376517-200210000-00004. PubMed  
735 PMID: 12391740.
- 736 43. Lobo EH, Frølich A, Kensing F, Rasmussen LJ, Livingston PM, Grundy J, et al.  
737 mHealth applications to support caregiver needs and engagement during stroke recovery: A  
738 content review. *Research in nursing & health*. 2021;44(1):213-25. Epub 2020/12/21. doi:  
739 10.1002/nur.22096. PubMed PMID: 33341958.
- 740 44. van Engelenburg S, Janssen M, Klievink B. Designing context-aware systems: A  
741 method for understanding and analysing context in practice. *Journal of Logical and Algebraic*  
742 *Methods in Programming*. 2019;103:79-104. doi: 10.1016/j.jlamp.2018.11.003.
- 743

## 744 **Supporting Information**

745 **S1 File. User Needs and Associated Comments**