Genomic epidemiology reveals the impact of national and international restrictions

measures on the SARS-CoV-2 epidemic in Brazil

Marta Giovanetti[1,2]*[&], Svetoslav Nanev Slavov[3,4]*, Vagner Fonseca[1,5,6,7]*, Eduan Wilkinson[6,7]*,

Houriiyah Tegally[6,7]*, José Salvatore Leister Patané[4]*, Vincent Louis Viala[4], James Emmanuel San[6,7],

Evandra Strazza Rodrigues[3], Elaine Vieira Santos[3], Flavia Aburjaile[2,8], Joilson Xavier[1,8], Hegger

Fritsch[1,8], Talita Emile Ribeiro Adelino[1,8], Felicidade Pereira[9], Arabela Leal[9], Felipe Campos de Melo

Iani[8], Glauco de Carvalho Pereira[8], Cynthia Vazquez[10], Gladys Mercedes Estigarribia Sanabria[11,12,13],

Elaine Cristina de Oliveira[14], Luiz Demarchi[15], Julio Croda[16], Rafael dos Santos Bezerra[3], Loyze Paola

Oliveira de Lima[4], Antonio Jorge Martins[4], Claudia Renata dos Santos Barros[4], Elaine Cristina

Marqueze[4], Jardelina de Souza Todao Bernardino[4], Debora Botequio Moretti[4], Ricardo Augusto

Brassaloti[17], Raquel de Lello Rocha Campos Cassano[17], Pilar Drummond Sampaio Corrêa Mariani[18], João

Paulo Kitajima[19], Bibiana Santos[19], Rodrigo Proto-Siqueira[20], Vlademir Vicente Cantarelli[21], Stephane

Tosta[2,9], Vanessa Brandão Nardy[9], Luciana Reboredo de Oliveira da Silva[9], Marcela Kelly Astete

Gómez[9], Jaqueline Gomes Lima[9], Adriana Aparecida Ribeiro[8], Natália Rocha Guimarães[8], Luiz Takao

Watanabe[14], Luana Barbosa Da Silva[14], Raquel da Silva Ferreira[14], Mara Patricia F. da Penha[22], María

José Ortega[10], Andrea Gómez de la Fuente[10], Shirley Villalba[10], Juan Torales[10], María Liz Gamarra[10],

Carolina Aquino[10], Gloria Patricia Martínez Figueredo[11,12,13], Wellington Santos Fava[16], Ana Rita C.

Motta-Castro[16], James Venturini[16], Sandra Maria do Vale Leone de Oliveira[16], Crhistinne Cavalheiro

Maymone Gonçalves[23], Maria do Carmo Debur Rossa[24], Guilherme Nardi Becker[24], Mayra Marinho

Presibella[24], Nelson Quallio Marques[24], Irina Nastassja Riediger[24], Sonia Raboni[25], Gabriela Mattoso

Coelho[26], Allan Henrique Depieri Cataneo[26], Camila Zanluca[26], Claudia N Duarte dos Santos[26], Patricia

Akemi Assato[27], Felipe Allan da Silva da Costa[27], Mirele Daiana Poleti[28], Jessika Cristina Chagas

Lesbon[28], Elisangela Chicaroni Mattos[28], Cecilia Artico Banho[29], Lívia Sacchetto[29], Marília Mazzi

Moraes[29], Rejane Maria Tommasini Grotto[27,30], Jayme A. Souza-Neto[27], Maurício Lacerda Nogueira[29],

Heidge Fukumasu[28], Luiz Lehmann Coutinho[17], Rodrigo Tocantins Calado[3], Raul Machado Neto[4], Ana

Maria Bispo de Filippis[1], Rivaldo Venancio da Cunha[31], Carla Freitas[5], Cassio Roberto Leonel Peterka[32],

Cássia de Fátima Rangel Fernandes[33], Wildo Navegantes de Araújo[34], Rodrigo Fabiano do Carmo Said[34],

Maria Almiron[34], Carlos Frederico Campelo de Albuquerque e Melo[34], José Lourenço[35,36], Tulio de

Oliveira[6,7,37,38], Edward C. Holmes[39], Ricardo Haddad[4], Sandra Coccuzzo Sampaio[4]^&, Maria Carolina

Elias4^&, Simone Kashima[3]^&, Luiz Carlos Junior de Alcantara[1,2]^&, Dimas Tadeu Covas[3,4]^&.

*These authors jointly contributed to this work

^These authors jointly supervised this work

Corresponding Authors&: sandra.coccuzzo@butantan.gov.br, carolina.eliassabbaga@butantan.gov.br,

skashima@hemocentro.fmrp.usp.br, luiz.alcantara@ioc.fiocruz.br, dimas.covas@butantan.gov.br;

giovanetti.marta@gmail.com

[1] Laboratório de Flavivírus, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil; [2]Laboratório de Genética Celular e

Molecular, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais,

Brazil; [3]University of São Paulo, Ribeirão Preto Medical School, Blood Center of Ribeirão Preto, Ribeirão Preto,

SP, Brazil; [4]Butantan Institute, São Paulo, Brazil; [5]Coordenação Geral de Laboratórios de Saúde Pública/Secretaria

de Vigilância em Saúde, Ministério da Saúde (CGLAB/SVS-MS) Brasília, Distrito Federal, Brazil; [6]KwaZulu-Natal

Research Innovation and Sequencing Platform (KRISP), School of Laboratory Medicine and Medical Sciences,

College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa; [7]Centre for Epidemic Response

and Innovation (CERI), School of Data Science and Computational Thinking, Stellenbosch University;

Stellenbosch, South Africa; [8]Laboratório Central de Saúde Pública do Estado de Minas Gerais, Fundacão Ezequiel

Dias, Belo Horizonte, Minas Gerais, Brazil; [9]Laboratorio Central de Saude Publica da Bahia–LACEN-BA,

Salvador, Bahia, Brazil; [10]Laboratorio Central de Salud Pública, Asunción, Paraguay; [11]Universidad Nacional del

Caaguazú, Instituto Regional de Investigación en Salud; [12]Laboratorio de Biología Molecular, Hospital Regional de

Coronel Oviedo; [13]Ministerio de Salud Pública y Bienestar Social; [14]Laboratório Central de Saúde Pública do Estado

de Mato Grosso, Cuiabá, Brazil; [15]Laboratório Central de Saúde Pública do Estado de Mato Grosso do Sul, Campo

Grande, Mato Grosso do Sul, Brazil; [16]Universidade Federal de Mato Grosso do Sul; [17]University of São Paulo,

Centro de Genômica Funcional da ESALQ, Piracicaba, SP, Brazil; [18]NGS Soluções Genômicas, Piracicaba, SP,

Brazil; [19]Mendelics Análise Genômica; [20]Instituto de Biologia Molecular, Laboratório Antonello, Rio Grande do

Sul, Brazil; [21]Universidade Federal de Ciências da Saúde de Porto Alegre (UFCSPA), Universidade Feevale, Grupo

Exame Laboratórios, Rio Grande do Sul, Brazil; [22]Vigilância em Saúde do Estado de Mato Grosso, Cuiabá, Brazil;

[23]Secretaria de Saúde do Estado do Mato Grosso do Sul; [24]Laboratório Central do Estado do Paraná (Lacen/PR);

[25]Hospital de Clínicas da Universidade Federal do Paraná, Curitiba, PR; [26]Laboratório de Virologia Molecular -

Instituto Carlos Chagas/Fiocruz PR, Curitiba, PR; [27]São Paulo State University (UNESP), School of Agricultural

Sciences, Department of Bioprocesses and Biotechnology, Botucatu, Brazil; [28]Department of Veterinary Medicine,

School of Animal Science and Food Engineering, University of Sao Paulo, Pirassununga, São Paulo, Brazil;

[29]Laboratório de Pesquisas em Virologia, Departamento de Doenças Dermatológicas, Infecciosas e Parasitárias,

Faculdade de Medicina de São José do Rio Preto; [30]Molecular Biology Laboratory, Applied Biotechnology

Laboratory, Clinical Hospital of the Botucatu Medical School, Brazil; [31]Fundação Oswaldo Cruz, Bio-Manguinhos,

Rio de Janeiro, Rio de Janeiro, Brazil; [32]Coordenacão Geral das Arboviroses, Secretaria de Vigilancia em

Saúde/Ministério da Saúde (CGARB/SVS-MS), Brasília, Distrito Federal, Brazil; [33]Departamento de Imunização e

Doenças Transmissíveisa/Secretaria de Vigilância em Saúde, Ministério da Saúde, Brasília, Distrito Federal, Brazil;

[34]Organizacão Pan-Americana da Saúde/Organizacão Mundial da Saúde, Brasília, Distrito Federal, Brazil;

[35]Department of Zoology, Peter Medawar Building, University of Oxford, Oxford, UK; [36]Biosystems and Integrative

Sciences Institute, Universidade de Lisboa, Lisboa, Portugal; [37]Centre for the AIDS Programme of Research in

South Africa (CAPRISA), Durban, South Africa; [38]Department of Global Health, University of Washington, Seattle,

WA, USA; [39]Sydney Institute for Infectious Diseases, School of Life and Environmental Sciences and School of

Medical Sciences, University of Sydney, Sydney, NSW, Australia.

Abstract

Brazil has experienced some of the highest numbers of COVID-19 cases and deaths globally and

from May 2021 made Latin America a pandemic epicenter. Although SARS-CoV-2 established

sustained transmission in Brazil early in the pandemic, important gaps remain in our

understanding of virus transmission dynamics at the national scale. Here, we describe the

genomic epidemiology of SARS-CoV-2 using near-full genomes sampled from 27 Brazilian

states and a bordering country - Paraguay. We show that the early stage of the pandemic in

Brazil was characterised by the co-circulation of multiple viral lineages, linked to multiple

importations predominantly from Europe, and subsequently characterized by large local

transmission clusters. As the epidemic progressed under an absence of effective restriction

measures, there was a local emergence and onward international spread of Variants of Concern

(VOC) and Variants Under Monitoring (VUM), including Gamma (P.1) and Zeta (P.2). In

addition, we provide a preliminary genomic overview of the epidemic in Paraguay, showing

evidence of importation from Brazil. These data reinforce the usefulness and need for the

implementation of widespread genomic surveillance in South America as a toolkit for pandemic

monitoring that provides a means to follow the real-time spread of emerging SARS-CoV-2

variants with possible implications for public health and immunization strategies.

Introduction

At the end of 2019, a novel respiratory pathogen designated the Severe Acute Respiratory

Syndrome Coronavirus 2 (SARS-CoV-2) emerged in the city of Wuhan, China. Since its

identification, the virus has spread rapidly on a global scale causing an unprecedented

Coronavirus disease 2019 (COVID-19) pandemic (declared on March 11[th] 2020) which has

overwhelmed many health care systems[1]. By the mid-February 2022, more than 415 million

cases of COVID-19, with more than 5.84 million associated deaths, have been reported

globally[2]. Brazil has become the epicentre of the COVID-19 epidemic in the Americas, with a

total of 21.2 million cases and a death toll exceeding 591,000 reported cases by the end of

September of 2021, making it one of the countries hardest hit by COVID-19[3]. Critically,

however, a lack of genome sequence data from Brazil has limited our ability to fully understand

transmission dynamics at the national scale.

To assess how population restriction measures might have played a role in shaping viral

diversity and evolution during the SARS-CoV-2 epidemic in Brazil, we performed a

phylogenetic and phylogeographic analysis of genomic data from 27 Brazilian states and one

neighbouring country – Paraguay – collected up to September 2021. We show that the early

spread of SARS-CoV-2 in Brazil resulted from initial founder events associated with

international travel (mainly from Europe), which then spread through regional mobility pathways

during the recession of the first epidemic wave. From this, Brazil emerged as an exporter of

SARS-CoV-2 variants to other countries. We further describe the emergence and spread of key

viral Variants of Concern (VOCs; for example, Gamma) and Variants Under Monitoring

(VUMs; for example, Zeta) in Brazil, highlighting how their emergence may have contributed to

a more severe second wave in the country.

Results

COVID-19 transmission dynamics in Brazil

The first confirmed infection of SARS-CoV-2 in Brazil was on February 26th 2020 in the State of

São Paulo (SP), in a traveller returning from Italy (**Fig. 1A**). On March 17th 2020 the first

COVID-19-related death, a 61-year old male, was reported in the same state[4,5]. Four days later,

all Brazilian states reported at least one confirmed case of COVID-19 and the Brazilian Ministry

of Health (BR-MoH) declared an outbreak of large-scale community transmission of the virus[6].

By April 10th, 2020 the virus had already reached remote locations such as the Yanomami

indigenous community located in the state of Roraima, in northern Brazil[6] (**Fig. 1A**).

After the World Health Organization (WHO) declared the outbreak of SARS-CoV-2 as a

public health emergency of international concern on January 30th 2020 the Brazilian government

introduced restriction measures to mitigate viral spread (**Fig. 1A**)[7]. The primary measure

involved social isolation, followed by the closure of schools, universities, and non-essential

businesses[8]. Additional measures included the mandatory use of personal protective masks[9], the

cancellation of events expected to attract large numbers of people and tourists, and only services

considered as essential such as markets and pharmacies remained open[8,10]. However, while the

epidemic was growing, restriction measures were progressively eased to mitigate negative

impacts on the economy. Notably, even during periods of restriction, travel between Brazilian

states largely remained possible, facilitating SARS-CoV-2 transmission throughout the

country[11]. The latter was likely linked to the emergence of new and more contagious viral

lineages documented as VOC - Gamma (lineage P.1) - and VUM - Zeta (lineage P.2), which may

also have contributed to a more severe second wave (**Fig. 1B**)[11,13,14,15].

The COVID-19 death toll in Brazil steadily rose from March 2021. It reached a daily

total of 4,250 on April 2021, the highest number of daily fatalities from COVID-19 worldwide

(**Fig. 1B**). Signs of collapse of the health system were reported in numerous cities among

different regions of the country. The situation worsened considerably following the emergence of

multiple VOCs and VUMs in parallel with a slow progress of the vaccination campaign[16].

Vaccination in Brazil began on January 17[th] 2021 when the Instituto Butantan imported the first

6 million doses of CoronaVac (a whole virus inactivated vaccine) in a collaboration with Sinovac

Biotech (**Fig. 1A**)[17,18]. As of February 16[th] 2022 approximately 71.8% of the Brazilian

population have been vaccinated with the first dose of any of the vaccines currently available

(CoronaVac, AstraZeneca, Pfizer and Janssen), and only 22% had been fully vaccinated (i.e. a

single dose of Janssen or both doses of the other vaccines)[19].

By analysing the total number of COVID-19 notified cases to the end of September 2021,

we observed that the Brazilian region with the highest population density (Southeast) also

contained the highest number of the cases registered in the country, with the state of São Paulo

documenting the largest number of cases (n=4,369,410) in that period (**Fig**. **1C**). However, when

we consider the incidence rate (number of reported cases/population) by state, we found that the

Midwest, the least populated region in Brazil, had the highest incidence rate, with 13,604.23

cases/100K inhabitants[1].

SARS-CoV-2 genomic data

A total of 3,866 near-full genomes sequences from SARS-CoV-2 RT-qPCR positive

samples were obtained as part of this study. SARS-CoV-2 sequencing spanned February 2020 to

June 2021, sampled from 8 of the 27 Brazilian states (São Paulo=3309; Rio Grande do Sul=48;

Paraná=55; Minas Gerais=80; Mato Grosso do Sul=36; Mato Grosso=51; Bahia=224) and one

neighbouring country, Paraguay (n=63). Almost half of the sequences were from Southeast

Brazil, consistent with it reporting most cases natiionally (**Fig. 1C**)[6]. The sequenced samples

were collected from 2,023 females and 1,843 males (**Table S1 and S2**) with a median age of

41.72 years (range: 1 to 90 years of age). All tested samples contained sufficient viral genetic

material (≥2ng/µL) for library preparation. For positive samples, PCR cycle threshold (Ct)

values were on average 19.93 (range: 10.75 to 30). Sequences had a median genome coverage of

95% (range: 80 to 99.99) and average genome coverage was typically higher for samples with

lower Ct values (**Fig. S1**). Epidemiological information and sequencing statistics of the

generated sequences from Brazil and Paraguay are detailed in **Tables S1** and **S2,** respectively.

Sequences were assigned to 39 different PANGO-lineages based on the proposed dynamic

nomenclature for SARS-CoV-2 lineages (**Fig. S1, Table S1 and S2**) and have been submitted to

GISAID following the WHO guidelines (**Tables S1** and **S2**) (Pangolin version 3.1.7, August

2021).

Phylogenetic inference and lineage diversity

The rapid spread of SARS-CoV-2, together with the reported circulation of several VOCs

and VUMs in Brazil prompted the intensification of genomic surveillance by the National

Network for Pandemic Alert of SARS-CoV-2 at the end of December 2020. As of June 30[th] 2021

some 17,135 SARS-CoV-2 genomes from Brazil had been deposited in the GISAID database

from all 27 Brazilian states (**Fig. 2A**). The states with the highest number of sequenced genomes

were São Paulo (n=9,600) and Rio de Janeiro (n=2,031). Although genomic surveillance began

as soon as the first confirmed infections were detected in Brazil, by the end of June 2021 there

was still a paucity of genomic data from some states such as Roraima (n=29), Acre (n=29),

Rondônia (n=37), Tocantins (n=27), Piauí (n=19) and the Federal District (n=33) (**Fig. 2A**). Half

of all Brazilian genomes were deposited in early 2021, suggesting intensified surveillance in the

second wave following the detection of Gamma (and other VOCs (e.g. Alpha/B.1.1.7) and

VUMs (e.g. Zeta) throughout the country (**Fig. 2B**).

To obtain a better understanding of the dynamics of SARS-CoV-2 spread in Brazil, we coupled epidemiological data with phylodynamic analysis for a data set comprising 25,288 available globally representative genomes, including the new genomes obtained in this study (n=3,866) sampled from December 26th, 2019 to June 28th 2021 (**Fig. 2C, Fig. S2**). A date-stamped phylogeny of these data indicated that most of the Brazilian sequences were interspersed with those introduced from several countries (**Fig. 2C, 2D**). This pattern further indicated that the co-circulation of multiple SARS-CoV-2 lineages over time was linked to multiple importations followed by large local transmissions concomitant with a high number of infections (**Fig. 2C, 2D**).

Using an ancestral location state reconstruction on the dated phylogeny we were able to infer the number of viral imports and exports between Brazil and the rest of the world, and between individual Brazilian regions (hereafter referred as the North, Northeast, Midwest, Southeast and South regions) (**Fig. 2D-F**). The bulk of imported introductions (estimated to be 114 independent ones) were largely from Europe (**Fig. 2D**), occurring before the implementation of restriction measures (April 2020) when the epidemic was rapidly progressing (**Fig. 2D**, **E**). However, at least 33 introduction events are inferred to have occurred during enforcement of preventive measures up to August 2020 (**Fig. 2D**, **E**), and hence before those measures were loosened. Finally, although Brazil was a major virus importer, there were approximately 10 times more inferred exportation events out of Brazil than viral introductions into Brazil (**Fig. 2E**).

Our estimates of viral movement within Brazil further suggested that the Southeast region was the largest contributor of viral exchanges to other regions, comprising approximately 40% of viral movements from one geographical region to another, followed by the North region that contributed to approximately 25% of all viral movements. Although these estimates are in line

with epidemiological data, this observation is also likely to be influenced by these two regions

having the greatest number of sequences available for analysis.

Spatiotemporal spreading of Brazilian VOC (Gamma) and VUM (Zeta)

We next focused on the two Brazilian variants that evolved from the B.1.1.28 lineage and

grew into large transmission clusters during the second wave of the epidemic from January 2021

- the VOC (Gamma/P.1) and VUM (Zeta/P.2). To assess the detailed evolution of these lineages

over time we performed a spatiotemporal phylogeographic analysis using a molecular clock

model.

The Gamma VOC was first sampled in Brazil in early January 2021[12,20]. It displayed an

unusual number of lineage-defining mutations in the S protein, including three designated that

may impact transmission, immune escape, and virulence - N501Y, E484K and K417T[21–23]. In

line with previous estimates[12,20], our phylogeographic analysis suggested that the Gamma variant

emerged around November 21st 2020 (95% highest posterior density, 12-29 November 2020) in

Manaus (Amazonas state) in Northern Brazil and spread extensively among Brazilian regions

(**Fig. 3A, C**). Our data reveals multiple introductions of this lineage from the Amazonas state to

Brazil's southeastern, northeastern and midwestern states (**Fig. 3A, C**). By mid-January 2021,

the southeastern and northern regions had also acted as source populations for the introduction of

this variant into the southern region (**Fig 3A, C**).

Zeta (P.2) is defined by the presence of the S:E484K mutation in the receptor binding

domain (RBD) and other lineage-defining mutations outside the S protein[15,24]. Although it was

first described in samples from October 2020 in the state of Rio de Janeiro, our phylogeographic

reconstruction suggests that the variant originated in Paraná state in South Brazil late August

2020 (95% highest posterior density, 19 August 2020 to 03 September 2020) (**Fig. 4B**). Since

then, Zeta has spread multiple times to much of the southeastern, northeastern, midwestern and

northern Brazilian regions (**Fig. 4D**). Together, our results further suggest that the transmission

dynamics roughly followed patterns of population density, moving most often between the most

populous localities (**Fig. A, B**).

By estimating the pattern of migration flows we also examined the potential role of Brazil

as an exporter of the Gamma and Zeta variants to the rest of the world (**Fig. 3**). While the North

region seeded approximately 47% of all Gamma infections into other regions, consistent with it

being where this lineage originated, there is strong evidence both from phylogeographic analysis

(**Fig. 3A**) and ancestral state reconstruction (**Fig. 3C**) that there was considerable subsequent

transfer of Gamma between all regions. Zeta had a different dispersal pattern to Gamma, with

73% of all Zeta movements originating in the Southeast and South regions, consistent with our

phylogeographic reconstruction that this is the geographic source of this lineage (**Fig. 3**).

Our analysis further revealed that Brazil has contributed to the international spread of

both variants with at least 316 and 32 exportation events to the rest of the world detected for

Gamma and Zeta variants, respectively (**Fig. 3E**). Consistent with importations, most exports

were to South America (65%) and Europe (14%), followed by Asia (11%), North America (5%),

Africa (2.5%) and Oceania (2.5%) with an increase between January 2021 and March 2021

coinciding with the second wave of infections in Brazil and some relaxation of international

travel restrictions (**Fig. 3E**). As shown elsewhere, these results demonstrate that under relaxation

of travel restrictions SARS-CoV-2 lineages can spread to a diverse range of international

locations[25-30].

Cross-border SARS-CoV-2 transmission from Brazil to Paraguay

To explore the burden of the Brazilian SARS-CoV-2 pandemic on other South American

countries, we provide a preliminary overview of the SARS-CoV-2 epidemic in Paraguay. The

first COVID-19 confirmed case was documented in Paraguay on March 7[th] 2020 in a 32-year-old

man from San Lorenzo, Central Department. Thirteen days later, the first death and the first case

of community transmission were also confirmed. COVID-19 cases in Paraguay rose sharply in

March (**Fig. 4A**) resulting in 100% occupancy of intensive care beds, prompting the government

to declare a strict quarantine to mitigate the spread of the virus[31,32]. By the end of June 2021 a

total number of 460,000 confirmed cases and 15,000 coronavirus-related deaths had been

reported in Paraguay[31].

The COVID-19 epidemic in Paraguay can generally be characterized by three phases:

phase I staring from March 10[th] 2020 characterized by restriction measures; phase II since May

4[th], 2020, also called "Intelligent/Smart Quarantine" with a gradual return to work and social

activities; and phase III implemented since October 5[th] 2020, known as the "COVID way of

living", characterized by the relaxation of the restrictions measures and the reopening of national

borders and resumption of international flights[32].

Since the beginning of the epidemic there has been a paucity of whole genomes

sequences from Paraguay, with only n=165 whole genome sequences available on GISAID by

the end of July 2021, about 0.0003% of known cases. This seriously impacts the ability to

characterize the molecular epidemiology of SARS-CoV-2 at a regional level. In collaboration

with the Pan-American Health Organization and the National Public Health Laboratory of

Asunción in Paraguay we obtained a total of 63 near complete genome sequences sampled

between July 2020 and June 2021, representing ~ 40% of the currently available genomes from

this country. The selection of the samples was based on the Ct value (≤30) and availability of

epidemiological metadata, such as date of sample collection, sex, age and municipality of

residence. Thus, by applying these inclusion criteria only 63 positive samples were considered

suitable for this study. As expected, we observed the co-circulation of multiple SARS-CoV-2

lineages **(Fig. 4B)**, linked to multiple importations and subsequently characterized by large

transmission clusters.

Importantly, our phylogenetic analysis revealed that most of the SARS-CoV-2 variants

currently circulating in Paraguay, including lineages (B.1.1.28; B.1.1.33; Zeta and Gamma),

originally emerged in Brazil (Fig. 4A and Fig.4B), thus suggesting cross-border transmission

from Brazil to Paraguay (**Fig. 4B, C**). This reinforces the importance of nonpharmaceutical

measures in containing and preventing the spread of viral strains into neighbouring countries.

As of July 31[th], 2021 a total of 78% of available genomic sequences from Paraguay were

linked to infections caused by Brazilian variants, with the Gamma VOC being the most prevalent

lineage in the country. As genome sequencing is not widespread, it is difficult to determine how

widely these variants have spread within Paraguay and to other Latin American countries.

However, the abundant COVID-19 cases in Brazil, a country that shares borders with ten

countries, suggests that this risk is likely to be high.

Discussion

Genome sequencing combined with epidemiological assessment has been widely used as

a tool to track the spread and evolution of SARS-CoV-2, enabling public health authorities to

tailor their control strategies. In addition, genomic data can reveal key properties of emerging

viruses, including precise epidemiological behaviour that patient data alone cannot capture.

However, the utility of genomic data is marred by sparse sampling and geographic inequities[33].

Brazil is an important case study, but SARS-CoV-2 sequences from Brazil are only available

from a tiny fraction of the number of confirmed cases into the country, curbing their utility in a

public health setting and limiting effective control strategies.

To help overcome these limitations, we report genomic data obtained by sequencing

3,866 SARS-CoV-2 infection cases confirmed by RT-qPCR from patients residing in 8 of the 27

Brazilian federal states and the neighbouring country of Paraguay. These were analysed together

with n=13,328 and n=102 (up to 30, June 20[th] 2021) publicly available complete genomes

sequences from both countries, respectively. By combining epidemiological and genomic data,

we show how the interplay between the implementation of restriction measures and sustained

SARS-CoV-2 transmission have shaped the Brazilian epidemic over 20 months, including the

dramatic resurgences in case numbers linked with the emergence of VOCs and VUMs. In

particular, we show that multiple independent importations of SARS-CoV-2, predominantly

from Europe, had occurred in Brazil during the early phase of the epidemic (up to April 2020).

We further detected multiple (n=33) international introductions during periods characterized by

the enforcement of preventive measures, demonstrating that these can be ineffective to prevent

importation, as previously reported in other countries[34].

Our analysis shows that viral importations were present throughout the entirety of the

study period, except between May and September 2020 and further suggest that Brazil played

much more as a viral exporter, resulting in 10 times more inferred exportations events than viral

introduction into the country (**Fig. 2E**). Importantly, this was also a period linked to the

emergence of Brazilian VOCs and VUMs. Although we have not measured passenger traffic to

and from Brazil, the viral migration pattern observed appears to be in line with results obtained

from other studies combining viral genetic data with epidemiological and travel data, and provide evidence that the dispersion of respiratory viruses is dictated by human mobility and population density[26-30].

We also provide the first preliminary overview of the SARS-CoV-2 epidemic in Paraguay, revealing a high virus connectivity with Brazil. This spread was likely facilitated by airplane and road networks that form major transport trails linking Brazil to other parts of South America. The rapid unhampered spread of such VOCs and VUMs into South American countries, such as Paraguay, indicates that the land-border controls to curb the international spread of the virus have been largely ineffective and highlights the inherent challenges in screening cross-border travellers to contain the spread of the virus within this region. Our data further suggest that national and international travel restrictions lifted at certain points during the Brazilian epidemic were likely responsible for both virus introduction from international localities and within-country transmission, with infected travelers acting as carriers.

More broadly, our study highlights the utility of SARS-CoV-2 genome sequencing in COVID-19 outbreak investigation and the need for more comprehensive country-wide studies of the epidemiology and spread of emerging viral strains. Since the disproportion on the number of genome sequences available for each Brazilian state/region, our results further highlight the need of more public investments to strength the genomic capacity across the country.

Additionally, the current co-circulation of VUMs and VOCs in Brazil, together with the slow vaccine rollout, has important implications for public health in this high populous and regionally important country[35]. Such epidemiological conditions create the perfect environment for the continued evolution of SARS-CoV-2, potentially enabling the emergence of novel variants of altered phenotype.

Materials and Methods

Ethics statement

This research was approved by the Ethics Review Committee of the Pan American Health Organization (PAHOERC.0344.01), the Federal University of Minas Gerais (CEP/CAAE: 32912820.6.1001.5149), the University of São Paulo (CEP/FZEA: 4.780.992) and the Blood Center of Ribeirão Preto (CEP/HCRP-FMRP: 50367721.7.1001.5440), and by the Paraguayan Ministry of Public Health and Social Welfare (MSPyBS/ S.G. no. 0944/18). The availability of these samples for research purposes during outbreaks of national concern is allowed under the terms of the 510/2016 Resolution of the National Ethical Committee for Research – Brazilian Ministry of Health (CONEP - Comissão Nacional de Ética em Pesquisa, Ministério da Saúde) that authorizes, without the necessity of an informed consent, the use of clinical samples collected in the Brazilian Central Public Health Laboratories to accelerate knowledge building and contribute to surveillance and outbreak response. The samples processed in this study were obtained anonymously from material exceeding the routine diagnosis in Brazilian public health laboratories that belong to the public network within BrMoH.

Genomic surveillance network

As a part of the National Network for Pandemic Alert of SARS-CoV-2, we performed genomic monitoring to rapidly understand the spread of SARS-CoV-2 at the national and cross border levels. This was performed in collaboration with the Central Laboratory of Public Health (LACEN) from the states of Bahia (BA), Mato Grosso (MT), Mato Grosso do Sul (MS), Minas Gerais (MG), as well as Blood Center of Ribeirão Preto, the Fiocruz Paraná (PR), the Butantan

Institute from the state of São Paulo (SP) state, the University of São Paulo and the Central

Laboratory of Health of Paraguay.

Sample collection and molecular diagnostic assays

Convenience clinical samples from patients with suspected SARS-CoV-2 infection from patients

residing in 8 of the 27 Brazilian federal states and from Asunción, Paraguay, and collected

between July 2020 and June 2021 were provided for diagnostic and genome sequencing

purposes. Viral RNA was extracted from nasopharyngeal swabs using an automated protocol and

tested for SARS-CoV-2 by multiplex real-time PCR assays: (i) the Allplex 2019-nCoV Assay

(Seegene) targeting the envelope (E), the RNA dependent RNA polymerase (RdRp) and the

nucleocapsid (N) genes; (ii) the Charité: SARS-CoV2 (E/RP) assay (Bio-Manguinhos/Fiocruz)

targeting the E gene, and (iii) the GeneFinder COVID-19 Plus RealAmp Kit (Osang Healthcare,

South Korea) supplied by the BrMoH, Butantan Institute and the Pan-American Health

Organization (OPAS).

cDNA synthesis and whole genome sequencing

Samples were selected for sequencing based on the Ct value (≤30) and availability of

epidemiological metadata, such as date of sample collection, sex, age and municipality of

residence. The preparation of SARS-CoV-2 genomic libraries was performed using both the

Illumina COVIDSeq test following the manufacturer's instructions and nanopore sequencing

using the ARTIC Network primal scheme (https://github.com/artic-network/artic-

ncov2019/tree/master/primer_schemes/nCoV-2019/V3)[36]. The normalized libraries were loaded

onto a 300-cycle MiSeq Reagent Kit v2 and run on the Illumina MiSeq instrument (Illumina, San

Diego, CA, USA).

Due to regional characteristics and the accessibility of resources of the different Brazilian

and Paraguayan laboratories, we also applied field SARS-CoV-2 sequencing using the Oxford

Nanopore MinION technology. In this case the SuperScript IV Reverse Transcriptase kit

(Invitrogen) was initially used for cDNA synthesis following the manufacturer's instructions.

The cDNA generated was subjected to multiplex PCR sequencing using the Q5 High Fidelity

Hot-Start DNA Polymerase (New England Biolabs) and a set of specific primers designed by the

ARTIC Network for sequencing the complete SARS-CoV-2 genome (Artic Network version

3)[36]. PCR conditions have been previously reported in[36]. All experiments were performed in a

biosafety level-2 cabinet. Amplicons were purified using 1x AMPure XP Beads (Beckman

Coulter) and quantified on a Qubit 3.0 fluorimeter (Thermofisher Scientific) using Qubit™

dsDNA HS Assay Kit (Thermofisher Scientific). DNA library preparation was performed using

the Ligation Sequencing Kit LSK109(Oxford Nanopore Technologies) and the Native Barcoding

Kit (NBD104 and NBD114, Oxford Nanopore Technologies). Sequencing libraries were loaded

into a R9.4 flow cell (Oxford Nanopore Technologies). In each sequencing run, we used negative

controls to prevent and check for possible contamination with less than 2% mean coverage.

Generation of consensus sequences from Illumina and nanopore

The genome assembly pipeline for Illumina reads involved: (i) read trimming and filtering using

Trimmomatic[37]; (ii) minimap2[38]for read mapping against the reference strain (Wuhan-Hu-1

genome reference - NCBI accession NC_045512.2); (iii) samtools[39] for sorting and indexing;

(iv) Pilon[40] for improving the indel detection; (v) bwa mem[41] for remapping against Pilon's

generated consensus; (vi) samtools mpileup to generate alignment quality values; (vii) seqtk[42] to

generate a quasi-final genome version; (viii) bwa mem for a 3rd round of remapping reads

against the quasi-final genome; and (ix) samtools depth to assess position depths given the *.bam

file from the previous step (nucleotide positions with read depth < 5 are denoted as "N").

Oxford Nanopore sequencing raw files were basecalled using Guppy v3.4.5 and barcode

demultiplexing was performed using qcat. Consensus sequences were generated by *de novo*

assembling using Genome Detective[43] that uses DIAMOND to identify and classify candidate

viral reads in broad taxonomic units, using the viral subset of the Swissprot UniRef protein

database. Candidate reads were next assigned to candidate reference sequences using NCBI

blastn and aligned using AGA (Annotated Genome Aligner) and MAFFT. Final contigs and

consensus sequences are then available as FASTA files.

Data quality control and global data set collection

To ensure the quality of the genome sequences generated in this study and to guarantee the

highest possible phylogenetic accuracy, only genomes >29,000bp and <1% of ambiguities were

considered (n=3,866). Multiple sequence alignment was performed with MAFFT[44] and a

preliminary phylogenetic tree was inferred using the maximum likelihood method in IQ-TREE2,

employing the GTR+I model of nucleotide substitution. Prior to further phylogenetic analysis

our data set was also assessed for both sequences with low data quality (e.g. with assembling

issues, sequencing and alignment errors, data annotation errors and sample contamination) and

for molecular clock signal (i.e. temporal structure) using TempEst v1.5.3[45].

We appended the 3,866 genome sequences newly generated under this project with an

extensive reference data set of SARS-CoV-2 sequences sampled globally collected since the start

of the outbreak, including n=13,328 near-complete genomes from Brazil and n=102 from

Paraguay (sampled up to June 30, 2021). A unique set of external references (n=7,992) were

obtained by including all the non-Brazilian and non-Paraguayan sequences from the global and

South American NextStrain builds (date of access; June 30[th], 2021). These global references

were further supplemented by the inclusion of a random sample of P.1 and P.2 lineages from

around the world. This approach was taken to ensure we captured sufficient global sequence

diversity while at the same time enriching our data set with South American sequences with

whom Brazil and Paraguay share borders. Additionally, during our down sampling strategy, we

took into account that the different sequencing efforts employed by different Brazilian states

resulted in differing numbers of available genomes for each state/region. For this reason, we

selected reference sequences considering both the number of reported cases for each state and the

number of available genome sequences, aiming at building a more representative data set (**Fig.**

S3).

Phylogenetic analysis

Sequences were aligned using MAFFT[44] and submitted to IQ-TREE2 for maximum likelihood

(ML) phylogenetic analysis[46] employing the general time reversible (GTR) model of nucleotide

substitution and a proportion of invariable sites (+I) as selected by the ModelFinder application.

Branch support was assessed using the approximate likelihood-ratio test based on the bootstrap

and the Shimodaira–Hasegawa-like procedure (SH-aLRT) with 1,000 replicates.

The raw ML tree topology was then used to estimate the number of viral transmission

events between various Brazilian regions and the rest of the world. TreeTime[47] was used to

transform this ML tree topology into a dated tree using a constant mean rate of 8.0 x 10^{-4}

nucleotide substitutions per site per year, after the exclusion of outlier sequences. A migration

model was fitted to the resulting time-scaled phylogenetic tree in TreeTime, mapping country

and regional locations to tips and internal nodes[47]. Using the resulting annotated tree topology

we were able to count the number of transitions (i.e. virus importations and exportations) within

different Brazilian regions and between Brazil and the rest of the world. Importantly, this

analysis was not dependent on a monophyletic clustering of SARS-CoV-2 in Brazil. To provide

a measure of confidence in the time and source of viral transitions, we performed the discrete

ancestral state reconstruction on 10 bootstrap replicate trees (**Fig. S4-S6**).

Lineage classification

We used the dynamic lineage classification as specified in the Phylogenetic Assignment of

Named Global Outbreak LINeages (Pangolin version 3.1.7) protocol[48]. This was aimed at

identifying the most epidemiologically important lineages of SARS-CoV-2 circulating within

South America (Brazil and Paraguay). Both VOCs and VUMs were designated based on the

World Health Organization framework as of July, 2021.

Phylogeographic reconstruction

VOCs and VUMs that emerged in Brazil (i.e. P.1 and P.2) were identified as monophyletic

groups on the time-resolved phylogenetic trees (**Fig.2 and Fig. S7**). Genome sequences from

these lineages were then extracted to infer continuous phylogeography histories using the

Markov chain Monte Carlo (MCMC) method in BEAST v1.10.4[49], employing the HKY + Γ

_nucleotide substitution model and a strict molecular clock in BEAST v.1.10.4 that assumes constant

evolutionary rates throughout the phylogeny.

Due to the huge size of the largest clusters we down-sampled them to <600 taxa per clade

to infer phylogeographies within a manageable time frame. Accordingly, we retained the ten

earliest sequences from each unique sampling location within Brazil and a random distribution

over the remaining samples from each location (n=531 for P.1 and n=428 for P.2). Briefly,

sequences from the subsampled cluster were aligned using MAFFT and preliminary ML trees

were inferred in IQ-TREE2 as described above. Prior to phylogeographic analysis, each lineage

was also assessed for molecular clock signal using the root-to-tip regression method available in

TempEst v1.5.3[45] following the removal of potential outliers that may violate the molecular

clock assumption. We accepted temporal structure when the correlation coefficient was > 0.2[50].

Linear regression of root-to-tip genetic distances against sampling dates indicated that the SARS-

CoV-2 sequences from Gamma and Zeta evolved in a relatively clock-like manner (r^2= 0.5;

coefficient correlation= 0.45 for Gamma and r^2= 0.5; coefficient correlation= 0.40 for Zeta) (**Fig.**

S7).

We modelled the phylogenetic diffusion and spread of these lineages within Brazil by

analysing localized transmission (between Brazilian regions) using a flexible relaxed random

walk (RRW) diffusion model[51,52] that accommodates branch-specific variation in rates of

dispersal with a Cauchy distribution and a jitter window site of 0.01[52]. The choice of Cauchy

distribution was based on recent studies demonstrating that it is successful for this kind of

analyses based on genomes of SARS-CoV-2 and its variants[4,50,54-56]. For each sequence, latitude

and longitude coordinates were attributed. MCMC analyses were set up in BEAST v1.10.4,

running in duplicate for 100 million interactions and sampling every 10,000 steps in the chain.

Convergence for each run was assessed in Tracer v1.7.1 (ESS for all relevant model parameters

>200). Maximum clade credibility trees for each run were summarized using TreeAnnotator after

discarding the initial 10% as burn-in. While the sampling is relatively homogeneous among

sampled locations, the phylogeographic reconstruction will of course remain sampling-

dependent. Hence, differences in sampling effort may have impacted the estimated transition

frequencies between locations. Despite this caveat, the phylogeographic analysis still provides

important information on the history and dynamics of dispersal of the lineages sampled, which

can in turn provide insights into the connectivity of locations along the transmission network.

Finally, we used the R package "seraphim"[55] to extract and map spatiotemporal information

embedded in the posterior trees. Note that a transmission link on the phylogeographic map can

denote one or more transmission events depending on the phylogeographic inference.

SARS-CoV-2 time series from Brazil

Data from weekly notified and laboratory confirmed cases of infection by SARS-CoV-2 in

Brazil, were supplied by the Brazilian Ministry of Health, as made available by the COVIDA

network at https://github.com/wcota/covid19br. For convenience, the geographical locations

were aggregated by Brazilian macro regions: North, Northeast, Southeast, South, and Midwest.

References

1. Coronavirus Disease (COVID-19) Situation Reports.

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports.

2. COVID-19 Map. *Johns Hopkins Coronavirus Resource Center*

https://coronavirus.jhu.edu/map.html.

3. COVID-19 Situation Reports - PAHO/WHO | Pan American Health Organization.

https://www.paho.org/en/covid-19-situation-reports.

4. Candido DS, et al. Evolution and epidemic spread of SARS-CoV-2 in Brazil. *Science*. **4**,

369(2020)

5. Xavier J, et al. The ongoing COVID-19 epidemic in Minas Gerais, Brazil: insights from

epidemiological data and SARS-CoV-2 whole genome sequencing. *Emerg Microbes Infect*.

9,1824(2020).

6. Coronavírus Brasil. https://covid.saude.gov.br/.

7. Cota, W. Monitoring the number of COVID-19 cases in Brazil. *Wesley Cota - Physics and*

Complex Networks https://covid19br.wcota.me/en/.

8. COVID-19 outbreak in Brazil: adherence to national preventive measures and impact on

people's lives, an online survey | BMC Public Health | Full Text.

https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-021-10222-z.

9. Pinto Neto, O. *et al.* Mathematical model of COVID-19 intervention scenarios for São

Paulo—Brazil. *Nat Commun* **12**, 418 (2021).

10. Monteiro de Oliveira, M., Fuller, T. L., Brasil, P., Gabaglia, C. R. & Nielsen-Saines, K.

Controlling the COVID-19 pandemic in Brazil: a challenge of continental proportions. *Nat*

Med **26**, 1505–1506 (2020).

11. Candido DDS. et al. Routes for COVID-19 importation in Brazil. *J Travel Med* **18**, 27

(2020).

12. Faria NR, et al. Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus,

Brazil. *Science* **21**,372 (2021).

13. Naveca FG, et al. COVID-19 in Amazonas, Brazil, was driven by the persistence of endemic

lineages and P.1 emergence. COVID-19 in Amazonas, Brazil, was driven by the persistence

of endemic lineages and P.1 emergence. *Nat Med*. **27**,1230 (2021).

14. Sant'Anna FH, et al. Emergence of the novel SARS-CoV-2 lineage VUI-NP13L and massive

spread of P.2 in South Brazil. *Emerg Microbes Infec.* **10**, 1431(2021).

15. Voloch, C. M. *et al.* Genomic characterization of a novel SARS-CoV-2 lineage from Rio de

Janeiro, Brazil. *J Virol* JVI.00119-21 (2021).

16. de Oliveira Andrade, R. Covid-19 is causing the collapse of Brazil's national health service.

BMJ **370**, m3032 (2020).

17. Vacinação contra a Covid-19 no Brasil - #PÁTRIAVACINADA.

https://www.gov.br/saude/pt-br/vacinacao.

18. Vacinas contra a COVID-19 - OPAS/OMS | Organização Pan-Americana da Saúde.

https://www.paho.org/pt/vacinas-contra-covid-19.

19. Brazil – COVID19 Vaccine Tracker. https://covid19.trackvaccines.org/country/brazil/.

20. Naveca, F. G. et al. COVID-19 in Amazonas, Brazil, was driven by the persistence of

endemic lineages and P.1 emergence. *Nat Med* **27**, 1230 (2021).

21. Weisblum, Y. et al. Escape from neutralizing antibodies by SARS-CoV-2 spike protein

variants. *eLife* **9**, e61312 (2020).

22. Nonaka, C. K. V. et al. Genomic Evidence of SARS-CoV-2 Reinfection Involving E484K

Spike Mutation, Brazil. *Emerging Infectious Diseases journal*. **5**, 27 (2021).

23. CDC. Coronavirus Disease 2019 (COVID-19). *Centers for Disease Control and Prevention*

https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-

emerging-variants.html (2020).

584 24. Sant'Anna, F. H. *et al.* Emergence of the novel SARS-CoV-2 lineage VUI-NP13L and

585   massive spread of P.2 in South Brazil. *Emerging Microbes & Infections* **10**, 1431–1440

586   (2021).

587 25. Kraemer MUG, et al. The effect of human mobility and control measures on the COVID-19

588   epidemic in China. *Science.* **1**, 368. (2020).

589 26. Lemey P, et al. Unifying viral genetics and human transportation data to predict the global

590   transmission dynamics of human influenza H3N2. *PLoS Pathog*. **5**, 29 (2014).

591 27. Lu L, et al. Quantifying predictors for the spatial diffusion of avian influenza virus in China.

592   *BMC Evol Biol*. **67**, 198. (2017).

593 28. Ganasegeran K, et al. Influence of Population Density for COVID-19 Spread in Malaysia:

594   An Ecological Study. *Int J Environ Res Public Health*. **4**, 76 (2021).

595 29. Magee D, et al. Combining phylogeography and spatial epidemiology to uncover predictors

596   of H5N1 influenza A virus diffusion. *Arch Virol*. **31**, 96 (2015).

597 30. du Plessis L, et al. Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the

598 UK. *Science*. **12**, 371 (2021).

599 31. PAHO Paraguay COVID-19 Response. https://paho-covid19-response-

600   who.hub.arcgis.com/pages/paho-paraguay-covid-19-response.

601 32. Paraguay: WHO Coronavirus Disease (COVID-19) Dashboard With Vaccination Data.

602   https://covid19.who.int.

603 33. Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data – from

604   vision to reality. *Euro Surveill* **22**, 30494 (2017).

605 34. Jorge DCP, et al. Assessing the nationwide impact of COVID-19 mitigation policies on the

606   transmission rate of SARS-CoV-2 in Brazil. *Epidemics*. **35**,100.

35. Victora PC, et al. Estimating the early impact of vaccination against COVID-19 on deaths

among elderly people in Brazil: Analyses of routinely-collected data on vaccine coverage

and mortality. *EClinicalMedicine*. **38**,101.(2021)

36. Quick, J. nCoV-2019 sequencing protocol. (2020) doi:10.17504/protocols.io.bbmuik6w.

37. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics*. **1**, 30 (2014).

38. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. **15**,34 (2018).

39. Danecek P, et al. Twelve years of SAMtools and BCFtools. *Gigascience*. 16, 10 (2021).

40. Walker BJ, et al. Pilon: an integrated tool for comprehensive microbial variant detection and

genome assembly improvement. *PLoS One*. **19**,9 (2014).

41. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. **15**, 25 (2009).

42. Shen W, et al. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File

Manipulation. *PLoS One*. 11,10 (2016).

43. Vilsker M, et al. Genome Detective: an automated system for virus identification from high-

throughput sequencing data. *Bioinformatics*. **1**, 35 (2019).

44. Katoh K, et al. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in

Performance and Usability. *Mol Biol Evo*l. **30**, 772 (2013)

45. Rambaut, A., Lam, T. T., Max Carvalho, L. & Pybus, O. G. Exploring the temporal structure

of heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol* **2**, vew007

(2016).

46. Minh BQ, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference

in the Genomic Era. *Mol Biol Evol*. **1**, 37 (2020).

47. Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood phylodynamic

analysis. *Virus Evol* **4**, vex042 (2018).

48. Rambaut A, et al. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist

genomic epidemiology | *Nat Microbiol*. **5**, 1403 (2020).

49. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian

phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evol*. **2**,298 (2018).

50. Tegally, H., Wilkinson, E., Lessells, R.J. et al. Sixteen novel lineages of SARS-CoV-2 in

South Africa. *Nat Med.* **27**, 440 (2021).

51. Lemey, P., Rambaut, A., Welch, J. J. & Suchard, M. A. Phylogeography takes a relaxed

random walk in continuous space and time. *Mol Biol Evol* **27**, 1877–1885 (2010).

52. Dellicour S, et al. Relax, Keep Walking - A Practical Guide to Continuous Phylogeographic

Inference with BEAST. *Mol Biol Evol*. **29**, 38 (2021).

53.Dellicour, S., Rose, R., Faria, N. R., Lemey, P. & Pybus, O. G. SERAPHIM: studying

environmental rasters and phylogenetically informed movements. *Bioinformatics* **32**, 3204–3206

(2016).

54. Tegally H, et al. Detection of a SARS-CoV-2 variant of concern in South Africa. Nature. doi:

10.1038/s41586-021-03402-9;

55. Viana R, et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern

Africa. Nature, doi: 10.1038/s41586-022-04411-y.

56. Dellicour S, Hong SL, Vrancken B, Chaillon A, Gill MS, Maurano MT, Ramaswami S,

Zappile P, Marier C, Harkins GW, Baele G, Duerr R, Heguy A. Dispersal dynamics of SARS-

CoV-2 lineages during the first epidemic wave in New York City. PLoS Pathog. 2021

20;17(5):e1009571. doi: 10.1371/journal.ppat.1009571.

Acknowledgments: We thank all those who have kindly deposited and shared genome data on GISAID (Table S3). The authors acknowledge the technical support of Luciana de Araujo Pimenta, Luiz Aurelio de Campos Crispin, Gabriela Mauric Frossard Ribeiro, Glaucia Maria Rodrigues Borges, Mariane Evaristo e Josiane Serrano Borges from the National Network for Pandemic Alert of SARS-CoV-2 and the contribution of all employees of General Coordination of Public Health Laboratories and professionals of Public Health Laboratories of Brazil and Network for Pandemic Alert of Emerging SARS-CoV-2 Variants for their contribution towards the sequencing effort and for their commitment and work during the fight of the COVID-19 pandemic.

Funding: This study was financed by São Paulo Research Foundation (FAPESP) (Grants Number: 2020/10127-1 and 2020/05367-3, 2013/08135-2), the Central Public Health Laboratories, Blood Center of Ribeirão Preto and supported by the Brazilian Ministry of Health and the Pan American Health Organization PAHO/WHO (APO21-00010098). This work was supported in part through National Institutes of Health USA grant U01 AI151698 for the United World Arbovirus Research Network (UWARN), the CRP- ICGEB RESEARCH GRANT 2020 Project CRP/BRA20-03, Contract CRP/20/03, the Oswaldo Cruz Foundation VPGDI-027-FIO-20-2-2-30, the Brazilian Ministry of Health (SCON2021-00180), the CNPq (426559/2018-5) and the Faperj E-26/202.930/2016. MG and LCJA are supported by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) (E-26/202.248/2018(238504); E26/202.665/2019(247400)). JX is supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) - Finance Code 001. JL is supported by a lectureship by the Department

of Zoology of the University of Oxford. VF is supported by Coordenação de Aperfeiçoamento de

Pessoal de Nível Superior (CAPES) (88882.349290/2019-01) and a Flagship grant from the

South African Medical Research Council (MRC-RFA-UFSP-01-2013/UKZN HIVEPI). ECH is

funded by an ARC Australian Laureate Fellowship (FL170100022). MLN is supported by

FAPESP (Grant 20/04836-0) and partially supported by a NIH grant to CreateNeo

(U01AI151807).

Author contributions:

<u>Molecular screening and produced SARS-CoV-2 genomic data:</u> MG, SNS, VLV, ESR, EVS,

FA, JX, HF, TERA, LPOL, AJM, CRSB, ECM, JSTB, DBM, RAB, RLRCC, PDSCM, JPK, BS,

RPS, VVC, ST, VBN, LROS, MKAG, JQL, AAR, NRG, LTW, LBS, RSF, MPFP, MJO, SGF,

SV, CA, GPMG, WSF, ARMC, JV, SMVLO, CCMG, MCDR, GNB, MPG, NQM, INR, SR,

GMC, AHDC, CZ, CDS, PAA, FASC, MDP, JCCL, ECM, CAB, LS, MMM, RMTG, JASN,

MLN, HF, LLC, RTC, RMN; <u>Collected samples and curated metadata:</u> CF, CRLP, CFRF, WN,

RFCS, MA, CCAM, RH, SCS, MCE, SK, LCJA, DTC, JASN, HF, LLC, FP, AL, FCMI, GCP,

CV, GMES, ECO, LD, JC; <u>Analysed the data:</u> MG, VF, EW, HT, JSLP, EJS, RB, JL; <u>Helped</u>

<u>with study design and data interpretation:</u> MG, VF, EW, HT, EH, JL, TdO, SCS, MCE, SK,

LCJA, DTC; <u>Wrote the initial manuscript, which was reviewed by all authors:</u> MG, SNS, SCS,

MCE, SK, LCJA, DTC.

Competing interests: The authors declare no competing interests.

Data and materials availability: All sequences that were generated and used in the present

study are listed in table S1-S3 (accessible on the GitHub repository) along with their GISAID

sequence IDs, dates of sampling, the originating and submitting laboratories and main authors.

All input files (e.g. alignments or XML files), all resulting output files and scripts used in the

study   are   shared   publicly   on   GitHub

(https://github.com/genomicsurveillance/Genomic_epidemiology_reveals_how_restriction_meas

ures_shaped_the_SARS-CoV-2_epidemic_in_Brazil).

ADDITIONAL INFORMATION

Figure S1. Sequencing statistics and lineage assignment of the SARS-CoV-2 genomes

generated in this study. (A) Genome coverage plotted against RT-qPCR cycle threshold value.

Colours represent genomes obtained from Brazil (green) and Paraguay (orange), respectively.

(B) Number of genomes obtained in this study from Brazil and Paraguay. Colours represent

different lineages.

Figure S2. Fully annotated Brazilian SARS-CoV-2 time tree. Time resolved maximum

likelihood phylogeny containing 17,135 high quality Brazilian SARS-CoV-2 near-full-genome

sequences (n=3,866 generated in this study) analysed against a backdrop of global reference

sequences. Variants of interest (VUM) and concern (VOC) are highlighted.

Figure S3. Number of total cases plotted against the number of sequences available for each

location Brazilian state/region.

Figure S4. Sensitivity of the viral introduction analysis to geographic sampling biases. A, B)

Sensitivity analysis, by performing the discrete ancestral state reconstruction on 10 bootstrap

replicate trees, showing how the proportion of international imports and exports into and from

Brazil from external locations did not vary in the time and source of viral transitions. **C-K)**

Equivalent sensitivity analysis showing how the number of viral exchanges within Brazilian

regions by counting the state changes from the root to the tips did not vary in the time and source

of viral transitions.

Figure S5. Sensitivity of the Gamma viral introduction analysis to geographic sampling

biases. A, B) Sensitivity analysis, by performing the discrete ancestral state reconstruction on 10

bootstrap replicate trees, showing how the proportion of international imports and exports of

Gamma variant into and from Brazil from external locations did not vary in the time and source

of viral transitions. **C-K)** Equivalent sensitivity analysis showing how the number of exchanges

of the Gamma variant between Brazilian regions did not vary in the time and source of viral

transitions.

Figure S6. Sensitivity of the Zeta viral introduction analysis to geographic sampling biases.

A, B) Sensitivity analysis, by performing the discrete ancestral state reconstruction on 10

bootstrap replicate trees, showing how the proportion of international imports and exports of

Zeta variant into and from Brazil from external locations did not vary in the time and source of

viral transitions. **C-K)** Equivalent sensitivity analysis showing how the number of exchanges of

the Zeta variant between Brazilian regions did not vary in the time and source of viral transitions.

Figure S7. Temporal Signal. (A) Root-to-tip regression plots for Brazilian lineages of concern

and of interest (P.1 – Gamma and P.2- Zeta).

Supplementary Table 1. Information on the n=3803 SARS-CoV-2 Brazilian samples sequenced

as part of this study.

Supplementary Table 2. Information on the n=63 SARS-CoV-2 from Paraguay samples

sequenced as part of this study.

Supplementary Table 3. GISAID acknowledgment table.

FIGURE LEGENDS

Fig. 1. Key events following the first confirmed infection of SARS-CoV-2 in Brazil. (A)

Timeline of SARS-CoV-2 key events in Brazil. The Brazilian map was colored according to

geographical macro region: North (red), Northeast (green), Southeast (purple), Midwest (light

blue) South (light orange). (B) Epidemic curve showing the progression of reported daily viral

infection numbers in Brazil from the beginning of the epidemic (grey) and deaths (red) in the

same period, with restriction phases indicated along the bottom. (C) Map of cumulative SARS-

CoV-2 cases per 100,000 inhabitants in Brazil up to June 2021.

Fig. 2. Phylogenetic analysis and SARS-CoV-2 lineage dynamics in Brazil. (A) Map of Brazil

with the number of sequences in GISAID as of 30th June 2021. (B) Temporal sampling of

sequences in Brazilian states through time with VOCs highlighted and annotated according to

their PANGO lineage assignment. (C) Time resolved maximum likelihood phylogeny containing

high qualitynear-full-genome sequences from Brazil (n=3866) obtained from this study, analysed

against a backdrop of global reference sequences (n=25,288). Variants under monitoring (VUM)

and concern (VOC) are highlighted on the phylogeny. (D) Sources of viral introductions into

Brazil characterized as external introductions from the rest of the world. (E) Sources of viral

exchanges (imports and exports) in and outside Brazil. (F) Number of viral exchanges within

Brazilian regions by counting the state changes from the root to the tips of the phylogeny in

panel C.

Fig. 3. Spatiotemporal spread of VOC and VUM in Brazil. (A) Phylogeographic

reconstruction of the spread of the Gamma VOC in Brazil. Circles represent nodes of the

maximum clade credibility phylogeny and are colored according to their inferred time of

occurrence. Shaded areas represent the 80% highest posterior density interval and depict the

uncertainty of the phylogeographic estimates for each node. Differences in population density

are shown on a dark-white scale; B) Phylogeographic reconstruction of the spread of the Zeta

VUM across Brazil. Circles represent nodes of the maximum clade credibility phylogeny and are

colored according to their inferred time of occurrence. Shaded areas represent the 80% highest

posterior density interval and depict the uncertainty of the phylogeographic estimates for each

node. Differences in population density are shown on a dark-white scale; In both Panels (A) and

(B) solid curved lines denote the links between nodes and the directionality of movement is anti-

clockwise along the curve (as shown in the "dispersal direction" sublegends). (C) Number of

exchanges of the Gamma variant between Brazilian regions (N=North; NE=Northeast;

MD=Midwest; SE=Southeast; S=South); (D) Number of exchanges of the Zeta variant between

Brazilian regions; (E) Sources of viral export of the VOC and VUM from Brazil to the rest of the

world.

Fig. 4. The SARS-CoV-2 epidemic in Paraguay. (A) Epidemic curve showing the progression

of reported viral infection numbers in Paraguay from the beginning of the epidemic (grey) and

deaths (red) in the same period; (B) Progressive distribution of the top 20 PANGO lineages in

Paraguay over time; (C) Time resolved maximum likelihood tree containing (n=63) high quality

near-complete genome sequences from Paraguay obtained in this study analysed against a

backdrop of global reference sequences. VUM and VOCs are highlighted on the phylogeny.

Small circles indicate genome sequences from Brazil. Bigger circles indicate sequences from

Paraguay. New genomic sequences from Paraguay obtained in this study are highlighted with a

red border.

A

26 Feb
1st confirmed case
in South America,
61 yo male returning
traveller from Italy

10 April
The virus had reached
remote locations;
a Yanomami teen died
of it in Roraima

20 April
Several cities started
to ease social isolation
guidelines in favor
of contact tracing.

30 April
Brazil overtook China's
official number of
confirmed cases,
surpassing 87,000

21 Mar
All Brazilian states reported
at least one confirmed
case of COVID-19, BR-MoH
declares large-scale community
transmission in the country

17 Mar
1st confirmed death for
COVID-19 , 61 yo male

11 Mar
SARS-CoV-2 confirmed as pandemic
by World Health Organization

Oct-Nov
Zeta (P.2) Variant is identified
in Rio de Janeiro

31 December:
Alpha (B.1.1.7) Variant is identified
in São Paulo

13 January:
Discovery of a new
variant in Manaus,
being called Gamma (P.1)

17 January:
Anvisa has authorized the
emergency use of CoronaVac
and the AstraZeneca vaccines

14 May
Delta (B.1.617.2) Variant is identified
in São Paulo

9 June:
Brazil registered
50 M people vaccinated

7 April:
For the first time,
Brazil has recorded
over 4,000 confirmed
COVID-19 deaths in 24 hours

04 April
Beta (B.1.351) Variant is identified
in São Paulo

| Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Jan | Feb | Mar | Apr | May | Jun |

2020   *2021*

Legend

Restriction measures

First confirmed case

First confirmed death

Vaccination campaign

First Brazilian VOI detected

First Brazilian VOC detected

B

Cases — Deaths ▌ Genome

Phase I   Phase II

cases   *deaths*

100k — 2.500
50k — 1.250
0 — 0

Mar 20 Apr 20 May 20 Jun 20 Jul 20 Aug 20 Sep 20 Oct 20 Nov 20 Dec 20 Jan 21 Feb 21 Mar 21 Apr 21 May 21 Jun 21 Jul 21

Month

C

Total Cases
(millions)
4
3
2
1

Population
(millions)
40
30
20
10

Incidence
(per 100k)
15000
10000
5000

A Gamma Spatiotemporal spreading

Population Density / km²
- 0 - 5
- 5 - 12
- 12 - 18
- 18 - 25
- 25 - 35
- 35 - 55
- 55 - 115
- > 115

Nov 2020
Jan 2021
Feb 2021
Mar 2021
Apr 2021

B Zeta Spatiotemporal spreading

Population Density / km²
- 0 - 5
- 5 - 12
- 12 - 18
- 18 - 25
- 25 - 35
- 35 - 55
- 55 - 115
- > 115

Sep 2020
Nov 2020
Feb 2021
Mar 2021

C
Origin: South, Midwest, Northeast, Southeast, North
Destination: Northeast, South, Midwest, Southeast

Gamma connection: North, Northeast, Midwest, South, Southeast

D
Origin: Northeast, North, Midwest, South, Southeast
Destination: North, South, Northeast, Midwest, Southeast

Zeta connection: North, Northeast, Midwest, South, Southeast

E
Number of exportation events

Lineage
- Gamma
- Zeta

Destination: Africa, Asia, Europe, North America, Oceania, South America