<u>Title Page</u>

Manuscript Type: Original Article

Manuscript Title: Effects of environmental polycyclic aromatic hydrocarbon exposure

and pro-Inflammatory activity on Type 2 Diabetes Mellitus.

Author: Shweta Srivastava

Author's Affiliation: Environmental Health Institute, University of Louisville, Louisville

KY, USA.

Corresponding Author Address:
Shweta Srivastava Ph.D., M.Sc.
302 E. Muhammad Ali Blvd.,
University of Louisville
Louisville KY 40202
Tel.: 502-852-2477
Email: shweta.srivastava.1@louisville.edu
15

Disclosure: Author has no conflicts of interest.

Abbreviations: AUD: alcohol use disorder, BMI: body mass index, CS: clinically
significant, HbA1C: Hemoglobin A1c or Glycated hemoglobin, HD: heavy drinkers,
HOMA: homeostatic model assessment for Types 2 Diabetes, TChol: total cholesterol,
HDL: high density lipids, LDL: low density lipids, MD: moderate drinkers, PC:
Polyaromatic Compounds, SD: social drinkers, SEQN: Respondent sequence number,
RIDSTATR: Interview/Examination status, RIAGENDR: Gender, RIDAGEYR: Age in
years at screening, RIDRETH3: Race/Hispanic origin w/ NH Asian, RIDEXPRG:
Pregnancy status at exam, BMXWT: Weight (kg), BMXHT: Standing Height (cm),
BMXBMI: Body Mass Index (kg/m**2), ALQ101: Had at least 12 alcohol drinks/1 yr.?,
ALQ110: Had at least 12 alcohol drinks/lifetime?, ALQ120Q: How often drink alcohol
over past 12 mos., ALQ120U: # days drink alcohol per wk., mos., yr., ALQ130: Avg #
alcoholic drinks/day - past 12 mos., ALQ141Q: # days have 4/5 drinks - past 12 mos.,
ALQ141U: # days per week, month, year?, ALQ151: Ever have 4/5 or more drinks every
day?, ALQ160: # days have 4/5 or more drinks in 2 hrs., LBXGH: Glycohemoglobin (%),
URXP01: 1-Hydroxynaphthalene (ng/L), URXP02: 2-Hydroxynaphthalene (ng/L),
URXP03: 3-Hydroxyfluorene (ng/L), URXP04: 2-Hydroxyfluorene (ng/L), URXP06: 1-
Hydroxyphenanthrene (ng/L), URXP10: 1-Hydroxypyrene (ng/L), URXP25: 2 & 3-

Hydroxyphenanthrene (ng/L), LBXTC: Total Cholesterol (mg/dL), LBDHDD: Direct HDL-Cholesterol (mg/dL), LBXTR: Triglyceride (mg/dL), LBDLDL: LDL-cholesterol (mg/dL), LBXHSCRP: HS C-Reactive Protein (mg/L), LBXSCR: Creatinine, refrigerated serum (mg/dL), LBXMOPCT: Monocyte percent (%).

Proprietorship: This article is the work of the University of Louisville.

Acknowledgements: The author thanks the NHANES for the dataset.

Funding and Support: This study has been supported by the Environment Health Institute of the University of Louisville, Louisville KY, USA.

Highlights:

1   BMI and Age are demographic risk factors for Diabetes Mellitus Type 2 (DMT2) in high PAH exposure

2   Acute proinflammatory activity characterized by CRP, is augmented by elevated monocyte levels and 1-HN and 2-HN independently

3   Prevalence of higher average drinks over time is observed with high PAH exposure

4   Pathway of DMT2 shows sexual dimorphism, with males showing association with triglycerides, and females with CRP independently with HbA1C

5   The arrangement of CRP, absolute monocyte levels, serum triglycerides and average drinks over time predict the HbA1C levels in individuals with high PAH exposure.

ABSTRACT

Background: Polycyclic aromatic hydrocarbons (PAHs) are formed due to incomplete combustion and known for their potential impact and persistence in the environment. PAHs exposure have been linked to cause adverse health effect including cancer and genetic mutations. The understanding of metabolic effects of PAH exposure are still less clear especially in the presence of pro-inflammatory stress like alcoholism or diabetes.

Objective: The aim of this article is to understand the metabolic effects of PAH exposure by analyzing the clinical biomarkers. This study has also accessed the interactive impact of PAH and other proinflammatory factors, like alcohol intake on the metabolic syndrome, especially Type 2 Diabetes Mellitus (T2DM).

Methods: All the data in this study are retrieved from CDC NHANES (2015-16). We investigated urinary levels of hydroxylated PAH metabolites (OH-PAHs) along with demographic, clinical and laboratory data. Questionnaire data for alcohol use and diabetes status were also included along with laboratory data. Laboratory measures included in the study were levels of PAHs, glycohemoglobin, glucose, cholesterol, lipids, triglyceride, complete blood count, lymphocytes, and monocytes. Generalize linear model Univariate factorial ANOVA was used to evaluate the group differences (both between the groups; as well as across all the groups) in the demographics, PAH exposure, drinking patterns, clinical data, and biomarker levels. Linear regression model was used to analyze the association of biomarkers, PAH exposure and drinking data. Multivariable regression model was used for multi-independent model to assess comorbidity association and their effect sizes on the clinical outcomes.

Results: BMI (p=0.002), and age (≤0.001) are independent demographic risk factors for T2DM in high PAH exposure. Acute proinflammatory activity characterized by CRP, is augmented by elevated monocyte levels (p≤0.001) and stepwise addition of 1-HN (p=0.005), and 2-HN (p=0.001) independently. Prevalence of highest average drinks over time is observed in the high PAH exposure; with males drinking almost twice compared to females in Gr.3. Pathway response of T2DM shows sexual dimorphism; with males showing association with triglycerides (p≤0.001), and females with CRP (p=0.015) independently with HbA1C. The arrangement of CRP, absolute monocyte levels, serum triglycerides and average drinks over time predict the HbA1C levels (adjusted R^2=0.226, p≤0.001) in individuals with high PAH exposure.

Discussion: In this large dataset investigation on humans, the adverse effects of high exposure of PAHs identified candidate demographic risk factors. Preclinical experimental studies on mice have suggested that PAHs exposure induces lipid metabolic disorders in a time-dependent manner, which we found in humans too. Sexual dimorphism is observed in alcohol drinking with males drinking more in the high PAH exposure group. Alcohol drinking as an independent factor associated with the DMT2 indicator, HbA1C in individuals with high PAH exposure.

Keywords: 1-HN, Alcohol, CRP, HbA1C, PAH, T2DM

INTRODUCTION

Polycyclic aromatic hydrocarbons (PAHs) are ubiquitous environmental pollutants linked to many adverse health effects [1, 2]. Polyaromatic Hydrocarbons (PAHs) are a group of chemicals that occur naturally in coal, crude oil, and gasoline. They also are produced when coal, oil, gas, wood, garbage, and tobacco are burned [3, 4]. PAH can also be present in food like charcoal grilled meat and processed food [5, 6]. National Institute for Occupational Safety and Health (NIOSH) recommends workplace air exposure levels of PAHs at less than 0.1 mg/m^3 [7].

Upon entering the body, PAHs are metabolized and eliminated via urine. Urinary concentrations of PAH metabolites, specifically mono-hydroxylated PAHs (OH-PAHs), have been used as biomarkers of human exposure to select PAHs, including naphthalene, fluorene, phenanthrene, and pyrene [8, 9].

PAHs can form small airborne particles and cause a pronounced pulmonary inflammatory response and lung cancer [10, 11]. PAHs have been found to reach to heart through circulation causing cardiotoxicity such as atherosclerosis, cardiac hypertrophy, arrhythmias, and contractile dysfunction [12, 13]. The transport of PAHs to systemic circulation can also significantly affect hepatic metabolism [14-16] since liver is major metabolic organ of PAHs. Inhaled Benzo[a]pyrene are metabolized to generate reactive oxygen species (ROS) in the liver, directly contributing to the hepatoxicity [17].

PAHs have been found to increase likelihood of Type 2 Diabetes Mellitus (T2DM) by 78-124% [18]. This risk is confounded by the fact that some PAHs increase

independent risk factors of T2DM such as obesity, alcohol intake, elevated blood

pressure and blood lipid [19, 20]. However, there are gaps in the understanding of the

clinical presentation or altered mechanisms that can clearly ascertain the role of PAHs

on T2DM. Its adverse effects on T2DM and its clinical marker, HbA1C are still under

investigation. Importantly, interaction of heavy or binge alcohol intake and PAHs on the

exacerbation of metabolic syndrome is not clear.

Majority of the research done so far are mainly focused on respiratory toxicity of

inhaled PAHs. In the present study, the 2015–2016 NHANES data for seven urinary

metabolites of PAH exposure namely (with their code in parenthesis for NHANES

tracking): 1-hydroxynaphthalene (URXP01), 2-hydroxynaphthalene (URXP02), 2-

hydroxyfluorene (URXP04), 3-hydroxyfluorene (URXP03), 1-hydroxyphenanthrene

(URXP06), 2- & 3-hydroxyphenanthrene (URXP25), and 1-hydroxypyrene (URXP10)

was used to assess their effects on T2DM. Primary aim of this study was to characterize

the exposure effect of PAHs on T2DM; and examine the modifying effects of alcohol

intake, blood pro-inflammatory mediators and lipids. A model was also developed to

describe the interaction of high PAHs, modifying risk factors on T2DM. Lastly, the role of

demographic confounders was also explored.

METHODOLOGY

Study Participants, questionnaires, and examination data

Population data were taken from the 2015–2016 NHANES provided by the

National Center for Health Statistics (NCHS) (https://wwwn.cdc.gov/Nchs/Nhanes/2015-

2016/DEMO_I.htm). All the laboratory measurements were performed by CDC's

Division of Laboratory Sciences at the National Center for Environmental Health. Survey

data were collected at the mobile examination center interview room on the day of the

health exam by CDC. The demographics data extracted for the study includes

information on the Survey participant's household interview and examination status,

gender, age, and pregnancy status. Examination data includes body measures and

blood pressure. Questioner data has information about drinking habit and known

diabetes status.

Laboratory Data

All the laboratory methods for blood and urine samples were performed by NHANES

that has been described thoroughly at:

(https://wwwn.cdc.gov/nchs/nhanes/continuousnhanes/labmethods.aspx?BeginYear=20

15). Laboratory measured data in the study are levels of PAHs, Glycohemoglobin,

Glucose, Cholesterol, Lipids, Triglyceride, complete blood count, Lymphocytes,

Monocytes, and hepatitis conditions.

Analytical method for urinary PAH measurement

The analytical procedure followed by NHANES involves enzymatic hydrolysis of

glucuronidated/ sulfated OH-PAH metabolites in urine, extraction by on-line solid phase

extraction, and separation and quantification using isotope dilution high performance

liquid chromatography-tandem mass spectrometry (on-line SPE-HPLC-MS/MS) [8].

Selection of eligible participants

After data collection from the NHANES website, samples where data for any of the

following were missing were excluded: missing examination status, age below 18 years,

pregnancy, missing BMI, missing laboratory data for Glycohemoglobin or Poly aromatic

hydrocarbons (PAHs). Individuals with missing alcohol drinking data for past 12 months

were also excluded.

The specific analytes measured in this method are OH-PAHs, namely URXP01, URXP02,

URXP03, URXP04, URXP06, URXP25, and URXP10 (In the NHANES database). PAH

metabolites data below lower limit of detection (LLOD) of the analytical method levels

were excluded (Fig. 1).

Statistics

Participants were categorized for this investigation based on their PAH

metabolite exposure range as per NHANES data (Supplement Table 1). Population

based on each metabolite levels was categorized into 3 groups: 1 below geometric

mean exposure range, 2 geometric mean exposure range and 3 higher than geometric

mean exposure of the reference range of URXP01 calculated by NHANEs (Fourth

National Report on Human Exposure to Environmental Chemicals Update (cdc.gov)). In

this study, we used the exposure levels of 1-HN (stated as URXP01 in the NHANES

database) for the analyses purpose.

Male with 14≥ drinks/ week, and females with 8≥ drinks/ week are heavy drinkers

as per NIAAA classification (Drinking Levels Defined | National Institute on Alcohol

Abuse and Alcoholism (NIAAA) (nih.gov). Low risk or moderate drinking are who drank

2≤ drinks/ day as males, and females as 1≤ drinks/ day (<u>Drinking too much alcohol can</u>

<u>harm your health. Learn the facts | CDC</u>).

Univariate factorial ANOVA was used to evaluate the group differences (both

between the groups; as well as across all the groups) in the demographics, PAH

exposure, drinking patterns, clinical data, and biomarker levels in context. Linear

regression model was used to analyze the association of biomarkers and clinical, PAH

exposure and drinking data. Demographic data were used as covariates, if found to be

significantly different in the between-group analyses. Multivariable regression model

was used for multi-independent model to assess comorbidity association and their effect

sizes on the clinical outcomes. IBM SPSS version 27 (IBM[®], Armonk, New York USA)

was used to compute all the statistical analysis and correlations in the study. Microsoft

365 version (Microsoft Corp., Redmond, Washington USA) was used for data

processing and Table/s development. Figures were processed using GraphPad Prism

(GraphPad, San Diego, CA USA). Data in table/s are presented as Mean and standard

deviation (M±SD). Statistical significance was set at $p < 0.05$.

RESULTS

Demographics, Drinking Profile and Diabetic Indexes by PAH Exposure Level

Age was significantly higher in Gr. 3 compared to Gr. 1 (Table 1). Notably, all the

individuals in each group showed overweight to borderline obese, especially Gr. 1 and

Gr. 2. Gr. 3 individuals drank on an average daily more than the rest of the two group

individuals, and significantly higher (p=0.015) compared to the Gr. 1 (Table 1).

Association of Diabetic Indexes and demographic and drinking Profile by PAH Exposure

HbA1C and age were very significantly associated (R=0.331, p≤0.001) (Fig. 2a)

in Gr. 3 (high exposure group of 1-Hydroxynapthelene). HbA1C and BMI were also very

significantly (p=0.002) associated (R=0.230) (Fig. 2b) in Gr.3. In this univariate

regression model, diabetic index (HbA1C) as a dependent variable showed unique

relationship with candidate demographic indices exclusively in the high exposure group

of 1-Hydroxynapthelene.

Association of Inflammatory Markers and PAH

Univariate analysis of regression showed significant prediction (R=0.141,

p=0.001) of C-Reactive protein by the absolute Monocyte level in Gr. 3. Multivariable

independent variable regression model was used to include the contributing role of

candidate PAHs that potentially are involved in the inflammation. CRP, marker of acute

pro-inflammation, as a dependent variable and absolute monocyte count (or numbers)

(AMC) as independent variable showed augmented effects of association with the

candidate PAHs; 1-HN (R=0.142, p=0.005) (Fig. 3a), 2-HN (R=0.149, p=0.001) (Fig.

3b); respectively.

To estimate the role of alcohol drinking, which is also a cause of inflammation,

we reviewed the levels of "Average Drinks Per Day" (ADPD) in all the groups (Fig. 4a).

We found that Gr. 3 individuals tend to drink more than the rest two groups. In Gr. 3,

males drank significantly higher than the females (Fig. 4b). Gr. 3 males showed

significant association of HbA1C and triglycerides (Fig. 4c). On the other hand, Gr. 3

females showed significant association of HbA1C and CRP (Fig. 4d).

With this finding on inflammation and immune activity, we evaluated the overall

impact on the Diabetic index, HbA1C in context of CRP, candidate lipid/s, drinking

measure/s and immune activity in the Gr. 3. HbA1C was significantly predictable with

CRP and AMC (Fig. 5), although CRP by itself showed only a close to trend level of

significance for direct relationship with HbA1C. This arrangement further augmented its

effects with simultaneously increasing order of statistical significance in a stepwise

manner with Triglycerides and Average Drinks Per Day were included respectively as

added independent variable/s in this multivariable prediction model.

DISCUSSION

Findings from this investigation support the pathological role of the high exposure

of PAH in the metabolic disorders; such as T2DM. Human studies have been performed

earlier [21], and this study assessed the proinflammatory status and factors, risk factors

and level of exposure to extend the present knowledge in a very large population. The

focus of this study was 1-HN (1-hydroxynaphthalene, stated as URXP01 in NHANES

database) and individuals exposed to this PAH yielded some noteworthy results.

Age and BMI stood out as the demographic measures that showed propensity

with T2DM diabetic index HbA1C in individuals with high PAH exposure of 1-HN, while

individuals with normal or low level of exposure did not. Similar study has been reported

for PAH exposure, obesity for the risk assessment of T2DM in patients with lung

disease [22].

Some studies report a close association of inflammation and PAH exposure [23]

as well in the 2003-2008 wave of NHANES database [24]. We found similar assessment

in the 2015-2015 database of NHANES database, especially in the context of the 1-HN

PAH, which was the scope of this study. CRP and inflammation associated blood cells

(absolute monocyte count) showed a significant relationship in individuals with high

exposure levels in this study. Interestingly, inflammation and role of PHAs along with the

oxidative stress has also been investigated in adolescents and pregnant women[24, 25].

One of the noteworthy findings in this study was that the individuals with high 1-

HN exposure drank the most compared to the other exposure groups (normal/low).

Males with high exposure of 1-HN had higher levels of average drinking than their

female counterparts suggesting trends of sexual dimorphism. Almost a 2-fold higher

average drinking in high PAH exposed males might be uniquely worrisome for a

possibly unidentified role of PAH. Males having more average drinks is aligned with the

findings of the NAS [26].

Earlier studies on mice have suggested that PAHs exposure induces lipid

associated metabolic disorders in a time-dependent manner [27]. Similarly, CRP level

observed in new-onset DM people diagnosed with HbA1c criterion has been recently

reported [28]. However, there has always been a gap in how to address such findings in

humans in context of PAH exposure and other risk factors. In this study, males with high

1-HN exposure showed close association of HbA1C and triglycerides, suggesting sex-

specific exposure associated response to metabolic conditions [29]. Females with high

1-HN exposure on the other hand showed that HbA1C was closely associated with pro-

inflammatory activity (CRP), suggesting propensity to the risks of chronic conditions

[30]. Such a sexual dimorphism in high PAH exposure is a healthcare concern for

metabolic disorder that needs more extensive evaluation; both mechanistically focus on

biological pathways as well clinically with well-developed research questions.

In this study, several important clinical and biological measures involved in pro-

inflammatory, pathological activity were observed to be higher as well as connected to

each other. Thus, it was important to know their overall effect on the HbA1C levels in

individuals with high PAH exposure [21, 31]. Indeed, the effect size on the HbA1C levels

grew in a stepwise manner in context of candidate markers with potential role in

inflammation and injury. As described in the multivariable model, role of several markers

(representing several pathways that can be affected) on HbA1C, support evaluating

PAH exposed individuals with comprehensive clinical assessment and varied tests to

understand the complexity involved [32]. A study on a larger subject size such as this

one, and its results may advance the understanding in the scientific literature on the

adverse and harmful effects of PAH exposure and its interaction with the risk factors

that are attributed to the metabolic disorders.

None-the-less, this study has several limitations that also need to be discussed.

This is a single timepoint observational study, and assessment of longitudinal data is

not in the scope of this investigation. No biological mechanism was tested in an

experimental model in this evaluation to validate the corresponding clinical

observations. Thus, no causations/ etiologies were identified or were the aims in this

study. Risk factors, like BMI and age were identified in high exposure group of

individuals. Aging and obesity associated with PAH exposure could be the important

newer directions of research that can offshoot from the findings of this study. The focus

of this study was limited to one PAH, 1-HN. However other PAHs are equally important

to investigate and can also be evaluated as future directions as independent projects.

This study also did not evaluate any other metabolic conditions.

In conclusion, this study identifies that the BMI and age likely are candidate

demographic risk factors for Diabetes Mellitus Type 2 (T2DM) in individuals with high

PAH exposure. Acute proinflammatory activity characterized by C-Reactive Protein, an

acute pro-inflammatory marker, is augmented when elevated monocyte levels are

observed in relevance of and 1-HN; and 2-HN independently. Prevalence of increased

average drinks over time is also observed with high PAH exposure, especially in males.

With high PAH exposure, T2DM shows sexual dimorphism. Males exposed with higher

levels of PAH show association with triglycerides, whereas females with CRP

independently predicted HbA1C. Importantly, the levels of CRP, absolute monocyte

levels (suggesting its activation), serum triglyceride levels, and average alcoholic drinks

predict the HbA1C levels robustly in individuals with high PAH exposure.

REPRINT REQUEST: Shweta Srivastava Ph.D., M.Sc.; Delia Baxter II, Rm. 428,
Environment Health Institute; 580 S. Preston Street, University of Louisville; Louisville
KY 40202. Email: shweta.srivastava.1@louisville.edu.

TABLES

TABLE 1. Demographic, drinking, PAH, Lipids, blood cell values, C-reactive protein and diabetic
markers are tabulated. Statistical significance was set at p≤0.05. Data presented as Mean±SD.

Measures	Group 1 (low) 1-Hydroxynaphthalene			Group 2 (Normal) 1-Hydroxynaphthalene			Group 3 (High) 1-Hydroxynaphthalene			all groups p-value
	Males (n= 287; %)	Females (n= 218; %)	Total (n = 505)	Males (n= 6; %)	Females (n= 7; %)	Total (n= 13)	Males (n= 186; %)	Females (n= 131; %)	Total (n= 317)	
Age (years) [c]	45.57 ± 17.76	43.46 ± 17.58	44.66 ± 17.69	43.17 ± 22.41	44.71 ± 13.22	44.00 ± 17.24	49.94 ± 16.52	48.98 ± 15.88	49.54 ± 16.24	NS
BMI (kg/m^2) [b#, c]	29.88 ± 6.33	31.507 ± 8.69	30.584 ± 7.48	29.23 ± 8.05	34.00 ± 7.37	31.80 ± 7.76	27.55 ± 5.66	29.83 ± 7.31	28.50 ± 6.48	≤0.001
Average Drink per Day in past 1 year	3.21±2.71	2.09±1.42	2.74±2.32	3±1.67	1.86±.69	2.38±1.32	3.87± 2.80	2.21±1.68	3.22±2.65	≤0.037
PAH metabolite level (ng/gm of creatinine)										
1-hydroxynaphthalene (ng/g of creatinine) **	1164.49 ± 788.29	1098.06 ±750.47	1135.81±772.14	3726.81±147.24	3760.83±180.34	3745.13±160.02	52465.91±329582.65	165450.57±845589.68	99156.73±600710.65	NA
2-hydroxynaphthalene (ng/g of creatinine) [c]	5893.71±7543.00	7866.85±6303.15	6745.48±7095.74	5561.87±2065.07	8140.95±2963.36	6950.61±2821.07	13022.58±8308.54	18245.61±13760.66	15180.99±11178.08	≤0.001
2-hydroxyfluorene (ng/g of creatinine) [a, b#, c]	234.74±349.28	225.86±279.47	230.91±320.74	589.49±226.21	336.84±329.75	453.45±304.75	1169.67±1609.27	1184.38±1138.77	1175.75±1431.68	≤0.001
3-hydroxyfluorene (ng/g of creatinine) [a, b, c]	105.74±116.42	95.19±145.33	101.18±129.66	279.80±89.09	212.44±260.72	243.53±196.25	730.24±761.15	786.40±830.78	753.45±789.87	≤0.001
1-hydroxyphenanthrene (ng/g of creatinine) [a#, c]	129.40±123.75	135.78±117.36	132.15±120.96	286.67±277.73	117.16±58.92	195.40±203.99	278.06±528.98	278.04±294.85	278.05±446.74	≤0.001
2- & 3-hydroxyphenanthrene (ng/g of creatinine) [c]	168.42±273.48	144.62±93.45	158.15±215.27	370.29±401.85	153.63±69.61	253.63±286.96	473.01±880.40	422.63±614.16	452.19±780.75	≤0.001
1-hydroxypyrene (ng/g of creatinine) [a, c]	137.14±148.64	153.18±100.39	144.07±130.16	537.42±693.53	131.80±77.49	319.01±497.71	419.58±602.28	415.68±461.92	417.97±547.85	≤0.001
Cholesterol and Triglycerides										
Total Cholesterol (mg/dL)	188.11±43.81	183.94±38.79	186.33±41.75	171.83±54.52	193.86±48.22	183.69±50.31	187.70±40.62	192.67±39.51	189.76±40.18	NS
HDL (mg/dL) [c]	47.79±14.64	55.23±16.92	50.96±16.06	45.17±11.50	61.71±14.03	54.08±15.07	51.45±19.34	57.33±16.62	53.88±18.47	0.061 [#]
LDL (mg/dL) [c]	112.30±35.54	105.86±30.42	109.59±33.56	88.50±72.83	132.67±27.57	115.00±47.87	115.67±39.88	112.16±32.13	114.36±37.08	NS
Triglycerides (mg/dL) [a#]	106.16 ±63.16	99.82 ±67.97	103.48 ±65.15	201.50 ±118.09	132.33 ±63.71	160.00 ±83.37	129.01 ±103.99	108.10 ±54.03	121.25 ±89.12	0.045
Candidate/ Target Blood Cell Measures										
Lymphocyte Absolute number (1000 cells/uL) [c]	2.07±0.66	2.40±0.85	2.21±0.76	2.02±0.47	2.09±0.39	2.06±0.41	2.23±0.77	2.53±0.75	2.35±0.77	0.015

Monocyte Absolute number (1000 cells/uL) ^{c}	0.60±0.20	0.57±0.18	0.59±0.19	0.66±0.38	0.54±0.10	0.59±0.25	0.63±0.19	0.61±0.20	0.62±0.19	*0.071*
Blood Glucose and diabetes measures										
Glycohemoglobin (%) $^{c\#}$	5.74±1.07	5.71±1.12	5.73±1.09	5.45±0.73	6.23±1.75	5.87±1.38	5.92±1.16	5.78±0.97	5.86±1.09	NS
Glucose, refrigerated serum (mg/dL)	104.96±43.22	99.57±32.45	102.66±39.06	108.17±31.19	94.33±12.08	101.25±23.68	107.67±47.28	101.98±38.97	105.32±44.06	NS
Inflammation markers										
HS C-Reactive Protein (mg/L) $^{c\#}$	2.93±5.17	5.35±7.08	3.96±6.17	5.58±5.21	3.31±5.41	4.45±5.20	4.82±10.72	5.30±7.54	5.02±9.53	NS

344   Footnote: BMI: Body Mass Index. HDL: High density Lipoprotein. LDL: Low Density Lipoprotein. [a] Significant difference between
345   group 1 and 2. [b] Significant difference between group 2 and 3. [c] Significant difference between group 1 and 3. # Trend level of
346   significance. [a]~, [b]~, [c]~ #: Trend level of significant difference between two groups. NS: Not Significant. NA: Not Applicable. **
347   comparisons were not shown since the groups were distributed by the levels of 1-hydroxynapthalene, thus the differences were all
348   significant.

Figure Legends

Figure 1. Flow Chart illustrating the selection of eligible participants. Eligible participants and those included in the analysis for the associations between urinary PAH levels and other clinical parameters.

Figure 2. Association of HbA1C and demographic parameters in high exposure group 3 of the study. Fig. 2a: Association of HbA1C and age. Fig. 2b: Association of HbA1C and Body Mass Index (BMI). Raw data plotted on x- and y-axes. Effect sizes/correction index is noted along with the p-value. Statistical significance was set at p≤0.05.

Figure 3. Association of C-Reactive Protein with absolute monocyte numbers in Group 3 of highly exposed subjects in multivariable regression model with PAH. 3a. 1-Hydroxynaphthalene. 3b. 2-Hydroxynaphthalene. Effect sizes/correction index is noted along with the p-value. Statistical significance was set at p≤0.05.

Figure 4. Alcohol drinking and PAH exposure; and interaction of diabetes and pro-inflammatory markers as sexual dimorphism in Gr. 3. Fig. 4a: Gr. 3 individuals drank more than the other groups. Fig. 4b: Males drank more than the females in Gr. 3. Gr. 3: Association of triglycerides and HbA1C in Gr. 3 males. Fig. 4d: Association of CRP and HbA1C in Gr. 3 females. Group 1: below geometric mean exposure, group 2: geometric mean exposure and group 3: higher than geometric mean exposure of the reference range. Statistical significance was set at p<0/05. Data presented as Mean ± Standard Error.

Figure 5. Stepwise multivariable regression analysis model illustrates the increasing effects on HbA1C as diabetic index by the pro-inflammatory activity presented as CRP,

371 absolute monocyte levels, serum triglyceride along with average drinking per day in the

372 Gr. 3. Statistical significance was set at $p < 0/05$. R^2 is noted along with the p-values.

References:

1. *Toxicological profile for Polycyclic Aromatic Hydrocarbons (PAHs).* , P.H.S. Department
of Health and Human Services, Editor. 1995: Atlanta, GA: U.S.
2. *Polycyclic aromatic hydrocarbons—occurrence in foods, die-tary exposure, and health*
effects. , S.C.o.F. uropean Commission, Editor. 2002.
3. Maliszewska-Kordybach, B., *Sources, concentrations, fate and effects of polycyclic*
aromatic hydrocarbons (PAHs) in the environment. Part A: PAHs in air. Polish journal of
environmental studies, 1999. **8**: p. 131-136.
4. Li, Z., et al., *PM2. 5 associated PAHs and inorganic elements from combustion of*
biomass, cable wrapping, domestic waste, and garbage for power generation. Aerosol
and air quality research, 2019. **19**(11): p. 2502-2517.
5. Farhadian, A., et al., *Effects of meat preheating and wrapping on the levels of polycyclic*
aromatic hydrocarbons in charcoal-grilled meat. Food chemistry, 2011. **124**(1): p. 141-
146.
6. Singh, L., J.G. Varshney, and T. Agarwal, *Polycyclic aromatic hydrocarbons formation*
and occurrence in processed food. Food Chemistry, 2016. **199**: p. 768.
7. Health, N.I.f.O.S.a. *What are the Standards and Regulations for PAHs Exposure?* 2013;
Available from: https://www.atsdr.cdc.gov/csem/polycyclic-aromatic-
hydrocarbons/standards_and_regulations_for_exposure.html.
8. Wang, Y., et al., *Quantification of urinary mono-hydroxylated metabolites of polycyclic*
aromatic hydrocarbons by on-line solid phase extraction-high performance liquid
chromatography-tandem mass spectrometry. Analytical and bioanalytical chemistry,
2017. **409**(4): p. 931-937.
9. Jongeneelen, F.J., *Benchmark guideline for urinary 1-hydroxypyrene as biomarker of*
occupational exposure to polycyclic aromatic hydrocarbons. Ann Occup Hyg, 2001.
45(1): p. 3-13.
10. Nemmar, A., *Possible mechanisms of the cardiovascular effects of inhaled particles:*
systemic translocation and prothrombotic effects. Toxicology Letters, 2004. **149**(1-3): p.
243-253.
11. Moorthy, B., C. Chu, and D.J. Carlin, *Polycyclic Aromatic Hydrocarbons: From*
Metabolism to Lung Cancer. Toxicological Sciences, 2015. **145**(1): p. 5-15.
12. Marris, C.R., et al., *Polyaromatic hydrocarbons in pollution: a heart-breaking matter.* The
Journal of Physiology, 2020. **598**(2): p. 227-247.
13. Ranjbar, M., et al., *Urinary Biomarkers of Polycyclic Aromatic Hydrocarbons Are*
Associated with Cardiometabolic Health Risk. PLOS ONE, 2015. **10**(9): p. e0137536.
14. Sarma, S.N., et al., *A metabolomics study on effects of polyaromatic compounds in oil*
sand extracts on the respiratory, hepatic and nervous systems using three human cell
lines. Environ Res, 2019. **178**: p. 108680.
15. Curtis, L.R., et al., *Reduced cytochrome P4501A activity and recovery from oxidative*
stress during subchronic benzo[a]pyrene and benzo[e]pyrene treatment of rainbow trout.
Toxicology and Applied Pharmacology, 2011. **254**(1): p. 1-7.

16.   Ataria, J.M., K. O'Halloran, and R. Gooneratne, *Hepatic and immune biological effect*
assays in C57BL/6 mice to measure polycyclic aromatic hydrocarbon bioavailability
under laboratory exposures with increasing environmental relevance. Environmental
Science and Pollution Research - International, 2007. **14**(4): p. 256-265.
17.   Ramesh, A., et al., *Comparative Metabolism, Bioavailability, and Toxicokinetics of*
Benzo[a]pyrene in Rats After Acute Oral, Inhalation, and Intravenous Administration.
Polycyclic Aromatic Compounds, 2002. **22**(3-4): p. 969-980.
18.   Alshaarawy, O., et al., *Urinary polycyclic aromatic hydrocarbon biomarkers and diabetes*
mellitus. Occupational and Environmental Medicine, 2014. **71**(6): p. 437-441.
19.   Yang, L., et al., *Dose-response relationship between polycyclic aromatic hydrocarbon*
metabolites and risk of diabetes in the general Chinese population. Environmental
Pollution, 2014. **195**: p. 24-30.
20.   Yang, L., et al., *Association of polycyclic aromatic hydrocarbons metabolites and risk of*
diabetes in coke oven workers. Environmental Pollution, 2017. **223**: p. 305-310.
21.   Zhang, H., et al., *Association of internal exposure to polycyclic aromatic hydrocarbons*
with inflammation and oxidative stress in prediabetic and healthy individuals.
Chemosphere, 2020. **253**: p. 126748.
22.   Hou, J., et al., *Exposure to polycyclic aromatic hydrocarbons and central obesity*
enhanced risk for diabetes among individuals with poor lung function. Chemosphere,
2017. **185**: p. 1136-1143.
23.   Alshaarawy, O., et al., *Polycyclic aromatic hydrocarbon biomarkers and serum markers*
of inflammation. A positive association that is more evident in men. Environmental
research, 2013. **126**: p. 98-104.
24.   Farzan, S.F., et al., *Urinary polycyclic aromatic hydrocarbons and measures of oxidative*
stress, inflammation and renal function in adolescents: NHANES 2003–2008.
Environmental research, 2016. **144**: p. 149-157.
25.   Ferguson, K.K., et al., *Urinary polycyclic aromatic hydrocarbon metabolite associations*
with biomarkers of inflammation, angiogenesis, and oxidative stress in pregnant women.
Environmental science & technology, 2017. **51**(8): p. 4652-4660.
26.   Rehm, J., T.K. Greenfield, and J.D. Rogers, *Average volume of alcohol consumption,*
patterns of drinking, and all-cause mortality: results from the US National Alcohol
Survey. American Journal of Epidemiology, 2001. **153**(1): p. 64-71.
27.   Li, F., et al., *Hepatotoxic effects of inhalation exposure to polycyclic aromatic*
hydrocarbons on lipid metabolism of C57BL/6 mice. Environment International, 2020.
134: p. 105000.
28.   Tutuncu, Y., et al., *A comparison of hs-CRP levels in new diabetes groups diagnosed*
based on FPG, 2-hPG, or HbA1c criteria. Journal of diabetes research, 2016. **2016**.
29.   Zaidi, S.M.H., A. Ghafoor, and F.A. Randhawa, *HbA1c as an indirect marker of*
hypertriglyceridemia in type 2 diabetes mellitus. Journal of Ayub Medical College
Abbottabad, 2015. **27**(3): p. 601-603.
30.   Shanahan, L., J. Freeman, and S. Bauldry, *Is very high C-reactive protein in young*
adults associated with indicators of chronic disease risk? Psychoneuroendocrinology,
2014. **40**: p. 76-85.
31.   Khosravipour, M. and H. Khosravipour, *The association between urinary metabolites of*
polycyclic aromatic hydrocarbons and diabetes: A systematic review and meta-analysis
study. Chemosphere, 2020. **247**: p. 125680.
32.   Mumtaz, M., et al., *ATSDR evaluation of health effects of chemicals. IV. Polycyclic*
aromatic hydrocarbons (PAHs): understanding a complex problem. Toxicology and
industrial health, 1996. **12**(6): p. 742-971.

Figure 1

Figure 2

Fig. 2a.

Fig. 2b.

Figure 3

Fig. 3a.

Fig. 3b.

Figure 4

Fig. 4a.

Fig. 4b.

Fig. 4c.

Gr. 3 Males

R=0.422; p≤0.001

Fig. 4d.

Gr. 3 Females

R=0.213; p=0.015

Figure 5