Effects of Glucagon-like Peptide-1 Receptor Agonists on Cardiovascular and Renal Outcomes: A Systematic Review, Meta-analysis, and Meta-regression Analysis

Short title: Cardiovascular and renal efficacy of GLP-1RAs

Satoshi Yoshiji, MD[1,2,3,4], Hiroto Minamino, MD[1,4], Daisuke Tanaka, MD, PhD[1], Shunsuke Yamane MD, PhD[1], Norio Harada, MD, PhD[1], and Nobuya Inagaki, MD, PhD[1]

[1]Department of Diabetes, Endocrinology and Nutrition, Graduate School of Medicine, Kyoto University, Kyoto, Japan

[2]Department of Human Genetics, McGill University, Montréal, Québec, Canada

[3]Kyoto-McGill International Collaborative Program in Genomic Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

[4]Japan Society for the Promotion of Science, Tokyo, Japan

Correspondence

Nobuya Inagaki MD, PhD,

Department of Diabetes, Endocrinology and Nutrition, Graduate School of Medicine,

Kyoto University, 54 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto, 606-8507, Japan

E-mail: inagaki@kuhp.kyoto-u.ac.jp

Tel: +81-075-751-3560

Fax: +81-075-751-4244

Abstract: 347 words.

Main text: 2981 words.

Number of Table: 1

Number of Figures: 4.

Supplementary material: 4 Tables and 5 Figures.

Abstract

Background: Cardiovascular and renal effects of glucagon-like peptide-1 receptor agonists (GLP-1RAs) have been inconsistent in cardiovascular outcome trials, and factors associated with the efficacy of GLP-1RAs remain to be clarified.

Objective: To evaluate cardiovascular and renal outcomes with GLP-1RAs and associations between these outcomes and the magnitude of HbA1c or weight reduction.

Data Sources and Study selection: We searched PubMed/MEDLINE for randomized, placebo-controlled trials of GLP-1RAs published until 11 August 2021 and selected trials reporting major adverse cardiovascular events (MACE) as the primary outcome.

Data Extraction and Synthesis: We extracted data of 60,080 individuals from eight studies (ELIXA, LEADER, SUSTAIN-6, EXSCEL, HARMONY, PIONEER 6, REWIND, and AMPLITUDE-O) and conducted a meta-analysis with a random-effects model to calculate pooled hazard ratios (HRs) of primary and secondary outcomes. Furthermore, we performed a univariable meta-regression analysis of these outcomes with HbA1c or weight reduction.

Main outcomes and Measures: The primary outcome was MACE (a composite of cardiovascular mortality, nonfatal stroke, and nonfatal myocardial infarction). Secondary outcomes included components of MACE, all-cause mortality,

hospitalization due to heart failure, and renal outcomes. Safety outcomes were severe

hypoglycemia, pancreatitis, pancreatic cancer, and retinopathy.

Results: GLP-1RAs reduced MACE (HR 0.86; 95% CI 0.80, 0.93; $P < 0.001$) and

secondary outcomes including cardiovascular mortality (0.87; 0.80 0.94; $P = 0.001$),

nonfatal myocardial infarction (0.90; 0.83, 0.98; $P = 0.020$) nonfatal stroke (0.83; 0.76,

0.92; $P < 0.001$), and the composite renal outcome (0.80; 0.73, 0.87; $P < 0.001$). No

increase in the incidence of safety outcomes was observed with GLP-1RAs.

Meta-regression analysis showed that HbA1c reduction was associated with logarithm

of HR (log-HR) of MACE ($P = 0.045$; $R^2 = 0.64$) and the composite renal outcome ($P =$

0.045; $R^2 = 0.80$), whereas weight reduction was not associated with any outcome.

Every 1.0% HbA1c reduction was associated with a decrease in log-HR of MACE and

the composite renal outcome by 0.27 and 0.35, respectively.

Conclusions: GLP-1RAs showed favorable effects on cardiovascular and renal

outcomes. Furthermore, reduction in HbA1c, but not body weight, was associated with

cardiovascular and renal risk reduction during the treatment with GLP-1RAs.

Introduction

Major adverse cardiovascular events (MACE) and chronic kidney disease are leading causes of morbidity and mortality in individuals with type 2 diabetes. Prevention and management of these disorders play pivotal roles in the treatment of type 2 diabetes[1, 2]. To date, a series of randomized, placebo-controlled clinical trials on cardiovascular and renal outcomes with glucagon-like peptide-1 receptor agonists (GLP-1RAs) have been reported, including ELIXA, LEADER, SUSTAIN-6, HARMONY, PIONEER 6, REWIND, and AMPLITUDE-O[3-10]. These trials evaluated the safety and efficacy of GLP-1RAs on the primary composite MACE outcome (cardiovascular mortality, nonfatal stroke, or nonfatal myocardial infarction) as well as secondary outcomes, including individual components of MACE, hospitalization due to heart failure, all-cause mortality, and renal outcomes. In these trials, GLP-1RAs consistently showed cardiovascular and renal safety, but cardiovascular and renal efficacy were inconsistent across trials: LEADER, SUSTAIN-6, and REWIND reduced MACE compared to placebo, whereas ELIXA and EXSCEL did not reduce MACE. Notably, AMPLITUDE-O reduced MACE for the first time as a trial with an exendin-based GLP-1RA[10].

Regarding the mechanism of action for cardiovascular and renal efficacy, it has

been reported that reducing HbA1c or body weight contributes to cardiovascular, renal,

and mortality risk reduction[11-16]. However, although cardiovascular outcome trials of

GLP-1RAs reported effective reductions in HbA1c and body weight, it is still unclear

whether HbA1c or body weight reduction is associated with the cardioprotective and

renoprotective profile.

To examine overall cardiovascular and renal effects as well as safety profiles of

GLP-1RAs, we performed a systematic review and meta-analysis of cardiovascular

outcome trials of GLP-1RAs, including AMPLITUDE-O. Furthermore, to evaluate

whether the magnitude of HbA1c or body weight reduction is associated with

cardiovascular and renal outcomes, we conducted a univariable meta-regression

analysis of cardiovascular and renal outcomes with HbA1c or weight reduction.

Methods

Data Sources and Searches

We performed a data search in accordance with the Preferred Reporting Items for

Systematic Reviews and Meta-Analysis (PRISMA) statement[17]. We searched the

PubMed/MEDLINE database for eligible randomized, placebo-controlled trials of

GLP-1RAs on MACE in individuals with type 2 diabetes, published until 11 August

2021. We placed no restrictions on the study population or language. The search terms

were described in Supplementary Figure 1. The protocol for this study is registered in

PROSPERO (CRD42021273058).

Study Selection

We selected studies reporting MACE as the primary outcome. For selected trials, we

also evaluated secondary outcomes and safety outcomes. SY and HM independently

conducted the search and solved any discordance through discussion.

We selected randomized, placebo-controlled trials of GLP-1RAs reporting major

adverse cardiovascular events (MACE) as the primary outcome. We excluded studies

not reporting MACE or a design of interest.

Data Extraction and Quality Assessment

We extracted summary statistics of the eligible studies from published articles of the

trials and their supplementary materials. We did not seek individual-level data. For

primary and secondary outcomes, we obtained relevant hazard ratios (HRs) and 95%

confidence intervals (95% CIs). For safety outcomes, odds ratios (ORs) and 95% CIs

were obtained. We evaluated the risk of bias for eligible studies with the Cochrane Risk

of Bias Tool[18]. SY and HM conducted the data extraction and quality assessment.

Data Synthesis and Analysis

For the present meta-analysis, the primary outcome was the HR of MACE. The

secondary outcomes were the HRs of individual components of MACE, all-cause

mortality, hospitalization due to heart failure, composite renal outcome, and renal

function outcome. Definitions of the composite renal outcome and the renal function

outcome were not standardized across the trials; detailed definitions of these renal

outcomes for each trial are provided in Supplementary Table 1. Definitions of the safety

outcomes, including severe hypoglycemia, pancreatitis, pancreatic cancer, and

retinopathy, are also described in Supplementary Table 2.

For the meta-analysis, we used the inverse variance method with a

random-effects model to obtain a weighted average of HRs or ORs with 95% CIs. We

used the DerSimonian-Laird estimator for the assessment of between-study variance[19].

We used the I^2 index and Cochran's Q test to measure the degree of heterogeneity. The

degree of heterogeneity based on the I^2 index was defined as low (I^2 index $\leq$ 25%),

moderate (I^2 index 26–50%), and high (I^2 index $>$ 50%)[20]. P values of $<$ 0.05 for

Cochran's Q test represent significant heterogeneity between the studies.

Regarding meta-regression analysis, normalized HbA1c and body weight

reduction of each GLP-1RA were calculated from head-to-head clinical as described

previously[21]. We defined HbA1c reduction and body weight reduction of liraglutide as

1.0% and 1.0 kg, respectively, and normalized those of other GLP-1RAs against

liraglutide. We delineated liraglutide as a normalizer because liraglutide was compared

head-to-head against all the other GLP-1RAs[22-28]. Using either normalized HbA1c

change or body weight change as an independent variable, we conducted univariable

meta-regression analyses for cardiovascular and renal outcomes with the log-HR as a

dependent variable. P values of < 0.05 represented significance.

We performed all statistical analysis with R (version 4.0.2, R Foundation for

Statistical Computing, Vienna, Austria) and the R package "meta" version 4.18.2.

Results

Study identification

Among 212 articles screened with the prespecified screening strategy, eight randomized,

placebo-controlled cardiovascular outcome trials which defined MACE as the primary

outcome were identified. Data from the following eight trials with a total of 60,080

individuals were analyzed in the present meta-analysis: ELIXA, LEADER, SUSTAIN-6,

EXSCEL, HARMONY, PIONEER 6, REWIND, and AMPLITUDE-O (Table 1). The

study selection process based on PRISMA is presented in Supplementary Figure 1.

According to the Cochrane risk of bias tool[18], the risk of bias for all included trials was

evaluated as low (Supplementary Table 3).

ELIXA, EXSCEL, and AMPLITUDE-O evaluated GLP-1RAs with an

exendin-based structural backbone, whereas LEADER, SUSTAIN-6, HARMONY, and

PIONEER 6 examined those with a human GLP-1 backbone. For all trials other than

ELIXA, the primary outcome was MACE, which was the time to the first occurrence of

cardiovascular mortality, nonfatal stroke, or nonfatal myocardial infarction. For ELIXA,

the primary outcome was extended MACE, which was MACE with an additional

component, i.e., hospitalization due to unstable angina. However, hospitalization due to

unstable angina accounted for less than 2.5% of the primary outcome in ELIXA. Given

that hospitalization for unstable angina had minimal influence on the primary outcome,

we included ELIXA in the meta-analysis. HARMONY and PIONEER 6 did not report

renal outcomes and were excluded from the analysis for renal outcomes.

Baseline characteristics

The baseline characteristics of the eight trials are summarized in Table 1. The median

duration of follow-up was the shortest in PIONEER 6 (1.3 years) and the longest in

REWIND (5.4 years). The average age ranged from 59.9 years in ELIXA to 66.2 years

in REWIND. Of the 60,080 individuals, 21930 (37%) were women, and 45640 (76%)

were of white ethnicity. Average body mass index varied from 30.1 kg/m^2 (ELIXA) to

32.8 kg/m^2 (SUSTAIN-6), and average baseline HbA1c ranged from 7.3% (REWIND)

to 8.9% (AMPLITUDE-O), and average estimated glomerular filtration rate (eGFR)

from 74 mL/min per 1.73 m^2 (PIONEER 6) to 80 mL/min per 1.73 m^2 (LEADER and

SUSTAIN-6). The proportion of individuals with previously established cardiovascular

disease was 77% (46,105 out of 60,080 individuals) across the eight trials and ranged

from 31% (REWIND) to 100% (ELIXA and HARMONY).

Meta-analysis

Overall, GLP-1RAs reduced MACE (HR 0.86; 95% CI 0.80, 0.93; $P < 0.001$) with

moderate heterogeneity ($I^2 = 44.5\%$, $P = 0.082$) (Figure 1). As for the individual

components of MACE, GLP-1RAs reduced cardiovascular mortality (HR 0.87; 95% CI

0.80, 0.94; $P = 0.001$) with low heterogeneity ($I^2 = 12.8\%$, $P = 0.330$), nonfatal

myocardial infarction (HR 0.90; 95% CI 0.83, 0.98; $P = 0.020$) with moderate

heterogeneity ($I^2 = 27.4\%$, $P = 0.210$), and nonfatal stroke (HR 0.83; 95% CI 0.76, 0.92;

$P < 0.001$) with low heterogeneity ($I^2 = 0\%$, $P = 0.639$). Normalized HbA1c and body

weight reduction used in the meta-regression analysis were presented in Supplementary

Table 4.

As for other secondary outcomes (Figure 2), GLP-1RAs reduced all-cause

mortality (HR 0.88; 95% CI 0.82, 0.94; $P < 0.001$) with low heterogeneity ($I^2 = 10.5\%$,

$P = 0.349$), hospitalization due to heart failure (HR 0.89; 95% CI 0.82, 0.98; $P = 0.013$)

with low heterogeneity ($I^2 = 2.5\%$, $P = 0.410$), composite renal outcome (HR 0.80; 95%

CI 0.73, 0.87; $P < 0.001$) with moderate heterogeneity ($I^2 = 45.0\%$, $P = 0.106$), and

renal function outcome (HR 0.84; 95% CI 0.73, 0.97; $P = 0.016$) with moderate

heterogeneity ($I^2 = 31.4\%$, $P = 0.200$).

Subgroup analysis

In a subgroup analysis of MACE stratified by the prior history of established

cardiovascular disease (Supplementary Figure 2), GLP-1RAs reduced MACE in

individuals with established cardiovascular disease (HR 0.85; 95% CI 0.79, 0.92; $P <$

0.001) with moderate heterogeneity ($I^2 = 48.0\%$, $P = 0.060$), but not in those without

established cardiovascular disease (HR 0.94; 95% CI 0.83, 1.06; $P = 0.303$) with

moderate heterogeneity ($I^2 = 0.0\%$, $P = 0.514$).

In another subgroup analysis of MACE stratified by the structural backbones of

GLP-1RAs (Supplementary Figure 3), exendin-based GLP-1RAs did not reduce MACE

(HR 0.90; 95% CI 0.78, 1.04; P = 0.163) and had a high and significant heterogeneity

among studies (I^2 = 67.2%, P = 0.047). On the other hand, human GLP-1-based

GLP-1RAs reduced MACE (HR 0.84; 95% CI 0.79, 0.90; P < 0.001) with low

heterogeneity (I^2 = 0.0%, P = 0.513).

Safety outcomes

Regarding safety outcomes (Supplementary Figure 4), GLP-1RAs did not increase ORs

for severe hypoglycemia (OR 0.87; 95% CI 0.71, 1.08; P = 0.208), pancreatitis (OR

1.16; 95% CI 0.90, 1.60; P = 0.384), pancreatic cancer (OR 0.97; 95% CI 0.64, 1.48; P

= 0.892), or retinopathy (OR 1.07; 95% CI 0.92, 1.25; P = 0.400).

Meta-regression analysis

The meta-regression analysis of the eight pooled trials showed that normalized HbA1c

reduction was associated with the log-HR of MACE (slope -0.27; 95% CI -0.53, -0.01;

P = 0.045; R^2 = 0.64), whereas normalized body weight reduction had no association

with the log-HR of MACE (slope = -0.05; 95% CI -0.17, 0.07; P = 0.390; R^2 < 0.01)

(Figure 3). Among secondary outcomes, normalized HbA1c reduction was associated

with the log-HR of the composite renal outcome (slope = -0.35; 95% CI -0.69, -0.01; P

= 0.045; R^2 = 0.80) (Figure 4). Although the associations between normalized HbA1c

reduction and other secondary outcomes were not significant, all outcomes except

hospitalization due to heart failure showed negative slopes for the regression line, which

was directionally concordant with MACE. As for body weight, normalized body weight

reduction was not associated with any secondary outcome; the 95% CIs were wide and

ranged from negative to positive values (Figure 4 and Supplementary Figure 5).

Discussion

Systematic review and meta-analysis

The present meta-analysis and meta-regression analysis included a total of 60,080

individuals from eight trials of GLP-1RAs, including AMPLITUDE-O. Overall,

GLP-1RAs reduced MACE, its components, cardiovascular mortality, nonfatal

myocardial infarction, nonfatal stroke, all-cause mortality, hospitalization due to heart

failure, composite renal outcome, renal function outcome. Regarding safety outcomes,

the occurrence of severe hypoglycemia, pancreatitis, pancreatic cancer, or retinopathy

was not increased with GLP-1RAs.

In a subgroup analysis stratified by a prior history of established cardiovascular

disease (Supplementary Figure 2), GLP-1RAs reduced MACE in individuals with

established cardiovascular disease but not in those without established cardiovascular

disease. Taking into account the anti-atherosclerotic properties of GLP-1RAs[29], it is

conceivable that cardiovascular benefits of GLP-1RAs were more prominent in

individuals with established cardiovascular disease. Our findings support current

guidelines which recommend GLP-1RAs for the treatment of diabetes in individuals

with established cardiovascular diseases or in those who are at high risk of

cardiovascular diseases[2, 30, 31].

In terms of individual study details, all the studies, except for ELIXA and

EXSCEL, reported a reduction in MACE. Notably, in AMPLITUDE-O, efpeglenatide

reduced MACE for the first time as an exendin-based GLP-1RA. Given the findings

from AMPLITUDE-O, we conducted the subgroup analysis of MACE according to the

structural backbone of GLP-1RAs (Supplementary Figure 3). Whereas human

GLP-1-based GLP-1RAs reduced MACE (HR 0.84; 95% CI 0.79, 0.90; $P < 0.001$) with

low heterogeneity ($I^2 = 0.0\%$, $P = 0.513$), exendin-based GLP-1RAs did not reduce

MACE (HR 0.90; 95% CI 0.78, 1.04; $P = 0.163$). The difference could be due to the

large heterogeneity of exendin-based GLP-1 trials ($I^2 = 67.2\%$, $P = 0.047$). Individually,

ELIXA was different from the other two trials (EXSCEL and AMPMLITUDE-O) in the

sense that it recruited individuals with a recent acute coronary syndrome[3]. It should also

be noted that EXSCEL reported poor adherence to the drug (around 40% of premature

discontinuation of the trial) and that lixisenatide, used in ELIXA, has the shortest

half-life of 2–4 hours, which is markedly shorter than that of other GLP-1RAs[32].

Therefore, in ELIXA and EXSCEL, multiple factors such as study design, adherence to

the drug, and half-life could have influenced the cardiovascular outcomes. On the other

hand, AMPLITUDE-O achieved a reduction of MACE and other secondary outcomes

despite the shortest follow-up periods of 1.8 years (2.1 years for ELIXA and 3.2 years

for EXSCEL)[3, 6, 10], showing that exendin-based GLP-1RAs can also exert

cardioprotective and renoprotective effects.

In terms of safety outcomes of GLP-1RAs (Supplementary Figure 4), our

meta-analysis showed that the occurrence of severe hypoglycemia, pancreatitis,

pancreatic cancer, or retinopathy did not increase, although definitions of safety

outcomes were not standardized across trials (Supplementary Table 2). Individually, the

incidence of retinopathy was increased in SUSTAIN-6, which may be attributable to the

rapid lowering of blood glucose[33]. Currently, a trial on the long-term effects of

semaglutide on diabetic retinopathy (FOCUS; ClinicalTrials.gov NCT03811561)[34] is

underway, which is expected to be completed in 2024.

Meta-regression analysis

Regarding the underlying mechanism of GLP-1RA efficacy, factors associated with the

cardioprotective and renoprotective profiles of GLP-1RAs remain unclear[29]. Therefore,

to evaluate whether reduction in HbA1c or body weight (two key clinical variables

decreased by GLP-1RAs) is associated with the cardioprotective and renoprotective

effects of GLP-1RAs, we conducted a meta-regression analysis of the log-HR of MACE

and other secondary outcomes with normalized reduction of HbA1c or body weight.

As for the primary composite MACE outcome (Figure 4), the results showed

that the log-HR of MACE decreased by 0.27 for every 1.0% normalized HbA1c

reduction. In contrast, log-HR of MACE was not associated with normalized body

weight reduction. In individual studies, albiglutide caused minimal weight reduction but

still showed a reduction in log-HR of MACE. On the other hand, lixisenatide, which

showed greater normalized body weight reduction than albiglutide and dulaglutide, did

not reduce cardiovascular risk.

The present findings that only HbA1c reduction but not body weight reduction

is associated with cardiovascular risk were consistent with a recent post-hoc mediation

study of the LEADER trial[35]. The mediation analysis showed that HbA1c was

potentially the largest mediator of cardiovascular efficacy of liraglutide, with percentage

mediation being 41–83%, whereas body weight was calculated to have a much smaller

percentage mediation of 4–14%. Considering findings from our meta-regression

analysis and the mediation analysis, it seems likely that HbA1c reduction can be a

marker of cardiovascular efficacy of GLP-1RAs. However, whether HbA1c reduction

constitutes a direct contributor or represents unmeasured contributors remains to be

clarified. From a different standpoint, previous intervention trials, including

DCCT/EDIC[11, 12] and UKPDS[13-15], which evaluated the safety and efficacy of intensive

glucose-lowering therapies, also supported the association of HbA1c lowering with

cardiovascular risk reduction. However, these studies required more than ten years of

follow-up to detect cardiovascular risk reduction, which is far longer than those of the

trials in the present analysis. These discrepancies indicate that not only HbA1c lowering

but pleiotropic effects of GLP-1RAs beyond glucose-lowering, such as amelioration of

endothelial damage and chronic inflammation[36, 37], may contribute to cardiovascular

risk reduction.

As for secondary outcomes, normalized HbA1c reduction was also associated

with the composite renal outcome. On the other hand, normalized body weight

reduction had high variability among studies and was not associated with any secondary

outcome (Figure 4 and Supplementary Figure 5). Consistent with our findings that

reduction in HbA1c but not body weight is associated with the composite renal outcome,

another mediation analysis of LEADER and SUSTAIN 6 trials on the composite renal

outcome suggested that HbA1c is potentially the greatest mediator of renal benefits,

whereas body weight has little to no effect[38]. In contrast to the association of HbA1c

reduction with the renal composite outcome, we did not find a significant association

with the renal function outcome (the narrower renal endpoint). This discrepancy may be

explained by the definitions of these outcomes: the renal composite outcome included

new-onset macroalbuminuria, which is predominantly prevented by GLP-1RAs[39],

whereas renal function outcome did not (Supplementary Table 1). However, we must

note that cardiovascular outcome trials were not designed to detect renal events as the

primary outcome. In this regard, FLOW trial (NCT03819153)[40], which evaluates the

effects of semaglutide on hard renal endpoints as the primary outcome in individuals

with chronic kidney disease, is expected to provide more insights into the renal effects

of the GLP-1RAs.

Limitations

This study has several limitations. First, we used only summary data for

meta-analysis and did not use individual-level data. Second, although heterogeneity

between studies was generally low–moderate, some differences between the trials

generated notable heterogeneity. For example, the unique recruiting criteria of ELIXA

(all individuals had a recent history of acute coronary syndrome)[3] likely contributed to

high heterogeneity among studies with exendin-based GLP-1RAs. Third, although we

conducted the meta-analysis and meta-regression analysis on various secondary

outcomes in addition to the primary outcome, the included trials were not sufficiently

powered to detect the differences in the secondary outcomes. This may have reduced

the statistical power for meta-regression analysis of the secondary outcomes.

Conclusions

In conclusion, the present systematic review and meta-analysis showed that GLP-1RAs

reduce MACE, its components, all-cause mortality, hospitalization due to heart failure,

the composite renal outcome, and the renal function outcome. Furthermore, the

meta-regression analysis showed that reduction in HbA1c, but not body weight, is

associated with cardiovascular and renal risk reduction during the treatment with

GLP-1RAs.

Data sharing

All the data were extracted from publicly available sources and included in the present

article.

Acknowledgements

SY and HM are supported by the Japan Society for the Promotion of the Science.

Author contributions

SY designed the study and analyzed the data. SY and HM collected and interpreted the

data and wrote the first draft of the manuscript. HM and DT verified the analysis. DT,

SYM, NH, and NI contributed to data interpretation and critical review of the

manuscript. NI is the guarantor of the study.

Funding

None.

Conflict of interests

NI received research funds from Terumo Corp., Drawbridge, Inc., and asken Inc. NI

received speaker honoraria from Kowa Co., Ltd., MSD K.K, Astellas Pharma Inc.,

Novo Nordisk Pharma Ltd., Ono Pharmaceutical Co., Ltd., Nippon Boehringer

Ingelheim Co., Ltd., Takeda Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Corp.,

Sumitomo Dainippon Pharma Co., Ltd., Sanofi K.K., Eli Lilly Japan K.K.; received

scholarship grant from Kissei Pharmaceutical Co., Ltd., Sanofi K.K., Daiichi-Sankyo

Co., Ltd., Mitsubishi Tanabe Pharma Corp., Takeda Pharmaceutical Co., Ltd., Japan

Tobacco Inc., Kyowa Kirin Co., Ltd., Sumitomo Dainippon Pharma Co., Ltd., Astellas

Pharma Inc., MSD K.K., Ono Pharmaceutical Co., Ltd., Sanwa Kagaku Kenkyusho Co.,

Ltd., Nippon Boehringer Ingelheim Co., Ltd., Novo Nordisk Pharma Ltd., Novartis

Pharma K.K., Life Scan Japan K.K. NI is an advisory board member of Novo Nordisk.

NI and NH were investigators for SUSTAIN and PIONEER programs. SY, HM, DT,

and SYM declare no conflict of interests.

References

1. American Diabetes Association. Cardiovascular disease and risk management: standards of medical care in diabetes—2021. *Diabetes Care.* 2021;44:S125-S150.

2. Buse JB, Wexler DJ, Tsapas A, et al. 2019 Update to: Management of hyperglycemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care.* 2020;43:487-493.

3. Pfeffer MA, Claggett B, Diaz R, et al. Lixisenatide in patients with type 2 diabetes and acute coronary syndrome. *New Engl J Med.* 2015;373:2247-2257.

4. Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and cardiovascular outcomes in type 2 diabetes. *New Engl J Med.* 2016;375:311-322.

5. Marso SP, Bain SC, Consoli A, et al. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *New Engl J Med.* 2016;375:1834-1844.

6. Holman RR, Bethel MA, Mentz RJ, et al. Effects of once-weekly exenatide on cardiovascular outcomes in type 2 diabetes. *New Engl J Med.* 2017;377:1228-1239.

7. Hernandez AF, Green JB, Janmohamed S, et al. Albiglutide and cardiovascular outcomes in patients with type 2 diabetes and cardiovascular disease (Harmony Outcomes): a double-blind, randomised placebo-controlled trial. *Lancet.* 2018;392:1519-1529.

8.   Husain M, Birkenfeld AL, Donsmark M, et al. Oral semaglutide and

cardiovascular outcomes in patients with type 2 diabetes. *New Engl J Med.*

2019;381:841-851.

9.   Gerstein HC, Colhoun HM, Dagenais GR, et al. Dulaglutide and cardiovascular

outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled

trial. *Lancet.* 2019;394:121-130.

10.   Gerstein HC, Sattar N, Rosenstock J, et al. Cardiovascular and renal outcomes

with efpeglenatide in type 2 diabetes. *New Engl J Med.* 2021;385:896-907.

11.   Nathan DM, Genuth S, Lachin J, et al. The effect of intensive treatment of

diabetes on the development and progression of long-term complications in

insulin-dependent diabetes mellitus. *New Engl J Med.* 1993;329:977-986.

12.   Nathan DM, Cleary PA, Backlund JY, et al. Intensive diabetes treatment and

cardiovascular disease in patients with type 1 diabetes. *New Engl J Med.*

2005;353:2643-2653.

13.   UKPDS Group. Intensive blood-glucose control with sulphonylureas or insulin

compared with conventional treatment and risk of complications in patients with type 2

diabetes (UKPDS 33). *Lancet.* 1998;352:837-853.

14.   UKPDS Group. Effect of intensive blood-glucose control with metformin on

complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet.* 1998;

352:854-865.

15.   Holman RR, Paul SK, Bethel MA, et al. 10-year follow-up of intensive glucose

control in type 2 diabetes. *New Engl J Med.* 2008;359:1577-1589.

16.   Ma C, Avenell A, Bolland M, et al. Effects of weight loss interventions for

adults who are obese on mortality, cardiovascular disease, and cancer: systematic

review and meta-analysis. *BMJ.* 2017;359:j4849.

17.   Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an

updated guideline for reporting systematic reviews. *BMJ.* 2021;327:n71.

18.   Higgins JPT, Altman DG, Gotzsche PC, et al. The Cochrane collaboration's

tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928-d.

19.   DerSimonian R, Laird N. Meta-analysis in clinical trials revisited. *Contemp*

Clin Trials .2015;45:139-145.

20.   Huedo-Medina TB, Sanchez-Meca J, Marin-Martinez F, Botella J. Assessing

heterogeneity in meta-analysis: Q statistic or I^2 index? *Psychol Methods.*

2006;11:193-206.

21.   Taylor SI. GLP-1 receptor agonists: differentiation within the class. *Lancet*

Diabetes Endocrinol. 2018;6:83-85.

22.   Buse JB, Nauck M, Forst T, et al. Exenatide once weekly versus liraglutide

once daily in patients with type 2 diabetes (DURATION-6): a randomised, open-label

study. *Lancet.* 2013;381:117-124.

23.   Dungan KM, Povedano ST, Forst T, et al. Once-weekly dulaglutide versus

once-daily liraglutide in metformin-treated patients with type 2 diabetes (AWARD-6): a

randomised, open-label, phase 3, non-inferiority trial. *Lancet.* 2014;384:1349-1357.

24.   Pratley RE, Nauck MA, Barnett AH, et al. Once-weekly albiglutide versus

once-daily liraglutide in patients with type 2 diabetes inadequately controlled on oral

drugs (HARMONY 7): a randomised, open-label, multicentre, non-inferiority phase 3

study. *Lancet Diabetes Endocrinol.* 2014;2:289-297.

25.   Nauck M, Rizzo M, Johnson A, Bosch-Traberg H, Madsen J, Cariou B.

Once-saily liraglutide cersus lixisenatide as add-on to metformin in type 2 diabetes: A

26-week randomized controlled clinical trial. *Diabetes Care.* 2016;39:1501-1509.

26.   Pratley R, Amod A, Hoff ST, et al. Oral semaglutide versus subcutaneous

liraglutide and placebo in type 2 diabetes (PIONEER 4): a randomised, double-blind,

phase 3a trial. *Lancet.* 2019;394:39-50.

27.   Rosenstock J, Sorli CH, Trautmann ME, et al. Once-weekly efpeglenatide

dose-range effects on glycemic control and body weight in patients with type 2 diabetes

on metformin or drug naive, referenced to liraglutide. *Diabetes Care.* 2019;42:1733-1741.

28. Capehorn MS, Catarig AM, Furberg JK, et al. Efficacy and safety of once-weekly semaglutide 1.0 mg vs once-daily liraglutide 1.2 mg as add-on to 1–3 oral antidiabetic drugs in subjects with type 2 diabetes (SUSTAIN 10). *Diabetes Metab.* 2020; 46:100-109.

29. Caruso I, Cignarelli A, Giorgino F. Heterogeneity and similarities in GLP-1 receptor agonist cardiovascular outcomes trials. *Trends Endocrinol Metab.* 2019;30:578-589.

30. Das SR, Everett BM, Birtcher KK, et al. 2020 Expert consensus decision pathway on novel therapies for cardiovascular risk reduction in patients with type 2 Diabetes. *J Am Coll Cardiol.* 2020;76:1117-1145.

31. Cosentino F, Grant PJ, Aboyans V, et al. 2019 ESC guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J.* 2020;41:255-323.

32. Sharma D, Verma S, Vaidya S, Kalia K, Tiwari V. Recent updates on GLP-1 agonists: current advancements & challenges. *Biomed Pharmacother.* 2018;108:952-962.

33.   Sharma A, Parachuri N, Kumar N, et al. Semaglutide and the risk of diabetic

retinopathy—current perspective. *Eye*. 2021; Published online 9 August.

https://doi.org/10.1038/s41433-021-01741-5.

34.   Novo Nordisk. A research study to look at how semaglutide compared to

placebo affects diabetic eye disease in people with type 2 diabetes. Identifier:

NCT03811561. https://clinicaltrials.gov/ct2/show/NCT03811561. Accessed 11 Augest

2021.

35.   Buse JB, Bain SC, Mann JFE, et al. Cardiovascular risk reduction with

liraglutide: An exploratory mediation analysis of the LEADER Trial. *Diabetes Care*.

2020;43:1546-1552.

36.   Rowlands J, Heng J, Newsholme P, Carlessi R. Pleiotropic effects of GLP-1

and analogs on cell signaling, metabolism, and function. *Front Endocrinol (Lausanne)*.

2018; 9: 672.

37.   Drucker DJ. The cardiovascular biology of glucagon-like peptide-1. *Cell Metab*.

2016;24:15-30.

38.   Mann JFE, Buse JB, Idorn T, et al. Potential kidney protection with liraglutide

and semaglutide: Exploratory mediation analysis. *Diab Obes Metab*.

2021;23:2058-2066.

39.   Vitale M, Haxhi J, Cirrito T, Pugliese G. Renal protection with glucagon-like

peptide-1 receptor agonists. *Curr Opin Pharmacol.* 2020;54:91-101.   1.

American Diabetes Association. Cardiovascular disease and risk management:

standards of medical care in diabetes—2021. *Diabetes Care.* 2021;44:S125-S150.

2.   Buse JB, Wexler DJ, Tsapas A, et al. 2019 Update to: Management of

hyperglycemia in type 2 diabetes, 2018. A consensus report by the American Diabetes

Association (ADA) and the European Association for the Study of Diabetes (EASD).

Diabetes Care. 2020;43:487-493.

3.   Pfeffer MA, Claggett B, Diaz R, et al. Lixisenatide in patients with type 2

diabetes and acute coronary syndrome. *New Engl J Med.* 2015;373:2247-2257.

4.   Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and

cardiovascular outcomes in type 2 diabetes. *New Engl J Med.* 2016;375:311-322.

5.   Marso SP, Bain SC, Consoli A, et al. Semaglutide and cardiovascular outcomes

in patients with type 2 diabetes. *New Engl J Med.* 2016;375:1834-1844.

6.   Holman RR, Bethel MA, Mentz RJ, et al. Effects of once-weekly exenatide on

cardiovascular outcomes in type 2 diabetes. *New Engl J Med.* 2017;377:1228-1239.

7.   Hernandez AF, Green JB, Janmohamed S, et al. Albiglutide and cardiovascular

outcomes in patients with type 2 diabetes and cardiovascular disease (Harmony

Outcomes): a double-blind, randomised placebo-controlled trial. *Lancet.*

2018;392:1519-1529.

8. Husain M, Birkenfeld AL, Donsmark M, et al. Oral semaglutide and

cardiovascular outcomes in patients with type 2 diabetes. *New Engl J Med.*

2019;381:841-851.

9. Gerstein HC, Colhoun HM, Dagenais GR, et al. Dulaglutide and cardiovascular

outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled

trial. *Lancet.* 2019;394:121-130.

10. Gerstein HC, Sattar N, Rosenstock J, et al. Cardiovascular and renal outcomes

with efpeglenatide in type 2 diabetes. *New Engl J Med.* 2021;385:896-907.

11. Nathan DM, Genuth S, Lachin J, et al. The effect of intensive treatment of

diabetes on the development and progression of long-term complications in

insulin-dependent diabetes mellitus. *New Engl J Med.* 1993;329:977-986.

12. Nathan DM, Cleary PA, Backlund JY, et al. Intensive diabetes treatment and

cardiovascular disease in patients with type 1 diabetes. *New Engl J Med.*

2005;353:2643-2653.

13. UKPDS Group. Intensive blood-glucose control with sulphonylureas or insulin

compared with conventional treatment and risk of complications in patients with type 2

diabetes (UKPDS 33). *Lancet.* 1998;352:837-853.

14.   UKPDS Group. Effect of intensive blood-glucose control with metformin on

complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet.* 1998;

352:854-865.

15.   Holman RR, Paul SK, Bethel MA, et al. 10-year follow-up of intensive glucose

control in type 2 diabetes. *New Engl J Med.* 2008;359:1577-1589.

16.   Ma C, Avenell A, Bolland M, et al. Effects of weight loss interventions for

adults who are obese on mortality, cardiovascular disease, and cancer: systematic

review and meta-analysis. *BMJ.* 2017;359:j4849.

17.   Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an

updated guideline for reporting systematic reviews. *BMJ.* 2021;327:n71.

18.   Higgins JPT, Altman DG, Gotzsche PC, et al. The Cochrane collaboration's

tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928-d.

19.   DerSimonian R, Laird N. Meta-analysis in clinical trials revisited. *Contemp*

Clin Trials .2015;45:139-145.

20.   Huedo-Medina TB, Sanchez-Meca J, Marin-Martinez F, Botella J. Assessing

heterogeneity in meta-analysis: Q statistic or I^2 index? *Psychol Methods.*

2006;11:193-206.

21.   Taylor SI. GLP-1 receptor agonists: differentiation within the class. *Lancet*

Diabetes Endocrinol. 2018;6:83-85.

22.   Buse JB, Nauck M, Forst T, et al. Exenatide once weekly versus liraglutide

once daily in patients with type 2 diabetes (DURATION-6): a randomised, open-label

study. *Lancet.* 2013;381:117-124.

23.   Dungan KM, Povedano ST, Forst T, et al. Once-weekly dulaglutide versus

once-daily liraglutide in metformin-treated patients with type 2 diabetes (AWARD-6): a

randomised, open-label, phase 3, non-inferiority trial. *Lancet.* 2014;384:1349-1357.

24.   Pratley RE, Nauck MA, Barnett AH, et al. Once-weekly albiglutide versus

once-daily liraglutide in patients with type 2 diabetes inadequately controlled on oral

drugs (HARMONY 7): a randomised, open-label, multicentre, non-inferiority phase 3

study. *Lancet Diabetes Endocrinol.* 2014;2:289-297.

25.   Nauck M, Rizzo M, Johnson A, Bosch-Traberg H, Madsen J, Cariou B.

Once-saily liraglutide cersus lixisenatide as add-on to metformin in type 2 diabetes: A

26-week randomized controlled clinical trial. *Diabetes Care.* 2016;39:1501-1509.

26.   Pratley R, Amod A, Hoff ST, et al. Oral semaglutide versus subcutaneous

liraglutide and placebo in type 2 diabetes (PIONEER 4): a randomised, double-blind,

phase 3a trial. *Lancet.* 2019;394:39-50.

27.   Rosenstock J, Sorli CH, Trautmann ME, et al. Once-weekly efpeglenatide

dose-range effects on glycemic control and body weight in patients with type 2 diabetes

on   metformin   or   drug   naive,   referenced   to   liraglutide. *Diabetes   Care.*

2019;42:1733-1741.

28.   Capehorn MS, Catarig AM, Furberg JK, et al. Efficacy and safety of

once-weekly semaglutide 1.0 mg vs once-daily liraglutide 1.2 mg as add-on to 1–3 oral

antidiabetic drugs in subjects with type 2 diabetes (SUSTAIN 10). *Diabetes Metab.*

2020; 46:100-109.

29.   Caruso I, Cignarelli A, Giorgino F. Heterogeneity and similarities in GLP-1

receptor   agonist   cardiovascular   outcomes   trials. *Trends   Endocrinol   Metab.*

2019;30:578-589.

30.   Das SR, Everett BM, Birtcher KK, et al. 2020 Expert consensus decision

pathway on novel therapies for cardiovascular risk reduction in patients with type 2

Diabetes. *J Am Coll Cardiol.* 2020;76:1117-1145.

31.   Cosentino F, Grant PJ, Aboyans V, et al. 2019 ESC guidelines on diabetes,

pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD.

Eur Heart J. 2020;41:255-323.

32.   Sharma D, Verma S, Vaidya S, Kalia K, Tiwari V. Recent updates on GLP-1

agonists: current advancements & challenges. *Biomed Pharmacother.* 2018;108:952-962.

33.   Sharma A, Parachuri N, Kumar N, et al. Semaglutide and the risk of diabetic retinopathy—current perspective. *Eye.* 2021; Published online 9 August. https://doi.org/10.1038/s41433-021-01741-5.

34.   Novo Nordisk. A research study to look at how semaglutide compared to placebo affects diabetic eye disease in people with type 2 diabetes. Identifier: NCT03811561. https://clinicaltrials.gov/ct2/show/NCT03811561. Accessed 11 Augest 2021.

35.   Buse JB, Bain SC, Mann JFE, et al. Cardiovascular risk reduction with liraglutide: An exploratory mediation analysis of the LEADER Trial. *Diabetes Care.* 2020;43:1546-1552.

36.   Rowlands J, Heng J, Newsholme P, Carlessi R. Pleiotropic effects of GLP-1 and analogs on cell signaling, metabolism, and function. *Front Endocrinol (Lausanne).* 2018; 9: 672.

37.   Drucker DJ. The cardiovascular biology of glucagon-like peptide-1. *Cell Metab.* 2016;24:15-30.

38.   Mann JFE, Buse JB, Idorn T, et al. Potential kidney protection with liraglutide

and   semaglutide: Exploratory   mediation   analysis. *Diab   Obes   Metab.*

2021;23:2058-2066.

39.   Vitale M, Haxhi J, Cirrito T, Pugliese G. Renal protection with glucagon-like

peptide-1 receptor agonists. *Curr Opin Pharmacol.* 2020;54:91-101.

40.   Novo Nordisk. A Research study to see how semaglutide works compared to

placebo in people with type 2 diabetes and chronic kidney disease. Identifier:

NCT03819153. https://clinicaltrials.gov/ct2/show/NCT03819153. Accessed 11 August

2021.

Figure legends

Figure 1. Effects of GLP-1RAs on the primary composite MACE outcome and its

components.

GLP-1RAs = glucagon-like peptide-1 receptor agonists. MACE = major adverse

cardiovascular events.

Figure 2. Effects of GLP-1 RAs on all-cause mortality, hospitalization due to heart

failure, and renal outcomes.

GLP-1RAs = glucagon-like peptide-1 receptor agonists.

Figure 3. Univariable meta-regression analysis of normalized reduction in HbA1c (A)

or body weight (B) with the logarithm of hazard ratio (log-HR) for the primary

composite MACE outcome.

The size of each trial's circle is inversely proportional to the variance of hazard ratio.

MACE = major adverse cardiovascular events.

Figure 4. Univariable meta-regression analysis of normalized reduction in HbA1c

reduction (A–D) or body weight (E–H) with the logarithm of hazard ratio (log-HR) for

all-cause mortality (A, E), nonfatal myocardial infarction (B, F), nonfatal stroke (C, G),

and composite renal outcome (D, H).

The size of each trial's circle is inversely proportional to the variance of hazard ratio.

Table 1.

Trial name	ELIXA	LEADER	SUSTAIN-6	EXSCEL	Harmony Outcomes	REWIND	PIONEER 6	AMPLITUDE-O
GLP-1 receptor agonists	Lixisenatide	Liraglutide	Semaglutide 0.5 or 1 mg/week	Exenatide QW	Albiglutide 30 or 50 mg/week	Dulaglutide	Oral semaglutide	Efpeglenatide
dose	20 μg/day	1.8 mg/day		2 mg/week		1.5 mg/week	14 mg/day	4 or 6 mg/week
Median year of follow-up (years)	2.1	3.8	2.1	3.2	1.5	5.4	1.3	1.8
Patients	6068	9340	3297	14752	9463	9901	3138	4076
Age	59.9 ± 9.7	64.2 ± 7.2	64.6 ± 7.4	62.0 (56.0–68.0)	64.1 ± 8.7	66.2 ± 6.5	66.0 ± 7.0	64.5 ± 8.2
Sex (%female)	1861 (31%)	3337 (36%)	1295 (39%)	5603 (38%)	2894 (31%)	4589 (46%)	1007 (32%)	1344 (33%)
Body mass index (kg/m2)	30.1 ± 5.6	32.5 ± 6.3	32.8 ± 6.2	31.8 (28.2–36.2)	32.3 ± 5.9	32.3 ± 5.7	32.3 ± 6.5	32.7 ± 6.2
White ethnicity	4576 (75%)	7238 (78%)	2736 (83%)	11175 (76%)	6583 (70%)	7498 (76%)	2300 (72%)	3534 (87%)
Duration of diabetes (years)	9.2 ± 8.2	12.8 ± 8.0	13.9 ± 8.1	12.0 (7.0–18.0)	14.1 ± 8.7	10.5 ± 7.2	14.9 ± 8.5	15.4 ± 8.8
HbA1c (%)	7.7 ± 1.3	8.7 ± 1.6	8.7 ± 1.5	8.0 (7.3–8.9)	8.7 ± 1.5	7.3 ± 1.1	8.2 ± 1.6	8.9 ± 1.5
Systolic blood pressure (mmHg)	129 ± 17	136 ± 18	136 ± 17	135 (124–145)	135 ± 17	137 ± 17	136 ± 18	135 ± 16
eGFR (mL/min/1.73 m2)†	78 ± 21	80	80 (61–92)	77 (61–92)	79 ± 25	75 (61–91)	74 ± 21	72 ± 22
Previous cardiovascular disease (%)	6068 (100%)	7598 (81%)	2735 (83%)	10782 (73%)	9463 (100%)	3114 (31%)	2695 (85%)	3650 (90%)
Previous heart failure (%)	1358 (22%)	1667 (18%)	777 (24%)	2389 (16%)	1922 (20%)	853 (9%)	288 (12%)	737 (18%)
Other glucose-lowering therapies								
Insulin	2374 (39%)	4169 (45%)	1913 (58%)	6368 (46%)	5597 (59%)	2363 (24%)	1930 (61%)	2560 (63%)
Metformin	4021 (66%)	7144 (76%)	2414 (73%)	11295 (77%)	6969 (74%)	8037 (81%)	2463 (77%)	2985 (72%)
Sulfonylurea	2004 (33%)	4733 (51%)	1410 (43%)	5401 (37%)	2725 (29%)	4552 (46%)	1027 (77%)	1036 (25%)
Thiazolidinedione	95 (2%)	575 (6%)	76 (2%)	579 (4%)	194 (2%)	168 (2%)	118 (4%)	–
DPP4 inhibitor	–	6 (<1%)	5 (<1%)	2203 (15%)	1437 (15%)	564 (6%)	2 (<1%)	–
SGLT2 inhibitor	–	–	5 (<1%)	77 (1%)	575 (6%)	620 (6%)	305 (10%)	618 (15%)

Continuous variables are presented as mean ± standard deviation (except for EXSCEL), and

categorical variables are presented as numbers (%). Continuous data on EXSCEL are presented as

median (Interquartile range, IQR).

[†]eGFR data on SUSTAIN-6, EXSCEL, and REWIND are expressed as median (IQR). The standard

deviation of eGFR in LEADER is not available.

GLP-1 = glucagon-like peptide-1, Exenatide QW = Exenatide once weekly, HbA1c = glycated

haemoglobin, eGFR = estimated glomerular filtration rate.

Figure 1.

Figure 2.

All-cause mortality	Treatment Events/Total (%)	Placebo Events/Total (%)		Weight	Hazard ratio (95% CI)
ELIXA	211/3034 (7%)	223/3034 (7%)		11.2%	0.94 (0.78–1.13)
LEADER	381/4668 (8%)	447/4672 (10%)		19.3%	0.85 (0.74–0.97)
SUSTAIN-6	62/1648 (4%)	60/1649 (4%)		3.3%	1.05 (0.74–1.50)
EXSCEL	507/7356 (7%)	584/7396 (8%)		25.0%	0.86 (0.77–0.97)
HARMONY	196/4731 (4%)	205/4732 (4%)		10.5%	0.95 (0.79–1.16)
PIONEER 6	23/1591 (1%)	45/1592 (3%)		1.7%	0.51 (0.31–0.84)
REWIND	536/4949 (11%)	592/4952 (12%)		25.0%	0.90 (0.80–1.01)
AMPLITUDE-O	111/2717 (4%)	69/1359 (5%)		4.5%	0.78 (0.58–1.06)
Overall	**2027/30694 (7%)**	**2225/29386 (8%)**		**100.0%**	**0.88 (0.82–0.94)**
					***P* < 0.001**

Heterogeneity: $I^2 = 10.5\%$, $P = 0.349$

Hospitalization due to heart failure	Treatment Events/Total (%)	Placebo Events/Total (%)		Weight	Hazard ratio (95% CI)
ELIXA	122/3034 (4%)	127/3034 (4%)		12.6%	0.96 (0.75–1.23)
LEADER	218/4668 (5%)	248/4672 (5%)		22.8%	0.87 (0.73–1.05)
SUSTAIN-6	59/1648 (4%)	54/1649 (3%)		5.8%	1.11 (0.77–1.61)
EXSCEL	219/7356 (3%)	231/7396 (3%)		22.0%	0.94 (0.78–1.13)
HARMONY	79/4731 (2%)	111/4732 (2%)		9.5%	0.71 (0.53–0.94)
PIONEER 6	21/1591 (1%)	24/1592 (2%)		2.3%	0.86 (0.48–1.55)
REWIND	213/4949 (4%)	226/4952 (5%)		21.5%	0.93 (0.77–1.12)
AMPLITUDE-O	40/2717 (1.5%)	31/1359 (2.3%)		3.5%	0.61 (0.38–0.98)
Overall	**977/30694 (3%)**	**1052/29386 (4%)**		**100.0%**	**0.89 (0.82–0.98)**
					***P* = 0.013**

Heterogeneity: $I^2 = 2.5\%$, $P = 0.410$

Composite renal outcome	Treatment Events/Total (%)	Placebo Events/Total (%)		Weight	Hazard ratio (95% CI)
ELIXA	172/2647 (6%)	203/2639 (8%)		12.8%	0.84 (0.68–1.02)
LEADER	268/4668 (6%)	337/4672 (7%)		17.4%	0.78 (0.67–0.92)
SUSTAIN-6	62/1648 (4%)	100/1649 (6%)		6.3%	0.64 (0.46–0.88)
EXSCEL	366/6256 (6%)	407/6222 (7%)		19.5%	0.88 (0.76–1.01)
REWIND	848/4949 (17%)	970/4952 (20%)		27.1%	0.85 (0.77–0.93)
AMPLITUDE-O	335/2717 (13%)	250/1359 (18%)		16.8%	0.68 (0.57–0.79)
Overall	**2051/22885 (9%)**	**2267/21493 (11%)**		**100.0%**	**0.80 (0.73–0.87)**
					***P* < 0.001**

Heterogeneity: $I^2 = 45.0\%$, $P = 0.106$

Renal function outcome	Treatment Events/Total (%)	Placebo Events/Total (%)		Weight	Hazard ratio (95% CI)
ELIXA	41/3031 (1%)	35/3032 (1%)		8.1%	1.16 (0.74–1.83)
LEADER	87/4668 (2%)	97/4672 (2%)		16.5%	0.89 (0.67–1.19)
SUSTAIN-6	18/1648 (1%)	14/1649 (1%)		3.7%	1.28 (0.64–2.58)
EXSCEL	246/6456 (4%)	273/6458 (4%)		29.6%	0.88 (0.74–1.05)
REWIND	169/4949 (3%)	237/4952 (5%)		25.9%	0.70 (0.57–0.85)
AMPLITUDE-O	121/2717 (5%)	76/1359 (6%)		16.2%	0.77 (0.57–1.02)
Overall	**682/22885 (3%)**	**732/21493 (3%)**		**100.0%**	**0.84 (0.73–0.97)**
					***P* = 0.016**

Heterogeneity: $I^2 = 31.4\%$, p = 0.200

Figure 3.

(A) Primary composite MACE outcome and HbA1c reduction

Slope = -0.27
(95% CI -0.53, -0.01),
P = 0.045

Lixisenatide
Exenatide QW
Dulaglutide
Liraglutide
Albiglutide
Oral semaglutide
Semaglutide
Efpeglenatide

Log-HR of primary component MACE outcome

Normalized reduction in HbA1c

(B) Primary composite MACE outcome and body weight reduction

Slope = -0.05
(95%CI -0.17, 0.07),
P = 0.390

Lixisenatide
Exenatide QW
Liraglutide
Dulaglutide
Albiglutide
Oral semaglutide
Efpeglenatide
Semaglutide

Log-HR of primary component MACE outcome

Normalized reduction in body weight

Figure 4.

(A) Cardiovascular mortality and HbA1c reduction
Slope = -0.13
(95% CI -0.55, 0.29),
P = 0.548

(B) Nonfatal myocardial infarction and HbA1c reduction
slope = -0.17
(95% CI -0.55, 0.20),
P = 0.368

(C) Nonfatal stroke and HbA1c reduction
Slope = -0.46
(95% CI -0.94, 0.02),
P = 0.063

(D) Composite renal outcome and HbA1c reduction
Slope = -0.35
(95% CI -0.69, -0.01),
P = 0.045

(E) Cardiovascular mortality and body weight reduction
Slope = -0.17
(95% CI -0.55, 0.20),
P = 0.368

(F) Nonfatal myocardial infarction and body weight reduction
Slope = -0.02
(95% CI -0.19, 0.16),
P = 0.242

(G) Nonfatal stroke and body weight reduction
Slope = -0.07
(95% CI -0.24, 0.09),
P = 0.388

(H) Composite renal outcome and body weight reduction
Slope = -0.14
(95% CI -0.30, 0.01),
P = 0.067