Establishment of a teaching hospital-based dementia consultation service for rurally-based regional district general hospitals

Björn H. Schott[1,2,3,6,*,#], Jakob Christian Voetlause[1,*], Juliana Lisa Amoah[1], Alexander
Kratzenberg[1,2], Michael Belz[1], Tobias Knipper[1], Charles Timäus[1], Carmen Beskow[1], Catherine
M. Sweeney-Reed[5,6], Jens Wiltfang[1,2,4], and Katrin Radenbach[1,#]

[1]Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Göttingen,

Germany

[2]German Center for Neurodegenerative Diseases, Göttingen, Germany

[3]Leibniz Institute for Neurobiology, Magdeburg, Germany

[4]Neurosciences and Signaling Group, Institute of Biomedicine (iBiMED), Department of Medical

Sciences, University of Aveiro, Aveiro, Portugal

[5]Neurocybernetics and Rehabilitation, Dept. of Neurology, Otto von Guericke University Magdeburg,

Magdeburg, Germany

[6]Center for Behavioral Brain Sciences, Otto von Guericke University Magdeburg, Magdeburg,

Germany

[*]These authors contributed equally to this work

15
[#] **Correspondence:**
Dr. Katrin Radenbach & Dr. Björn Schott
Department of Psychiatry and Psychotherapy
University Medical Center Göttingen
Von Siebold-Str. 5
37075 Göttingen, Germany
katrin.radenbach@med.uni-goettingen.de
bjoernhendrik.schott@med.uni-goettingen.de

Keywords: dementia, delirium, consultation service, rural hospitals, demographic change,
rural health care

Abstract

Objective: The treatment of patients with dementia poses a considerable challenge to regional district
general hospitals, particularly in rural areas. Here we report the establishment and initial evaluation of
a dementia-specific consultation service provided by a teaching hospital-based Psychiatry Department
to regional district general hospitals in surrounding smaller towns.

Methods: The consultation service was provided to patients with pre-existing or newly suspected
dementia, who were in acute hospital care for concurrent conditions. An evaluation of 61 consultations
– 49 on-site and 12 via telemedicine – was performed to assess the needs of the participating hospitals
and the specific nature of the referrals to the consultation service.

Results: Suspected dementia or cognitive dysfunction was the primary reason for consultation requests
(>50% of cases). Other common requests concerned suspected delirium, behavioral symptoms, and
therapeutic recommendations. During the consultations, a diagnosis of dementia was reached in 52.5%
of cases, with other common diagnoses including delirium and depression. Recommendations related
to pharmacotherapy were given in 54.1% of consultations. Other recommendations included referral
for outpatient neurological or psychiatric follow-up, further diagnostic assessment, or assessment in a
memory clinic. Geriatric psychiatric inpatient treatment was recommended in only seven cases (11.5
%).

Conclusions: Our initial evaluation demonstrates the feasibility of providing a dementia-specific
consultation service in rural areas. The service has the potential to reduce acute transfers to inpatient
geriatric psychiatry and enables older patients with dementia or delirium to be treated locally by
helping and empowering rurally-based regional hospitals to manage these problems and associated
complications.

1 Introduction

Patients with dementia admitted to hospital for acute treatment of concurrent conditions are at increased risk of complications and adverse events such as falls, delirium, or severe behavioral symptoms, culminating in overall higher mortality (Inouye et al., 2014b; Dewing and Dijk, 2016; Blair et al., 2018). Advanced age is the primary risk factor for developing dementia, and the proportion of individuals affected by dementia is therefore increasing steeply in aging populations, such as in most European countries(Cao et al., 2020).

The situation of people with dementia in hospital is frequently complicated by other related conditions, which may present similarly, yet necessitate a different or more comprehensive treatment approach. A cardinal example is delirium, which is common following surgery (Ní Chróinín et al., 2021), and can be aggravated by inadequate medication (Inouye et al., 2014b). Delirium typically results in prolonged length of inpatient stay and a short-term 20-fold increase in mortality. A quarter of older patients with delirium die within three to four months, and 41% of previously self-caring individuals require discharge to a nursing facility. Symptoms are fully reversible in approximately 50% of cases (Inouye et al., 2014a), but recovery may depend on timely recognition and initiation of appropriate treatment. Another challenge in the care of the elderly, and especially patients with dementia, is the frequent polypharmacy, which refers to the use of four to five or even more medications at the same time. Even in the absence of obvious side-effects, polypharmacy must be regarded as a risk factor for the occurrence of complications like delirium or behavioral disorders. Such complications can arise because numerous pharmaceuticals, for example loop diuretics (Sica, 2004) or digitalis glycosides (Marvanova, 2016), are generally capable of triggering delirium in predisposed patients. Furthermore, psychotropic medications frequently used for behavioral disorders may themselves induce considerable somatic and psychiatric side-effects (Inouye et al., 2014a).

Several approaches have been proposed to improve the care for patients with dementia and to prevent delirium in acute care hospitals. Interventions to improve care in general include patient-centered care provided by hospital staff (Chenoweth et al., 2021) or the training of volunteers (Blair et al., 2018). To specifically reduce the incidence of delirium, both non-pharmacological (Burton et al., 2021) and pharmacological (Park et al., 2021) interventions show moderate effectiveness, but there is considerable variability concerning the benefit for the individual patient.

While these approaches are promising and are currently being further developed, their applicability may be particularly challenging in smaller, rural hospitals: In addition to the increasing average age of the population, younger individuals frequently concentrate in larger cities, leaving small towns and rural areas predominantly affected by the consequences of having an aging population (Milbert and Sturm, 2016). This trend leads to a disproportionately increasing number of patients with dementia in rural hospitals, and at the same time, contributes to the lack of qualified staff to support specialized units for dementia treatment and care, such as neurology or (geriatric) psychiatry departments (van den Berg et al., 2011; Mathur et al., 2019). Here we propose and evaluate the feasibility of an out-reach consultation service provided by a specialized institution within rural surroundings to alleviate this problem.

We report the establishment of a dementia-specific consultation service to improve the care for patients with dementia treated in rurally-based regional hospitals. The service as described here was provided by the Department of Psychiatry and Psychotherapy of the University Medical Center Göttingen, Germany, and was available for patients treated in local regional hospitals upon request by their treating clinicians. An initial evaluation was performed to assess the needs and questions

94　commonly raised by the treating hospitals as well as recommendations commonly made by the
95　providers of the service.

2　Methods

2.1.　Establishment of a dementia consultation service

2.1.1.　Preparatory work

100　The proposal for establishing a dementia-specific consultation service for rurally-based regional
101　hospitals was initially developed as part of an initiative funded by the Department of Social Affairs,
102　Health and Equality of the State of Lower Saxony (2019). The study was approved by the Ethics
103　Committee of the University Medical Center Göttingen (approval number 13/12/19) and was registered
104　with the German Clinical Trials Register (registration number DRKS00022366).

2.1.2.　Setting and participating hospitals

The consultation service was provided by the Department of Psychiatry and Psychotherapy of the
University Medical Center Göttingen (UMG). The university city of Göttingen, with a population of
118,911 (December 2019, https://www.statistik.niedersachsen.de), is surrounded by the rural counties
of Göttingen (county), Goslar, and Northeim. We identified nine hospitals without in-house Psychiatry
and/or Neurology Departments in the assigned service area of the Department of Psychiatry and
Psychotherapy of the UMG.

2.1.3.　Team

113　The team consisted of two senior residents in psychiatry (T.K. and C.T.), a nurse with long-standing
114　experience in geriatric psychiatry and dementia care (C.B.), as well as a project coordinator (A.K.) and
115　two research students (J.C.V. and J.L.A.). Organizational and clinical supervision was carried out by
116　two senior psychiatrists specialized in geriatric psychiatry and/or consultant liaison psychiatry (K.R.
117　and B.H.S.).

2.1.4.　Legal issues and service agreements

The service agreement was drawn up in collaboration with the UMG's legal department. After a
formal presentation of the project was held in a partner hospital expressing interest, the written service
agreements were sent to the respective hospital for review if further interest was expressed. Upon
mutual agreement, the service agreements were signed by the Medical Director of the Department of
Psychiatry and Psychotherapy of the UMG and the board members (UMG and partner hospital).
Service agreements were established with five partner hospitals, including four general care hospitals
and one hospital specialized in interdisciplinary diabetes care. Agreements were signed between
18.05.2020 and 21.07.2020.

2.2.　Implementation of the consultation service

2.2.1.　Request and performance of consultations

Information regarding the process of requesting consultations, including contact options, was
provided to the hospitals as a flowchart during the initial project visit. Consultations could be requested
by phone, e-mail, or fax. They included history and neuropsychiatric examination and were performed
either on-site or via telemedical service, and were followed by recommendations regarding diagnostic
and therapeutic measures. Further diagnostic assessment for the differential diagnosis of a dementia

syndrome or its complications was recommended where applicable, as well as specific therapies and
support from Social Services. Specific attention was paid to complex polypharmacy, and advice was
given by physicians on how to avoid possible pharmacological risk factors for deterioration of the
patient's condition (e.g., by avoiding potentially delirium-inducing substances).

An educational program complemented the consultation service. The program included both
scheduled talks on selected topics concerning dementia treatment and care as well as hands-on
teaching. A nurse with long-standing experience in geriatric psychiatry and dementia care supported
the nursing staff on-site by providing advice on the nursing management of patients with dementia,
including means of communication and improving the hospital environment to meet the special needs
of these patients. Advice was provided both in general and on a case-by-case basis. A detailed
evaluation of the educational program and the local expertise form part of a separate, ongoing, study.

2.2.2. Standardized documentation of consultations

A standardized form was used for systematic clinical documentation of the consultations, including
the questions leading to consultation, a detailed medical and psychiatric history, and the
recommendations made by the providing physician. The questionnaire was available in both digital
and paper form (see Appendix). The Mini-Mental State Examination (MMSE) was additionally
administered to a total of 40 patients. While primarily designed to meet the clinical needs of the
requesting institutions, these standardized forms also provided the basis for the statistical evaluation of
the consultations reported here.

2.3. Preliminary evaluation

2.3.1. Data extraction and statistical analysis

A central aspect of the current study was the evaluation of the questions raised by the partner
institutions when requesting our consultation service for an individual patient. The circumstances
leading to initiation of the consultations were also considered. Since the wording of questions arising
under comparable circumstances varied, we performed a content analysis using inductive
categorization to enable quantitative analyses (Mayring and Fenzl, 2014). As consultation forms were
primarily completed to meet the needs of the referring clinicians, the categories were defined according
to the existing content. For example, the category *Suspected Dementia* included questions regarding
dementia, memory impairment, and cognitive decline, whereas questions concerning confusion and
disorientation were also assigned to the category *Suspected Delirium*. Furthermore, questions relating
to drug intolerance and to recommendations on pharmacotherapy were included in a single category
(*Pharmacological Issues*): referring clinicians typically requested advice regarding a potential change
or discontinuation of medication in both areas.

Given the preliminary nature of the current evaluation, we report our findings primarily as
frequency data. Where inference statistics were applicable, X^2 tests were used to evaluate frequency
deviations, and t-tests were applied to continuous data. Owing to the relatively small sample sizes,
unequal variances were assumed in independent-sample t-tests (i.e., Welch's t-tests were used).

2.3.2 Follow-up consultations

We additionally considered separately the requests for follow-up consultations, to gain insights into
the medium-term outcome of the implementation of the program. Due to the low number of follow-up
consultations, they could only be considered qualitatively.

3   Results

3.1. Consultations performed across participating institutions

The consultations analyzed took place from 10.07.2020 to 14.03.2021. Within the study period, 65
consultations were performed, four of which were follow-up consultations provided for patients who
had already been examined previously, in order to follow the subsequent course of their condition and
to potentially adapt diagnostics and therapy accordingly. The sample thus consists of 65 consultations
for 61 patients. The following general evaluation of the data is based on the number of patients (N =
61).

Despite limitations imposed by the ongoing COVID-19 pandemic during the study period, the
majority of consultations (N = 49, 80.3%) were performed on-site (Figure 1A). The number of requests
varied considerably between the participating hospitals (N = 5; Figure 1B), significantly deviating from
an equal distribution (X^2 (4) = 77.93, p < .001), with more than half of the consultations requested by
one hospital (Figure 1B).

3.2. Demographic and clinical characteristics of the patients

3.2.1. Demographics of the patients

Among the patients in our sample, more than half were 80 years or older (Figure 1C), and the
number of women in the age groups of 80 years and above was higher compared to men (Figure 1D).
The average age of women among the patients was 82.5 years (men: 73.1 years, t (27.7) = 2.94, p =
.007). While this asymmetry was generally in line with the higher life expectancy among women, it
should be noted that the proportion of men (n = 19) was still lower than would be expected based on
the general population (2021).

3.2.2. Conditions leading to hospital admission and indications for consultation requests

The conditions leading to acute admission to the partner hospitals were categorized according to
the relevant medical specialty or the specific reason for admission. In absolute numbers, the most
common reasons for patients' admission to the partner hospitals were problems classified under
Surgery (Figure 2A), with falls being the most frequently reported event leading to admission (N = 15,
24.6%). Other common reasons were categorized under Internal Medicine, Neurology or Psychiatry,
and planned adaptation or modification of pharmacotherapy. The latter arose particularly at the partner
hospital specializing in the treatment of diabetes. Notably, in 17 cases (27.9%), the primary reasons
for admission to the partner hospitals were neuropsychiatric problems, although the partner hospitals
had neither neurology nor psychiatry departments.

The most common symptoms reported to the service by the partner institutions when requesting a
consultation could be classified as being related to *Behavioral Symptoms*, *Risk of Falls*, *Cognitive
Impairment*, and *Sleep Disorders* (Figure 2B). Following data extraction, the questions were classified
as indications for consultation and assigned to the following categories: *Suspected Dementia*,
Suspected Delirium, *Behavioral Symptoms in Dementia*, *Pharmacological Issues* and *Suspected
Depression* (Figure 2B).

In line with the focus of our consultation service on dementia, over half of consultation requests
(50.8%, Figure 2C) were related to suspected dementia and/or cognitive impairment, and over a fifth
(23.0%) specifically concerned behavioral symptoms in dementia. Questions relating to
pharmacotherapy comprised both suspected adverse effects and requests for recommendations
regarding pharmacological treatment.

3.2.3. Pre-existing conditions and risk factors

A pre-existing confirmed diagnosis of dementia was reported in only nine consultation requests
(14.8%) (Figure 3A). More broadly, a previous history of cognitive deficits was reported in 22 patients
(36.1%), suggesting that dementia or cognitive impairment in older adults may be underdiagnosed
among patients in rural primary care hospitals.

Beyond dementia and cognitive dysfunction, 38 patients (64.4%) had a history of at least three
chronic medical conditions, and 23 (39.0%) had six or more diagnoses of a chronic disease.
Considering the importance of delirium in the differential diagnosis for dementia and as a risk factor
for morbidity and mortality in older adults (Inouye et al., 2014a), risk factors for developing delirium
were specifically explored. Acute or chronic pain, hearing or visual impairment, recent surgical
procedures, and infections were noted most frequently (Figure 3B).

3.3. Consultations

3.3.1. Diagnostic assessment

Mirroring the focus of the consultation requests on suspected dementia, the group of dementias
(ICD-10: F00 – F03) constituted the diagnoses made most frequently by the physicians performing the
consultations (Figure 4A). Further diagnoses typically associated with cognitive symptoms included
delirium (ICD-10: F05.x, F1x.4) and mild cognitive impairment (MCI; ICD-10: F06.7), a risk state for
dementia (Schmidtke and Hermeneit, 2008). Notably, among the dementia subtypes, vascular
dementias were most frequently recorded as diagnoses (Figure 4B), despite dementia in Alzheimer's
disease being the most common form of dementia in the population (2016). The MMSE (Folstein et
al., 1975) was performed in 40 patients (Figure 4C), and the average MMSE score was lower in patients
with a diagnosis of dementia (18.86 ± 5.16) than in those with other diagnoses (22.75 ± 5.01; t (21.7)
= 22.23, p = 0.016). As expected, polypharmacy was highly prevalent among the patients referred for
consultation, with a median number of 10 different medications (range: $2 - 23$; mean = 10.31 ± 5.01;
Figure 4D). There was, however, no difference in the number of medications as a function of dementia
diagnosis or delirium (all p >0.468).

3.3.2. Recommendations

The physicians performing the consultations made a wide range of recommendations (Figure 5A).
Recommendations regarding pharmacotherapy given most frequently (54.1% of patients). Among
these, initiation of novel pharmacological treatment, including medication to be administered as
required (*pro re nata*, PRN), was most commonly recommended, followed by discontinuation or
reduction, and change of medication, which were recommended with equal frequency (Figure 5B).
Non-pharmacological measures to treat behavioral symptoms were recommended for six patients
(9.8%), but it should be noted that assistance with non-pharmacological interventions was primarily
provided by the geriatric nurse (C.B.), who interacted with the nursing staff of the participating
hospitals largely independently and provided advanced training in non-pharmacological treatment of
dementia patients.

Further diagnostic assessment, referral for outpatient neuropsychiatric treatment, or consultation at
a memory clinic were also frequently recommended. For seven patients (11.5%), transfer to a geriatric
psychiatric inpatient ward was recommended. Follow-up consultations were recommended for seven
patients but were subsequently requested and performed for four patients only.

3.3.3. Follow-up consultations

Follow-up consultations were requested for three of the patients, with one patient receiving two
follow-up visits. In two cases, follow-up visits were requested for patients who had initially been seen
by the consultation service for suspected dementia and/or post-stroke depression after the patients had
shown nocturnal episodes of confusion suggestive of delirium.

4   Discussion

The main goal of the study, the establishment of a combined outreach and telemedical dementia
consultation service for rurally-based regional district general hospitals, was successfully achieved
despite the obstacles posed by the ongoing COVID-19 pandemic. The requests received from the
participating hospitals emphasize the need for the dissemination of expertise in dementia treatment and
care among smaller hospitals in rural areas.

4.1.   Implementation and evaluation of the consultations

The basic goal of the project, to improve the general care of people with dementia in rurally-based
regional hospitals, was achieved: Agreements were signed with five partner hospitals, and
consultations were performed for all five institutions, albeit with considerably varying frequency.
Notably, over half of the consultations were requested by one of the participating hospitals. This
tendency is probably best explained by the fact that this hospital has a geriatric department, which
treats a large number of people with dementia, and the staff in this department may be more sensitive
to the presence of cognitive or behavioral symptoms.

Across all participating hospitals, consultations were predominantly requested in relation to
previously diagnosed or suspected dementia or associated behavioral problems (Figure 2C),
confirming the need for a dementia-specific consultation service. Other frequent diagnoses included
delirium, which is often difficult to distinguish cross-sectionally from dementia (Inouye et al., 2014b),
and depressive syndromes, which are an important part of the differential diagnosis, especially in
individuals with subjectively reported cognitive symptoms (Arie, 1983). We note the relatively low
number of consultation requests (14.8 %) for patients with a pre-existing confirmed diagnosis of
dementia. Although not explicitly stated in the information provided to the referring hospitals
introducing the consultation service, our findings suggest that the service was considered to be useful
primarily for establishing a new diagnosis, whereas patients who had previously received a diagnosis
were likely to be already under regular medical care. Indeed, the predominant reason for requesting a
consultation was for diagnosis, with suspected dementia (50.8 %), suspected delirium (24.6 %), and
suspected depression (8.2 %) accounting for over three-quarters of the consultation indications.

Somewhat unexpectedly, the diagnosis of vascular dementia was the most frequently diagnosed
form of dementia during the consultations. Given that dementia due to Alzheimer's disease is the most
common form in older adults (2016), we cannot exclude the possibility that some patients diagnosed
with vascular dementia may nevertheless have had amyloid pathology and would thus have to be
correctly classified as having mixed dementia. This differential diagnosis was, however, difficult to
make as part of the consultation and should be performed, for example, in a memory clinic (Wallin et
al., 2016). Alternatively, or more likely additionally, the proportionately high prevalence of vascular
dementia in this patient group might reflect local risk profiles. The risk of major cardiovascular disease
in Lower Saxony is higher than the average in Germany (Dornquast, 2016). Interestingly, however, a
study previously conducted in the same region identified an under-diagnosis of Alzheimer's dementia,
with a corresponding over-diagnosis of an underlying vascular cause for dementia symptoms (Stoppe
et al., 1994). Although a subsequent study found a reduction in the effect over the intervening years,

the tendency nonetheless persisted (Maeck et al., 2007), which may contribute to the proportions
among the diagnoses in our cohort.

More than 70% of the patients treated were receiving eight or more medications (Figure 4D),
emphasizing the importance of considering the risks associated with polypharmacy in older patients,
and particularly those with dementia, or more generally, cognitive impairment (Inouye et al., 2014a;
Blair et al., 2018). Accordingly, in more than 50% of the consultations, an adjustment of
pharmacotherapy was recommended, which commonly involved dose reduction or discontinuation, as
well as replacement with medication with a more favorable side-effect profile (Figure 5). We note that
the number of medications taken by a given patient was independent of patient age, degree of cognitive
impairment, and diagnosis reached. Although such dependencies may become apparent in a larger
study population, the finding underlines the importance of service provision for this patient group that
is not limited to tight criteria but rather is based on communication with colleagues at the partner
hospitals.

A direct transfer to a (geriatric) psychiatric inpatient unit was recommended in only 11.5% of cases.
This low referral rate indicates that acute transfers of people with dementia to psychiatric departments
– often associated with great distress for those affected – can possibly be at least partially avoided by
providing adequate outreach consultation services. While further neurological or psychiatric
assessment was indeed frequently recommended, it can often take place in an outpatient setting. The
frequent recommendation of further diagnostic assessment, referral for outpatient neuropsychiatric
treatment, or consultation at a memory clinic reflect the fact that dementia and cognitive impairment
are chronic conditions that cannot be adequately treated in a single hospital stay. Access to the proposed
consultation service meant that these unmet needs could be identified and seamlessly addressed by
appropriate specialist facilities following the acute hospital stay, as opposed to requiring a subsequent
separate assessment in the community before the necessary referral could be made.

The follow-up findings support the notion that the occurrence of delirium in patients with pre-
diagnosed dementia is a common and clinically relevant complication warranting appropriate
treatment. While the low number of follow-up consultations does not allow us to generalize from this
observation, it is in line with the frequent occurrence of delirium in hospitalized patients with pre-
existing dementia (Han et al., 2022).

4.2. Partnership with participating hospitals

Implementing a partnership with external hospitals (partly under private ownership) requires
establishment of continuous communication. Individual approaches and intense exchange were
essential in building a system of shared patient care. Gaining the interest of potential partners in the
project was best achieved through personal contact, preferably on-site. Establishing contact via letter
or e-mail remained more challenging, presumably due to the limited time capacities of the respective
chief and senior physicians. Once the partnership had been established, it was essential to have fixed
contact persons at the respective partner hospitals, who were aware of the possibility of dementia-
specific consultations and could both request consultations and distribute information regarding our
service. The establishment of a routine with fixed appointments and contact persons proved helpful for
regular consultation requests. Telemedical consultations were not possible at all sites due to limited
Wi-Fi and other technical factors. The successful implementation of this service in 12 cases, however,
provides support for the feasibility of such an approach, which may grow in relevance during the
current COVID-19 pandemic as well as future potential pandemics, in which this patient group is at
high risk of serious consequences of infection, due to both advanced age and comorbidity. A specialist
consultation service can provide diagnostic assessment and treatment plans without increasing social
contact and infection risk.

The role of the nurse in providing non-pharmacological interventions underlines the requirement
for a multidisciplinary service to address the clinical needs of this patient group, as highlighted in
previous studies (Blair et al., 2018; Burton et al., 2021).

4.3.Limitations

A full evaluation of this consultation service requires a systematic, in-depth, long-term follow-up
study. Future work should include a comprehensive quantitative and qualitative assessment of the
impact of the implementation of the service. We suggest that the feasibility of the program
implementation, as demonstrated here, provides the basis for such future work.

As only three patients received repeated visits, we could not derive quantitative information
regarding the potential impact of the program on the longitudinal course of individual patients'
treatment. In a larger-scale future study, a greater number of patients would be expected to receive
repeated consultations, which would enable collection of more representative information on medium-
as well as long-term outcomes.

4.4. Conclusion and perspectives

Our findings indicate that provision of a dementia-specific consultation service for rurally-based,
regional district general hospitals is feasible for university-based geriatric psychiatric units. We suggest
that the concept of the dementia consultation service provided by a tertiary referral center (e.g.,
university hospital) to surrounding regional hospitals may provide a helpful addition to improve care
for patients with dementia in rural areas. The service has the potential to reduce acute transfers to in-
patient geriatric psychiatry. It enables older patients with dementia or delirium to be treated locally by
assisting and empowering rurally-based regional hospitals to manage these problems and associated
complications. Future studies are required to provide a more extensive evaluation of these effects. As
our project was tailored to meet the capacities and needs of the German healthcare system, future
research should also investigate to what extent the concept is transferrable to other geographic regions
with comparable organization of hospital care.

5 Conflict of Interest

378 The authors have no conflict of interest, financial or otherwise, to declare.

6 Author Contributions

B.H.S., J.W., and K.R. conceived and designed the study and acquired funding; A.K. coordinated
the consultation service; T.K. and C.A.T. performed the medical consultations; C.B. performed the
nursing consultations; B.H.S. and K.R. supervised consultations; J.C.V., J.L.A., and A.K. performed
data curation; B.H.S., J.C.V., and M.B. analyzed the data; B.H.S. and J.C.V. wrote the first draft of the
manuscript; C.M.S.R., B.H.S., M.B. and K.R. interpreted the analyses and edited the manuscript; all
authors contributed to editing and critically revising the manuscript.

7 Funding

387   This work was supported by the State of Lower Saxony, Department of Social Affairs, Health and
388 Equal Opportunities (funding reference: 3SL_G.Demenz-04-2019). B.H.S. holds a grant from the
389 European Union and the State of Saxony-Anhalt (Research Alliance *Autonomy in Old Age*). The
390 funders had no role in the design or conduct of the study.

8 Acknowledgments

392   The authors would like to thank the staff of all participating hospitals. We thank Thomas Voigt and
393 Matthias Siegert from the legal department of the UMG for support drafting the agreements and Petra
394 Arndt, Hermann Esselmann, Ute Gotthardt, Cornelia Hönicke, Tobias Urbanczyk, and David Zilles for
395 help with project administration. We further thank Hermann Esselmann for helpful comments on this
396 manuscript.

9 Ethics Statement

This work was approved by the Ethics Committee of the University Medical Center Göttingen
(approval number 13_12_19). Informed consent to be examined and treated by a physician from the
University Medical Center Göttingen (UMG) was obtained from the patients or their caregivers by the
participating hospitals. Additional consent to participate in research was waived by the Ethics
Committee of the UMG, as all consultations were performed according to the current S3 Guidelines
Dementia, and all data were processed and analyzed in a fully anonymized way.

10 Data Availability Statement

405 Data from this study are available in anonymized form from the corresponding authors upon
406 reasonable request.

11   References

Alzheimer's Association (2016). 2016 Alzheimer's disease facts and figures. Alzheimers Dement 12, 459-509.

Niedersächsisches Ministerium für Soziales Gesundheit und Gleichstellung (2016) [Online]. Available: https://www.ms.niedersachsen.de/startseite/service_kontakt/presseinformationen/demenzkranke-im-krankenhaus-besser-versorgen-land-foerdert-modellprojekte-mit-15-millionen-euro-175936.html.

Statistisches Bundesamt (2021) [Online]. Available: https://www.destatis.de/DE/Themen/Querschnitt/Demografischer-Wandel/_inhalt.html.

Arie, T. (1983). Pseudodementia. Br Med J (Clin Res Ed) 286, 1301-1302.

Blair, A., Anderson, K., and Bateman, C. (2018). The "Golden Angels": effects of trained volunteers on specialling and readmission rates for people with dementia and delirium in rural hospitals. Int Psychogeriatr 30, 1707-1716.

Burton, J.K., Craig, L.E., Yong, S.Q., Siddiqi, N., Teale, E.A., Woodhouse, R., Barugh, A.J., Shepherd, A.M., Brunton, A., Freeman, S.C., Sutton, A.J., and Quinn, T.J. (2021). Non-pharmacological interventions for preventing delirium in hospitalised non-ICU patients. Cochrane Database Syst Rev 7, Cd013307.

Cao, Q., Tan, C.C., Xu, W., Hu, H., Cao, X.P., Dong, Q., Tan, L., and Yu, J.T. (2020). The Prevalence of Dementia: A Systematic Review and Meta-Analysis. J Alzheimers Dis 73, 1157-1166.

Chenoweth, L., Williams, A., Fry, M., Endean, E., and Liu, Z. (2021). Outcomes of Person-centered Care for Persons with Dementia in the Acute Care Setting: A Pilot Study. Clin Gerontol, 1-15.

Dewing, J., and Dijk, S. (2016). What is the current state of care for older people with dementia in general hospitals? A literature review. Dementia (London) 15, 106-124.

Dornquast, C.K., Lars E.; Neuhauser, Hannelore K.; Willich, Stefan N.; Reinhold, Thomas; Busch, Markus A. (2016). Regional differences in the prevalence of cardiovascular disease—results from the German Health Update (GEDA) from 2009–2012. Dtsch Arztebl Int 113, 704-711.

Folstein, M.F., Folstein, S.E., and Mchugh, P.R. (1975). "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. J Psychiatr Res 12, 189-198.

Han, Q.Y.C., Rodrigues, N.G., Klainin-Yobas, P., Haugan, G., and Wu, X.V. (2022). Prevalence, Risk Factors, and Impact of Delirium on Hospitalized Older Adults With Dementia: A Systematic Review and Meta-Analysis. J Am Med Dir Assoc 23, 23-32.e27.

Inouye, S.K., Marcantonio, E.R., and Metzger, E.D. (2014a). Doing Damage in Delirium: The Hazards of Antipsychotic Treatment in Elderly Persons. Lancet Psychiatry 1, 312-315.

Inouye, S.K., Westendorp, R.G., and Saczynski, J.S. (2014b). Delirium in elderly people. Lancet 383, 911-922.

Maeck, L., Haak, S., Knoblauch, A., and Stoppe, G. (2007). Early diagnosis of dementia in primary care: a representative eight-year follow-up study in Lower Saxony, Germany. Int J Geriatr Psychiatry 22, 23-31.

Marvanova, M. (2016). Drug-induced cognitive impairment: Effect of cardiovascular agents. Ment Health Clin 6, 201-206.

Mathur, S., Walter, S., Grunwald, I.Q., Helwig, S.A., Lesmeister, M., and Fassbender, K. (2019). Improving Prehospital Stroke Services in Rural and Underserved Settings With Mobile Stroke Units. Front Neurol 10, 159.

Mayring, P., and Fenzl, T. (2014). "Qualitative Inhaltsanalyse," in Handbuch Methoden der empirischen Sozialforschung, eds. N. Baur & J. Blasius. (Wiesbaden: Wiesbaden), 543–556.

Milbert, A., and Sturm, G. (2016). Binnenwanderungen in Deutschland zwischen 1975 und 2013. Informationen zur Raumentwicklung 2, 121-144.

Ní Chróinín, D., Francis, N., Wong, P., Kim, Y.D., Nham, S., and D'amours, S. (2021). Older trauma patients
are at high risk of delirium, especially those with underlying dementia or baseline frailty. Trauma Surg
Acute Care Open 6, e000639.

Park, S.K., Lim, T., Cho, H., Yoon, H.K., Lee, H.J., Lee, J.H., Yoo, S., Kim, J.T., and Kim, W.H. (2021).
Comparative effectiveness of pharmacological interventions to prevent postoperative delirium: a network
meta-analysis. Sci Rep 11, 11922.

Schmidtke, K., and Hermeneit, S. (2008). High rate of conversion to Alzheimer's disease in a cohort of amnestic
MCI patients. Int Psychogeriatr 20, 96-108.

Sica, D.A. (2004). Diuretic-related side effects: development and treatment. J Clin Hypertens (Greenwich) 6,
532-540.

Stoppe, G., Sandholzer, H., Staedt, J., Winter, S., Kiefer, J., Kochen, M.M., and Rüther, E. (1994). Diagnosis of
dementia in primary care: results of a representative survey in lower Saxony, Germany. Eur Arch Psychiatry
Clin Neurosci 244, 278-283.

Van Den Berg, N., Grabe, H.J., Freyberger, H.J., and Hoffmann, W. (2011). A telephone- and text-message
based telemedical care concept for patients with mental health disorders--study protocol for a randomized,
controlled study design. BMC Psychiatry 11, 30.

Wallin, A., Nordlund, A., Jonsson, M., Lind, K., Edman, Å., Göthlin, M., Stålhammar, J., Eckerström, M., Kern,
S., Börjesson-Hanson, A., Carlsson, M., Olsson, E., Zetterberg, H., Blennow, K., Svensson, J., Öhrfelt, A.,
Bjerke, M., Rolstad, S., and Eckerström, C. (2016). The Gothenburg MCI study: Design and distribution
of Alzheimer's disease and subcortical vascular disease diagnoses from baseline to 6-year follow-up. J
Cereb Blood Flow Metab 36, 114-131.

12   Figures

Figure 1: Basic conditions of the consultation service. A: Proportions of consultations performed
on-site or via telemedicine. B: Distribution of the consultation requests among the participating
hospitals. C: Age distribution among the patients for whom consultations were requested. D: Age
distribution, separated by gender.

478
479

480   **Figure 2: Clinical history and indications for consultations.** A: Reasons leading to initial
481   admission of the patients. B: Symptoms reported by requesting institutions. C: Specific indications for
482   consultation. Total numbers sum to >100% as multiple items could be reported for one patient.

483
484

Figure 3: Clinical history related to dementia and risk for delirium. A: Proportions of patients
with recorded history of dementia (left) or cognitive deficits (right), respectively. B: Risk factors for
developing delirium. Total numbers sum to >100% as multiple items could be reported for one patient.

Figure 4: Diagnostic assessment during consultations. A: Primary diagnoses noted by the
consulting physicians. B: Proportions of dementia subtypes among patients diagnosed with any form
of dementia. Medium grey denotes the (suspected) presence of Alzheimer's pathology. Hatch fill
denotes (suspected) vascular pathology. C: MMSE scores among the patients investigated during
consultations. D: Number of pharmacological agents taken by the patients.

Figure 5: Therapeutic recommendations. A: General recommendations. B: Pharmacological
recommendations. Total numbers add to > 100% as multiple recommendations could be given for one
patient.

500
501
502