Projecting the transition of COVID-19 burden towards the young population

while vaccines are rolled out: a modelling study

Jun Cai[1]*, Juan Yang[1,2,3]*, Xiaowei Deng[1], Cheng Peng[1], Xinhua Chen[1], Qianhui

Wu[1], Hengcong Liu[1], Juanjuan Zhang[1,2,3], Wen Zheng[1], Junyi Zou[1], Zeyao Zhao[1],

Marco Ajelli[4,5]†, Hongjie Yu[1,2,3]†

1. Department of Infectious Diseases, Huashan Hospital, School of Public Health, Fudan

University

2. Key Laboratory of Public Health Safety, Ministry of Education, Shanghai, China

3. Shanghai Institute of Infectious Disease and Biosecurity, Fudan University, Shanghai, China

4. Department of Epidemiology and Biostatistics, Indiana University School of Public Health,

Bloomington, IN, USA

5. Laboratory for the Modelling of Biological and Socio-technical Systems, Northeastern

University, Boston, MA, USA

*These authors contributed equally to this work.

†These authors are joint senior authors contributed equally to this work.

Corresponding authors: Hongjie Yu, Shanghai Institute of Infectious Disease and

Biosecurity, School of Public Health, Fudan University, Shanghai 200032, China

E-mail: yhj@fudan.edu.cn

Word count (main text): 3,210

Word count (abstract): 261

Abstract

SARS-CoV-2 infection causes most cases of severe illness and fatality in older age groups. In China, over 85% of individuals aged $\geq$12 years have been vaccinated against COVID-19 (albeit with vaccines developed against historical lineages), while children aged 0–11 years are currently not eligible for vaccination (as of September 2021). The aim of this work is to assess whether the importation of Delta variant infections will shift the COVID-19 burden from adults to children. We developed an age-structured susceptible-infectious-removed model of SARS-CoV-2 transmission dynamics to simulate the epidemics triggered by the importation of Delta variant infections and project the age-specific incidence of SARS-CoV-2 infections, cases, hospitalisations, intensive care unit (ICU) admissions, and deaths. In the context of the vaccination programme targeting individuals aged ≥12 years (as of September 2021), and in the absence of non-pharmaceutical interventions, the importation of Delta variant infections could lead to widespread transmission and substantial disease burden in mainland China, even with vaccination coverage as high as 97% across the currently eligible age groups. The symptomatic SARS-CoV-2 infections and hospitalisation are projected to shift towards children and young adolescents, with 13% of symptomatic infections and 30% of hospitalisations occurring in those aged 0–11 years. Extending the vaccination roll-out to include children aged 3–11 years is estimated to dramatically decrease the burden of symptomatic infections and hospitalisations within this age group (54% and 81%, respectively), but would have a low impact on protecting infants (aged 0–2 years). Our findings highlight the need to strengthen vaccination efforts by simultaneously extending the target population and elevating vaccine effectiveness.

Author summary

Older age groups have the highest risk of severe illness and fatality from SARS-CoV-2 infection. While the vaccination coverage in China is highly skewed towards older age groups. No study has quantified to what extent the spread of Delta variant infections and lack of vaccination in younger age groups will shift the COVID-19 burden towards younger age groups and how this will affect the return to normal. To this end, we developed an age-structured transmission model to mimic the epidemics triggered by imported Delta variant infections in China. In the context of the vaccination programme targeting individuals aged ≥12 years as of September 2021, and in the absence of non-pharmaceutical interventions, the importation of Delta variant infections would lead to substantial disease burden, which will shift towards the 0–11-year age group. Extending the vaccination to children aged 3–11 years, for whom the currently available vaccines have been licensed, is projected to dramatically decrease symptomatic infections and hospitalisations within this age group but would have a low impact in protecting infants aged 0–2 years. This study highlights the value of extending vaccination to children aged 3–11 years and protecting infants 0–2 years by vaccinating their contacts.

Introduction

The coronavirus disease 2019 (COVID-19) pandemic is still raging worldwide. The highly contagious Delta variant (B.1.617.2) of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has become the dominant strain across the world[1],[2]. The global circulation of the Delta variant has led to a resurgence of COVID-19 cases worldwide, including in areas with high vaccination coverage[3, 4].

A clear positive correlation has been shown between severe illness and fatality from SARS-CoV-2 infection and increasing age[5-7]. COVID-19 vaccines are highly efficacious in preventing severe illness[8-10] Due to the age-prioritised vaccination strategies adopted by most countries[11], vaccine coverage tends to be higher in older adults[12]. However, since the summer of 2021, while the Delta variant continued its spread, the number of COVID-19 cases and hospitalisations have trended upwards in many countries with an increasing fraction of children and adolescents[13-16].

China has been able to avoid the widespread local transmission of SARS-CoV-2 since March 2020[17]. However, the importations of Delta variant infections have caused several local outbreaks, the largest of which occurred in Nanjing city and led to over 1,000 reported cases and spilled over to >25 cities[18]. The most recent outbreak started in September 2021 in Putian City, where it spread to a second location, Xiamen City, and caused local transmission. Similar to what was observed in other countries, this outbreak is characterized by a high proportion (57/129) of notified infections in children younger than 12 years[19].

To prevent the spread of infection in younger age groups, in late July 2021, China extended the vaccination programme to include adolescents (aged 12–17 years). As of September 15, 2021, 1.99 billion and 0.17 billion doses were administered to adults (18+ years) and adolescents (12–17 years), respectively, corresponding to 92% and 88% of these two target populations[20]. Two inactivated COVID-19 vaccines have

already obtained emergency use approval for administration in individuals aged ≥3 years. The vaccination roll-out is expected to include children aged 3–11 years at a later stage[21]. No vaccine has yet received approval for use in children aged ≤2 years.

The vaccines used in mainland China were developed using historical SARS-CoV-2 lineages and have proven to be highly effective in protecting against severe illness caused by the Delta variant[22-24]. However, their effectiveness in preventing infection appears to be reduced[25]. Moreover, vaccination coverage is highly skewed towards older age groups. It is thus legit to ask whether the spread of Delta variant infections and lack of vaccination in younger age groups will shift the COVID-19 burden towards younger age groups and how this can affect the return to normal. To fill this gap, we developed an age-structured SARS-CoV-2 transmission model to project age-specific epidemiological outcomes, should an epidemic start to unfold. In particular, the model is used to forward simulate alternative vaccination strategies and allows tracking the number of infections, cases, hospitalisations, intensive care unit (ICU) admissions, and deaths by age.

Methods and methods

SARS-CoV-2 transmission and vaccination model

We developed an age-structured (16 age groups, Table S2) stochastic compartmental susceptible-infectious-removed (SIR) model to simulate the transmission of SARS-CoV-2 (Delta variant) and assess the health impact of age-targeted vaccination campaigns (Fig S1). Detailed information about the model and adopted parameters are described in S1 Text. A summary is presented here.

The model is calibrated to represent the Chinese population and considers the age-mixing patterns quantified prior to the COVID-19 pandemic[26]. Based on contact tracing data in the Hunan province of China[27], children aged under 15 years were

found to have a lower susceptibility to SARS-CoV-2 infection than other age groups, which was confirmed by several independent studies[28] (Table S1). A sensitivity analysis considering homogeneous susceptibility across age groups is presented in Fig S3. Simulations were initiated with 40 imported infections[29] on November 1, 2021 and run forward for 1 year. We considered a basic reproductive number $R_0 = 6$ for the Delta variant[30-32]. It is still unclear whether the Delta variant has a shorter generation time than the original lineage, but a first analysis conducted in Singapore found no significant difference[33], and as such, we set the average generation time of the Delta variant to be 7 days, in line with estimates for the wild type[34].

Vaccination strategy and vaccine effectiveness

We considered two vaccination strategies: (1) vaccination of individuals aged ≥12 years (here referred to as the "adults+adolescents" vaccination strategy), in agreement with the vaccination programme in place in China as of September 2021; (2) as in strategy (1), but the target population is extended to include children aged 3–11 years starting from November 1, 2021–this strategy is referred to here as the "adults+adolescents+children" vaccination strategy. We consider that susceptible individuals are only eligible for vaccination, mimicking the programme implemented in China. Moreover, we accounted for individuals with contraindications against vaccination and pregnant women as they were excluded from the Chinese vaccination programme (Table S2)[35]. We simulated the daily distribution of vaccine doses according to the observed vaccination capacity (S1 Text).

We considered a two-dose vaccine with a 21-day interval between doses. Fourteen days after the second dose, the vaccine efficacy in protecting against an infection caused by the Delta variant was set at 54.3%[35]. The considered vaccine effectiveness (VE) against clinical endpoints such as symptomatic COVID-19, hospitalisations, ICU admissions, and deaths are reported in Table 1. We further explored a range of higher VE values for the sensitivity analyses (Fig S7). We considered VE to be homogeneous across age groups[36-38]. We considered a

"leaky" vaccine in which all vaccinated individuals are exposed to a lower risk of

infection, which is 1-VE times that of non-vaccinated individuals[39]. We assumed

that both the vaccine and natural infection confer protection for longer than the study

period (i.e., 1 year)[40].

Model of COVID-19 burden

The number of infections was produced by the transmission model previously

described. To estimate the COVID-19 burden, we rescaled the number of infections to

obtain estimates of the cumulative incidence of symptomatic cases, hospitalisations,

ICU admissions, and deaths under different vaccination strategies (S1 Text).

Integrating different studies in Scotland[41], Canada[42], and England[43, 44], we

estimated that the increased risk associated with the Delta variant was 182% (97–

317%) for hospitalisation, 287% (98–399%) for ICU admission, and 137% (50–

230%) for death compared to the original lineage, respectively (Table 1). We also

assumed that the age-specific probability of developing symptoms upon Delta

infection is similar to that of the historical lineage. This choice was inspired by the

finding that no significant difference was found for the Alpha variant as compared to

historical lineages[45]. To compare the severity of disease across different age groups,

we calculated the rate ratios as the incidence rate per age group dividing the overall

incidence rate for each health outcome under different vaccination strategies.

Data analysis

For each scenario, 200 stochastic simulations were performed. The outcomes of these

simulations determined the distribution of the cumulative number of symptomatic

infections, hospitalisations, ICU admissions, and deaths by age. We defined 95%

credible intervals as quantiles of the estimated distributions of 0.025 and 0.975.

Table 1. Parameters regulating the transmissibility, vaccine effectiveness/efficacy, and
COVID-19 burden

Parameter description	Estimated value for the SARS-CoV-2 Delta variant
Epidemiology	
Basic reproduction number (R_0)	6[30-32]
Vaccine effectiveness/efficacy	
Against infection (%) (ϵ)	54.3[35]
Against symptomatic disease (%) (ϵ^{symp})	69.5 (42.8, 96.3)[22]
Against hospitalisations (%) (ϵ^{hosp})	87.5 (86.7, 88.2)[8] [a]
Against ICU admissions (%) (ϵ^{icu})	93[22-24] [b]
Against deaths (%) (ϵ^{death})	93[22-24]
Disease burden [c]	
Proportion of infections that develop symptoms for the wild-type (r_a^{symp})	18.1%, 22.4%, 30.5%, 35.5%, and 64.6% for 0–19, 20–39, 40–59, 60–79, and ≥80 years[7]
Proportion of infections that require hospitalisation for the wild-type (r_a^{hosp})	7.24%, 6.54%, 10.16%, 12.00%, and 21.83% separately for 0–19, 20–39, 40–59, 60–79, and ≥80 years[6, 7]
Proportion of infections that require ICU for the wild-type (r_a^{icu})	0.1014%, 0.2747%, 0.4266%, 0.7617%, 0.8999%, 2.8198%, and 5.1312% separately for 0–19, 20–39, 40–49, 50–59, 60–64, 65–79, and ≥80 years[6, 7, 46]
Infection fatality ratio for the wild-type (r_a^{death})	0.0923%, 0.1456%, 0.7259%, 3.7346%, and 6.7959% separately for 0–19, 20–39, 40–59, 60–79, and ≥80 years[6, 7]
Risk ratio of symptomatic infection associated with the Delta variant compared to the wild-type ($\Delta_{WT \rightarrow Delta}^{symp}$)	1[45] [d]
Risk ratio of hospitalisation associated with the Delta variant compared to the wild-type ($\Delta_{WT \rightarrow Delta}^{hosp}$)	2.82 (1.97, 4.17)[41-44]
Risk ratio of ICU admission associated with the Delta variant compared to the wild-type ($\Delta_{WT \rightarrow Delta}^{icu}$)	3.87 (1.98, 4.99)[42]
Risk ratio of death associated with the Delta variant compared to the wild-type ($\Delta_{WT \rightarrow Delta}^{death}$)	2.37 (1.50, 3.30)[42]

[a] Although the main variants of concern detected in Chile during the study period from February 2 to May 1, 2021
were the Alpha and Gamma variants, we assumed that the effectiveness reported in reference [8] applies to the
Delta variant.

[b] Pooling the effectiveness of inactivated COVID-19 vaccines against severe illness caused by the Delta variant
estimated in the recent outbreaks of the Delta variant in Guangdong[22, 23] and Jiangsu[24], China, we assume
conservative effectiveness in preventing serious illness.

[c] The risks of different clinical outcomes associated with the Delta variant are expressed as increased risks
compared to the wild-type.

[d] Based on the conclusion that no significant change in reported symptoms associated with the Alpha variant
compared to wild-type is found in reference[45], we assumed that the probability of developing symptoms by age
for the Delta variant is the same as the wild-type.

Results

Baseline scenario

In the baseline scenario, we assumed that 40 individuals infected with the Delta variant were introduced in the population on November 1, 2021, and the basic reproduction number R_0 was 6 in the absence of interventions and immunity[30-32]. Given the actual vaccination rates, we project that the "adults+adolescents" vaccination strategy could reach a 97% coverage of the target population (83% of the total population) by November 1, 2021 (Fig 1A).

Model simulations suggest that the importation of Delta variant infections in November 2021 would still have the potential to generate a major epidemic wave in the absence of non-pharmaceutical interventions (NPIs). Such an epidemic is estimated to cause 1,693 (95% CI, 1,596–1,738) symptomatic infections, 750 (95% CI, 686–776) hospitalisations, 48 (95% CI, 45–50) ICU admissions, and 38 (95% CI, 35–40) deaths per 10,000 individuals over 1 year (Figs 1C and 2). These figures correspond to 11–21 fold the disease burden of the initial epidemic in Wuhan, China[6]. Under the adopted vaccination programme, 13% of symptomatic infections and 30% of hospitalisations were estimated to occur in children younger than 12 years (Fig S6), who are ineligible for COVID-19 vaccination as of September 2021.

Two inactivated COVID-19 vaccines have been licensed for use in children aged 3–17 years[47, 48]. However, vaccination has not been implemented among children aged 3–11 years as of September 2021. Here, we simulate a scenario where the vaccine is offered to children aged 3–11 years (representing 11.0% of the population) from November 1, 2021 (i.e., the "adults+adolescents+children" vaccination strategy). As compared to the "adults+adolescents" vaccination strategy, this alternative strategy is estimated to reduce the incidence of symptomatic cases, hospitalisations, ICU admissions, and deaths by 6% (95% CI, 3–12%), 20% (95% CI, 17–27%), 8% (95% CI, 4–15%), and 5% (95% CI, 1–13%), respectively (Fig 2). Despite the beneficial

mitigation effect of this strategy, extending vaccination to children is estimated not to

be enough to suppress viral circulation. Indeed, 5.2 million deaths (95% CI, 4.8–5.4)

(approximately 0.3% of the Chinese population) are projected under this vaccination

programme, should NPIs not be implemented (Fig S6).

We used the model to simulate a counterfactual scenario to estimate what would have

happened if a vaccination programme had not been implemented. Compared with this

scenario, the "adults+adolescents" vaccination strategy led to a 47% and 54% increase

in the rate ratios of symptomatic infections in children aged 0–2 and 3–11 years,

respectively. We recall that the rate ratio is defined as the incidence rate per age group

divided by the overall incidence rate[49]. A higher increase is observed in the rate

ratios of hospitalisation for these two population groups: 178% and 191% for children

aged 0–2 and 3–11 years, respectively. At the same time, the rate ratios in adults aged

≥60 years is estimated to decrease by 25%. Similar patterns are observed in the rate

ratios of ICU admissions and deaths (Fig 3). Extending the vaccination to children

aged 3–11 years is projected to dramatically decrease the burden of symptomatic

infections and hospitalisations within the same age group by 54% and 81%,

respectively (Fig S6). However, due to the strong age-assortativity of contact patterns

of this age group (i.e., individuals aged 3–11 years primarily mix with other

individuals of the same age) (Fig S4A), extending the vaccination to children does not

strongly impact the COVID-19 burden in other age groups (Fig S6). No evident effect

is projected on the 0–2-year age group (Fig 3 and S6).

Under either the "adults+adolescents" or "adults+adolescents+children" vaccination

strategies, the mean incidence of symptomatic cases among unvaccinated individuals

is estimated to be 1.5–2.0 fold that of vaccinated individuals, with larger differences

observed in terms of the incidence of severe clinical outcomes (3.4–5.0 fold for

hospitalisation, 5.5–8.8 fold for ICU admission, and 4.1–8.8 fold for death) (Fig 2).

Improving VE, in the absence/presence of NPIs

In our model, VE in preventing ICU admissions and deaths caused by a Delta variant infection was set at 93%, based on real-world VE studies in Guangdong and Jiangsu, China (Table 1)[22-24]. Should the VE against the infection of the Delta variant be increased to >84% from 54.3% used in the baseline[35], the "adults+adolescents+children" vaccination strategy is estimated to reduce the number of deaths by 99% even in the absence of any NPIs, leading to <40,000 deaths over 1 year, similar to the annual excess respiratory disease deaths associated with influenza in China[50] (Fig S7).

We leveraged the developed model to explore the impact of implementing different levels of NPIs on the COVID-19 burden while the "adults+adolescents+children" vaccination strategy is in place and considering different levels of VE against infection. When VE against infection is set at 54.3% (baseline scenario), stringent NPIs capable of maintaining the net reproduction number R_e<2.5 are needed, in combination with the vaccination programme, to reduce the yearly COVID-19 death toll to a level similar to that of seasonal influenza[50] (<50,000 deaths). Should a new vaccine with higher (>75%) VE against the infection be adopted, NPIs could be relaxed. Moderate NPIs able to keep R_e<4.3 in combination with the vaccination programme is estimated to be sufficient to decrease the annual number of deaths to less than 40,000 (Fig S7).

Sensitivity analysis on susceptibility to infection and contact patterns by age

In the baseline analysis, age-varying susceptibility to SARS-CoV-2 infection was considered. That is, using adults aged 15–64 years as a reference group, children have a 42% lower risk of infection than adults (Table S1)[27]. No estimates are available for susceptibility to Delta variant infections by age. As such, we performed an analysis assuming the same susceptibility to infection across all age groups. The obtained results are consistent with those obtained in the baseline analysis, with average variations lower than 12% (Fig S3).

Another feature able to shape the epidemiology of COVID-19 is the contact pattern of

the population. In the baseline analysis, we used pre-pandemic mixing patterns to

represent a situation close to the objective of returning to a pre-COVID-19 pandemic

lifestyle[26] (Fig S4A). However, whether mixing patterns will ever return to be the

same after the pandemic remains to be seen. As such, we tested the robustness of our

findings by considering an alternative contact matrix estimated in Shanghai, China,

right after the end of the lockdown in March 2020[51] (Fig S4B). The obtained results

are consistent with those obtained with the pre-pandemic contact matrix, with mean

variations in the estimated burden lower than 6% (Fig S5).

Discussion

Our modelling study indicates that, under a vaccination programme targeting

individuals aged ≥12 years, symptomatic SARS-CoV-2 infections and hospitalisation

would shift towards children and young adolescents in the case of a new COVID-19

wave caused by the Delta variant. These modelling results obtained for China are

backed up by empirical evidence from other countries, where an upsurge of COVID-

19 cases and hospitalisations among children and adolescents has been reported since

the summer of 2021[13, 14, 49]. Evidence of this epidemiological shift towards

children and adolescents comes from the analyses of local outbreaks in Putian city,

China, in September 2021, where 44.2% of the reported cases occurred in those aged

0–11 years[19].

The numbers depicted by the model are related to the relatively low vaccine efficacy

of the vaccines in use in China against Delta variant infection (54.3%[35]) as of

September 2021. Model projections show that, in the absence of NPIs, even when

97% of the population aged ≥12 years is fully vaccinated, the importation of Delta

variant infections could lead to a new COVID-19 wave and substantial health burden.

In particular, the estimated number of deaths over 1 year was 38 per 10,000

individuals as compared, for instance, with 3.6 deaths per 10,000 individuals in the

first epidemic wave in Wuhan in early 2020[6]. Extending vaccination to children

would mitigate the number of symptomatic cases and severe outcomes but would not

be enough to suppress transmission. Relying on NPIs such as border control

screenings, border quarantine, isolation, and mask wearing remains key in this phase

of widespread circulation of highly transmissible SARS-CoV-2 variants across the

globe. This finding is consistent with studies in France, the UK, and the US where,

despite the distribution of more effective vaccines (e.g., mRNA vaccines and

adenoviral vector vaccines), the transmission is far from being suppressed as of

September 2021[52-54].

To return to normal (i.e., no NPIs), the combination of high vaccine coverage,

including children and adolescents, and a highly efficacious vaccine that prevents

infection appear to be key to reducing COVID-19 burden to levels closer to those of

seasonal influenza. To reach this target, our modelling results suggest that the

vaccination programme would need to be extended to children aged 3–11 years, and

vaccine efficacy against the infection would need to be >84%, in line with the efficacy

estimated for the mRNA-1273 (Moderna) vaccine[9]. China is currently attempting to

develop mRNA vaccines[55], vaccines that specifically target the Delta variant[56],

multi-valent vaccines, and universal vaccines[57-59]. However, whether such

vaccines will become available in the short term is unknown. Moreover, the waning of

vaccine-induced immunity may also worsen the situation[54, 60-62]. For this reason,

several countries recommend booster shots, at least for certain high-risk groups[63,

64]. The debate on whether to prioritise the administration of booster doses to older

age groups or extend the vaccination programme to those aged 3–11 years merits

further modelling investigations. Currently, the best way to protect infants (aged 0–2

years) appears to be by vaccinating their contacts, such as family members.

Although the inactivated vaccines in use in China as of September 2021 show a

relatively low effectiveness against mild and moderate disease caused by Delta variant

infection, they appear to be highly effective against severe outcomes[8, 22, 23]. In the

context of a constantly changing virus and the evolving epidemiology of the

pandemic, it is of great importance to continue to ramp up efforts to increase

vaccination coverage by using currently available vaccines. Promoting equitable

access to COVID-19 vaccines is critical to decreasing viral circulation and the

likelihood of the emergence of new variants of concern.

Our findings should be interpreted by considering the following limitations. First, we

assumed that vaccine protection lasted longer than the time of our simulations (1

year). Recent immunological studies have shown that the neutralising antibody titres

of inactivated vaccines decline to a low level at 6 months after the second vaccine

dose[60, 61], which may indicate waning of vaccine efficacy. However, the

correlation with protection from SARS-CoV-2 infection has yet to be proven. Second,

we estimated the disease burden potentially caused by the importation of Delta variant

infections in the absence or presence of NPIs. The impact of NPIs has been modelled

through a simple reduction in transmissibility occurring homogenously with age. We

still lack evidence regarding whether this is the case or whether NPIs lead to age-

related effects. Moreover, our analysis is not suited to pinpoint which NPIs need to be

performed to reach the considered levels of transmission reduction. Further studies

exploring this direction are required.

In conclusion, in the absence of NPIs, the importation of Delta variant infections

could lead to widespread transmission and substantial disease burden in mainland

China, even with vaccination coverage as high as 97% of those aged ≥12 years. Our

findings highlight the need to strengthen vaccination efforts by simultaneously

extending the target age groups of the vaccination campaign, and elevating VE via

booster vaccination and developing new highly efficacious vaccines. Although

vaccination is key to dramatically reducing severe outcomes of SARS-CoV-2

infections and the overall burden of the COVID-19 pandemic, vaccination

programmes targeting individuals aged ≥12 years can potentially shift the COVID-19

burden towards younger ages. This highlights the importance of extending

vaccination to children aged 3–11 years and protecting infants by vaccinating their

contacts.

Supporting information

S1 Supporting information. Extended details on methods and additional analyses.

Author Contributions

Conceptualization: Hongjie Yu.

Data curation: Jun Cai, Juan Yang, Xiaowei Deng, Xinhua Chen, Qianhui Wu, Hengcong Liu, Juanjuan Zhang, Wen Zheng, Junyi Zou, Zeyao Zhao.

Formal analysis: Jun Cai, Juan Yang, Xiaowei Deng, Cheng Peng.

Funding acquisition: Hongjie Yu.

Investigation: Jun Cai, Juan Yang, Xiaowei Deng, Peng Cheng, Marco Ajelli, Hongjie Yu.

Methodology: Jun Cai, Xiaowei Deng.

Software: Jun Cai, Xiaowei Deng.

Supervision: Hongjie Yu.

Validation: Jun Cai, Xiaowei Deng, Hengcong Liu.

Visualization: Jun Cai, Xiaowei Deng, Cheng Peng, Juan Yang.

Writing-original draft: Jun Cai, Juan Yang.

Writing-review & editing: Marco Ajelli, Hongjie Yu.

Competing interests

H.Y. received research funding from Sanofi Pasteur, GlaxoSmithKline, Yichang HEC Changjiang Pharmaceutical Company, Shanghai Roche Pharmaceutical Company, and SINOVAC Biotech Ltd. M.A. received research funding from Seqirus. Except for research funding from SINOVAC Biotech Ltd, which is related to the data analysis of clinical trials of immunogenicity and safety of CoronaVac, the others are not related to COVID-19. All the other authors have no competing interests.

Data Availability Statement

The data and code that support the findings of this study will be made available on

GitHub upon the acceptance of this manuscript.

Funding

This study was supported by grants from the Key Program of the National Natural

Science Foundation of China (82130093) and the National Institute for Health

Research (NIHR) (grant no. 16/137/109) using UK aid from the UK Government to

support global health research. The views expressed in this publication are those of

the author(s) and not necessarily those of the NIHR or the UK Department of Health

and Social Care.

Figures

Fig 1. Time series of vaccine coverage and daily incidence of new SARS-CoV-2

Delta variant infections. A Age-specific vaccine coverage over time for the

"adults+adolescents" vaccination strategy. The vaccination programme was initiated

on November 30, 2020 (as first officially reported in China). The vertical dotted lines

represent the (simulated) seeding of the infection (November 1, 2021). The subpanel

reports a table showing the age-specific coverage. The line corresponds to the mean

value, while the shaded area represents the 95% quantile intervals (CI). **B** As A, but

for the "adults+adolescents+children" vaccination strategy. **C** Simulated daily

incidence of new SARS-CoV-2 Delta variant infections per 10,000 individuals for the

two strategies (mean and 95% CI).

Fig 2. Disease burden due to an epidemic caused by the introduction of SARS-CoV-2 Delta variant infections in China. A Cumulative number of symptomatic cases per 10,000 individuals after one simulated year by vaccination strategy (AA = "adults+adolescents" vaccination strategy, AAC = "adults+adolescents+children" vaccination strategy), vaccination status, and age group. The vaccinated group are those individuals who are administrated with two doses. **B** As A, but for the incidence of hospitalisations. **C** As A, but for the incidence of ICU admissions. **D** As A, but for mortality. The horizontal dotted lines in **A**, **B**, and **D** represent the rates of symptomatic cases, hospitalisations, and deaths, respectively, of the first pandemic wave of COVID-19 in Wuhan, China[6].

441 **Fig 3. Rate ratios of symptomatic cases, hospitalisations, ICU admissions, and**

442 **deaths due to SARS-CoV-2 Delta variant infections in China by age group and**

443 **vaccination strategy.** NV = no vaccination, AA = "adults+adolescents" vaccination

444 strategy, AAC = "adults+adolescents+children" vaccination strategy.

References

1.	WHO Team Emergency Respinse. Weekly epidemiological update on COVID‑19 - 10 August 2021 2021 [updated 2021-08-10; cited 2021 2021-08-12]. Available from: https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---10-august-2021.

2.	CDC. Delta Variant: What We Know About the Science 2021 [updated 2021-08-19; cited 2021 2021-08-20]. Available from: https://www.cdc.gov/coronavirus/2019-ncov/variants/delta-variant.html.

3.	WHO. WHO Coronavirus (COVID-19) Dashboard 2021 [updated 2021-09-07; cited 2021 2021-09-07]. Available from: https://covid19.who.int/.

4.	Our World in Data. Coronavirus (COVID-19) Vaccinations 2021 [updated 2021-09-06; cited 2021 2021-09-07]. Available from: https://covid19.who.int/.

5.	O'Driscoll M, Ribeiro Dos Santos G, Wang L, Cummings DAT, Azman AS, Paireau J, et al. Age-specific mortality and immunity patterns of SARS-CoV-2. Nature. 2021;590(7844):140-5. doi: 10.1038/s41586-020-2918-0.

6.	Yang J, Chen X, Deng X, Chen Z, Gong H, Yan H, et al. Disease burden and clinical severity of the first pandemic wave of COVID-19 in Wuhan, China. Nature Communications. 2020;11(1):5411. doi: 10.1038/s41467-020-19238-2.

7.	Poletti P, Tirani M, Cereda D, Trentini F, Guzzetta G, Sabatino G, et al. Association of Age With Likelihood of Developing Symptoms and Critical Disease Among Close Contacts Exposed to Patients With Confirmed SARS-CoV-2 Infection in Italy. JAMA Network Open. 2021;4(3):e211085-e. doi: 10.1001/jamanetworkopen.2021.1085.

8.	Jara A, Undurraga EA, González C, Paredes F, Fontecilla T, Jara G, et al. Effectiveness of an Inactivated SARS-CoV-2 Vaccine in Chile. New England Journal of Medicine. 2021. doi: 10.1056/NEJMoa2107715.

9.	Tang P, Hasan MR, Chemaitelly H, Yassine HM, Benslimane F, Al Khatib HA, et al. BNT162b2 and mRNA-1273 COVID-19 vaccine effectiveness against the Delta (B.1.617.2) variant in Qatar. medRxiv. 2021:2021.08.11.21261885. doi: 10.1101/2021.08.11.21261885.

10.	Chemaitelly H, Yassine HM, Benslimane FM, Al Khatib HA, Tang P, Hasan MR, et al. mRNA-1273 COVID-19 vaccine effectiveness against the B.1.1.7 and B.1.351 variants and severe COVID-19 disease in Qatar. Nature Medicine. 2021. doi: 10.1038/s41591-021-01446-y.

11.	Noh EB, Nam H-K, Lee H. Which Group Should be Vaccinated First?: A Systematic Review. Infect Chemother. 2021;53(2):261-70.

12.	CDC. Demographic Characteristics of People Receiving COVID-19 Vaccinations in the United States 2021 [updated 2021-09-04; cited 2021 2021-09-07]. Available from: https://covid.cdc.gov/covid-data-tracker/#vaccination-demographic.

13.	Siegel DA. Trends in COVID-19 Cases, Emergency Department Visits, and Hospital Admissions Among Children and Adolescents Aged 0–17 Years—United States, August 2020–August 2021. MMWR Morbidity and Mortality Weekly Report. 2021;70.

14.	Delahoy MJ. Hospitalizations Associated with COVID-19 Among Children and Adolescents—COVID-NET, 14 States, March 1, 2020–August 14, 2021. MMWR Morbidity and Mortality Weekly Report. 2021;70.

15.	Office for National Statistics of the UK. Coronavirus (COVID-19) latest insights 2021 [updated 2021-09-16; cited 2021 2021-09-17]. Available from:

https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19latestinsights/Overview.

16. Ian Mount, Bernhard Warner, Dunn K. The kids are (not) alright: Europe sounds the alarm as Delta variant soars among teens and 20-somethings 2021 [updated 2021-07-08; cited 2021 2021-09-17]. Available from: https://fortune.com/2021/07/08/kids-vulnerable-covid-delta-variant-vaccinated-europe/.

17. National Health Commission. Notification on situation of COVID-19 epidemic in China 2021 [updated 2021-09-17; cited 2021 2021-09-17]. Available from: http://www.nhc.gov.cn/xcs/yqtb/list_gzbd.shtml.

18. Xinhua Net. China goes all out to contain Delta outbreak 2021 [updated 2021-08-02; cited 2021 2021-08-22]. Available from: http://www.xinhuanet.com/english/2021-08/02/c_1310103377.htm.

19. News Office of Putian City. The 19th press conference on epidemic prevention and control held in Putian 2021 [updated 2021-09-16; cited 2021 2021-09-17]. Available from: http://www.putian.gov.cn/zwgk/xwfbhs/202109/t20210916_1647826.htm.

20. The State Council Information Office PRC. Press conference held on situation regarding prevetion and control of the COVID-19 epidemic and vaccination 2021 [updated 2021-09-16; cited 2021 2021-09-17]. Available from: http://www.nhc.gov.cn/xcs/s3574/202109/f89d117481b24572a346c19fde76a396.shtml.

21. The State Council Information Office PRC. Press conference held on situation regarding prevetion and control of the COVID-19 epidemic and vaccination 2021 [updated 2021-08-27; cited 2021 2021-08-27]. Available from: http://www.nhc.gov.cn/xcs/fkdt/202108/f211dd79672642b7908ffb7575d020a1.shtml.

22. Kang M, Yi Y, Li Y, Sun L, Deng A, Hu T, et al. Effectiveness of Inactivated COVID-19 Vaccines Against COVID-19 Pneumonia and Severe Illness Caused by the B. 1.617. 2 (Delta) Variant: Evidence from an Outbreak in Guangdong, China. SSRN. 2021.

23. Li X-n, Huang Y, Wang W, Jing Q-l, Zhang C-h, Qin P-z, et al. Efficacy of inactivated SARS-CoV-2 vaccines against the Delta variant infection in Guangzhou: A test-negative case-control real-world study. Emerging Microbes & Infections. 2021:1-32. doi: 10.1080/22221751.2021.1969291.

24. Wang J, Huang P, Yi Y, Zhu M, Li J, Yi C, et al. Effectiveness of inactive COVID-19 vaccines against severe illness in B.1.617.2 (Delta) variant-infected patients in Jiangsu, China. medRxiv. 2021:2021.09.02.21263010. doi: 10.1101/2021.09.02.21263010.

25. Chen X, Wang W, Chen X, Wu Q, Sun R, Ge S, et al. Prediction of long-term kinetics of vaccine-elicited neutralizing antibody and time-varying vaccine-specific efficacy against the SARS-CoV-2 Delta variant by clinical endpoint. medRxiv. 2021:2021.09.23.21263715. doi: 10.1101/2021.09.23.21263715.

26. Zhang J, Klepac P, Read JM, Rosello A, Wang X, Lai S, et al. Patterns of human social contact and contact with animals in Shanghai, China. Scientific Reports. 2019;9(1):15141. doi: 10.1038/s41598-019-51609-8.

27. Hu S, Wang W, Wang Y, Litvinova M, Luo K, Ren L, et al. Infectivity, susceptibility, and risk factors associated with SARS-CoV-2 transmission under intensive contact tracing in Hunan, China. Nature Communications. 2021;12(1):1533. doi: 10.1038/s41467-021-21710-6.

28. Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward J, Hudson L, et al. Susceptibility

to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic Review and Meta-analysis. JAMA Pediatrics. 2021;175(2):143-56. doi: 10.1001/jamapediatrics.2020.4573.

29. Yang J, Marziano V, Deng X, Guzzetta G, Zhang J, Trentini F, et al. Despite vaccination, China needs non-pharmaceutical interventions to prevent widespread outbreaks of COVID-19 in 2021. Nature Human Behaviour. 2021;5(8):1009-20. doi: 10.1038/s41562-021-01155-z.

30. Scientific Advisory Group for Emergencies. SPI-M-O: Consensus Statement on COVID-19, 12 May 2021 2021 [updated 2021-05-14; cited 2021 2021-06-11]. Available from: https://www.gov.uk/government/publications/spi-m-o-consensus-statement-on-covid-19-12-may-2021.

31. Dagpunar J. Interim estimates of increased transmissibility, growth rate, and reproduction number of the Covid-19 B.1.617.2 variant of concern in the United Kingdom. medRxiv. 2021:2021.06.03.21258293. doi: 10.1101/2021.06.03.21258293.

32. Challen R, Dyson L, Overton CE, Guzman-Rincon LM, Hill EM, Stage HB, et al. Early epidemiological signatures of novel SARS-CoV-2 variants: establishment of B.1.617.2 in England. medRxiv. 2021:2021.06.05.21258365. doi: 10.1101/2021.06.05.21258365.

33. Pung R, Mak TM, Kucharski AJ, Lee VJ. Serial intervals in SARS-CoV-2 B.1.617.2 variant cases. The Lancet. 2021. doi: 10.1016/S0140-6736(21)01697-4.

34. Sun K, Wang W, Gao L, Wang Y, Luo K, Ren L, et al. Transmission heterogeneities, kinetics, and controllability of SARS-CoV-2. Science. 2020:eabe2424. doi: 10.1126/science.abe2424.

35. Liu H, Zhang J, Cai J, Deng X, Peng C, Chen X, et al. Herd immunity induced by COVID-19 vaccination programs to suppress epidemics caused by SARS-CoV-2 wild type and variants in China. medRxiv. 2021:2021.07.23.21261013. doi: 10.1101/2021.07.23.21261013.

36. Al Kaabi N, Zhang Y, Xia S, Yang Y, Al Qahtani MM, Abdulrazzaq N, et al. Effect of 2 Inactivated SARS-CoV-2 Vaccines on Symptomatic COVID-19 Infection in Adults: A Randomized Clinical Trial. JAMA. 2021. doi: 10.1001/jama.2021.8565.

37. Palacios R, Batista AP, Albuquerque CSN, Patiño EG, Santos JdP, Tilli Reis Pessoa Conde M, et al. Efficacy and safety of a COVID-19 inactivated vaccine in healthcare professionals in Brazil: the PROFISCOV study. 2021.

38. Han B, Song Y, Li C, Yang W, Ma Q, Jiang Z, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial. The Lancet Infectious Diseases. 2021. doi: 10.1016/S1473-3099(21)00319-4.

39. Halloran ME, Longini IM, Struchiner CJ, Longini IM. Design and analysis of vaccine studies: Springer; 2010.

40. Wang Z, Muecksch F, Schaefer-Babajew D, Finkin S, Viant C, Gaebler C, et al. Naturally enhanced neutralizing breadth against SARS-CoV-2 one year after infection. Nature. 2021;595(7867):426-31. doi: 10.1038/s41586-021-03696-9.

41. Sheikh A, McMenamin J, Taylor B, Robertson C. SARS-CoV-2 Delta VOC in Scotland: demographics, risk of hospital admission, and vaccine effectiveness. The Lancet. 2021. doi: 10.1016/S0140-6736(21)01358-1.

42. Fisman DN, Tuite AR. Progressive Increase in Virulence of Novel SARS-CoV-2 Variants in Ontario, Canada. medRxiv. 2021:2021.07.05.21260050. doi: 10.1101/2021.07.05.21260050.

43. Nyberg T, Twohig KA, Harris RJ, Seaman SR, Flannagan J, Allen H, et al. Risk of hospital

admission for patients with SARS-CoV-2 variant B.1.1.7: cohort analysis. BMJ. 2021;373:n1412. doi: 10.1136/bmj.n1412.

44. Twohig KA, Nyberg T, Zaidi A, Thelwall S, Sinnathamby MA, Aliabadi S, et al. Hospital admission and emergency care attendance risk for SARS-CoV-2 delta (B.1.617.2) compared with alpha (B.1.1.7) variants of concern: a cohort study. The Lancet Infectious Diseases. 2021. doi: 10.1016/S1473-3099(21)00475-8.

45. Graham MS, Sudre CH, May A, Antonelli M, Murray B, Varsavsky T, et al. Changes in symptomatology, reinfection, and transmissibility associated with the SARS-CoV-2 variant B.1.1.7: an ecological study. The Lancet Public Health. 2021;6(5):e335-e45. doi: 10.1016/S2468-2667(21)00055-4.

46. Moghadas SM, Shoukat A, Fitzpatrick MC, Wells CR, Sah P, Pandey A, et al. Projecting hospital utilization during the COVID-19 outbreaks in the United States. Proceedings of the National Academy of Sciences. 2020;117(16):9122. doi: 10.1073/pnas.2004064117.

47. Global Times. China approves emergency use of Sinovac's COVID-19 vaccine for children aged 3-17 2021 [updated 2021-06-05; cited 2021 2021-08-22]. Available from: https://www.globaltimes.cn/page/202106/1225430.shtml.

48. Global Times. China approves emergency use of Sinopharm COVID-19 vaccine for children aged 3-17 2021 [updated 2021-08-18; cited 2021 2021-08-22]. Available from: https://www.globaltimes.cn/page/202108/1231842.shtml.

49. Virginia Department of Health. COVID-19 Attack Rates by Vaccination Status and Age 2021 [updated 2021-08-23; cited 2021 2021-09-07]. Available from: https://www.vdh.virginia.gov/coronavirus/2021/08/23/covid-19-attack-rates-by-vaccination-status-and-age/.

50. Li L, Liu Y, Wu P, Peng Z, Wang X, Chen T, et al. Influenza-associated excess respiratory mortality in China, 2010–15: a population-based study. The Lancet Public Health. 2019;4(9):e473-e81. doi: https://doi.org/10.1016/S2468-2667(19)30163-X.

51. Zhang J, Litvinova M, Liang Y, Zheng W, Shi H, Vespignani A, et al. The impact of relaxing interventions on human contact patterns and SARS-CoV-2 transmission in China. Science Advances. 2021;7(19):eabe2584. doi: 10.1126/sciadv.abe2584.

52. Bosetti P, Kiem CT, Andronico A, Colizza V, Yazdanpanah Y, Fontanet A, et al. Epidemiology and control of SARS-CoV-2 epidemics in partially vaccinated populations: a modeling study applied to France. HAL. 2021.

53. Dyson L, Hill EM, Moore S, Curran-Sebastian J, Tildesley MJ, Lythgoe KA, et al. Possible future waves of SARS-CoV-2 infection generated by variants of concern with a range of characteristics. medRxiv. 2021:2021.06.07.21258476. doi: 10.1101/2021.06.07.21258476.

54. Rosenberg ES, Holtgrave DR, Dorabawila V, Conroy M, Greene D, Lutterloh E, et al. New COVID-19 Cases and Hospitalizations Among Adults, by Vaccination Status — New York, May 3–July 25, 2021. MMWR. 2021. Epub 2021-08-18. doi: http://dx.doi.org/10.15585/mmwr.mm7034e1.

55. Walvax Biotechnology Co. L. A Phase III Clinical Study of a SARS-CoV-2 Messenger Ribonucleic Acid (mRNA) Vaccine Candidate Against COVID-19 in Population Aged 18 Years and Above 2021 [updated 2021-05-17; cited 2021 2021-08-22]. Available from: https://clinicaltrials.gov/ct2/show/NCT04847102.

56. Global Times. Chinese producers quickly progressing on Delta variant vaccines, ready for clinical trials 2021 [updated 2021-08-06; cited 2021 2021-08-25]. Available from:

https://www.globaltimes.cn/page/202108/1230741.shtml.

57. He C, Yang J, He X, Hong W, Lei H, Chen Z, et al. A bivalent recombinant vaccine targeting
the S1 protein induces neutralizing antibodies against both SARS-CoV-2 variants and wild-type
of the virus. MedComm. 2021;n/a(n/a). doi: https://doi.org/10.1002/mco2.72.

58. Wang J. A bivalent recombinant vaccine: a promising strategy against both SARS-CoV-2
variants and wild type of the virus. Signal Transduction and Targeted Therapy. 2021;6(1):278. doi:
10.1038/s41392-021-00691-4.

59. Tan C-W, Chia W-N, Young BE, Zhu F, Lim B-L, Sia W-R, et al. Pan-Sarbecovirus Neutralizing
Antibodies in BNT162b2-Immunized SARS-CoV-1 Survivors. New England Journal of Medicine.
2021. doi: 10.1056/NEJMoa2108453.

60. Pan H, Wu Q, Zeng G, Yang J, Jiang D, Deng X, et al. Immunogenicity and safety of a third
dose, and immune persistence of CoronaVac vaccine in healthy adults aged 18-59 years: interim
results from a double-blind, randomized, placebo-controlled phase 2 clinical trial. medRxiv.
2021:2021.07.23.21261026. doi: 10.1101/2021.07.23.21261026.

61. Li M, Yang J, Wang L, Wu Q, Wu Z, Zheng W, et al. A booster dose is immunogenic and will
be needed for older adults who have completed two doses vaccination with CoronaVac: a
randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. medRxiv.
2021:2021.08.03.21261544. doi: 10.1101/2021.08.03.21261544.

62. Thomas SJ, Moreira ED, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Six Month Safety
and Efficacy of the BNT162b2 mRNA COVID-19 Vaccine. medRxiv. 2021:2021.07.28.21261159. doi:
10.1101/2021.07.28.21261159.

63. Ministry of Health Israel. The Ministry of Health Director General Has Approved the
Recommendation to Administer a Third Vaccine Doe to 50-Year-Olds and Older and to Other
Populations 2021 [updated 2019-08-13; cited 2021 2021-08-22]. Available from:
https://www.gov.il/en/departments/news/13082021-01.

64. U.S. Food and Drug Administration. Joint Statement from HHS Public Health and Medical
Experts on COVID-19 Booster Shots 2021 [updated 2019-08-18; cited 2021 2021-08-22].
Available from: https://www.fda.gov/news-events/press-announcements/joint-statement-hhs-
public-health-and-medical-experts-covid-19-booster-shots.

650