Using self-controlled case series to understand the relationship between conflict and

cholera in Nigeria and the Democratic Republic of Congo

Gina E C Charnley[1,2]*, Kévin Jean[3] , Ilan Kelman[4,5,6] , Katy A M Gaythorpe[1,2]**, Kris A

Murray[1,2,7]**

1. Department of Infectious Disease Epidemiology, School of Public Health, Imperial College

London, London, UK

2. MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, UK.

3. Maître de conférences, Laboratoire MESuRS - Cnam Paris, Paris, France.

4. Institute for Risk and Disaster Reduction, Faculty of Mathematical and Physical Sciences,

University College London, London, UK

5. Institute for Global Health, Faculty of Population Health, University College London, London, UK

6. University of Agder, Kristiansand, Norway.

7. MRC Unit The Gambia at London School of Hygiene and Tropical Medicine, Fajara, The

Gambia.

* Corresponding author

** Joint senior author

Email: g.charnley19@imperial.ac.uk

Abstract

Background Cholera outbreaks contribute significantly to diarrhoeal disease mortality, especially in low-income countries. Cholera outbreaks have several social and environmental risk factors and extreme conditions can act as catalysts for outbreaks. A social extreme with known links to infectious disease outbreaks is conflict, causing disruption to services, loss of income and displacement.

Methods Here, we explored this relationship in Nigeria and the Democratic Republic of Congo (DRC), by fitting publicly available cholera and conflict data to conditional logistic regression models. We used the self-controlled case series method in a novel application, to understand if an exposure period of excess risk (conflict), increased the relative incidence of cholera. We also adapted the percentage attributable fraction equations to understand the proportion of cholera outbreaks attributable to conflict onset.

Results There were 12,829 conflict events and 1,178 cholera outbreaks during the period analysed in Nigeria and the DRC. Conflict increased the risk of cholera in Nigeria by 3.6 times and 2.6 times for the DRC. Conflict was attributed to 19.7% and 12.3% of cholera outbreaks in Nigeria and the DRC, respectively. This was higher for some states/provinces, with a maximum increased risk of 7.5 times. Several states/provinces with the strongest positive relationship were also areas of high reported conflict or were neighbouring states/provinces.

Conclusion Our results help highlight the importance of rapid and sufficient assistance during social extremes and the need for pre-existing vulnerabilities such as poverty and access to healthcare to be addressed. In fragile states, conflict resolution should be a top priority to avoid excess risk for both cholera and other health and social implications.

Introduction

Diarrhoeal diseases are the eighth leading cause of death worldwide, with cholera contributing significantly, especially in low- and middle-income countries [1]. Over 94% of World Health Organization (WHO) reported cholera cases are in Africa and more research is needed to understand cholera dynamics on the continent [2]. Previous research has found several

environmental and socioeconomic links with cholera, including temperature, precipitation, poverty

and water, sanitation, and hygiene (WASH) [3-5]. Furthermore, extremes of these environmental

and social conditions can act as catalysts for outbreaks, such as droughts, floods, and conflicts [6-

8].

We will focus on the impacts of conflict on cholera outbreaks and compare the results across two

countries in Africa, Nigeria and the Democratic Republic of Congo (DRC). Several mechanisms

have been suggested through which conflict can lead to infectious disease outbreaks [9-12].

During conflicts, services can be disrupted including access to WASH, disruption of disease control

programmes and collapse of health systems (e.g., vaccination coverage). Those displaced by

conflict may also find it difficult to access healthcare [13-15]. Populations may not seek medical

treatment as they perceive it or healthcare facilities to be unsafe. For example, during the 2018

Ebola outbreak in the DRC healthcare facilities were attacked, dampening efforts to control the

virus [16]. Conflict can worsen pre-existing vulnerabilities including poverty, as conflicts can cause

loss of income, disruption to education, damage to livelihoods and displacement [17].

Nigeria and the DRC share social and environmental similarities, as well as experiencing cholera

outbreaks. Both have active conflicts including the Boko Haram Insurgency in northeastern Nigeria

[18] and political unrest in the eastern DRC [19]. They also have the second and third highest

numbers of estimated cholera cases per year in Africa, respectively [20], with the Kivu provinces

being the most active cholera foci in the world [21]. In addition, Nigeria and the DRC have a

tropical climate, poor access to WASH and a large proportion of the population living in poverty

(<$1.25/day) at 87.7% for the DRC and 62% for Nigeria [22], which are known cholera risk factors.

Few studies have investigated the impacts of conflict on cholera outbreaks, especially

quantitatively. Studies have commonly focused on cholera and conflict in Yemen [10,23-24], its

effect on vaccination efforts [25] or the impact of conflict on other diseases such as Ebola [16] and

COVID-19 [26]. Africa is also a chronically understudied continent in relation to cholera, despite

reporting a large proportion of global cases [2].

To bridge this research gap, we aimed to investigate if the incidence of cholera outbreak onset was

affected by conflicts in Nigeria and the DRC in the past 23 years. To test this, we used the Self-

Controlled Case Series (SCCS) method in a novel application and to understand its use in other

contexts [27]. The SCCS methods have previously been used to test the effectiveness of drug and

vaccine intervention on an individual [28-29] and population level [30]. Furthermore, we adapted

the recently developed percentage attributable fraction (PAF) equations to the work presented here

[30], to understand the proportion of cholera outbreaks attributable to conflict.

A sensitivity analysis provided further insight into the duration of the conflict effect on outbreaks

and a sub-national analysis highlighted priority areas for national cholera prevention. Based on

these results, we suggest mechanisms for which conflict is driving cholera and potential risk

factors, building on previous research in this area. We hope this information can be used to

strengthen disease prevention in conflict settings and reduce additional mortality and morbidity in

conflicts.

Methods

Datasets

Cholera data were compiled from a range of publicly available sources (WHO disease outbreak

news, ProMED, ReliefWeb, WHO regional office for Africa weekly outbreak and emergencies,

UNICEF cholera platform, EM-DAT, the Nigerian Centre for Disease Control, and a literature

search) in both English and French. The full compiled dataset is available in a GitHub repository

(https://github.com/GinaCharnley/cholera_data_drc_nga). An outbreak was defined by the onset of

a cholera case and the case definitions for the two countries are shown in S1 Information. Conflict

data were provided by the United Nations Office for the Coordination of Humanitarian Affairs

Humanitarian Data Exchange [31]. The data included sub-national conflict events, categorised by

event type including battles, explosions, protests, riots, strategic developments, and violence

against civilians.

The spatial granularity of the analysis was to administrative level 1 (states for Nigeria and

provinces for the DRC) and all data points that were reported on a finer spatial scale were

aggregated to the upper level. The study period was specified as January 1997 to May 2020, as

these were the first and last reports in the conflict data. The temporal scale was set to weekly, with

continuous weeks from epidemiological week 1 in 1997 to epidemiological week 20 in 2020 (1-

1,220). Continuous weeks was chosen for compatibility with the model and to include periods of

conflict that endured from one year into the next. Weeks was chosen, rather than days, to account

for reporting lags, as previous work has reported issues in the granularity of data and timeliness of

reporting, especially in humanitarian crises due to different sources of data and logistical difficulties

[32-34]. Additional information on the datasets used here are available in S2 Information.

Model Structure and Fitting

The SCCS method investigates the association between an exposure and an outcome event. The

aim is to estimate the effect, by comparing the relative incidence of the adverse events (outbreaks)

within an exposure period of hypothesised excess risk (conflicts), compared to all other times

(peace, according to the dataset used) [35-37].

Both the event and exposure were set as binary outcomes, either being present (1) or not (0).

The observation period was the full study period (1-1,220). The exposure period was the first week

after conflict onset and was reported as multiple onsets, not one long exposure period. The event

was defined by the week the cholera outbreaks was reported.

Each event and exposure that occurred in the same state/province were designated an

identification number and a pre-exposure, exposure, and post exposure period (see S1 and S2

Tables). The data was fit to conditional logistic regression models (function clogit(), R package

"survival") [38]. The model coefficient values were used to calculate incidence rate ratio (IRR), to

understand the magnitude in which conflict increased the risk of cholera outbreaks.

The datasets for each country were then split by state/province and the analysis repeated for each,

to understand if the significance of conflict on cholera outbreaks varied by sub-national location

and if conflict was more important in some states/provinces compared to others. All statistical

analyses were carried out in R version 3.6.2 and the threshold for significance was $p = <0.05$.

Percentage Attributable Fraction

The recently developed percentage attributable fraction (PAF) equations [30] were adapted to the

model results found here. The PAF values estimate the percentage of outbreaks that could be

attributed to conflict at a national level, using the datasets and the IRR values from the model

results (full equations in S3 Information).

Sensitivity Analysis

A sensitivity analysis was used to test different methods of defining the exposure end point (set to

one week in the main analysis) and to understand the impact this would have on the results. It

allowed for further understanding of how long after a conflict event the risk of cholera is

heightened. Full information is shown in S4 Information and S1 and S2 Figs.

Results

Conflict and Cholera Occurrence

The distribution of conflict and cholera in Nigeria and the DRC in the datasets used here is shown

temporally and spatially in Fig 1. The data shows an increase in reported conflict and cholera,

especially after 2010 (Fig 1a-d) and a large proportion of the cholera cases have been reported in

conflict-stricken areas (Fig 1e-f).

Fig 1. Changes in cholera and conflict. a, monthly cholera cases and deaths for Nigeria and **b**, the Democratic Republic of Congo. **c**, Monthly frequency of conflict events and fatalities for Nigeria and **d**, the Democratic Republic of Congo and **e** the number of conflict events and cholera cases as a percentage of the total number of national cases by administrative level 1 for Nigeria and **f**, the Democratic Republic of Congo. All figures are for the full datasets.

165 The total number of conflicts and outbreaks for each state/province during the study period is

166 shown below in Fig 2 and totaled 4,639 conflict and 396 cholera outbreaks for the DRC and in

167 Nigeria, 8,190 conflicts and 782 cholera outbreaks.

Fig 2. Percentage of events in each dataset including conflict and outbreaks, for **a**, Nigeria and **b**, the Democratic Republic of Congo by administrative level 1. FCT - Federal Capital Territory.

169 To be included in the analysis, the state/province had to report both outbreaks and conflicts during

170 the study period. As such, 22 provinces were included for the DRC and 36 states for Nigeria

171　(states and provinces excluded are shown in S5 Information). The temporal distribution of the

172　exposure periods and outbreaks included in the SCCS model for each state/province is shown in

173　Fig 3.

Fig 3. Swimmer plots. Showing the conflict exposure period in the SCCS model (1 week after the onset) and the outbreaks (black triangles) for each state/province for **a**, Nigeria and **b**, the Democratic Republic of Congo.

Model Output

Conflict significantly increased the risk of cholera outbreaks (IRR) in the past 23 years in Nigeria

and the DRC. Nigeria showed an effect of greater magnitude, increasing the risk of cholera

outbreaks by up to 3.6 times (IRR = 3.6, 95%CI = 3.3-3.9). Whereas for the DRC, the risk was

increased by 2.6 times (IRR = 2.6, 95%CI = 2.3-2.9).

Of the 36 Nigerian states included in the analysis, 24 showed significant associations between

conflict and cholera outbreaks. The strongest effect was found in Kebbi, Lagos, Osun, Borno and

Nasarawa, with IRR values ranging from 6.8 to 6.2 (Fig 4a).

Eleven out of 22 DRC provinces included in the analysis showed a significant relationship between

conflict and cholera. Tanganyika, Kasaï-Oriental, Maniema, Nord-Kivu and Kasaï found the

strongest values and some were the highest values found in the analysis. In Tanganyika, conflict

increased cholera risk by 7.5 times and 3.7 times for Kasaï (Fig 4b).

Percentage Attributable Fraction

Based on the IRR values from the model results (3.6 for Nigeria and 2.6 for the DRC), the onset of

a conflict event was attributable to 19.7% of cholera outbreaks in Nigeria and 12.3% for the DRC.

Sensitivity Analysis

The effect of conflict on cholera outbreaks at a national and sub-national level for both Nigeria and

the DRC decreased with increasing exposure period. By week 6 the change was minimal and

plateaued or increased (full results in S3 and S4 Figs).

Fig 4. Incidence rate ratio (IRR) for the effect of exposure to conflict within one week of the event and cholera at a sub-national level. For **a**, Nigeria and **b**, the Democratic Republic of Congo. Only results that were significant at the threshold p=<0.05 are plotted here.

Discussion

Conflict events in Nigeria increased the risk of cholera outbreaks by 3.6 times and 2.6 times for the

DRC. The percentage of cholera outbreaks attributable to the conflicts reported here was 19.7%

for Nigeria and 12.3% for the DRC. The states/provinces with the highest increased risk were

Kebbi, Nigeria at 6.9 times and Kasaï-Oriental, the DRC at 7.3 times. This showed that in some

states/provinces, the effect of conflict was much greater than the national level.

The sensitivity analysis showed a decrease in effect as the weeks progressed, with some

states/provinces seeing a plateau or increase around 6 weeks after the event. The decrease with

the lag duration may be a" diluting" effect, as the probability of an outbreak will increase across a

longer period. The states/provinces that increased after week 6, were often those with the

strongest initial effect, especially in the DRC. This larger initial effect may have a longer lasting

impact, potentially due to conflict severity.

States/provinces with the greatest increased risk often coincided with areas of high conflict. This

provides further evidence to the hypothesis that conflict may be a driver of cholera in Nigeria and

the DRC. States/provinces surrounding high conflict areas were also highly significant areas (e.g.,

Abia, Ogun, Osun, Maniema, and Tanganyika), showing a potential spill-over effect. The

states/provinces here were studied independently but a possible explanation may be people

fleeing areas of conflict or a cholera outbreak to neighbouring states, as displacement is a known

risk factor for disease outbreaks [12]. This is especially important for cholera, as a large proportion

of people will be asymptomatic but can still shed the pathogen into local reservoirs [39].

Cholera outbreaks can be explosive and self-limiting, due to the high number of asymptomatic

individuals, reducing the susceptible pool [39]. This potentially explains why the impacts of conflict

on cholera was seen just one week after the event. The incubation period of cholera is short [40],

making the effect within the first week found here biologically possible for the pathogen and a

realistic timeframe for elevated exposure to manifest in cases. Other examples of cholera cases

emerging within the first week after an adverse event include Cyclone Thane hitting the Bay of

Bengal [41], water supply interruption in the DRC [42] and Cyclone Aila in West Bengal [43]. This

provides further evidence of the need for quick and effective aid in humanitarian crisis to avoid

outbreaks and reduce mortality [44].

Healthcare facilities can suffer in periods of conflict and cholera outbreaks can overwhelm

systems, a potential cause of the relationship between conflict and cholera shown here. Care can

be inaccessible because of direct infrastructure damage or difficulties getting to the facilities due to

impromptu roadblocks [45]. Supplies may be stolen and/or unable to be delivered, including oral

rehydration solution (ORS), pathogen-sensitive antibiotics and oral cholera vaccines, which are

important for cholera outbreak control and mortality [46]. Finally, safety is a serious issue, both for

healthcare workers and patients and non-governmental (NGO) organisations can withdraw from

these areas, citing an inability to ensure the safety of their staff [47]. Steps need to be taken

globally to reduce this violence, such as using active clinical management for all patients to

enhance the acceptance of pathogen-specific treatment centres [48].

Conflict has the potential to worsen pre-existing vulnerabilities, which can exacerbate poverty,

another potential cause of the effect of conflict on cholera. The impacts of poverty can be far

reaching and is a known risk factor for cholera [5,49], along with other diseases [50]. For example,

poor urban settlements have faced the brunt of outbreaks including Zika, Ebola, typhoid, and

cholera, due to crowding and poor access to WASH [51]. Those in poorer communities may also

have more contacts and greater transmission, creating a vicious cycle [50]. Conflict can result in

loss of possessions, habitual residence, and an inability to find employment, reducing income

generation, savings and financial backstops [17]. In times of worsening poverty, people may not be

able to afford healthcare and basic medical supplies, especially in vulnerable groups. This

disruption to daily life can cause many more deaths than direct battlefield fatalities and leads to

stagnation in development [52].

A lack of WASH facilities is likely to have contributed to the positive relationship between cholera

and conflict found here. Although WASH and poverty were not directly evaluated, their effects are

likely to have been important. Conflict events can lead to disruption in sanitation and hygiene and

adverse events can act as catalysts in the interaction of contaminated water and the human

populations [53]. Displacement from conflict can cause issues in accessing WASH (e.g., latrine

access, soap availability) and several displacement camps have seen rapid cholera outbreaks,

including the DRC after the Rwandan genocide in 1994 [2]. If people are displaced due to conflict,

this may result in the use of water with unknown risk, or because alternatives are lacking, leading

to outbreaks.

A potential limitation of this analysis is the confounding effect of waterbodies in the most significant

states/provinces including the Lake Chad basin in Nigeria and the African Great Lakes Region in

the DRC. Water is considered fundamental in cholera transmission [54], although no study has yet

demonstrated a long-term persistence of toxigenic *Vibrio cholerae* in African lakes [55]. Several

reporting issues are likely to have affected the data used here, including underreporting,

overreporting and a reporting lag. Underreporting is a significant issue in global cholera estimates,

due to asymptomatic cases and disincentives to report [56]. This may be worsened in conflicts, due

displaced populations, or disruption to surveillance. Alternatively, during times of conflict health

surveillance can be enhanced by the government and/or NGOs [33]. Reporting delay is another

potential problem and some national reporting delays, have been found to range from 12 days for

meningococcal disease to 40 days for pertussis [33]. To help account for these issues, both the

event and the exposure was set to a binary outcome and weekly data was used, instead of daily.

In summary, our analysis shows a clear relationship between cholera and conflict in both Nigeria

and the DRC at both a national and sub-national level. Conflict increased the risk of cholera

outbreaks by up to 7.3 times in some states/provinces and almost 20% of cholera outbreaks being

attributable to conflict in Nigeria. This finding potentially holds in other countries and diseases and

highlights how the SCCS methodology could be used in different contexts. Cholera risks are likely

multi-factorial in both northeastern Nigeria and eastern DRC and several conditions need to be met

for emergencies to lead to cholera outbreaks. Sufficient and rapid support, along with enhanced

efforts to build community trust can reduce this access risk. Finding conflict resolution should be

the main priority in fragile states and pre-existing vulnerabilities need to be addressed, such as

poverty, expansion of affordable healthcare and improvements in WASH. By reducing these

vulnerabilities, communities will have greater resources to adapt to social extremes and could help

to reduce vulnerabilities both in times of conflict and peace.

Acknowledgements

This work was supported by the Natural Environmental Research Council [NE/S007415/1], as part

of the Grantham Institute for Climate Change and the Environment's (Imperial College London)

Science and Solutions for a Changing Planet Doctoral Training Partnership. We also acknowledge

joint Centre funding from the UK Medical Research Council and Department for International

Development [MR/R0156600/1]. We thank the organisations who published and those who

collected the data that was used here, including the Nigerian Centre for Disease Control and

Minstère de la Santé RDC. Finally, we acknowledge and thank Heather Whitaker (Open University)

and Yonas Weldeselassie (University of Warwick) for their assistance and advice in the SCCS

methodology.

Competing interests

The authors declare no competing interests.

References

1. World Health Organization. The top 10 causes of death. https://www.who.int/news-room/fact-

sheets/detail/the-top-10-causes-of-death (accessed September 9, 2018).

2. Nkoko DB, Giraudoux P, Plisnier PD, Tinda AM, Piarroux M, Sudre B et al. Dynamics of

cholera outbreaks in Great Lakes region of Africa, 1978–2008. Emerg Infect Dis. 2011; 17:

2026.

3.   D'Mello-Guyett L, Gallandat K, Van den Bergh R, Taylor D, Bulit G, Legros D, Maes P et al.

Prevention and control of cholera with household and community water, sanitation and hygiene

(WASH) interventions: a scoping review of current international guidelines. PloS One 2020; 15:

e0226549.

4.   Leckebusch GC, Abdussalam AF. Climate and socioeconomic influences on interannual

variability of cholera in Nigeria. Health Place. 2015; 34: 107-17.

5.   Talavera A, Perez EM. Is cholera disease associated with poverty?. J Infect Dev Ctries. 2009;

3: 408-11.

6.   Charnley GE, Kelman I, Green N, Hinsley W, Gaythorpe KA, Murray KA. Relationships

between drought and epidemic cholera in Africa. medRxiv 2021. doi:

https://doi.org/10.1101/2021.07.16.21260629.

7.   Gormley M. Untangling the causes of the 2016–18 cholera epidemic in Yemen. Lancet Glob

Health. 2018; 6: e600-1.

8.   Rieckmann A, Tamason CC, Gurley ES, Rod NH, Jensen PK. Exploring droughts and floods

and their association with cholera outbreaks in sub-Saharan Africa: a register-based ecological

study from 1990 to 2010. Am J Trop Med. 2018; 98: 1269-74.

9.   Al-Salem WS, Pigott DM, Subramaniam K, Haines LR, Kelly-Hope L, Molyneux DH et al.

Cutaneous leishmaniasis and conflict in Syria. Emerg Infect Dis. 2016; 22: 931.

10. Dureab FA, Shibib K, Al-Yousufi R, Jahn A. Yemen: cholera outbreak and the ongoing armed

conflict. J Infect Dev Ctries. 2018; 12: 397-403.

11. Kraemer MU, Pigott DM, Hill SC, Vanderslott S, Reiner RC, Stasse S et al. Dynamics of

conflict during the Ebola outbreak in the Democratic Republic of the Congo 2018–2019. BMC

Med. 2020; 18: 1-0.

12. Watson JT, Gayer M, Connolly MA. Epidemics after natural disasters. Emerg Infect Dis. 2007;

13: 1.

13. Charnley GE, Kelman I, Gaythorpe KA, Murray KA. Traits and risk factors of post-disaster

infectious disease outbreaks: a systematic review. Sci Rep. 2021; 11: 1-4.

14. Gayer M, Legros D, Formenty P, Connolly MA. Conflict and emerging infectious diseases.

Emerg Infect Dis. 2007; 13: 1625.

15. McPake B, Witter S, Ssali S, Wurie H, Namakula J, Ssengooba F. Ebola in the context of

conflict affected states and health systems: case studies of Northern Uganda and Sierra

Leone. Conf Health. 2015; 9: 1-9.

16. Wells CR, Pandey A, Mbah ML, Gaüzère BA, Malvy D, Singer BH et al. The exacerbation of

Ebola outbreaks by conflict in the Democratic Republic of the Congo. PNAS. 2019; 116:

24366-72.

17. Okunlola OC, Okafor IG. Conflict–Poverty Relationship in Africa: A Disaggregated Approach. J

Interdiscip Econ. 2020: 1-26.

18. Agbiboa D. The ongoing campaign of terror in Nigeria: Boko Haram versus the state. Stability.

2013; 2.

19. Council on Foreign Relations. Violence in the Democratic Republic of Congo.

https:/www.cfr.org/global-conflict-tracker/conflict/violence-democratic-republic-congo (accessed

September 9, 2018).

20. Ali M, Nelson AR, Lopez AL, Sack DA. Updated global burden of cholera in endemic countries.

PLoS Negl Trop Dis. 2015; 9: e0003832.

21. Bompangue D, Giraudoux P, Piarroux M, Mutombo G, Shamavu R, Sudre B et al. Cholera

epidemics, war and disasters around Goma and Lake Kivu: an eight-year survey. PLoS Negl

Trop Dis. 2009; 3: e436.

22. United Nations Statistical Division, 2015. Millennium Development Goal Indicators.

https://unstats.un.org/unsd/mdg/SeriesDetail.aspx?srid=580 (accessed September 9, 2018).

23. Blackburn CC, Lenze PE, Casey RP. Conflict and cholera: Yemen's man-made public health

crisis and the global implications of weaponizing health. Health Secur. 2020; 18: 125-31.

24. Federspiel F, Ali M. The cholera outbreak in Yemen: lessons learned and way forward. BMC

Public Health 2018; 18: 1-8.

25. Sato R. Effect of armed conflict on vaccination: evidence from the Boko haram insurgency in

northeastern Nigeria. Confl Health. 2019; 13: 1-0.

26. Tijjani SJ, Ma L. Is Nigeria prepared and ready to respond to the COVID-19 pandemic in its

conflict-affected northeastern states?. Int J Equity Health. 2020; 19: 1-4.

27. Farrington CP. Relative incidence estimation from case series for vaccine safety evaluation.

Biometrics 1995: 228-35.

28. Brauer R, Smeeth L, Anaya-Izquierdo K, Timmis A, Denaxas SC, Farrington CP et al.

Antipsychotic drugs and risks of myocardial infarction: a self-controlled case series study. Eur

Heart J. 2015; 36: 984-92.

29. Douglas IJ, Evans SJ, Pocock S, Smeeth L. The risk of fractures associated with

thiazolidinediones: a self-controlled case-series study. PLoS Med 2009; 6: e1000154.

30. Jean K, Raad H, Gaythorpe KA, Hamlet A, Mueller JE, Hogan D et al. Assessing the impact of

preventive mass vaccination campaigns on yellow fever outbreaks in Africa: A population-level

self-controlled case series study. PLoS Med. 2021; 18: e1003523.

31. HDX. The Humanitarian Data Exchange. https://data.humdata.org (accessed September 9,

2018).

32. Jajosky RA, Groseclose SL. Evaluation of reporting timeliness of public health surveillance

systems for infectious diseases. BMC Public Health 2004; 4: 1-9.

33. Ri S, Blair AH, Kim CJ, Haar RJ. Attacks on healthcare facilities as an indicator of violence

against civilians in Syria: An exploratory analysis of open-source data. PloS One 2019; 14:

e0217905.

34. Weidmann NB. A closer look at reporting bias in conflict event data. Am J Pol Sci. 2016; 60:

206-18.

35. Petersen I, Douglas I, Whitaker H. Self controlled case series methods: an alternative to

standard epidemiological study designs. BMJ 2016; 354.

36. Farrington CP, Whitaker HJ, Hocine MN. Case series analysis for censored, perturbed, or

curtailed post-event exposures. Biostatistics 2009; 10: 3-16.

37. Whitaker HJ, Paddy Farrington C, Spiessens B, Musonda P. Tutorial in biostatistics: the self-

controlled case series method. Stat Med. 2006; 25: 1768-97.

38. Therneau TM, 2021. A Package for Survival Analysis in R. https://CRAN.R-

project.org/package=survival (accessed 2 October 2021).

39. King AA, Ionides EL, Pascual M, Bouma MJ. Inapparent infections and cholera dynamics.

Nature 2008; 454: 877-80.

40. Azman AS, Rudolph KE, Cummings DA, Lessler J. The incubation period of cholera: a

systematic review. J Infect. 2013; 66: 432-8.

41. Fredrick T, Ponnaiah M, Murhekar MV, Jayaraman Y, David JK, Vadivoo S et al. Cholera

outbreak linked with lack of safe water supply following a tropical cyclone in Pondicherry, India,

2012. J Health Popul Nutr. 2015; 33: 31.

42. Jeandron A, Saidi JM, Kapama A, Burhole M, Birembano F, Vandevelde T et al. Water supply

interruptions and suspected cholera incidence: a time-series regression in the Democratic

Republic of the Congo. PLoS Med. 2015; 12: e1001893.

43. Bhunia R, Ghosh S. Waterborne cholera outbreak following cyclone Aila in Sundarban area of

West Bengal, India, 2009. Trans R Soc Trop. 2011; 105: 214-9.

44. Tauxe RV, Holmberg SD, Dodin A, Wells JV, Blake PA. Epidemic cholera in Mali: high mortality

and multiple routes of transmission in a famine area. Epidemiol Infect. 1988; 100: 279-289.

45. Sousa C, Hagopian A. Conflict, health care and professional perseverance: a qualitative study

in the West Bank. Glob Public Health. 2011; 6: 520-33.

46. Cartwright EJ, Patel MK, Mbopi-Keou FX, Ayers T, Haenke B, Wagenaar BH et al. Recurrent

epidemic cholera with high mortality in Cameroon: persistent challenges 40 years into the

seventh pandemic. Epidemiol Infect. 2013; 141: 2083-93.

47. MSF. DRC: Violent attacks against staff force MSF to end projects in Fizi territory, South Kivu.

https://www.msf.org/msf-forced-pull-out-eastern-drc-territory-following-violent-attacks

(accessed September 9, 2018).

48. Nguyen VK. An epidemic of suspicion—Ebola and violence in the DRC. N Engl J Med. 2019;

380: 1298-9.

49. Penrose K, Castro MC, Werema J, Ryan ET. Informal urban settlements and cholera risk in

Dar es Salaam, Tanzania. PLoS Negl Trop Dis. 2010; 4: e631.

50. Fallah MP, Skrip LA, Gertler S, Yamin D, Galvani AP. Quantifying poverty as a driver of Ebola

transmission. PLoS Negl Trop Dis. 2015; 9: e0004260.

51. Eisenstein M. Disease: poverty and pathogens. Nature 2016; 531: S61-3.

52. Trani JF, Bakhshi P, Noor AA, Lopez D, Mashkoor A. Poverty, vulnerability, and provision of

healthcare in Afghanistan. Soc Sci Med. 2010; 70: 1745-55.

53. Jutla A, Khan R, Colwell R. Natural disasters and cholera outbreaks: current understanding

and future outlook. Curr Environ Health Rep. 2017; 4: 99-107.

54. Birmingham ME, Lee LA, Ndayimirije N, Nkurikiye S, Hersh BS, Wells JG et al. Epidemic

cholera in Burundi: patterns of transmission in the Great Rift Valley Lake region. Lancet. 1997;

349: 981-5.

55. Bompangue D, Giraudoux P, Handschumacher P, Piarroux M, Sudre B, Ekwanzala M et al.

Lakes as source of cholera outbreaks, Democratic Republic of Congo. Emerg Infect Dis. 2008;

14: 798.

56. Elimian KO, Musah A, Mezue S, Oyebanji O, Yennan S, Jinadu A et al. Descriptive

epidemiology of cholera outbreak in Nigeria, January–November, 2018: implications for the

global roadmap strategy. BMC Public Health 2019; 19: 1-1.

Supporting information

S1 Information. Cholera case definitions according to the Nigerian Centre for Disease Control and the Ministère de la Santé Publique de la République démocratique du Congo.

NCDC:

Suspected case: Severe dehydration or death from acute watery diarrhoea in a patient aged 5 years or more. In an epidemic situation: A suspected case in any person aged 5 years or more with acute watery diarrhoea with or without vomiting.

Confirmed case: A suspected case in which *Vibrio cholerae* O1 or O139 has been isolated in the stool.

RDC Ministère de la Santé:

Suspected case: Severe dehydration or death following acute watery diarrhoea in a patient aged 5 years or more. In an epidemic situation: Acute watery diarrhoea with or without vomiting in a patient aged 1 year or more.

S2 Information. Dataset information.

Cholera data was compiled from a range of publicly available sources (WHO's disease outbreak news, ProMED, ReliefWeb, WHO's regional office for Africa weekly outbreak and emergencies, UNICEF cholera platform, EM-DAT, the Nigerian CDC and a literature search) in both English and French. A data charting form was used to enable a dynamic data entry process and collected data on date, geographic location, cases, deaths, hospitalisations, fatality rates, gender, age, oral cholera vaccinations, risk factors, aid and the source of the report. Data spanned from 1971-2021 for Nigeria and 1978-2021 for the DRC on a daily temporal scale and was provided at the finest spatial scale possible.

Conflict data was provided by the United Nations Office for the Coordination of Humanitarian Affairs's Humanitarian Data Exchange (HDX, 2020). The data included sub-national conflict events for both countries on a fine spatial scale, given to the exact location in longitude/latitude. This was reported on a daily temporal scale and spanned from 1997 to 2020. The data was also categorised by event type which included battles, explosions, protests, riots, strategic developments and violence against civilians. This was further sub-categorised within these groups and reported number of fatalities.

The study period was selected as Jan 1997 to May 2020, as these were the first and last reports in the conflict data. The spatial granularity of the analysis was to administrative level 1 (states for Nigeria and provinces for the DRC) and all data points that were reported on a finer spatial scale were attributed to the upper level. To be included in the analysis, the state/province had to report both outbreaks and conflicts during the study period, therefore 22 provinces were included for the DRC and 36 states for Nigeria.

S3 Information. Percentage attributable fraction (PAF) equations.

Estimate the number of outbreaks attributable to conflicts, A_i, for each province i. Using the formula:

$$A_i = \lambda_i d_i^{E+}(IRR - 1) \tag{1}$$

Where d_i^{E+} is the total duration of conflict exposure for the province i (if no conflict in province i, thus $d_i^{E+} = 0$), λ_i is the rate of outbreak occurrence in a Poisson process in the absence of conflict, and IRR is the incidence rate ratio associated with exposure to conflict. With N_i^{E-} being the number of outbreaks observed in the province i during the un-exposed period and T being the total period of observation, an estimator of λ_i is $\hat{\lambda}_i = N_i^{E-}/(T - d_i^{E+})$, which leads to:

$$\hat{A} = \sum_i \frac{N_i^{E-} d_i^{E+}}{(T - d_i^{E+})}(\hat{IRR} - 1) \tag{2}$$

Based on $\hat{A}$ and N, the total number of outbreaks observed, we can easily obtain the equivalent of the population attributable fraction, PAF, which corresponds to the proportion of the total number of outbreaks in both countries that are attributable to conflicts (this is equivalent to the PAF obtained in classical epidemiological studies, but here population refers to the "population of provinces"):

$$PAF = \frac{\hat{A}}{N} \tag{3}$$

S4 Information. Sensitivity analysis. Alternative exposure end points to identify the effect of lag.

Five alternative exposure periods were tested from the original exposure period (1 week after the

onset of exposure, lag 1) and were named lag periods due to the potential lag effect from conflict

onset to cholera outbreaks, these included:

1. Lag 2 - Week of conflict onset + 2 weeks

2. Lag 4 - Week of conflict onset + 4 weeks

3. Lag 6 - Week of conflict onset + 6 weeks

4. Lag 8 - Week of conflict onset + 8 weeks

5. Lag 10 - Week of conflict onset + 10 weeks

The sensitivity analysis was run on both a national and sub-national level and S1 and S2 Figs

show additional swimmer plots of lag 10 and line plots of the temporal trends.

S5 Information. Excluded events. States/provinces removed as they did not report conflict and

cholera in the study period (1997-2020).

Democratic Republic of Congo:

. Haut-Uele - 629 conflict events removed

. Kasaï-Central - 234 conflict events removed

. Lomani - 101 conflict events removed

. Tshuapa - 70 conflict events removed

Nigeria:

. Imo - 239 conflict events removed

578 **S1 Table. The layout of the data frame used to create the pseudo-dataset for the model.**

579 Each new 'end' period represents a different sensitivity analysis and each event and exposure are

580 given a reference number (indiv). The example shown here is for the Democratic Republic of

581 Congo Conflict dataset.

province	exday	eventday	start	end	start1	end0	end1	end2	end4	end6	end8	end10	indiv
Bas-Uele	3	374	1	542	3	3	4	5	7	9	11	13	1
Bas-Uele	4	374	1	542	4	4	5	6	8	10	12	14	2
Bas-Uele	6	374	1	542	6	6	7	8	10	12	14	16	3
Bas-Uele	7	374	1	542	7	7	8	9	11	13	15	17	4
Bas-Uele	9	374	1	542	9	9	10	11	13	15	17	19	5

583 **S2 Table. The pseudo-dataset created from the data in S1 Table for the first two reference**

584 **numbers.** This data can then be fit to the model (conditional logistic regression).

indiv	exday	eventday	start	end	event	exgr	interval	loginterval
1	3	374	1	3	0	0	2	0.693147180559945
1	3	374	3	13	0	1	10	2.30258509299405
1	3	374	13	542	1	0	529	6.2709884318583
2	4	374	1	4	0	0	3	1.09861228866811
2	4	374	4	14	0	1	10	2.30258509299405
2	4	374	14	542	1	0	528	6.26909628370626

S1 Figure. Swimmer plots showing the conflict dataset for lag 10 in the sensitivity

Analysis. In relation to outbreaks (black triangles) for Nigeria (NGA) and the Democratic Republic

of Congo (COD).

603 S2 Figure. Number of outbreak (orange) and conflict (purple) events by year in Nigeria and

604 the Democratic Republic of Congo over the full study period.

S3 Figure. Results of national sensitivity analysis. Incidence rate ratio (IRR) for the effect of

exposure to conflict within 1, 2, 4, 6, 8 and 10 weeks of the event and cholera for **a**, Nigeria and **b**,

the Democratic Republic of Congo. Only results that were significant at the threshold p=<0.05 are

plotted here. From week 1 to week 10 the risk decreased from 3.6 to 2.08 for Nigeria and from 2.6

to 1.5 for the DRC. This suggests that the risk of conflict on cholera is highest soon after the event

but remains a detectable association albeit at a lower level for potentially a long period of time after

the event.

 S4 Figure. Results of subnational sensitivity analysis. Incidence rate ratio (IRR) for the effect

 of exposure to conflict within 1, 2, 4, 6, 8 and 10 weeks of the event and cholera at administrative

 level 1. For **a**, Nigeria and **b**, the Democratic Republic of Congo. Only results that were significant

 at the threshold $p=<0.05$ are plotted here. Thirty Nigerian states and 13 DRC provinces were found

 to be significant for at least one of the lag periods and the most significant states predominately

 followed the trends of the national analysis. Values ranged from Kebbi at 6.9 to 4.0 times increased

 risk of cholera, to Gombe at 2.4 to 1.5 (Fig 6a).