Original Research

Ambient heat exposure and COPD hospitalisations in England: A

nationwide case-crossover study during 2007-2018.

Garyfallos Konstantinoudis,[1,*] Cosetta Minelli,[2] Ana Maria Vicedo Cabrera,[3,4] Joan Ballester,[5]

Antonio Gasparrini,[6,7,8] and Marta Blangiardo[1]

Affiliations:

[1]MRC Centre for Environment and Health, Department of Epidemiology and Biostatistics,

School of Public Health, Imperial College London, London, UK

[2]National Heart and Lung Institute, Imperial College London, London, United Kingdom

[3]Institute of Social and Preventive Medicine, University of Bern, Switzerland

[4]Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

[5]ISGlobal, Barcelona, Spain.

[6]Department of Public Health Environments and Society, London School of Hygiene Tropical

Medicine, London, UK

[7]Centre for Statistical Methodology, London School of Hygiene Tropical Medicine, London,

UK

[8]Centre on Climate Change and Planetary Health, London School of Hygiene Tropical

Medicine, London, UK

*Corresponding author:

Garyfallos Konstantinoudis, MRC Centre for Environment and Health, Department of

Epidemiology and Biostatistics, School of Public Health, Imperial College London, London,

UK

E-mail: g.konstantinoudis@imperial.ac.uk

Word count: 3488

Key Messages

What is the key question?

What is the short-term effect of heat exposure on COPD hospitalisation and which

contextual/societal factors affect population vulnerability?

What is the bottom line?

For every 1^{o}C increase in summer temperatures higher than 23.2^{o}C, the risk of COPD

hospitalisation increases by 1.47%, and populations in the North and in the South East are more

vulnerable.

Why read on?

This large nationwide study in England using individual data quantifies the effect of heat

exposure on COPD hospitalisations; these findings inform future policies regarding

preparedness and resilience of public health systems against the increasing COPD burden due

to the increasing temperatures.

Abstract

Background There is emerging evidence suggesting a link between ambient heat exposure and

chronic obstructive pulmonary disease (COPD) hospitalisations. Individual and contextual

characteristics can affect population vulnerabilities to COPD hospitalisation due to heat

exposure. This study quantifies the effect of ambient heat on COPD hospitalisations and

examines population vulnerabilities by age, sex, and contextual characteristics.

Methods Individual data on COPD hospitalisation at high geographical resolution (postcodes)

during 2007-2018 in England was retrieved from the small area health statistics unit. Maximum

temperature at 1 km×1km resolution was available from the UK Met Office. We employed a

case-cross over study design and fitted Bayesian conditional Poisson regression models. We

adjusted for relative humidity and national holidays, and examined effect modification by age,

sex, green space, average temperature, deprivation and urbanicity.

Results After accounting for confounding, we found a 1.47% (95% Credible Interval 1.19% to

1.73%), increase in the hospitalisation risk for every 1°C increase in temperatures above 23.2°C

(lags 0-2 days). We reported weak evidence of an effect modification by sex and age. We found

a strong spatial determinant of the COPD hospitalisation risk due to heat exposure, that was

alleviated when we accounted for contextual characteristics. 1 851 (95% CrI 1 576 to 2 079)

COPD hospitalisations were associated with temperatures above 23.2°C annually.

Conclusion Our study suggests that resources should be allocated to support the public health

systems, for instance through developing or expanding heat-health alerts, to challenge the

increasing future heat-related COPD hospitalisation burden.

Introduction

Chronic obstructive pulmonary disease (COPD) is the most prevalent chronic respiratory

disease worldwide, with point prevalence varying from 1.56% in Sub-Saharan Africa to 6.09%

in Central Europe, eastern Europe, and central Asia in 2007.[1] In England, COPD is a significant

cause of morbidity and mortality, leading to 115 000 emergency admissions and 24 000 deaths

per year.[2] The causes of acute exacerbation of COPD are established and include factors such

as sex, age, COPD severity and comorbidities.[3] Environmental triggers of COPD

hospitalisations such as air-pollution exposure have also been discussed extensively.[4] There is

emerging evidence suggesting a link between heat exposure and COPD hospitalisation, either

directly or through exacerbating the effects of factors such as ozone concentration that are

associated with these events.[5]

Several previous studies have examined the effect of high temperatures on COPD

hospitalisations, reporting higher rates with heat exposure[6-8] and heat waves.[9,10] The majority of

these studies are based on aggregated data (at the city or regional level),[6,9-11] whereas only a few

considered individual data.[7,8] Use of individual data allows investigation of possible effect

modification by individual factors such as age and sex, and it avoids ecological bias arising

when group-level associations do not reflect associations at the individual level.[12] Although

previous studies have assessed the vulnerability related to individual factors, such as age and sex,[6,8]

contextual characteristics, such as green space, average temperature, deprivation and urbanicity,

are still poorly characterised. Two of the previous studies have examined the spatial variation of

the temperature effect on COPD hospitalisation, using however very coarse geographical

resolution.[6,8]

In this nationwide study in England during 2007-2018, we investigated the effect of heat

exposure on COPD hospital admissions using a semi-ecological framework. We took advantage

of the individual data availability of the outcome and adopted a case-crossover study design that

naturally accounts for time-constant variables at the individual patient level. Thus, we were able to

account for factors like age, sex, comorbidities, deprivation as well as lifestyle characteristics such

as physical activity through the study design. We also adjusted for time-varying confounders,

such as air-pollution exposure and relative humidity and examined how the effect of temperature

is modified by age, sex and in space. Last, we assessed the extent to which contextual

characteristics, such as green space, deprivation, urbanicity and average temperature, contribute

to the observed spatial variation of the effect of temperature.

Methods

Study population

We included inpatient hospital admissions from COPD in England during 2007-2018 as

retrieved from Hospital Episode Statistics (HES) data held by the UK Small Area Health

Statistics Unit, provided by the Health and Social Care Information Centre. Age, postcode of

residence at time of the hospitalisation, and date of hospitalisation were available for each

record. We focused only on admissions with acute exacerbation of COPD as primary diagnosis.

We investigated the following diagnostic groups: J40–44 according to the International

Classification of Disease version 10 (ICD10).[13] The analysis is restricted to June, July and

August.

Exposure

Daily minimum and maximum temperatures were available at 1 km×1 km resolution from the

UK Met Office with methods described elsewhere.[14] In brief, the daily temperature in each

grid-cell was estimated based on inverse-distance-weighted interpolation of monitoring data,

also accounting for latitude and longitude, elevation, coastal influence, and proportion of urban

land use. To assign daily temperature to health records, the postcode centroids of each patient

were spatially linked to the 1km× 1km grid cell, applying a 100m fuzziness to the postcode

location to fulfil governance requirements. We focused on daily maximum temperature, as we

are interested in heat exposure, averaged over the day of hospitalisation and the preceding two

days (lags 0-2 days) to estimate the cumulative health effects.[15-17]

Covariates

We used hourly concentration of Ozone (O_3) and atmospheric particulate matter that has a

diameter of less than 2.5 μm ($PM_{2.5}$), as retrieved from the unified model produced by the Met

Office measured in $\mu g/m^3$. [18] The model outcome is then post-processed to correct for bias

using observational data. [18] For O_3 we calculated the daily mean of the 8 hours of maximum O_3,

whereas for $PM_{2.5}$ the daily mean concentration. The geographical resolution of the air-

pollutants is 12 km×12 km for the years 2007-2011 and 2km×2km during 2012-2019. We

adjusted for relative humidity (daily and at a 10 km×10 km grid) through a model that

integrates MetOffice data on daily observations from the meteorological stations and monthly

nationwide data as retrieved from HadUK,[14] see Online Supplement Text S1.1. All covariates

were included at lags 0-2 days, to match the exposure lags. O_3, $PM_{2.5}$ and relative humidity

were included as linear terms in the model. We also accounted for the effect of national

holidays through a dummy variable.

Spatial effect modifiers

We selected these spatial effect modifiers based on consistency with the literature,[19] data

availability in England and a priori hypotheses, see Online Supplement Text S1.3. As a

measure of green space we used the proportion of a region that is covered by green land such as

woodland, agricultural land, grassland and other natural vegetated land as classified in the Land

Cover Map 2015 (LCM15).[20]Deprivation is measured using the Index of Multiple Deprivation

(IMD) 2015, as retrieved from the Ministry of Housing, Communities and Local Government.[21]

We used the quintiles of IMD in our analysis. For these two modifiers, we selected the year

2015 as the most representative data point, among the ones available, for our study period.

Urbanicity (predominantly rural, urban with significant rural and predominantly urban) is based

on the Office for National Statistics (ONS) classification in 2011 (the most recent year for

which data was available at the time of analysis).[22] We also incorporated the average

temperature during 2007-2018, as a measure of adaptation on higher temperatures.[23] Green

space and average temperature were included as linear terms in the model. Due to power and

computational considerations, all spatial effect modifiers were included at the lower tier local

authority level (LTLA; Online Supplement Fig. S1).

Statistical methods

We used a time-stratified case-crossover design, commonly used for analysing the effect of

transient exposures.[24 25] The temperature on the day of COPD hospitalisation (event day), is

compared with the temperature on non-event days. In the case-crossover design a case serves as

its own control, thus this design automatically controls for factors that do not vary or vary

slowly over time, such as sex or deprivation. We selected non-event days on the same day of

week and calendar month as the event day to avoid the overlap bias.[26] Thus we could have

maximum 4 non-event days per event day.

We modelled the effect of temperature on event compared to non-event days by specifying

Bayesian hierarchical conditional Poisson models, with a fixed effect on the event/non-event

day grouping.[19 27] We accounted for recurrent hospitalisations by adding a random effect on

each patient. For the main analysis, we treated relative humidity and national holidays as

confounders and thus adjusted for them, but we did not adjust for air-pollutants because they

were treated as mediators, see directed acyclic graph on Online Supplement Text S1.3.[28] As the

effect of temperature on health is typically non-linear,[19] we used piecewise linear threshold

models, to allow more flexible fits, but retain ease of interpretation. We considered nationwide

thresholds, specified as the 50th, 55th …, 95th percentile of the daily temperatures. We selected

the threshold based on the WAIC, an estimate of predictive accuracy, with smaller values

indicating better fits.[29 30] We then ran additional models allowing the effect of heat exposure

(temperatures above the threshold) to vary by sex (male and female), age (0-64, 65-74, 75+)

and space (LTLA). We additionally included the air-pollutants in these models to examine the

sensitivity of the effect if the air-pollutants were confounders. For the spatial effect

modification, we used the Besag-York-Mollie prior that assumes local dependency among

adjacent LTLAs.[31] We fitted this model with and without the spatial effect modifiers, while

adjusting for confounders. The model is described in detail in the Online Supplement Text

S1.2. Results are reported as medians and 95% Credible Intervals (CrI; 95% probability that the

true values lies within this interval) of % increase in the hospitalisation risk for every 1°C

increase in temperatures above the threshold;[32] additionally, we report posterior probabilities of

a positive % increase. For the spatially varying risk, we also reported posterior probabilities

that the % hospitalisation risk is larger than the average % hospitalisation risk.

Population attributable fraction

To calculate the population attributable fraction, we extended[33] to incorporate the spatial

dimension of the effect of heat exposure. We first calculated the cumulative heat exposure -

COPD hospitalisation relative risk (RR_s) for the s-th LTLA. We could then calculate the

attributable fraction: $AF_s = (RR_s - 1)/RR_s$. Let n_s be the number of hospitalisations at days

above 23.2°C and N_s the total number of hospitalisations, then $AF_s(n_s/N_s)$ is the population

attributable fraction, i.e., the number of COPD hospitalisations attributable to summer heat

exposure. In our Bayesian formulation we were able to propagate all the random variable-

specific uncertainty in our estimates.

Sensitivity analyses

We repeated the main analysis for the lags 0, 1 and 2 independently. We also used b-splines to

model the temperature effect and examined the linearity assumption above the threshold.

All analyses are run in R-NIMBLE (Numerical Inference for Hierarchical Models Using

Bayesian and Likelihood Estimation).[34] The code for running the analysis is online available at

.

Results

Population

We retrieved 1 570 288 COPD hospital records during 2007-2018 in England. After removing

the duplicated records, the ones with place of residence outside England, the ones not occurred

in summer months and the ones for which we could not sample non-event days, we had 320

411 records available for the analysis Fig. 1.

Exposure, Covariates and Effect modifiers

The median maximum temperature across England has increased from 19.42°C in 2007 to

22.20°C in 2018, Online Supplement Table S1. The median maximum temperature exposure is

20.91°C at lag 0 for event and 20.39°C non-event days, 20.97°C for event and 20.94°C for non-

event days at lag 1, 20.92°C for event and 20.90°C for non-event days at lag 2 and 20.93°C for

event and 20.92°C for non-event days at lag 0-2, Online Supplement Table S2. The distribution

of the covariates across event and non-event days and the spatial distribution of the effect

modifiers at the LTLA level can be found on the Online Supplement, Table S2-5, and Fig. S2-

5.

WAIC analysis

In the model adjusted for relative humidity and national holidays, the 80th percentile of the

temperature (23.2°C) was the threshold minimising the WAIC, Online Supplement Table S6.

We found a 0.37% (95% CrI 0.09% to 0.65%) increase in the COPD hospitalisation risk for

every 1°C increase in temperatures below 23.2°C, Online Supplement Table S6. In contrast, the

effect above 23.2°C was higher, namely 1.46% (95% CrI 1.19% to 1.71%), Online Supplement

Table S6. All subsequent analyses were conducted using the 80-th percentile of the temperature

as the threshold.

Age and sex effect modification

In the unadjusted models, the percentage of risk increase in hospitalisations for every 1°C

increase above the threshold varies from 0.92% (95% CrI 0.25% to 1.63%) in females 64 years

old or younger to 1.56% (95% CrI 0.94% to 2.20%) in females aged 65-74, Fig. 2 and Online

Supplement Table S6. After adjusting for relative humidity and national holidays the effects are

slightly higher varying from 1.14% (95% CrI 0.39% to 1.84%) in females 64 years old or

younger to 1.75% (95% CrI 1.13% to 2.41%) in males 65-74 years old, Fig. 2 and Online

Supplement Table S7. Additionally adjusting for air-pollution substantially reduces the

observed effect, Fig. 2 and Online Supplement Table S7.

Spatial effect modification

The spatial variation of the effect of heat exposure on COPD hospitalisations is shown on Fig.

3. The risk of COPD hospitalisation is less than 1.31% for every 1°C increase in heat exposure

in South West, top left panel Fig. 3. In contrast, populations in the South East are more

vulnerable: the probability that the effect of heat exposure is larger than the national average

estimate ranges between 0.6 and 1, top right panel Fig. 3. After incorporating green space,

deprivation, urbanicity, and average temperature, the observed variation of the effect of

temperature is alleviated, bottom panels Fig. 3.

We found weak evidence that populations in areas with higher proportions of green space,

larger average temperature and higher level of urbanicity are more resilient to COPD

hospitalisations due to heat exposure, Table 1. If we increase an LTLA's proportion of green

space by 1%, the spatial effect of the heat exposure changes by -1.46% (95% CrI -6.99% to

4.39%), Table 1. For every 1°C increase in the average temperature per LTLA, the spatial effect

of the heat exposure changes by -0.41% (95% CrI -1.49% to 0.71%), Table 1. The spatial effect

of heat exposure in urban LTLAs with significant rural and predominantly urban LTLAs

changes by -0.79% (95% CrI -3.10% to 1.51%) and -1.57% (95% CrI -4.16% to 0.96%)

respectively compared with predominantly rural LTLAs, Table 1.

Population attributable burden

We found that 1 851 (95% CrI 1 576 to 2 079) COPD hospitalisations were associated with

temperatures above 23.2^{o}C annually. This accounts for 7.8% (95% CrI 6.7% to 8.8%) of the

total COPD hospitalisations during the summer months from 2007 to 2019. The proportion of

COPD hospitalisations attributable to temperatures above the threshold has a clear spatial

structure and is more than 8% in East Midlands, East of England, London and South East, while

it is below 5% in the South West (Fig. 4).

Sensitivity analysis

The lag with the highest influence was lag 1 with the risk of COPD hospitalisation being 1.37%

(95% CrI 1.14% to 1.58%) for every 1^{o}C increase in heat exposure. For lag 0 and lag 2 the

point estimate was still positive, but lower in magnitude, 0.71% (95% CrI 0.50% to 0.93%) and

1.01% (95% CrI 0.78% to 1.24%) respectively, likely due to the correlation with temperatures

at lag 1. The linearity assumption above the 23.2^{o}C threshold looks reasonable, Online

Supplement Fig. S6.

Discussion

This is the first nationwide case-crossover study in England investigating the short-term effects

of heat exposure on COPD hospitalisation. After accounting for confounding, the results

indicate that for every 1^{o}C increase in heat exposure the COPD hospitalisation risk increases by

1.47% (95% CrI 1.19% to 1.73%), with evidence that $PM_{2.5}$ and O_3 mediate this relationship.

We found weak evidence of an effect modification by sex and age. The attributable burden of

heat exposure has a clear spatial structure, with areas in East Midlands, East of England,

London and South East affected the most. Assuming a causal relationship, 7.8% (95% CrI 6.7%

to 8.8%) of COPD hospitalisations could be attributed to heat exposure during the summer

months between 2007 and 2018.

The main strength of our study is the availability of postal codes, exploiting the highest spatial

resolution available for linkage with the exposure and confounding factors. Such geographical

resolution is expected to minimise misclassification resulting from any spatial misalignment

between the outcome and exposure/confounder. The availability of individual data for the

outcome also minimises ecological bias,[12] while guaranteeing high statistical power due to the

population-based nature of the study. We ascertained hospital records from NHS digital

covering almost all hospitalisation occurred in the public sector in England during 2007-2018.

Our study has some limitations. First, residential temperature does not reflect the actual

temperature exposure of an individual, as individuals are exposed to different temperatures in

the course of the day. In addition to this, the outdoor temperature, as provided by Met Office

does not reflect the actual temperature exposure inside the house. Nevertheless, in line with

most of the studies in this field and given the lack of more precise individual exposure data, we

used residential temperature outdoors as a proxy for the individual exposure. To allow for

flexible fits, we used a linear threshold model. More complex relationships may need multiple

thresholds, however the WAIC analysis suggested that the linearity assumption suffices.

Although we adjusted for the main COPD hospitalisation environmental contributors, we could

not evaluate other potential confounders (e.g., seasonal allergies and pollen counts) due to the

lack of available data. Additionally, exposure to other air-pollutants, such as NO_2, SO_2, might

also confound the observed relationship; we decided to adjust for $PM_{2.5}$ and O_3 as they seem to

have a larger impact on COPD hospitalisation and to avoid potential collinearity with other

pollutants.

Our results can be compared with studies examining COPD hospital admissions and ambient

temperatures during the hottest months.[6 11 35 36] Our study is in line with a US study including

12.5 million participants that found a 4.7% (95% CrI 3.9 to 5.5%) increase in the COPD

hospitalisation rate at lag 0 for every 5.6^{o}C increase in the average daily temperature during

May-September.[6] Our study is also in line with a case-crossover study in Brazil that reported a

5% (95% CrI 4% to 6%) increase in the hospitalisation odds for every 5^{o}C increase in the

average temperature (0-3 lags) during the 4 hottest months.[11] In contrast, a study in New York

reported a 7.64% increase in the risk of COPD admissions for each 1^{o}C increase in daily mean

apparent temperature above 32^{o}C.[36] A study in 12 European cities, reported a 4.5% (95% CrI

1.9 to 7.3) and 3.1% (95% CrI 0.8 to 5.5) increase in total respiratory admissions (the majority

being COPD) in Mediterranean and North-Continental cities, respectively, for every for each

1^{o}C increase in the maximum apparent temperature (lag 0-3 days) above the 90th percentile.[37]

A study in Taiwan reported negative correlation between the average daily temperature and

emergency admissions with COPD, but a 14% increase in the emergency COPD admissions

when the diurnal temperature range is larger than 9.6^{o}C.[35]

We found weak evidence of an effect modification by age and sex, but discrepancies in

vulnerability in space. A previous study in Brazil, reported higher COPD hospitalization odds

for women and the older people.[11] In the models adjusted for relative humidity and national

holidays, in line with a previous study in the US,[6] the age group 65-74 was the most vulnerable.

Some spatial variability by regions or counties was also observed in previous studies in Brazil

and the US, potentially due to socioeconomic characteristics or exposure to higher average

summer temperatures.[6 11] In our study, green space, average temperature, deprivation and

urbanicity explained some of the observed variation in the observed spatial vulnerabilities, the

evidence of an effect was however inconclusive.

Some discrepancies of our results compared with previous studies can have multiple

explanations. Previous studies reporting higher effect estimates had available coarser

geographical resolution (city or county level), leading to inadequate adjustment for

confounding, as confounders, can vary in high geographical resolution.[11 35 37 38] Differences in

the definition of the outcome can also lead to the discrepancies as previous studies have used

the apparent temperature,[36 37] or diurnal temperature range,[35] while others, more in line with our

approach, the daily mean.[6 11] Decisions regarding the selection of the temperature threshold, the

warm-season months and the lags can also partly explain the observed difference in the effect

estimates. Most previous studies adjusted for air-pollution,[6 36 37] while we did not, as we

assumed that air-pollution is a mediator;[28] when we added air-pollution to model the effect of

heat exposure was much reduced.

Acute COPD episodes are associated with airways and systemic inflammation but also with

cardiovascular comorbidity and may be triggered by exposures to heat.[37] Exposure to ambient

heat can lead to heat dissipation through hyperventilation and may trigger dynamic

hyperinflation and dyspnoea in patients with pre-existing COPD.[6 11] The higher risk of COPD

hospitalisation in the 65-74 age group observed in our study could be explained by the inability

of this frail population to dissipate excess heat through circulatory adjustment, and exposure to

extreme temperatures increases their risk of developing pulmonary vascular resistance

secondary to peripheral pooling of blood or hypovolemia.[37] In addition, older populations are of

higher risk to have cardiovascular comorbidities, which are hypothesised to increase the risk of

COPD hospitalisations associated with heat exposure. Nevertheless, such evidence is

inconclusive.[37] We also reported a weak protective effect of higher average temperatures,

arguing towards protective adaptation to heat, possibly related to differences in housing stock

or behaviour during hot weather.[11] We observed weak evidence of increased resilience in

populations in more deprived areas and in areas with higher degrees of urbanicity. Although

this evidence is inconclusive, potential factors that could confound the observed effect include

differences in demographics, for instance ethnicity.

Previous studies examining future trends in COPD, population demographics and temperature

changes have predicted a higher COPD prevalence, a raise in the average age of the population

and increased global temperatures.[39-41] Resources should be allocated to support the

preparedness and resilience of public health systems, for instance through developing or

expanding heat-health alerts, to challenge the increasing heat exposure related COPD

hospitalisation burden.

Acknowledgements

We thank Hima Daby, Gajanan Natu, Eric Johnson and Bethan Davies for their help with data

acquisition, storage, preparation, and governance. All authors acknowledge infrastructure

support for the Department of Epidemiology and Biostatistics provided by the NIHR Imperial

Biomedical Research Centre (BRC). Hospital Episode Statistics data are copyright © 2021, re-

used with the permission of NHS Digital. All rights reserved. The Hospital Episode Statistics

data were obtained from NHS Digital.

Contributors

G.K and M.B. conceived the study. M.B. supervised the study. G.K. developed the initial study

protocol and discussed it with M.B., C.M., A.M.V.C., J.B and A.G.. G.K. developed the

statistical model, prepared the covariate data and led the acquisition of hospitalisation data.

M.B. validated the code. G.K. ran all the analysis and wrote the initial draft. All the authors

contributed in modifying the paper and critically interpreting the results. All authors read and

approved the final version for publication.

Funding

G.K. is supported by an MRC Skills Development Fellowship [MR/T025352/1]. M.B. is

supported by a National Institutes of Health, grant number [R01HD092580-01A1].

Infrastructure support for this research was provided by the National Institute for Health

Research Imperial Biomedical Research Centre (BRC). A.G. is supported by the Medical

Research Council-UK (Grant ID: MR/R013349/1), the Natural Environment Research Council

UK (Grant ID: NE/R009384/1) and the European Union's Horizon 2020 Project Exhaustion

(Grant ID: 820655). The work was partly supported by the MRC Centre for Environment and

Health, which is funded by the Medical Research Council (MR/S019669/1, 2019-2024). J.B.

gratefully acknowledges funding from the European Union's Horizon 2020 research and

innovation programme under grant agreement No 865564 (European Research Council

Consolidator Grant EARLY-ADAPT), support from the Spanish Ministry of Science and

Innovation through the "Centro de Excelencia Severo Ochoa 2019-2023" Program (CEX2018-

000806-S), and support from the Generalitat de Catalunya through the CERCA Program.

The work of the UK Small Area Health Statistics Unit is overseen by Public Health England

(PHE) and funded by PHE as part of the MRC-PHE Centre for Environment and Health also

supported by the UK Medical Research Council, Grant number: MR/L01341X/1), and the

National Institute for Health Research (NIHR) through its Health Protection Units (HPRUs) at

Imperial College London in Environmental Exposures and Health and in Chemical and

Radiation Threats and Hazards, and through Health Data Research UK (HDR UK). This paper

does not necessarily reflect the views of Public Health England, the National Institute for

Health Research or the Department of Health and Social Care.

Competing interests

The authors declare no competing interests.

Patient consent for publication

Not required.

Ethics

The analyses were covered by national research ethics approval from the London-South East

Research Ethics Committee (Reference 17/LO/0846). Data access was covered by the Health

Research Authority Confidentiality Advisory Group under section 251 of the National Health

Service Act 2006 and the Health Service (Control of Patient Information) Regulations 2002

(Reference 20/CAG/0028).

Data availability statement

SAHSU does not have permission to supply data to third parties. For reproducibility purposes we

have simulated data and provided the code used for the analysis in

https://github.com/gkonstantinoudis/COPDTempSVC.

References

1. England N. Overview of potential to reduce lives lost from Chronic Obstructive Pulmonary

Disease (COPD). *Department of Health* 2014

2. Soriano JB, Kendrick PJ, Paulson KR, et al. Prevalence and attributable health burden of

chronic respiratory diseases, 1990-2017: a systematic analysis for the Global Burden of

Disease Study 2017. *Lancet Resp Med* 2020;8(6):585-96 doi: 10.1016/S2213-

2600(20)30105-3

3. Alqahtani JS, Njoku CM, Bereznicki B, et al. Risk factors for all-cause hospital readmission

following exacerbation of COPD: a systematic review and meta-analysis. Eur Respir

Rev 2020;29(156) doi: 10.1183/16000617.0166-2019

4. DeVries R, Kriebel D, Sama S. Outdoor Air Pollution and COPD-Related Emergency

Department Visits, Hospital Admissions, and Mortality: A Meta-Analysis. Copd

2017;14(1):113-21. doi: 10.1080/15412555.2016.1216956

5. De Sario M, Katsouyanni K, Michelozzi P. Climate change, extreme weather events, air

pollution and respiratory health in Europe. Eur Respir J 2013;42(3):826-43. doi:

10.1183/09031936.00074712

6. Anderson B, Dominici F, Wang Y, et al. Heat-Related Emergency Hospitalizations for

Respiratory Diseases in the Medicare Population. Am J Epidemiol 2013;177:S64-S64

doi: 10.1164/rccm.201211-1969OC

7. Astrom DO, Schifano P, Asta F, et al. The effect of heat waves on mortality in susceptible

groups: a cohort study of a mediterranean and a northern European City. Environ

Health-Glob 2015;14 doi: 10.1186/s12940-015-0012-0

8. Zhao Q, Li SS, Coelho MDZS, et al. Ambient heat and hospitalisation for COPD in Brazil: a

nationwide case-crossover study. Thorax 2019;74(11):1031-36. doi: 10.1136/thoraxjnl-

2019-213486

9. Bobb JF, Obermeyer Z, Wang Y, et al. Cause-Specific Risk of Hospital Admission Related

to Extreme Heat in Older Adults. Jama-J Am Med Assoc 2014;312(24):2659-67. doi:

10.1001/jama.2014.15715

10. Monteiro A, Carvalho V, Oliveira T, et al. Excess mortality and morbidity during the July

2006 heat wave in Porto, Portugal. Int J Biometeorol 2013;57(1):155-67. doi:

10.1007/s00484-012-0543-9

11. Zhang Y, Liu X, Kong D. Effects of Ambient Temperature on Acute Exacerbations of

Chronic Obstructive Pulmonary Disease: Results from a Time-Series Analysis of

143318 Hospitalizations (vol 15, pg 213, 2020). Int J Chronic Obstr 2021;16:2129-31.

doi: 10.2147/Copd.S322834

12. Wakefield J. Ecologic studies revisited. Annu Rev Publ Health 2008;29:75-+. doi:

10.1146/annurev.publhealth.29.020907.090821

13. Rothnie KJ, Mullerova H, Thomas SL, et al. Recording of hospitalizations for acute

exacerbations of COPD in UK electronic health care records. Clin Epidemiol

2016;8:771-82. doi: 10.2147/Clep.S117867

14. Hollis D, McCarthy M, Kendon M, et al. HadUK-Grid-A new UK dataset of gridded

climate observations. Geosci Data J 2019;6(2):151-59. doi: 10.1002/gdj3.78

15. Gasparrini A, Armstrong B, Kovats S, et al. The effect of high temperatures on cause-

specific mortality in England and Wales. Occup Environ Med 2012;69(1):56-61. doi:

10.1136/oem.2010.059782

16. Hajat S, Armstrong BG, Gouveia N, et al. Mortality displacement of heat-related deaths - A

comparison of Delhi, Sao Paulo, and London. Epidemiology 2005;16(5):613-20. doi:

10.1097/01.ede.0000164559.41092.2a

17. Hajat S, Kovats RS, Lachowycz K. Heat-related and cold-related deaths in England and

Wales: who is at risk? Occup Environ Med 2007;64(2):93-100. doi:

10.1136/oem.2006.029017

18. Neal LS, Agnew P, Moseley S, et al. Application of a statistical post-processing technique

to a gridded, operational, air quality forecast. Atmos Environ 2014;98:385-93. doi:

10.1016/j.atmosenv.2014.09.004

19. Bennett JE, Blangiardo M, Fecht D, et al. Vulnerability to the mortality effects of warm

temperature in the districts of England and Wales. Nat Clim Change 2014;4(4):269-73.

doi: 10.1038/Nclimate2123

20. Rowland C, Morton D, Carrasco Tornero L, et al. Land Cover Map 2015 (1km percentage

aggregate class, GB). 2017 doi: 10.5285/711c8dc1-0f4e-42ad-a703-8b5d19c92247

21. McLennan D, Noble S, Noble M, et al. The English indices of deprivation 2019: technical

report. 2019

22. Statistics DR. The 2011 Rural-Urban Classification for Output Areas in England, 2018.

23. Arbuthnott K, Hajat S, Heaviside C, et al. Changes in population susceptibility to heat and

cold over time: assessing adaptation to climate change. Environ Health-Glob 2016;15

doi: 10.1186/s12940-016-0102-7

24. Lu Y, Zeger SL. On the equivalence of case-crossover and time series methods in

environmental epidemiology. Biostatistics 2007;8(2):337-44. doi:

10.1093/biostatistics/kxl013

25. Maclure M. The Case-Crossover Design: A Method for Studying Transient Effects on the

Risk of Acute Events. Am J Epidemiol 2017;185(11):1174-83. doi: 10.1093/aje/kwx105

26. Janes H, Sheppard L, Lumley T. Overlap bias in the case-crossover design, with application

to air pollution exposures. Stat Med 2005;24(2):285-300. doi: 10.1002/sim.1889

27. Armstrong BG, Gasparrini A, Tobias A. Conditional Poisson models: a flexible alternative

to conditional logistic case cross-over analysis. Bmc Med Res Methodol 2014;14 doi:

10.1186/1471-2288-14-122

28. Buckley JP, Samet JM, Richardson DB. Commentary: Does air pollution confound studies

of temperature? Epidemiology 2014;25(2):242-45 doi:

10.1097/EDE.0000000000000051

29. Gelman A, Hwang J, Vehtari A. Understanding predictive information criteria for Bayesian

models. Stat Comput 2014;24(6):997-1016. doi: 10.1007/s11222-013-9416-2

30. Watanabe S. Asymptotic Equivalence of Bayes Cross Validation and Widely Applicable

Information Criterion in Singular Learning Theory. J Mach Learn Res 2010;11:3571-

94.

31. Besag J, York J, Mollié A. Bayesian image restoration, with two applications in spatial

statistics. Annals of the institute of statistical mathematics 1991;43(1):1-20 doi:

10.1007/BF00116466

32. Navidi W, Weinhandl E. Risk set sampling for case-crossover designs. Epidemiology

2002;13(1):100-05. doi: Doi 10.1097/00001648-200201000-00016

33. Mansournia MA, Altman DG. Population attributable fraction. BMJ 2018;360 doi:

10.1136/bmj.k757

34. de Valpine P, Turek D, Paciorek CJ, et al. Programming With Models: Writing Statistical

Algorithms for General Model Structures With NIMBLE. J Comput Graph Stat

2017;26(2):403-13. doi: 10.1080/10618600.2016.1172487

35. Liang WM, Liu WP, Kuo HW. Diurnal temperature range and emergency room admissions

for chronic obstructive pulmonary disease in Taiwan. Int J Biometeorol 2009;53(1):17-

23. doi: 10.1007/s00484-008-0187-y

36. Lin S, Luo M, Walker RJ, et al. Extreme High Temperatures and Hospital Admissions for

Respiratory and Cardiovascular Diseases. Epidemiology 2009;20(5):738-46. doi:

10.1097/EDE.0b013e3181ad5522

37. Michelozzi P, Accetta G, De Sario M, et al. High Temperature and Hospitalizations for

Cardiovascular and Respiratory Causes in 12 European Cities. Am J Resp Crit Care

2009;179(5):383-89. doi: 10.1164/rccm.200802-217OC

38. Konstantinoudis G, Padellini T, Bennett J, et al. Long-term exposure to air-pollution and

COVID-19 mortality in England: A hierarchical spatial analysis. Environment

International 2021;146 doi: 10.1016/j.envint.2020.106316

39. Meehl GA, Stocker TF, Collins WD, et al. Global climate projections. Chapter 10. 2007

40. Nash A. National population projections: 2018-based. Office for National Statistics 2019

41. McLean S, Hoogendoorn M, Hoogenveen RT, et al. Projecting the COPD population and

costs in England and Scotland: 2011 to 2030. Sci Rep-Uk 2016;6 doi:

10.1038/srep31893

Tables

Table 1. Median, 95% credible intervals of the percentage change of the heat exposure related spatial

hospitalization risk due to green space, average temperature, index of multiple deprivation

and urbanicity and probability that this percentage change is higher than 0.

Effect modifier	Percentage change	Pr(% change>0)[1]
Green space[2]	-1.46 (-6.99 to 4.39)	0.30
Average temperature[3]	-0.41 (-1.49 to 0.71)	0.22
IMD[4]		
Q1	1	
Q2	0.81 (-1.16 to 3.08)	0.78
Q3	1.57 (-0.76 to 4.06)	0.91
Q4	0.75 (-1.68 to 3.36)	0.71
Q5	1.62 (-1.31 to 4.49)	0.85
Predominantly Rural	1	
Urban with significant rural	-0.79 (-3.10 to 1.51)	0.25
Predominantly urban	-1.57 (-4.16 to 0.96)	0.12

[1] Posterior probability that the percentage change is larger than zero.

[2] Green space is the proportion of a region covered by green land such as woodland, agricultural

land, grassland and other natural vegetated land.

[3] The average temperature is the mean summer temperature per LTLA during 2007-2018°C.

[4] Index of multiple deprivation. IMD is calculated based on the following domains: a. income,

b. employment, c. education, skills and training, d. health and disability, e. crime, f. barriers to

572   housing and services and g. living environment deprivation. Q1 denotes the most deprived

573   areas, whereas Q5 the least deprived.

Figures

Figures

Fig. 1 Flowchart of COPD hospitalisations.

Fig. 2 Percentage risk of COPD hospitalisation for every 1^{o}C increase in the temperatures

above 23.2^{o}C during the summer months between 2007 and 2018, for the unadjusted (left

panel), the model adjusted for relative humidity (RH) and national holidays (NL) (mid

panel) and the model additional adjusted for air-pollution (POL) (right panel). Results are

stratified by age (0-64, 65-74, 75+, total) and sex (male, female, total).

Fig. 3 Median spatial COPD hospitalisation risk for every 1^{o}C increase in the temperatures

above 23.2^{o}C and posterior probability that the risk is larger than the overall risk in England

during the summer months between 2007 and 2018. The top panels refer to the model without

incorporating contextual characteristics, whereas the panels below otherwise. All models

were fully adjusted.

Fig. 4 The percentage of COPD hospitalisations by lower tier local authorities attributed to

exposure to summer temperatures above 23.2^{o}C during 2007-2018 in England. This effect

assumes a causal relationship between heat exposure and COPD hospitalisation risk. The

island on the left is a zoomed version of London.

Spatial attributable fraction