Increased Risk for Type 2 Diabetes in Relation to Adiposity in Middle-Aged Black South African Men compared to Women

Journal:	*European Journal of Endocrinology*
Manuscript ID	EJE-21-0527
mstype:	Clinical Study
Date Submitted by the Author:	17-May-2021
Complete List of Authors:	Kufe, Clement; University of the Witwatersrand Faculty of Health Sciences, Paediatrics; WITS University Micklesfield, Lisa; South African Medical Research Council/University of the Witwatersrand, Developmental Pathways for Health Research Unit (DPHRU), Paediatrics Masemola, Maphoko; 1South African Medical Research Council/University of the Witwatersrand, Developmental Pathways for Health Research Unit (DPHRU), Paediatrics Chikowore, Tinashe; 1South African Medical Research Council/University of the Witwatersrand, Developmental Pathways for Health Research Unit (DPHRU), Paediatrics Kengne, Andre; Non-Communicable Disease Research Unit, South African Medical Research Council, Non-Communicable Disease Research Unit Karpe, Fredrik; University of Oxford, Oxford, UK, Oxford Centre for Diabetes, Endocrinology and Metabolism; Oxford Medical Research Centre, Oxford Radcliffe Hospitals Trust, OCDEM, Churchill Hospitsl, Oxford, UK, National Institute for Health Research Norris, Shane; South African Medical Research Council/University of the Witwatersrand, Developmental Pathways for Health Research (DPHRU), Paediatrics; School of Health and Human Development, University of Southampton, UK, Global Health Research Institute Crowther, Nigel; National Health Laboratory Service and University of Witwatersrand, Department of Chemical Pathology Olsson, Tommy; Umeå University, Department of Public Health and Clinical Medicine, Medicine Goedecke, Julia; South African Medical Research Council/University of the Witwatersrand, Developmental Pathways for Health Research Unit (DPHRU), Paediatrics; South African Medical Research Council, Non-Communicable Diseases Research Unit
Keywords:	Diabetes Mellitus, Insulin, Obesity

Increased Risk for Type 2 Diabetes in Relation to Adiposity in Middle-Aged Black South African Men compared to Women

Clement N. Kufe[1][§], Lisa K. Micklesfield[1], Maphoko Masemola[1], Tinashe Chikowore[1], Andre Pascal Kengne[2], Fredrik Karpe[3,4], Shane A. Norris[1,5], Nigel J. Crowther[6], Tommy Olsson[7], Julia H. Goedecke[1,2]

Authors affiliations:

[1]South African Medical Research Council/University of the Witwatersrand, Developmental Pathways for Health Research Unit (DPHRU), Department of Paediatrics, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

[2]Non-Communicable Diseases Research Unit, South African Medical Research Council, Cape Town, South Africa

[3]Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford, UK

[4]National Institute for Health Research, Oxford Biomedical Research Centre, Oxford Radcliffe Hospitals Trust, OCDEM, Churchill Hospital, Oxford, UK

[5]Global Health Research Institute, School of Health and Human Development, University of Southampton, UK

[6]Department of Chemical Pathology, National Health Laboratory Service and School of Pathology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

[7]Department of Public Health and Clinical Medicine, Medicine, Umeå University, Umeå, Sweden

Running title: Increased Risk for T2D from Adiposity

§ Corresponding author

Clement N. Kufe

SAMRC/WITS Developmental Pathways for Health Research Unit,

Department of Paediatrics,

School of Clinical Medicine,

Faculty of Health Sciences,

University of the Witwatersrand, Johannesburg, South Africa

Phone: +27 63 092 6943

E-mail: klekufe@gmail.com

Email addresses:

CNK: klekufe@gmail.com

LKM: lisa.micklesfield@wits.ac.za

MM: maphokomasemola@gmail.com

TC: tinashedoc@gmail.com

APK: andre.kengne@mrc.ac.za

FK: fredrik.karpe@ocdem.ox.ac.uk

SAN: shane.norris@wits.ac.za

NC: nigel.crowther@nhls.ac.za

TO: tommy.g.olsson@umu.se

JHG: julia.goedecke@mrc.ac.za

Word count: 3498 (3500 maximum)

Number of tables and figures: 4

Abstract

Aims: Despite a higher prevalence of overweight/obesity in black South African women compared to men, the prevalence of type 2 diabetes does not differ. We explored if this could be due to sex differences in insulin sensitivity, clearance and/or beta-cell function, and also sex-specific associations with total and regional adiposity.

Methods: This cross-sectional study included 804 black South African men (n=388) and women (n=416). Dual-energy x-ray absorptiometry was used to measure total and regional adiposity. Insulin sensitivity (Matsuda index), secretion (C-peptide index) and clearance (C-peptide/insulin ratio) were estimated from an oral glucose tolerance test.

Results: After adjusting for sex differences in fat mass index, men were less insulin sensitive and had lower beta cell function than women (p<0.001), with the strength of the associations with measures of total and central adiposity being greater in men than women (p<0.001 for interactions). Further, the association between total adiposity and type 2 diabetes risk was also greater in men than women (relative risk ratio (95% confidence interval): 2.05 (1.42–2.96), p<0.001 vs. 1.38 (1.03–1.85), p=0.031).

Conclusion: With increasing adiposity, particularly increased centralisation of body fat linked to decreased insulin sensitivity and beta cell function, black African men are at greater risk for type 2 diabetes than their female counterparts.

Key words: Body fat distribution, insulin sensitivity, insulin secretion, beta-cell function, basal and postprandial insulin clearance, ethnicity. **(Word count abstract: 207)**

Introduction

Type 2 diabetes (T2D) is a global health problem, with low-middle income countries

particularly affected. It is projected that sub-Saharan Africa will have the highest increase in

T2D compared to the rest of the world, and in 2019 South Africa (SA) had the highest estimated

number of people with diabetes (4.6 million) in the sub-Saharan region, and the highest age–

adjusted comparative prevalence of diabetes (12.7%) in adults (1), which is higher than the

global average (2). Within SSA and SA, the prevalence of T2D does not differ by sex, despite

large sexual dimorphism in obesity rates (3). For example, in SA the prevalence of T2D in

black SA men and women is similar (10.2% vs. 13.8%) (4) , but the prevalence of overweight

and obesity differs markedly (27.4% vs. 67.4%) (5).

The reason for this discrepancy in the association between overweight/obesity and diabetes

risk in men and women is not clear. Recent studies have started to explore the underlying

pathophysiology of T2D in Africans (6–10), and shown that black African women present with

a phenotype of low insulin sensitivity and hyperinsulinemia due to higher insulin secretion and

lower hepatic insulin clearance compared to white SA women (7) and black SA men (8).

However, the majority of these studies have been undertaken in premenopausal women

(6,7,10), with limited data in middle-aged men and women (8).

Notably, men typically have greater central fat mass (particularly visceral adipose tissue

(VAT)) and less peripheral subcutaneous adipose tissue (SAT) than women, which is

associated with a higher risk for T2D (8,11,12). However, the sex differences in the association

between whole body and regional adiposity, and T2D risk, including insulin sensitivity,

secretion and clearance, to our knowledge, has not been studied in African men and women.

Accordingly, the aims of this study were to compare insulin sensitivity, clearance and beta-cell

function between middle-aged black South African men and women who differ in obesity

prevalence, and to explore sex-specific associations with total and regional adiposity.

Methods

This cross-sectional study includes the analysis of the follow-up data that was part of a

longitudinal study designed to investigate the determinants of T2D risk in middle-aged black

SA men and women. Data collection for the baseline study, as part of the AWI-Gen (Africa

Wits-INDEPTH partnerships for Genomic Research) study (13), took place between 2011 and

2015 in black SA men (n=1027) and women (n=1008) residing in Soweto, South Africa (14).

Follow-up data, analysed for this study, was collected between January 2017 and August 2018

on a sample of 502 men and 527 women randomly selected from the original sample.

Participants living with HIV were excluded from this data analysis to avoid the confounding

effects of the virus and antiretroviral therapy on the outcomes. Complete data was available for

804 participants (388 men and 416 women) and complete oral glucose tolerance (OGTT) data

was available on 734 of these participants (Supplementary files: Figure 1).

The study was conducted in accordance with the tenets of the Helsinki declaration and was

approved by the Human Research Ethics Committee (HREC) of the University of the

Witwatersrand (M160604 and M160975). Prior to inclusion in the study all procedures and

possible risks were explained and all participants signed a consent form. Data collection took

place at the South African Medical Research Council/University of the Witwatersrand

Developmental Pathways for Health Research Unit at the Chris Hani Baragwanath Hospital in

Soweto, Johannesburg, South Africa.

Socio–demographic and medical questionnaire

Interviewer administered questionnaires were completed and captured onto REDCap (15).

Data collected included age, marital status (married/unmarried), current employment

(employed/not employed), highest educational level completed (no formal

schooling/elementary school, secondary school level, tertiary education), alcohol intake and

tobacco consumption (Yes/No), and self–reported diabetes and/or diabetes medication taken.

Menopausal stage was classified according to last menstrual period (16).

Anthropometry

Weight was measured to the nearest 0.1 kg using a TANITA digital scale (model: TBF-410,

TANITA Corporation, US). Height was measured to the nearest 0.1cm using a wall–mounted

stadiometer (Holtain, UK). Waist circumference (WC) and hip circumference (HC) were

measured to the nearest 0.1cm with a non–stretchable tape. For the WC, the tape was placed

horizontally between the iliac crest in the mid–axillary plane and the lowest rib margin. For

the HC, the tape was placed around the level of the greatest protrusion of the buttocks. Waist-

to-hip ratio (WHR) and BMI were calculated, and participants categorised according to the

World Health Organisation (WHO) criteria (17).

Body composition and body fat distribution measurements

Dual–energy X-ray absorptiometry (DXA) was used to measure whole body composition,

including sub-total (total body minus head to account for any artefacts that may influence the

DXA reading) fat mass (FM, kg and % body mass) and fat–free soft tissue mass (FFSTM), and

regional FM including trunk, arm and leg FM (QDR 4500A, Hologic Inc., Bedford, USA,

APEX software version 4.0.2). Fat mass index (FMI, sub-total fat mass kg/height^2) and FFSTM

index (FFSTM/height^2) were calculated. Regional fat distribution was expressed relative to

sub-total FM (%FM), with trunk fat (%FM) representing central fat distribution and arm and

leg fat (%FM) representing upper- and lower-body peripheral fat distribution, respectively.

Abdominal VAT and SAT areas were estimated from DXA (18).

Blood sampling and analysis

Blood samples (10 ml) were drawn after an overnight fast for the determination of glycated

haemoglobin (HbA1c), plasma glucose, serum insulin, C–peptide and follicle stimulating

hormone (FSH) concentrations. Participants then completed a standard 75g oral glucose

tolerance test (OGTT) over 2-hours during which blood samples (5 ml) were drawn at 30 min

intervals for the determination of glucose, insulin and C-peptide concentrations. Participants

with known diabetes and/or those with fasting blood glucose ≥11.1 mmol/l (n=76) (ACCU-

CHEK®, MedNet GmbH, Munster, Germany) did not complete the OGTT.

Plasma glucose concentrations were measured on the Randox RX Daytona Chemistry Analyser

(Randox Laboratories Ltd., London, UK). HbA1c concentrations were measured using the D–

10TM Haemoglobin Analyser (Bio–Rad Laboratories, Inc. USA). Serum insulin and C–peptide

concentrations were measured on the Immulite® 1000 Immunoassay System (Siemens

Chemiluminescent Healthcare GmbH, Henkestr, Germany). FSH was measured on serum

using the ARCHITECT Chemiluminescent Microparticle Immunoassay assay (Abbott

Laboratories, Abbott Ireland).

Based on the fasting plasma glucose (FPG) and 2–h OGTT glucose results, participants were

classified according to the WHO criteria (19). Participants with impaired fasting glucose and

impaired glucose tolerance were combined and described as having impaired glucose

metabolism (IGM).

Calculations from the OGTT

The homeostasis model assessment (HOMA-IR) was used to estimate fasting insulin resistance (20). The Matsuda Index (21), was used to estimate insulin sensitivity for participants with complete OGTT data (n=628), alongside the composite score (22) for participants who only had data for 0 and 120 minutes (n=106). These composite measures have been shown to compare well (22), and were significantly correlated in this study (r=0.874; p<0.001) to the Matsuda Index. Early phase insulin response to the OGTT was estimated using the insulinogenic index (IGI) (23). Participants without data at 30 minutes or whose insulin response was <0 were excluded from the analysis. Insulin secretion was calculated using the C-peptide index, the ratio of the increment in C-peptide relative to glucose in the first 30 minutes of OGTT (23). C-peptide is produced in equimolar quantities to endogenous insulin, and unlike insulin, there is negligible hepatic extraction of C-peptide, and hence the C-peptide index and the C-peptide to insulin ratio may serve as proxy measures of insulin secretion and clearance, respectively (24,25). Basal and postprandial insulin clearance were calculated as the ratio of fasting C–peptide to insulin, and the incremental area under the curve (iAUC) of C-peptide to iAUC insulin, calculated using the trapezoidal method, respectively. The oral disposition index (oDI), which reflects insulin secretion adjusted for the level of insulin sensitivity (26–28), was calculated as the product of the C-peptide index and Matsuda index (23) which demonstrated a hyperbolic relationship and was used as the measure of beta-cell function. These calculations were only performed in participants without known T2D and/or not taking medications for T2D, and who underwent an OGTT.

Statistical analysis

Data were analysed using Stata 15.1/IC (StataCorp, College Station, TX, USA). Variables are

summarised as percentages for categorical data, mean ± standard deviation (SD) for normally

distributed continuous data, and median (25th-75th percentile) if not normally distributed.

Normality was assessed using the Shapiro–Wilk test and Q-Q probability plots. Sex differences

were determined using Students t-test for normally distributed continuous data, Mann-Whitney

U and Kruskal–Wallis tests for skewed continuous data, and Chi-squared test for categorical

data. Sex differences in glucose and insulin measures are presented before and after adjusting

for FMI using one–way analysis of covariance (ANCOVA). Z-scores were derived for the total

and regional adiposity measures for the combined sample, as well as sex-stratified using

Fisher's Yates transformation (29). By using Z-scores we were able to compare the risk

magnitude per 1 SD change in total and regional adiposity measurements. Multinomial logistic

regression was used to explore the relationship between total and regional adiposity measures,

and IGM and T2D, using NGT as the reference, and including age, sex, smoking, alcohol

intake, education, and FMI (for regional measures), as covariates. All participants with known

(n=65) and newly diagnosed (n=42) diabetes were included in the multinomial analyses. We

explored sex*adiposity z-score interactions and only found a significant interaction for FMI.

Accordingly, the data (excluding FMI) were analysed in the combined sample and the relative

risk ratio (RRR) and 95% confidence intervals for IGM and T2D are presented. For the

continuous measures of insulin sensitivity (Matsuda index), clearance (fasting C–

peptide/insulin ratio) and beta-cell function (oDI), robust regressions were used to explore

associations with adiposity z-scores, including age, smoking, alcohol intake, education and

FMI (for regional adiposity measures) as covariates. Participants with known diabetes and/or

taking medication for diabetes and those without OGTT data were excluded from the robust

regression analyses. Due to significant sex interactions in most models, the analyses were

completed separately for men and women using sex-specific total and regional adiposity z-

scores. A p–value of <0.05 was considered significant.

Results

Socio-demographic and body composition characteristics

A total of 804 participants (48.3% men) with a mean age of 54.6±6.0 years were included

(Table 1). Men were younger than women and significantly more men were married than

women. Current employment status was not different between the sexes, however more men

than women (18.1 vs. 12.5%) had completed tertiary education.

Mean BMI was higher in women than men (p<0.001), and accordingly a larger proportion of

the women presented with obesity (70.2% vs. 26.6%) (Table 1). While waist circumference

was similar, men had higher WHR due to the higher hip circumference of the women. While

FFSTM was higher in men, FM (kg and %) and FMI were higher in women. When expressed

relative to FM, women had significantly greater leg FM, while men had more central FM

(trunk), but arm FM did not differ. Within the central depot, men had less VAT and SAT (both

p<0.001), but a higher VAT/SAT ratio.

Differences in glucose and insulin measures between men and women

Although fasting glucose and iAUC for glucose were not different between the sexes, HbA1C

and 2 h glucose were higher in women than men (Table 1). Fasting insulin and C-peptide, and

iAUC for insulin, were also higher in women than men. Accordingly, HOMA-IR was higher

and insulin sensitivity (Matsuda index) was lower in women compared to men, accompanied

by a higher insulin response (IGI) characterised by higher insulin secretion (C-peptide index)

and lower insulin clearance (basal and postprandial). However, the oDI, a measure of beta-cell

function, did not differ by sex.

When adjusting for differences in FMI (Table 2), there were no longer sex differences in

HbA1C, 2-hour glucose, insulin response, or basal and postprandial insulin clearance, while

insulin secretion remained higher in women. In contrast, fasting insulin and C-peptide, as well

as HOMA-IR were higher, and insulin sensitivity and beta-cell function were lower in men

compared to women.

The prevalences of NGT, IGM and T2DM were not significantly different between men and

women.

Associations between total and regional adiposity and risk for IGM and type 2 diabetes

There was a significant sex*FMI z-score interaction (p<0.001), such that the RRR for IGM and

T2D were greater for men than women (Figure 1A). Associations between regional adiposity

z-scores and risk for IGM and T2D did not differ by sex, and the RRR for the combined sample

are presented in Table 3. Trunk fat and VAT z-scores were associated with a higher risk for

both IGM and T2D, with every 1 SD increase in trunk fat and VAT being associated with a 4.8

fold and 2.6 fold increased risk for T2D, respectively. In contrast, higher leg fat z-score was

associated with a 58% and 79% lower risk for IGM and T2D, respectively, while a 1 SD higher

arm fat z-score was associated with a 2.2-fold greater risk for T2D only. SAT z-score was not

associated with IGM or T2D.

Sex-specific associations between total and regional adiposity z-scores and insulin

measures

There were significant sex*FMI z-score interactions for insulin sensitivity, clearance and beta-

cell function, with associations consistently being stronger in men than women (Figure 1B–

1D). There were also significant sex*regional adiposity interactions for most measures of

insulin sensitivity and response and therefore the results are presented separately for men and

women (Table 4). Lower insulin sensitivity was associated with higher central fat mass (trunk

fat and VAT), and lower leg fat in both men and women, but the associations with central fat

mass were stronger in men than women ($p<0.001$ for all interactions). In contrast, arm fat mass

was associated with lower insulin sensitivity in women only ($p<0.001$ for interaction). Beta-

cell function (oDI) was negatively associated with VAT in both men and women. In contrast

beta-cell function was positively associated with peripheral fat mass in women only ($p=0.040$

for interaction). Basal insulin clearance was negatively associated with trunk fat mass in both

men and women, with a stronger association in men ($p=0.017$ for interaction). In contrast, basal

insulin clearance was negatively associated with VAT and arm fat, and positively associated

with leg fat in women only, but the strength of the association did not differ significantly

between sexes. The associations for postprandial insulin clearance were similar to those for

basal insulin clearance (data not shown). As the women were at different phases of the

menopausal transition with 17.6% being premenopausal, 14.7% perimenopausal and 67.7%

being postmenopausal, we wanted to ascertain whether the associations presented above

differed by menopausal phase. The associations between total and regional adiposity and

insulin sensitivity, secretion and beta-cell function did not differ between menopausal groups.

In contrast, the associations between FMI, trunk, leg and arm z-scores and basal insulin

clearance differed by menopausal phase, being stronger in the pre- than peri- and

postmenopausal women (data not shown).

Discussion

The main and novel findings of this study were that in a sample of black men and women with

a mean age of 54.6 years, after adjustments for differences in body fat, insulin sensitivity,

secretion and beta-cell function were lower in black SA men compared to women, while insulin

clearance did not differ by sex. In line with this, the strength of the association between total

adiposity and T2D risk was greater in men compared to women. Although black SA women

have a higher prevalence of obesity (70.2 vs. 26.6%) and greater whole-body fatness (43.6 vs.

26.3%) than men, they present with a more 'favourable' body fat distribution, characterised by

less central fat mass and greater peripheral fat mass. This phenotype has been associated with

lower diabetes risk. This together with the greater impact of body fatness on diabetes risk could

explain the similar prevalence of diabetes in men and women (11.1 vs. 15.4%) despite the

lower adiposity in men in this study.

These findings also suggest that with increasing adiposity, black SA men will be at greater risk

for T2D than their female counterparts. We found that the association between total adiposity

and risk for T2D was higher in men than women (Figure 1A). Further we showed that with

increasing FMI the decline in insulin sensitivity was greater in men compared to women,

similar to earlier studies from SA (8,30), which was also associated with a more pronounced

decrease in beta-cell function in men compared to women.

The finding of similar T2D prevalence (1) despite marked differences in the prevalence of

obesity (3) between sexes are consistent and representative of South Africa and the SSA region.

In order to understand the sexual dimorphism in this relationship, it is obviously essential to

account for sex differences in body fatness as well as disentangle the sex-specific associations

between regional adiposity and T2D risk. After adjusting for differences in body fatness, men

had lower insulin sensitivity, insulin secretion and beta-cell function compared to women,

placing the men at higher risk for T2D. Indeed, a lower beta-cell function, estimated using the

oDI, has been shown to predict the development of T2D over a 10 year period in a Japanese

American cohort (26).

Black African women have been shown to present with hyperinsulinaemia compared to their

European counterparts, often beyond that required to maintain normoglycaemia (8,31).

Hyperinsulinemia in black African women has previously been attributed to alterations in both

insulin secretion and clearance, depending on age, and/or level of glycemia (10,32). Studies in

African American women have shown that decreased hepatic insulin clearance is the main

contributor to hyperinsulinemia (33). In contrast, we show that the higher IGI in women

compared to men, was associated with higher insulin secretion without differences in insulin

clearance. Due to limited longitudinal studies, it is not known whether the higher IGI in women

is protective or may actually cause insulin resistance (9).

It is well recognised globally and in South Africa that men have greater central body fat and

less lower body peripheral fat compared to women (8,30). Similarly, we showed that men had

greater trunk fat mass, a higher VAT/SAT ratio, and less leg and similar arm fat mass than

women. This adiposity phenotype is associated with greater diabetes risk (8,30,32,34). Indeed,

we showed that a 1SD increase in trunk z-score was associated with a more than two-fold

greater risk for IGM and nearly five times greater risk for T2D, and was also associated with

lower insulin sensitivity and lower basal insulin clearance. In contrast, peripheral fat is

typically associated with reduced risk for diabetes (30,34). We showed that a 1 SD increase in

leg z-score was associated with a 58% lower risk for IGM and a 79% lower risk for T2D, as

well as higher insulin sensitivity in both sexes. Notably, the strength of the inverse association

between central fat distribution and insulin sensitivity was greater in men compared to women.

Several studies in different populations have shown VAT to be more strongly associated with

insulin resistance, and therefore a greater risk for T2DM, in men than women (30,35–37). A

further a novel finding of the study was that the positive relationship between beta-cell function

and leg FM was weaker in men compared to women, suggesting lower 'protective' effect of

leg FM on beta-cell function in men compared to women. Accordingly, despite a lower

prevalence of overweight and obesity in men compared to women in our study, this

'unfavourable' regional fat distribution and the sex-specific relationships with insulin

sensitivity and beta-cell function places them at greater risk for future T2D.

This is the first study, to our knowledge, in black SA men and women with detailed measures

of insulin sensitivity, secretion and clearance, and beta-cell function, based on estimates from

an OGTT. We were also able to use DXA, which provides an accurate assessment of body

composition and regional adiposity. A limitation is the cross–sectional nature of the study

which does not allow us to infer causality. Although the sex differences in obesity and total

adiposity may be seen as a limitation, it reflects the status of obesity within South Africa and

the sub-Saharan African region (3), and adjustments for total body fatness and the calculation

of z-scores were used in the analyses to determine whether these sex differences in adiposity

were influencing the insulin- and glucose-related variables. There were no effects of the

menopausal transition *per se* on the association between adiposity and insulin sensitivity,

secretion and beta-cell function and therefore the sex differences reported cannot be explained

by menopausal status. Furthermore, the conclusions for this study are valid only for HIV

negative individuals.

In summary, for the same level of body fatness, black South African men are less insulin

sensitive and had lower insulin secretion and beta-cell function than women, with the strength

of the association between adiposity and T2D risk being greater in men compared to women.

This suggests that with increasing adiposity, particularly an increase in central adiposity, black

SA men face an increased risk for T2D in comparison with their female counterparts.

Longitudinal studies are required to confirm the results of this study.

Funding: The study was jointly funded by the South African Medical Research Council (MRC) from South African National Department of Health, MRC UK (via the Newton Fund) and GSK Africa Non-Communicable Disease Open Lab (via a supporting Grant project Number: ES/N013891/1) and South African National Research Foundation (Grant no: UID:98561).

Conflict of Interest: No potential conflicts of interest relevant to this article were reported

Authors' contributions: CNK, JHG and LKM designed the study and CNK analysed the data, drafted and revised the manuscript under the supervision of JHG and LKM. All authors reviewed/edited, read and approved all the drafts and the final version of the manuscript. The authors retained the control of the final content of the publication.

Acknowledgements: We are grateful to the participants as well as the following DPHRU staff for their input during data collection and entry: Melikhanya Soboyisi, Tshifhiwa Ratshikombo, Vukosi Mkansi, Sphume Thango, Mosadiapula Nakedi, Thabile Sibiya, Bonisiwe Mlambo, Caroline Makura, Dr Mamosilo Lichaba, Karabo Pearl Nkhahle and team, and Prof Michèle Ramsay for AWI-Gen study development and oversight.

References
1. IDF. Diabetes Atlas, 9th Edition 2019. 2019.
2. Kengne AP, Bentham J, Zhou B, Peer N, Matsha TE, Bixby H, et al. Trends in obesity and diabetes across Africa from 1980 to 2014: An analysis of pooled population-based studies. Int J Epidemiol. 2017;46(5):1421–32.
3. Ramsay M, Crowther NJ, Agongo G, Ali SA, Boua RP, Gómez-olivé FX, et al. Regional and sex-specific variation in BMI distribution in four sub-Saharan African countries : The H3Africa AWI-Gen study Regional and sex-specific variation in BMI distribution in four sub-Saharan. Glob Health Action. 2018;11(2).
4. Peer N, Steyn K, Lombard C, Lambert E V., Vythilingum B, Levitt NS. Rising Diabetes Prevalence among Urban-Dwelling Black South Africans. PLoS One. 2012;7(9).
5. NDoH, Stats SA S& I. South Africa Demographic and Health Survey 2016. 2019.
6. Goedecke JH, Dave JA, Faulenbach M V, Utzschneider KM, Lambert E V, West S, et al. Insulin response in relation to insulin sensitivity: An appropriate β-cell response in black South African women. Diabetes Care. 2009;32(5):860–5.
7. Goedecke JH, Keswell D, Weinreich C, Fan J, Victor H. Ethnic differences in hepatic and systemic insulin sensitivity and their associated determinants in obese black and white South African women. 2015;58(11):2647–52.
8. Goedecke JH, George C, Veras K, Peer N, Lombard C, Victor H, et al. Sex differences in insulin sensitivity and insulin response with increasing age in black South African men and women. Diabetes Res Clin Pract. 2016;122:207–14.
9. Goedecke JH, Olsson T. Pathogenesis of type 2 diabetes risk in black Africans: a South African perspective. J Intern Med. 2020;1–11.
10. Goedecke JH, Mtintsilana A, Dlamini SN, Pascal Kengne A. Type 2 diabetes mellitus in African women. Diabetes Res Clin Pract. 2017;123:87–96.
11. Bredella MA. Sex differences in body composition. Adv Exp Med Biol. 2017;1043:9–27.
12. Schorr M, Dichtel LE, Gerweck A V., Valera RD, Torriani M, Miller KK, et al. Sex differences in body composition and association with cardiometabolic risk. Biol Sex Differ. 2018;9(1):1–10.
13. Ramsay M, Crowther N, Tambo E, Agongo G, Baloyi V, Dikotope S, et al. H3Africa AWI-Gen Collaborative Centre: A resource to study the interplay between genomic and environmental risk factors for cardiometabolic diseases in four sub-Saharan African countries. Glob Heal Epidemiol Genomics. 2016;
14. Ali SA, Soo C, Agongo G, Alberts M, Amenga-Etego L, Boua RP, et al. Genomic and environmental risk factors for cardiometabolic diseases in Africa: methods used for Phase 1 of the AWI-Gen population cross-sectional study. Glob Health Action. 2018;11(sup2).
15. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)-A metadata-driven methodology and workflow process for providing translational research informatics support. J Biomed Inform. 2009;
16. Jaff NG, Snyman T, Norris SA, Crowther NJ. Staging reproductive aging using stages of reproductive aging workshop + 10 in black urban african women in the study of women entering and in endocrine transition. Menopause. 2014;21(11):1225–33.
17. WHO. Expert Committee on Physical Status: The use and interpretation of anthropometric physical status. World Health Organization technical report series. Geneva; 1995.
18. Micklesfield LK, Goedecke JH, Punyanitya M, Wilson KE, Kelly TL. Dual-Energy X-

Ray Performs as Well as Clinical Computed Tomography for the Measurement of Visceral Fat. Obesity. 2012;20(5):1109–14.

19. WHO. Classification of diabetes mellitus. Vol. 21, WHO. Geneva; 2019.

20. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and β-cell function from fasting plasma glucose and insulin concentrations in man. Diabetologia. 1985;28(7):412–9.

21. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing: Comparison with the euglycemic insulin clamp. Diabetes Care. 1999;22(9):1462–70.

22. DeFronzo RA, Matsuda M. Reduced time points to calculate the composite index. Vol. 33, Diabetes Care. 2010. p. 2010.

23. Tura A, Kautzky-Willer A, Pacini G. Insulinogenic indices from insulin and C-peptide: Comparison of beta-cell function from OGTT and IVGTT. Diabetes Res Clin Pract. 2006;72(3):298–301.

24. Polonsky KS, Rubenstein AH. C-peptide as a measure of the secretion and hepatic extraction of insulin. Pitfalls and limitations. Vol. 33, Diabetes. 1984. p. 486–94.

25. Leighton E, Sainsbury CA, Jones GC. A Practical Review of C-Peptide Testing in Diabetes. Vol. 8, Diabetes Therapy. Springer Healthcare; 2017. p. 475–87.

26. Utzschneider KM, Prigeon RL, Faulenbach M V., Tong J, Carr DB, Boyko EJ, et al. Oral Disposition index predicts the development of future diabetes above and beyond fasting and 2-h glucose levels. Diabetes Care. 2009;32(2):335–41.

27. Kahn SE, Prigeon RL, McCulloch DK, Boyko EJ, Bergman RN, Schwartz MW, et al. Quantification of the relationship between insulin sensitivity and β- cell function in human subjects: Evidence for a hyperbolic function. Diabetes. 1993;42(11):1663–72.

28. Mather KJ, Chen M, Hannon TS. Linearization of the Disposition Index equation allows evaluation of secretion-sensitivity coupling slopes. J Diabetes Complications. 2020;34(7):107589.

29. Armitage P BG. Statistical Methods in Medical Research, 2002. Blackwell Science Ltd: Oxford, UK. Blackwell Science Ltd: Oxford, UK; 2002.

30. Davidson FE, Matsha TE, Erasmus RT, Kengne AP, Goedecke JH. Associations between body fat distribution and cardiometabolic risk factors in mixed-ancestry South African women and men. Cardiovasc J Afr. 2019;30(6):321–30.

31. Lee CC, Haffner SM, Wagenknecht LE, Lorenzo C, Norris JM, Bergman RN, Stefanovski D, Anderson AM, Rotter JI, Goodarzi MO HA. Insulin Clearance and the Incidence of Type 2 Diabetes in Hispanics and African Americans. Diabetes Care. 2013;36:901–7.

32. Mtintsilana A, Micklesfield LK, Chorell E, Olsson T, Goedecke JH. Fat redistribution and accumulation of visceral adipose tissue predicts type 2 diabetes risk in middle-aged black South African women: a 13-year longitudinal study. Nutr Diabetes. 2019;9(1).

33. Kodama K, Tojjar D, Yamada S, Toda K, Patel CJ BA. Ethnic Differences in the Relationship Between Insulin Sensitivity and Insulin Response. A systematic review and meta-analysis. Diabetes Care. 2013;36(6):1789–96.

34. Keswell D, Tootla M, Goedecke JH. Associations between body fat distribution, insulin resistance and dyslipidaemia in black and white South African women. Cardiovasc J Afr. 2016;27(3):177–83.

35. Kautzky-willer A, Harreiter J, Pacini G. Sex and Gender Differences in Risk, Pathophysiology and Complications of Type 2 Diabetes Mellitus. 2016;37(June):278–316.

36. Geer EB, Shen W. Gender differences in insulin resistance, body composition, and

energy balance. Gend Med. 2009;

37.   De Mutsert R, Gast K, Widya R, De Koning E, Jazet I, Lamb H, et al. Associations of Abdominal Subcutaneous and Visceral Fat with Insulin Resistance and Secretion Differ between Men and Women: The Netherlands Epidemiology of Obesity Study. Metab Syndr Relat Disord. 2018;16(1):54–63.

Table 1: Socio-demographic, body composition, glucose and insulin measures in black South African men and women (n=804)

Variable	Men	Women	p-value
n (%)	388 (48.3%)	416 (51.7%)	-
Age (years)	54.2±6.2	55.0±5.8	0.047
Socio-demographic characteristics, row: n (%)			
Married, n (%)	219 (56.7)	186 (44.9)	0.001
Currently employed (n, %)	232 (47.3)	258 (52.6)	0.519
Currently smokes, n (%)	179 (46.1)	30 (7.2)	<0.001
Alcohol intake, n (%)			
Never	107 (27.6)	304 (73.1)	
Sometimes (monthly or less and 2-4 times a month)	163 (42.0)	93 (22.4)	
Often (2-3 times and 4 or more times a week)	118 (30.4)	19 (4.6)	<0.001
Educational attainment, n (%)			
No formal schooling/elementary school level	43 (11.1)	38 (9.2)	
Secondary school level	274 (70.8)	325 (78.3)	0.042
Tertiary education	70 (18.1)	52 (12.5)	
Body composition			
Height (cm)	171±6	158±6	<0.001
Weight (kg)	77.4±18.4	85.4±18.0	<0.001
BMI (kg/m^2)	26.4±6.0	34.0±7.0	<0.001
Waist circumference (cm)	96.1±15.4	97.4±13.1	0.191
Hip circumference (cm)	100.6±11.1	116.6±13.6	<0.001
WHR	0.95±0.06	0.84±0.10	<0.001
BMI categories, n (%)			
Underweight (<18.5 kg/m^2)	26 (6.7)	2 (0.5)	
Normal weight (18.50-24.99 kg/m^2)	145 (37.4)	31 (7.5)	<0.001
Overweight (25-29.99 kg/m^2)	115 (29.6)	90 (21.6)	
Obese (≥30.0 kg/m^2)	102 (26.3)	293 (70.4)	
DXA (n=763)			
Fat–free soft tissue mass (kg)	49.4±9.1	41.7±7.0	<0.001
Fat–free soft tissue mass index (kg/m^2)	16.8±2.8	16.6±2.6	0.270
Body fat mass (kg)	20.9±8.9	37.7±10.3	<0.001
Body fat (%)	26.3±6.2	44.0±4.8	<0.001
Fat mass index (kg/m^2)	7.1±3.0	15.1±4.1	<0.001
Trunk (% FM)	46.9±5.3	43.5±5.7	<0.001
Leg (% FM)	40.8±5.0	43.9±6.2	<0.001
Arm (% FM)	12.4±1.3	12.6±1.8	0.055
VAT (cm^2)	91.9±47.4	109.6±44.7	<0.001
SAT (cm^2)	215.8±129.5	474.6±144.3	<0.001
VAT/SAT	0.50±0.19	0.24±0.09	<0.001
Glucose and insulin measures (n=804)			
HbA1c (%)	5.8±1.1	6.3±1.4	<0.001
Fasting glucose (mmol/L)	5.3±1.5	5.5±2.0	0.057
2 h glucose (mmol/L) (n) (n=735)	6.1±2.6	6.6±2.7	0.009
iAUC for glucose (mmol/L) (n=735)	177 (72 – 297)	157 (79 – 262)	0.368
Fasting insulin (mIU/ml)	5.9 (2.3 – 11.9)	9.4 (5.2 – 15.2)	<0.001
iAUC for insulin (mIU/ml) (n=735)	4 132 (2526–7304)	4 692 (3080–7216)	0.021

Fasting C-peptide (ng/ml)	1.84±1.07	2.09±1.17	0.002
iAUC for C-peptide (ng/ml) (n=735)	641 (447 – 925)	638 (465 – 875)	0.915
HOMA–IR	1.37 (0.51 – 2.81)	2.11 (1.13 – 3.69)	<0.001
Matsuda index (mgl^2/mU min) (n=734)	7.1 (3.6 – 13.2)	5.0 (3.1 – 8.4)	<0.001
Insulinogenic Index (mIU/mmol) (n=624)	16.9 (8.3 – 33.0)	23.4 (12.7 – 43.4)	0.001
C-peptide Index (ng/mmol) (n=612)	2.25 (1.27 – 3.79)	2.73 (1.56 – 4.57)	0.002
oDI (mIU/mmol) (n=644)	13.74 (7.22 – 26.09)	13.40 (5.98 – 28.08)	0.917
Basal insulin clearance (ng/mIU)	0.28 (0.20 – 0.39)	0.20 (0.15 – 0.27)	<0.001
Postprandial insulin clearance (ng/mIU) (n=698)	0.18 (0.14 – 0.25)	0.14 (0.12 – 0.17)	<0.001
Glucose tolerance status[1]			
NGT: (FPG <6.1 / 2-h PG <7.8mmol/L)	263 (67.8)	261 (62.7)	
IGM: (FPG: 6.1–6.9 / IGT 2hPG: 7.8–11.0)	82 (21.1)	91 (21.9)	0.163
T2DM: (FPG ≥7/2-h PG≥11.1/diabetes medication)	43 (11.1)	64 (15.4)	

Values expressed as mean±SD, median (25[th] – 75[th] percentile) or n (percentage). BMI: body mass index, WHR: waist-hip ratio, DXA: dual x–ray absorptiometry, VAT: visceral adipose tissue, SAT: subcutaneous adipose tissue; HbA1c: glycated haemoglobin, iAUC: integrated area under the curve, HOMA-IR: Homeostatic Model Assessment of Insulin Resistance, NGT: normal glucose tolerance, IGM: impaired glucose metabolism, T2DM: type 2 diabetes mellitus.

Matsuda Index: measure of insulin sensitivity (21,22); Insulinogenic Index: measure of insulin response: $\Delta I_{30}/\Delta G_{30}$ (23); C-peptide Index: measure of insulin secretion: $\Delta CP_{30}/\Delta G_{30}$ (23); oDI: oral disposition index, measure of beta-cell function, calculated as product of C-peptide index and Matsuda index (26–28); Glycaemic tolerance status[1]: definition and diagnosis of diabetes and intermediate hyperglycemia: report of a WHO/IDF consultation

Table 2: Glucose and insulin measures in black SA men and women adjusted for FMI

Variable, n	Adjusted for FMI		
Glucose and insulin measures	**Men**	**Women**	**p-value**
HbA1c (%) (n=761)	6.0 (5.8, 6.1)	6.1 (6.0 – 6.3)	0.256
Fasting glucose (mmol/L) (n=763)	5.5 (5.3 – 5.7)	5.2 (5.0 – 5.5)	0.160
2 h glucose (mmol/L) (n=697)	6.5 (6.2 – 6.9)	6.0 (5.7 – 6.4)	0.116
iAUC for glucose (mmol/L) (n=697)	221 (198 – 244)	163 (140 – 185)	0.003
Fasting insulin (mIU/ml) (n=761)	12.3 (11.1 – 13.5)	7.8 (6.6 – 8.9)	<0.001
iAUC for insulin (mIU/ml) (n=697)	6387 (5842 – 6932)	4704 (4165 – 5244)	0.001
Fasting C-peptide (ng/ml) (n=761)	2.29 (2.16 – 2.41)	1.59 (1.47 – 1.72)	<0.001
iAUC for C-peptide (ng/ml) (n=698)	746 (631 – 861)	895 (781 – 1009)	0.123
HOMA–IR (n=761)	3.03 (2.62 – 3.44)	2.06 (1.66 – 2.45)	0.005
Matsuda Index (mgl^2/mU min) (n=696)	6.3 (5.5 – 7.2)	9.8 (8.9 – 10.6)	<0.001
Insulinogenic index (mIU/mmol) (n=659)	27.4 (16.6 – 38.3)	43.3 (32.3 – 54.2)	0.089
C-peptide index (ng/mmol) (n=613)	2.75 (1.43 – 4.07)	6.91 (5.60 – 8.22)	<0.001
oDI (mIU/mmol) (n=613)	14.0 (1.7 – 26.3)	62.3 (50.0 – 74.5)	<0.001
Basal insulin clearance (ng/mIU) (n=761)	0.27 (0.25 – 0.28)	0.28 (0.26 – 0.29)	0.448
Postprandial insulin clearance (ng/mIU) (n=698)	0.19 (0.18 – 0.21)	0.18 (0.17 – 0.19)	0.403

Data adjusted for fat mass index (FMI) presented as median (95% Confidence Interval);

HbA1c: glycated haemoglobin,

iAUC: integrated area under the curve,

HOMA-IR: Homeostatic Model Assessment of Insulin Resistance,

Matsuda Index: measure of insulin sensitivity (24, 25),

Insulinogenic Index: measure of insulin response: $\Delta I_{30}/\Delta G_{30}$ (26),

C-peptide Index: measure of insulin secretion: $\Delta CP_{30}/\Delta G_{30}$ (26),

oDI: oral disposition index, measure of beta-cell function, calculated as the product of the C-peptide index and Matsuda index (29-31)

Table 3: Associations between regional adiposity z-scores and risk for IGM and type 2 diabetes in men and women combined

Men and women	RRR	95% CI	p-value	Model R²
Trunk z-score (n=761)				
IGM	2.35	1.43 – 3.87	0.001	<0.001
Type 2 diabetes	4.76	2.68 – 8.45	<0.001	
Leg z-score (n=759)				
IGM	0.42	0.24 – 0.71	0.001	<0.001
Type 2 diabetes	0.21	0.11 – 0.39	<0.001	
Arm z-score (n=759)				
IGM	1.35	0.83 – 2.21	0.224	<0.001
Type 2 diabetes	2.19	1.25 – 3.81	0.006	
VAT z-score (n=753)				
IGM	1.76	1.35 – 2.28	<0.001	<0.001
Type 2 diabetes	2.58	1.92 – 3.48	<0.001	
SAT z-score (n=753)				
IGM	1.35	0.74 – 2.48	0.331	<0.001
Type 2 diabetes	1.37	0.68 – 2.75	0.376	

Results of multinomial logistic regression presented as relative risk ratios (RRR) and 95% confidence interval and represent risk of outcome with 1 SD increase in regional adiposity. Model used normal glucose tolerance (NGT) as the reference group compared to impaired glucose metabolism (IGM) and type 2 diabetes, adjusted for: age, smoking, alcohol intake, education attainment, FMI and sex

Table 4: Associations between regional adiposity z-scores and insulin sensitivity, basal insulin clearance and Beta-cell function

	ß	95% CI	p-value	Model R^2	Model p-value	ß	95% CI	p-value	Model R^2	Model p-value
	Men					Women				
	Insulin Sensitivity (n=344)					Insulin Sensitivity (n=349)				
FMI z-score[#]	-3.442	-4.011 to -2.873	<0.001	0.284	<0.001	-1.225	-1.631 to -0.819	<0.001	0.069	<0.001
Trunk z-score [#]	-2.680	-4.829 to -0.532	0.015	0.295	<0.001	-1.748	-2.545 to -0.950	<0.001	0.097	<0.001
Leg z-score	1.695	0.233 to 3.157	0.023	0.293	<0.001	1.218	0.454 to 1.982	0.002	0.086	<0.001
Arm z-score[#]	0.027	-1.667 to 1.722	0.975	0.284	<0.001	-0.893	-1.716 to -0.070	0.033	0.080	<0.001
VAT z-score[#]	-1.679	-2.540 to -0.818	<0.001	0.309	<0.001	-1.174	-1.657 to -0.691	<0.001	0.106	<0.001
SAT z-score[#]	-1.043	-2.961 to 0.876	0.286	0.286	<0.001	-0.693	-1.582 to 0.195	0.126	0.075	<0.001
	Beta-cell function (n=304)					Beta-cell function (n=306)				
FMI z-score[#]	-3.9594	-5.3600 - -2.5589	<0.001	0.065	<0.001	-2.6420	-4.0684 to -1.2156	<0.001	0.027	0.001
Trunk z-score	-4.0597	-9.2112 to 1.0917	0.122	0.068	<0.001	-4.3905	-7.2286 to -1.5524	0.003	0.040	<0.001
Leg z-score[#]	3.182	-0.3294 to 6.6940	0.076	0.071	<0.001	4.3120	1.7781 to 6.8460	0.001	0.044	<0.001
Arm z-score[#]	0.1387	-3.8845 to 4.1618	0.946	0.065	<0.001	-0.5693	-3.3668 to 2.2282	0.689	0.028	0.001
VAT z-score	-2.6180	-4.7016 to 0.5344	0.014	0.074	<0.001	-3.8567	-5.5820 to -2.1314	<0.001	0.051	<0.001
SAT z-score	1.6166	-2.9625 to 6.1957	0.488	0.066	<0.001	-1.8141	-4.7217 to 1.0936	0.220	0.029	0.001
	Basal Insulin Clearance (n=346)					Basal Insulin Clearance (n=351)				
FMI z-score[#]	-0.047	-0.062 to -0.033	<0.001	0.121	<0.001	-0.027	-0.035 to -0.018	<0.001	0.092	<0.001
Trunk z-score[#]	-0.055	-0.109 to -0.001	0.047	0.129	<0.001	-0.026	-0.044 to -0.009	0.002	0.108	<0.001
Leg z-score	0.006	-0.031 to 0.044	0.738	0.121	<0.001	0.017	0.001 to 0.033	0.036	0.099	<0.001
Arm z-score[#]	-0.015	-0.058 to 0.027	0.477	0.122	<0.001	-0.022	-0.038 to -0.005	0.011	0.104	<0.001
VAT z-score	0.001	-0.022 to 0.023	0.945	0.121	<0.001	-0.018	-0.028 to -0.008	0.001	0.111	<0.001
SAT z-score[#]	-0.033	-0.082 to 0.015	0.175	0.125	<0.001	-0.005	-0.024 to 0.013	0.555	0.093	<0.001

Beta coefficients for robust regression models for men and women, adjusted for age, smoking, alcohol intake, education attainment and FMI (except for FMI z-score), VAT: visceral adipose tissue, SAT: subcutaneous adipose tissue,
Insulin sensitivity estimated from the Matsuda Index (24, 25) and beta-cell function was estimated using the oral Disposition index calculated as the product of C-peptide and Matsuda Index (26), and basal insulin clearance calculated as the ratio of fasting C–peptide to fasting insulin
[#]$p<0.05$ for sex*z-score body fat interaction term

Figure 1: Bar Graph of the relative risk ratio (RRR) of impaired glucose metabolism (IGM) and type 2 diabetes mellitus compared to the normal glucose tolerant (NGT) in men and women, IGM: (RRR (95%CI): 1.70 (1.27–2.29), p<0.001 vs. 1.23 (0.95–1.60), p=0.115) and T2D: (2.05 (1.42–2.96), p<0.001 vs. 1.38 (1.03–1.85), p=0.031) for men and women, respectively (A); Sex-specific associations between FMI z-scores and insulin sensitivity (Matsuda index) (B), beta-cell function (oral disposition index) (C) and basal insulin clearance (D), modelled as predictive margins of sex with 95% CI