Cluster detection with random neighbourhood covering: application to invasive Group A Streptococcal disease

Massimo Cavallaro[1,2,3*], Juliana Coelho[3], Derren Ready[3,6], Valerie Decraene[3], Theresa Lamagni[3], Noel D. McCarthy[1,4,5], Dan Todkill[3,4], Matt J. Keeling[1,2]

[1] The Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research, University of Warwick, Coventry, CV4 7AL, United Kingdom.

[2] School of Life Sciences and Mathematics Institute, University of Warwick, Coventry, CV4 7AL, United Kingdom.

[3] National Infection Service, UK Health Security Agency, United Kingdom.

[4] Warwick Medical School, University of Warwick, Coventry, CV4 7AL, United Kingdom.

[5] Institute of Population Health, School of Medicine, Trinity College Dublin, University of Dublin, 2 Dublin, Ireland.

[6] Health Protection Research Unit in Behavioural Science and Evaluation at the University of Bristol, Bristol, UK

*m.cavallaro@warwick.ac.uk

Abstract

The rapid detection of outbreaks is a key step in the effective control and containment of infectious diseases. In particular, the identification of cases which might be epidemiologically linked is crucial in directing outbreak-containment efforts shaping the intervention of public health authorities. Often this requires the detection of clusters of cases whose numbers exceed those expected by a background of sporadic cases. Quantifying exceedances rapidly is particularly challenging when only few cases are typically reported in a precise location and time. To address such important public health concerns, we present a general method which can detect spatio-temporal deviations from a Poisson point process and estimate the odd of an isolate being part of a cluster. This method can be applied to diseases where detailed geographical information is available. In addition, we propose an approach to explicitly take account of delays in microbial typing. As a case study, we considered invasive group A Streptococcus infection events as recorded and typed by Public Health England from 2015 to 2020.

Introduction

Disease surveillance is the rapid collection and analysis of data to identify public health priorities, develop targeted response, and monitor the effects of interventions [1]. The

importance of surveillance in controlling the spread of diseases is highlighted by several
recent challenges, ranging from the fast evolving COVID-19 pandemic [2], to hospital-
acquired anti-microbial resistant infections [3], to the burden of chronic non-communicable
diseases [4].

For decades, statistical principles and methods have been leveraged by public health officials
to identify increases in diseases and raise early warnings where necessary. The effectiveness
of these methods has been improved by the availability of large digital databases, which can
store infection records and quickly be interrogated and processed [5]. Thanks to these
advances, syndromic surveillance is performed daily in the UK based on common symptoms
such as cough, diarrhoea, or fever through general practitioner consultations [6-8]. In
addition, the surveillance of microbiological diagnosis of specific pathogenic organisms is of
key public health importance. Most of the existing statistical models for surveillance are
tailored to dense temporal data streams, consisting of tens or hundreds of health records per
week or day [9-15], although methods based on the so-called scan statistics [16] have good
performance in detecting geographical areas with small confined outbreaks [17-25].

Streptococcus pyogenes, also known as group A Streptococcus (GAS), is a Gram-positive
human pathogen that can colonise the throat and skin of humans. Asymptomatic throat
carriage rates in the general population are reported to be around 7.5% in high-income
countries [26]. Complex host-pathogen interactions in GAS infections can cause a broad
range of symptom severity – from mild or asymptomatic infections to death. The most severe
disease occurs when these bacteria invade parts of the body where they are usually not
found (such as the blood, the muscles, or the lungs) and cause invasive infections, which
requires hospitalisation and can lead to death [27]. An important part of the epidemiological
surveillance of invasive GAS (iGAS) infections is the typing of collected bacterial isolates. A
widely used typing method for GAS is based on the sequence of the N-terminal part of the M-
protein gene (*emm*), which is a highly polymorphic gene and allows us to distinguish strains
of *S. pyogenes* which would otherwise look identical. So far, more than 200 *emm* types have
been identified in Europe and North America, with few common types contributing to a high
percentage of the total number of iGAS cases [28]. In the UK, iGAS infections are notifiable
and isolated from iGAS cases are sent to the national reference laboratory for *emm* typing,
enabling the identification of cases which share the same *emm* gene and which could denote
an outbreak. Non-invasive GAS isolates are rarely typed outside of an outbreak investigation.
iGAS disease is of particular concern due to the recent upsurge in the number of cases in
some countries [29] and potential for spread amongst marginalized groups [30] or in
community nursing teams [31]. In these latter two settings, outbreaks can take place across
wide geographic areas and are therefore more difficult to detect using standard public health
investigations.

To help identify clusters of iGAS which share the same *emm* type and are epidemiologically
linked by geography and in time, we developed the random neighbourhood covering
(RaNCover) approach. As an effective and automated algorithm, it would be complementary
to what local health protection or public health teams can offer in the initial stages of
outbreak detection, and may be of particular use with outbreaks which cross regional or
team borders that might otherwise be difficult to detect. The general problem RaNCover
addresses is to classify a single recorded infection as being part of a cluster or being

consistent with a baseline of sporadic cases. To this end, we consider a neighbourhood of the point location and time of the recorded case and appropriately draw a random collection of spatio-temporal cylinders to cover it. Based on the properties of this collection of covering cylinders it is possible to perform classification. RaNCover is specifically tailored to diseases where detailed spatio-temporal information is available and aggregating data in coarse-grained bins (which results in less detailed information) is not desirable. As proof of concept, we deployed our framework on *emm* typed iGAS infections recorded in England.

The manuscript is organised as follows. In section *Methods* we introduce the outbreak-detection principles, define the RaNCover approach, illustrate it on simulated data, introduce the *emm* typed iGAS data, and derive the endemic baseline for this disease. The latter is used to generate synthetic data and perform additional simulation experiments for validation. In the section *Results,* RaNCover is retrospectively and prospectively applied to the iGAS data and the detected clusters are discussed with the help of a dimensionality reduction technique, which allows us to conveniently present the results. In the same section, we tackle *emm* typing delay. We conclude with a *Discussion* section.

Methods

Null model and outbreak detection

A spatio-temporal disease pattern can often be decomposed into two components, endemic and epidemic — although such a distinction can be blurred [32]. The endemic component explains a baseline rate of cases with a persistent yet stable pattern, only reinforced by immigration [33]. The epidemic element can be thought of as realizations of processes with auto-regressive or self-exciting behavior, which arise from a common source, e.g., environmental exposure, increased susceptibility in the population, or invasion of a new pathogen. In other words, the presence of epidemic events is correlated with and reinforces the occurrence of successive events [32].

We argue that the most important characteristic common to all epidemic data is that observations are not independent. Therefore, to state whether cases are observed in atypical frequency, we must compare our observations with those expected if the cases were uncorrelated in both space and time. If the recorded case observations are consistent with such an uncorrelated baseline, no warning should be raised. This suggests that the first step of an outbreak detection system is to develop a naïve model for the observed cases — a *null model* — which does not consider any spatial or temporal correlations. The second step is then to detect (statistically significant) deviations of the observed pattern from that produced by the null model.

We let $X(A)$ denote the random variable that describes the number of cases detected in a region of space and time ($A \subset S$) under the null model (X) and define a threshold value θ_A such that

$$\text{Prob}(X(A) > \theta_A) = \alpha. \qquad (1)$$

A basic exceedance criterion therefore consists of triggering a warning flag if the number of cases observed over A exceeds a value θ_A. Here, α is the false-positive detection rate, which is the probability of triggering a warning by chance when no outbreak has occurred, and the set $\mathcal{S}$ represents the entire extent of the spatio-temporal data. The approach described so far is not new [5, 34] and can be regarded as the standard in statistical process control [35].

The random neighbourhood covering approach

We assume that the null model is an inhomogeneous Poisson point process with intensity $\lambda(x, y, t)$, where x and y are spatial coordinates and t is the time. We stress that the rate parameter λ is not necessarily constant in space or time. Indeed, even in the absence of localized outbreaks, we may expect many episodes in highly populated regions and at certain times of the year, as many diseases are known to be influenced by seasonality while weekends or holidays can impact the daily recording practice [10, 13] [36]. A Poisson point process across space and time ($\mathcal{S}$) is defined by the following two properties:

1. $X(A_1)$ and $X(A_2)$ are independent for disjoint subsets A_1 and A_2 of $\mathcal{S}$;
2. For any subset $A \subset \mathcal{S}$, $X(A)$ has Poisson distribution

$$\text{Prob}(X(A) = n) = \Lambda(A)^n e^{-\Lambda(A)}/n!, \qquad (2)$$

$$\Lambda(A) = \int_A d\xi\, \lambda(\xi),$$

where $\xi = (x, y, t)$ and $\Lambda(A)$ is referred to as the intensity measure [37]. Given the intensity function, it is straightforward to numerically obtain $\Lambda(A)$, find the αth quantile θ, and state whether the observed number n exceeds this value in the subset A. Those subsets such that the observed event number exceeds θ are tagged as outbreak subsets. For simplicity, it is convenient to treat the time t as an integer variable rounding it up to, e.g., days or weeks, thus $\mathcal{S}$ is $\mathbb{R}^2 \times \mathbb{N}$, its generic subset A is the union $\cup_{i \in \mathbb{N}} A_i$, $A_i = \{(x, y, t) \in A | t = i\}$, and the intensity measure can be easily computed as sum of two-dimensional integrals, i.e., $\Lambda(A) = \sum_i \int_{A_i} dx dy \lambda(x, y, i)$.

The intensity function λ represents the average trend of the endemic infections. We assume that it factorises into a spatial factor λ_0 and a temporal factor λ_{time}, i.e., $\lambda(x, y, t) = \lambda_0(x, y)\, \lambda_{\text{time}}(t)$, thus excluding any correlation between the space and time trends. We make the simplifying assumption that $\lambda_0(x, y)$ is proportional to the population density, reflecting the fact that the number of observed episodes in an area is likely to increase with its population size – although $\lambda_0(x, y)$ could be matched to historical infection data if these were plentiful. This population-based weighting is arguably the most important factor in the spatial distribution of the episodes, although, across geographical areas, other factors such as the climatic zone or disparities in access to health care could affect the baseline. The temporal factor encodes the disease seasonality, and other long-term trends, and is generally obtained by fitting a periodic function to available data.

The inequality (1) provides a basic exceedance criterion for flagging a subset A (which encompasses a geographical region for a certain time interval) but is far from being ideal. Indeed, a warning is raised with probability α even if the data is actually generated by the baseline model $X(A)$. Further, the aberration can only be tested on the subset A and

outbreaks diffused over a larger area are likely to be missed [38]. To address these issues we
consider, for each reported case (of given space-time coordinates $\mathbf{x} \in \mathcal{S}$), N overlapping
subsets $A_i \subset \mathcal{S}, \mathrm{i} = 1,2,\cdots,N$, each of which contains the case ($\mathbf{x} \in A_i$) and apply the criterion
of (1) to each one of them in order to test whether there is aberration from the baseline
Poisson point process (Fig 1). In the absence of any aberrations and in agreement with
property 2, the number of detected cases in any subset A_i has distribution (2). If its intensity
measures $\Lambda(A_i)$ are the same for all the subsets, we expect just a fraction α of them to exceed
the threshold θ_{A_i} (while in the presence of epidemic events the fraction is greater than α).
Throughout this manuscript, we stick to the standard false discovery rate $\alpha = 0.05$ for each
single subset. It is worth noting that, by considering the state of multiple subsets, we increase
robustness to random fluctuations and make highly unlikely to identify a cluster from
random case data. In the null model, we indeed expect just 5% of independent subset to have
a warning flag and even with just 20 subsets, the probability of generating a false flag is very
small (probability of 19 out of 20 subsets having a flag is 3.6×10^{-24}).

For a detected case $\mathbf{x}$, we refer to the proportion $w(\mathbf{x})$ of flagged subsets $A \ni \mathbf{x}$ as its *warning*
score. By taking this approach, we are able to go beyond the binary warning flag for an
outbreak event, and hence present $w(\mathbf{x}) \in [0,1]$ as a continuous warning score specific to a
detected case, high values of w corresponding to a high probability that an outbreak is
occurring. This suggests raising a warning if $w(\mathbf{x})$ is significantly greater than a chosen
threshold, where the statistical significance is tested by means of a binomial test for
proportions. Wilson intervals [39] are used as 2.5\%-97.5\% confidence intervals (CIs) for
the warning scores. As N increases, these CIs tighten and the test significancy improves thus
allowing practitioners to tune the parameter N and reach their desired significance level.

It is convenient to focus on subsets placed over detected cases, thus testing for aberration
only where necessary (Fig 1). This obviously implies that every subset always contains at
least one point event and its null-model prediction is shifted to $1 + X(A)$. We opt to select
cylinders of the same volume $\int_A \mathrm{d}\,\xi$. In order to allow cylinders of the same volume to enclose
outbreaks highly diffused in space or lasting long in time, we randomly generate cylinders of
variable radia ϱ and heights h, still subject to the constraint that their volume $\pi \varrho^2 h$ is
constant. The algorithm appears to be robust to the choice of this volume (see simulation
experiments); its value is heuristically set to avoid excessively small volumes, expected to
contain only one event, and excessively large volumes, which would contain more cases but
might not represent well local information.

It is worth noting that a cylinder can accommodate many point events and nearby events are
likely to be enclosed by the same cylinder. Consequently, the warning scores are correlated
and their values reflect the neighbourhood structure of the detected cases, as demonstrated
in the case study of section *Results*. Also, in simulation experiments performed for validation,
clusters are detected as groups of cases with high warning scores. To summarise, for a single
infection case $\mathbf{x}$, RaNCover returns a warning score $w(\mathbf{x})$ that measures to what extent its
presence alters the local count statistics from those of a Poisson point process of intensity λ.
An appropriate baseline intensity needs to be estimated or given as an input.

We now illustrate the method in action by means of synthetic data and baseline. To this end,
we assume the baseline intensity $\lambda = \lambda_0(x,y)\lambda_{\mathrm{time}}(t)$ with

$$\lambda_0(x, y) = \left(C + D\, e^{-E((x-0.5)^2+(y-0.5)^2)}\right), \quad (3)$$

$$\lambda_{\text{time}}(t) = \left(\frac{A}{2}(\sin(t) + 1) + B\right) \qquad (4)$$

over a unit square region with parameters $(A, B, C, D, E) = (7, 0.9, 0.5, 20, 80)$ where x and y are spatial coordinates, while t is the time index ranging from 1 to 20. This describes a spatial distribution of cases peaked at the centre of the region, modulated by a periodic temporal trend. For each value of t we simulate a spatial Poisson point process with intensity λ, which yields a random pattern of points representing endemic baseline episodes. An outbreak in a delimited polygonal region is simulated by randomly generating additional Poisson point events with intensity $50\,\lambda_{\text{time}}(t)$ for $t = 5,6,7,8,15,16,17,18$ and zero elsewhere. Spatial Poisson processes are generated using the thinning method [40]. The resulting point pattern is illustrated in Fig 2A for the first 9 time steps. To identify the events corresponding to the simulated outbreak in this point pattern we apply RaNCover as follows. We draw $N = 10000$ random covering cylinders of equal volume, with radia $\rho \cong (0.10, 0.12, 0.14, 0.17, 0.23)$ and height $h = (1,2,3,4,5)$. The chosen volume is $\pi\rho^2 h \cong 0.17$, which corresponds to about 5 expected baseline events per cylinder. The centre coordinates are randomly sampled with replacement from the episode locations and times, and randomly shifted whilst ensuring that each cylinder contains its assigned episode location. We then flag all the cylinders according to the basic exceedance criterion with $\alpha = 0.05$. The classification performed by the algorithm is summarised by the warning scores, computed using this set of cylinders. These warning scores indeed appear to capture the synthetic outbreak, as illustrated in Fig 2B.

Performance metrics

Excellent performance is confirmed by the receiver operator characteristic (ROC) curves (Fig S1 A) and the area under the curve (ROC-AUC 0.90, 95%CI 0.85-0.95 by means of bootstrap). The warning scores obtained upon increasing the volume of the cylinders by 40% achieve close performances (ROC-AUC 0.92, 95%CI 0.87-0.96) and are strongly correlated with the first replicate output (Pearson $\rho = 0.95$, P=1×10^{-83}), thus suggesting that the result is robust to the choice of the cylinder size (Fig S1 B-C).

In a more realistic surveillance scenario, the warning scores are prospectively updated as soon as new cases are detected and included into the dataset, which we take to occur at every point in time. Observing the individual warning scores varying as new observations are included allows us to assess timeliness. This measures the time elapsed between the case detection day and the day at which its warning score stabilises around its final value (corresponding to the retrospectively computed warning score). To simulate this, we iteratively apply the procedure detailed above to all the observations occurred until time τ, $\tau = 1, 2, \dots, 20$ and observe how the warning scores of cases detected previous to the update time τ vary with τ, while new cases are included and assessed. Fig 3 shows this process in action, the three panels show the warning scores for outbreak and endemic cases (as defined by the Poisson intensity function) that are detected at times t=5, t=6, and t=7. For this simulation, the values of the warning scores appear to converge after only one or two updates, which suggests that the method can reach excellent timeliness.

Defining iGAS background rate and reporting delay

Our study cohort consists of 10,820 iGAS records comprising 465 different *emm* subtypes, with the three most prevalent types (viz., `1.0`, `89.0`, and `12.0`) accounting for more than 38% of all records, while other *emm* types occur one order of magnitude less frequently (Fig 4). A total of 235 different *emm* subtypes were recorded only once in the cohort.

We estimated the endemic baseline for iGAS as follows. As streptococcal isolates of different *emm* types are thought to belong to different endemic populations, they cannot form a single outbreak even when they occur in the same geographical location and at the same time [29]. To encode such mutual independence, the baseline intensity $\lambda_M(x, y, t)$ specific to the *emm* type M is obtained by multiplying $\lambda(x, y, t)$ by the relative frequency λ_m of M in the population, computed retrospectively until time τ. In other words, the null-model intensity function is assumed to obey

$$\lambda_M(x, y, t) = \lambda_m \, \lambda_0(x, y) \, \lambda_{\text{time}}(t). \quad (5)$$

The temporal and spatial factors (λ_{time} and λ_0, respectively) remain the same for all *emm* types. As the incidence of iGAS infection is subject to seasonality [27], $\lambda_{\text{time}}(t)$ is chosen to be periodic with a period T of one year, viz.,

$$\lambda_{\text{time}}(t|\vartheta_\tau) = a + b\,t + (c + b\,t)\sin(2\pi d/T + 2\pi t/T), \quad (6)$$

where $\vartheta_\tau := (a, b, c, d)$ are parameters that must be retrospectively estimated from the data until time τ. To determine these parameters, we use the principle of maximum likelihood estimation (MLE). We denote by $t_1, t_2, \ldots, t_n$ the ordered times of all cases detected until τ, $0 < t_1 < t_2 < \cdots < t_n < \tau$, regardless of their *emm* types and geographical locations. The parameter τ identifies the time interval used to estimate the temporal trend of the baseline; for iGAS this has to be long enough to capture seasonality, with longer τ arguably producing better estimates. The likelihood function $L(\vartheta_\tau)$ under the Poisson null model is given by

$$\log L(\vartheta_\tau) = -\int_0^\tau \lambda_{\text{time}}(u|\vartheta_\tau)\mathrm{d}u + \sum_{i=0}^n \log \lambda_{\text{time}}(t_i|\vartheta_\tau).$$

The value of ϑ_τ that maximises the likelihood was recursively obtained by maximising over a, c, d using the Nelder-Mead method with b held fixed and then finding the global maximum over b with the other parameters held fixed.

The episodes in the database are labelled by patient postcode location. For each postcode, the estimated population size is obtained from the 2011 Census in England and Wales by the Office for National Statistics (ONS), which allows us to tabulate the values of the spatial baseline intensity $\lambda_0(x, y)$. The corresponding spatial coordinates are taken to be the postcode centroid locations, also retrieved from the ONS. This completes the definition of the endemic intensity function $\lambda(x, y, t)$.

The available iGAS records also include the week at which the original sample was taken, which, on average, occurred a week earlier than the laboratory *emm* typing. The time elapsed between the sample date and the *emm*-type report is referred to as the reporting delay. We

generalise RaNCover to include untyped cases by introducing the baseline intensity $\lambda_U :=$
$\lambda_u(t; \tau)\lambda_{\text{time}}(t)\lambda_0(x, y)$ of untyped cases at each update time τ, where $\lambda_u(t; \tau)$ is the average
fraction of cases detected at time t and *emm* typed after time τ. For a generic spatio-temporal
subset A, we refer to the number of typed observations, the typed baseline prediction, the
number of untyped observation, and the untyped baseline prediction as $n_M(A)$, $X_M(A)$,
$n_u(A)$, and $X_u(A)$, respectively. The next step is to define a strategy based on the standard
exceedance criterion (1). We flag A if $n_M(A)$ exceeds the threshold value $\theta_{M,A}$ or $n_u(A)$
exceeds the value $\theta_{u,A}$, i.e.,

$$(n_M(A) > \theta_{M,A}) \vee (n_u(A) > \theta_{u,A}),$$

with thresholds defined by $\text{Prob}(X_M(A) > \theta_{M,A}) = \alpha$ and $\text{Prob}(X_u(A) > \theta_{u,A}) = \alpha$,
respectively. $X_M(A)$ and $X_u(A)$ are assumed to be Poisson point processes of intensities
$(1 - \lambda_u(t; \tau))\lambda_M(x, y, t)$ and $\lambda_U(x, y, t; \tau)$, respectively. The warning score of an untyped case
$\mathbf{x}$ is the fraction $w(\mathbf{x})$ of subsets $A \ni \mathbf{x}$ flagged accordingly to this criterion.

Ethics statement

288 The data used in this study were supplied from the UK Health Security Agency (UKHSA)
289 under strict data protection protocols agreed between the University of Warwick and
290 UKHSA. The ethics of the use of these data for these purposes was agreed by UKHSA.

Results

Simulation study

Benchmarking surveillance systems benefits from realistic simulations of health data [41,
42]. As described in *Methods*, we assume that, in the absence of outbreaks, the detected cases
are a realization of a generative Poisson point process with factorised intensity function
$\lambda_0(x, y)\lambda_{\text{time}}(t)$. Accordingly, we simulated the point process with intensity defined by
equations (2) and (3) and parameters as estimated in section *Methods*, thus yielding a
synthetic baseline of 4915 endemic cases (distributed over a period of 100 weeks in
England) that should not raise any public-health concerns. The temporal intensity function
of equation (6) fitted to all data in the study period is illustrated in Fig 5. We simulated an
outbreak by including additional (epidemic) point events localised in 19 selected postcode
areas (arbitrarily chosen around the town of St. Albans, north of Greater London) during a
chosen time interval (weeks 40 to 59). In order to imitate the initial spread of infections and
their subsequent control, the total number of epidemic events per week is initially low,
increases and peaks around week 50, and finally decreases until the number of epidemic
cases drops to zero (Fig 6 A). Aggregating the cases over their geographic locations clearly
shows the temporal trend as illustrated in Fig S2 A for the endemic cases and in Fig S2 B for
both endemic and epidemic cases. Based on these time series, it is virtually impossible to
detect excess cases during the outbreak. Analogously, aggregating over the time yields the
geographical distributions of cases, which mainly occur alongside the most populated area
(Fig S2 C) and show no appreciable changes if the outbreak in St. Albans is included (Fig S2

D). Yet, whether a single case is part of the outbreak or not is clearly labelled in the synthetic
dataset, thus making it ideal for evaluation purposes; this typically is not the case in real data.

The set of these simulated data was used to test whether RaNCover can detect the epidemic
components in it. RaNCover for retrospective analysis returns a warning score $w(\mathbf{x})$ for each
case $\mathbf{x}$. We set $\alpha = 0.05$, $N = 1000000$, and cylinder volumes $\cong 2605\ km^2 \times week$, which
overall yielded tight CIs (Fig S3). In Fig 6B, each warning score is plotted vs the case detection
time t. In this figure, the markers are coloured in blue if the case was simulated from the
baseline and red if simulated from the outbreak. It is possible to appreciate that the baseline
events have low warning scores within statistical fluctuations. The bulk of the outbreak
events always had warning scores close to one and can be unambiguously discriminated
from the baseline based on $w(\mathbf{x})$. Unsurprisingly, few cases at the tails of outbreak can be
confused with background, but their warning scores reach relatively large values ($w(\mathbf{x}) >$
0.8). The ROC-AUC is 0.99 (95% CI 0.98-1.00), thus suggesting excellent performances. In Fig
6C, we show that the method can also identify the geographic location of the outbreak with
precision. The result appears robust to the choice of the cylinder volume, as suggested by the
fact that the warning scores calculated from a second replicate where cylinder volume was
increased by 40% are strongly correlated with the first replicate (Fig 6D).

As a further test we sequentially calculated the warning scores prospectively, using only the
data simulated until the week τ, $\tau = 40,41,\cdots,59$. As time progresses and new cases are
detected, the warning scores are updated, thus allowing us to observe how quickly the
outbreak is detected. The warning-score traces for the epidemic cases are plotted in Fig 6E,
where it is possible to appreciate that, for the episode that occurred at t=40 (which is the
first outbreak event and appears isolated at first, see Fig 6A), the warning score is initially
very low and increases when new outbreak cases occur (until it converges to a higher stable
value). Similarly, the other warning scores become stable after few steps.

For comparison, we performed cluster detection with SaTScan™ [43] for retrospective
space-time analysis using the discrete Poisson model [17]. SaTScan™ is a mature software
that computes the scan statistics and is written in Java [43]. For our simulation experiment,
the cylinder with the optimal likelihood ratio encloses 16 weeks (from 43 to 58) and 33
postcodes (Fig 6C, inset). These SaTScan™ results are widely consistent with ours; however,
it is worth noting that SaTScan™ took around 3 hours to evaluate this sample dataset on an
Intel Core i5™ CPU (excluding default Monte Carlo hypothesis testing and not accounting for
seasonality, which would further increase the computational burden), thus significantly
longer than the 5 minutes circa required by our R [44] implementation of RaNCover. As
SaTScan™ gives cylindrical boundaries of a detected cluster, these do not necessarily
separate the outbreak cases from the baseline and, within these boundaries, there are cases
correctly labelled only by RaNCover (Fig. 6C).

Application to iGAS disease: Cluster analysis and data-point embedding

Warning scores for all episodes of selected *emm* types are retrospectively generated
following the procedure detailed in *Methods*, with $N = 20,000$ cylinders per *emm* type with
respect to the baseline intensity of equation (5). The spatial and temporal factors are
obtained from census data and MLE fit, respectively. For each *emm* type considered, the type-

specific factor λ_m is the relative frequencies of cases of that *emm* type (as illustrated in Fig 4
at each update time). As the average baseline intensity governs the cylinders' volumes, these
are differentially set for each *emm* type, the rarer the *emm* type, the larger the volumes.

RaNCover consistently assigns high warning scores to neighbour cases and is thus able to
reveal clusters which may belong to outbreaks. As each event is described by its spatial
coordinates (e.g., latitude and longitude) and its date, the neighbourhood must defined in
both space and time. We use the t-distributed stochastic neighbour embedding (t-SNE)
method to plot these cases (identified by three dimensions, i.e., latitude, longitude, and
detection time) in a two-dimensional space in a way that preserves neighbouring identities
[45]. In other words, t-SNE is a general statistical-learning method for dimensionality
reduction which here is used to pin a two-dimensional point for each point in the three-
dimensional space, in such a way that nearby events in space and time appear next to each
other in the two-dimensional map. Using a visualisation method based on neighbourhoods
rather than on absolute distances is very useful in our context, where the local prevalence of
each *emm* type is important. Indeed, for *emm* types with low prevalence, recorded episodes
that are distant in space and time can be correctly identified as being part of the same
outbreak. Conversely, common *emm* types already established in the host population may
show more clusters in the same geographical area and time scale. We illustrate the results
for 4 rare *emm* types (94.0, 108.1, 44.0, and 33.0), and a common *emm* type that did not
appear to cluster (12.0).

emm type 94.0

emm type 94.0 accounts for 466 invasive cases recorded over the 2015-2020 study period.
Its frequency λ_m relative to all iGAS episodes stabilised in early 2018 after a phase of early
growth (Fig 4B). Applying t-SNE to the episodes' spatial and temporal coordinates yielded
the scatter plot of Fig 7A, which shows a cluster with episodes of high warning scores ($w >$
0.95) in bright red. These episodes are localised in both time (Fig 7B) and space (Fig 7C),
and, specifically, may correspond to a small outbreak in the towns of Bournemouth and
Weymouth (South West region) (Fig 7D) in 2017 (before λ_m reaches its plateau, see Fig 4B).
Fig 7 also suggests the presence of other clusters of cases in the North West region, yet with
$w < 0.95$, which might have triggered investigations but are not retrospectively tagged as of
concern with a warning cut off at 0.95.

emm type 108.1

emm type 108.1, with 270 invasive cases recorded over the whole study period, was less
common than the former. Yet this *emm* type is concerning as its prevalence has been steadily
increasing since 2018 (Fig 4B) and in 2019 it accounted for more confirmed outbreaks than
any other *emm* type [46]. For this *emm* type, RaNCover raised many warnings as the expected
baseline is lowered by the absence of episodes prior to 2018 (Fig 8). The clusters with the
highest warning scores include episodes in South East and South West, West Midlands, East
Midland, and North West. Interestingly, in Yorkshire and the Humber region, we detect the
presence of two clusters both localised in the city of Leeds but separated in time. Another
cluster localised around the river Humber was also detected.

emm type 44.0

emm type 44.0 totals only 163 recorded cases and its relative prevalence is nearly constant in time, except for a small increase in mid-2019 (Fig 4B). Consistently, RaNCover only detected a single cluster of episodes between March and August 2019 localised around the town of Chelmsford in the East of England (Fig 9). The cases in the South East and London are geographically distant but occurred at the same time as the bulk of the main cluster and are therefore mapped nearby the main cluster in Fig 9C.

emm type 33.0

With a total of only 61 invasive cases recorded, the *emm* type 33.0 is uncommon in England. Its relative prevalence has been linearly growing since 2019 (Fig 4B). Similar to 108.1, RaNCover yielded high warning score for almost all detected cases. t-SNE embedding shows that these belong to a single cluster which corresponds to cases diffused in central England with epicenter in the West Midlands (Fig 10).

emm type 12.0

At the other end of the spectrum, the *emm* type 12.0 is relatively common (782 cases over the study period) and its trend has been decreasing since mid-2016. The application of RaNCover highlighted very few points which might raise concern with the majority of the episodes being assigned low warning scores and none being assigned a warning score higher than 0.95 (Fig 11).

Application to iGAS disease: Prospective analysis and *emm* typing delay

The cluster analysis detailed in the previous subsection is performed retrospectively and is based on cases labelled by the week at which the reference laboratory identified the pathogen's *emm* type. As anticipated for synthetic data (Figs 3 and 6 E), it is also possible to observe single-case warning scores varying in time as new cases are *emm* typed and the outbreak progresses. As an example, we focus on the *emm* type 44.0, which appeared to have caused an outbreak in the East of England (Fig 9) with 23 cases retrospectively flagged at the end of the study period (circled red markers in Fig 9 A-B). The warning scores computed at each week are illustrated in Fig 12A. As the first case of the cluster is detected, its score is high yet substantially below the threshold of 0.95. As more cases are detected, it approaches the threshold and sharply increases within the first four weeks (Fig 12B), during which nine cases are identified in the region.

It is possible to combine reporting delays in RaNCover. During the interval from the sampling to the *emm* typing date, we are aware that, at a precise location and time, a case of an *emm* type of concern *might* have occurred. Therefore, only incomplete information is available. Even so, it is desirable to account for this, since it may allow earlier outbreak detection. However, it is possible that an untyped case is not actually associated with the clusters of interest, thus including it into a surveillance system may substantially tamper the overall results. It is important to choose a suitable strategy, which we take to flag a subset if the untyped or typed observation exceeds their respective baselines. Prospectively, as the time progresses, new cases are observed and some previously observed cases are *emm* typed. In

such a scenario, at a given point in time, in addition to the typed iGAS cases and their
baselines (one for each *emm* type) one can collect the untyped cases and compute the
corresponding baseline. As an example, the number of cases per week and the continuous
baselines are plotted against time in Fig 12C for the *emm* type `44.0` -- similar snapshots can
be obtained for any other point in time. The number of untyped cases is highest in
correspondence of the update week and quickly decays to zero as the observation week goes
back in time. Using these two baselines and according to the procedure detailed in *Methods*,
we computed the warning scores relative to a potential *emm* `44.0` outbreak of the untyped
cases at each update week. More specifically, we are interested in the early warning scores
of those cases which are later found to belong to the 2019 cluster. The early warning scores
did not reach the threshold of 0.95, yet they are significantly larger than the baseline warning
scores thus potentially providing early warning signals as illustrated in Fig 12D.

Discussion

In this paper, we have presented the random neighbourhood covering (RaNCover) approach,
a statistical method that can identify outbreaks by detecting deviations from a baseline
Poisson point process. It is based on a simple and scalable approach, i.e., measuring the
expected and observed number of cases in random portions (viz., cylinders) of space and
time. While the statistics in a single random region are not highly informative, aggregating
information over many random regions permits the detection of infectious-disease clusters.
A strength of this approach is that it returns a warning score for each recorded episode. This
makes it ideal in application to diseases which are sparsely distributed, where it is
appropriate to assess a new health event as soon as it is recorded, rather than waiting until
enough cases are collected over a determined region or time interval.

Our method is akin to those based on the so-called scan statistics [16]. They both rely on
creating a large number of spatio-temporal sectors, each with a different set of detected
cases in it. By means of the scan statistics one can identify the sectors with the highest
likelihood ratio, which correspond to the regions where outbreaks are most likely to have
occurred. Conversely, in RaNCover, drawing and inspecting many cylinders covering each
event $\mathbf{x}$ permits defining a continuous warning score w specific to $\mathbf{x}$. Compared to existing
methods, it excels in execution speed and flexibility. One potential limitation of our approach
is that the values of the warning scores depend on the chosen volumes of the spatio-temporal
cylinders and are subject to fluctuations due to the random sampling of these sectors. In fact,
extensive simulations show that, performing replicates with different sector sizes, the
results remain consistent and fluctuations can be controlled by increasing the sampling.

RaNCover can be used for both retrospective and prospective analyses. Retrospective
validation was performed on simulated data using Receiver Operating Characteristic (ROC)
curves. Deriving a binary warning from w requires choosing a cut off, which also governs
sensitivity and specificity. For a given cut off, the sensitivity is defined here as the fraction of
simulation-outbreak cases for which an alarm is generated, while the specificity is defined
as the fraction of baseline cases that did not generate an alarm. The overall performance
quantified by the ROC curve (obtained by plotting the sensitivity against the specificity at
various cut-off settings) and the area under the curve (AUC or C-statistics) were excellent

(ROC-AUCs >0.9 in all simulation scenarios). Fine tuning the cut off also serves as a way of
reducing the number of false alarms or increasing the true positive rate, which can be useful
in real-world surveillance settings.

As a proof-of-concept, we retrospectively detected a number of potential outbreaks of iGAS
infections in England. For this disease, we estimated the baseline taking into account
population denominator data and iGAS seasonality. In the absence of a definitive and
comprehensive list of all real outbreaks that should be detected by public-health
surveillance, it is hard to fully benchmark any surveillance method. In order to circumvent
this latter point and confirm that our method consistently detects events in close proximity
(both in space and time), we proposed to visualise the data using t-distributed stochastic
neighbour embedding (t-SNE) approach. The use of this dimensionality reduction algorithm
on real iGAS data highlights events proximal in both space and time and confirms that
detected cases with high warning scores typically cluster together.

The timeliness was prospectively assessed by tracking how the individual warning scores
vary as new observations are included. It is worth noting that, in general, the timeliness can
be influenced by the speed of the outbreak growth [36]. It can also be influenced by the
outbreak type and its full assessment requires comprehensive testing with different
outbreak scenarios, as performed for established methods, e.g., in references [36, 41, 47, 48].
Reporting delays are also known to affect surveillance practice. These can differ substantially
between pathogens (which may require laboratory tests of different complexity) and
between laboratories; some identifying some specimens may need several steps in
succession; reporting delays may even vary over time, thus making relevant to address these
with a general approach [49-51]. It is desirable to leverage the information on an infection
case potentially caused by the pathogen of interest, as this can anticipate the complete
detection of a cluster upon *emm* typing. As an approach to this, we thus defined an early
warning score based on exceedances in both typed and untyped records in the
neighbourhoods of a case of interest. For cases that were later identified as part of a cluster,
these early scores were indeed higher than those in the background, thus suggesting good
predictive power.

Further work is planned to validate RaNCover in public health practice. This is based on the
compilation of iGAS outbreak incidents identified through a variety of consensus sources,
including whole-genome sequencing, contact tracing, and detailed public health
investigation, which can ultimately confirm the presence of epidemiological links.

In conclusion, the RaNCover method is a convenient clustering tool for the spatio-temporal
surveillance of communicable diseases such as iGAS infections, which typically are sparsely
distributed over a geographical area and yet pose major threats to people's health due to
their severity. For these diseases, traditional surveillance methods typically require
aggregating data over time and space. Consequently, they cannot fully leverage all the
detailed geographical information that is available. Similarly to SaTScan[(TM)], our method can
be used at arbitrary levels of geographical and temporal aggregation. RaNCover further
provides a continuous warning score for each detected case, rather than for coarse-grained
time windows or geographical regions, and yet remains able to highlight emerging clusters

of diseases. The method only requires deriving a null model for the baseline and can be used
for different types of diseases, regardless of the complexity of their host-pathogen dynamics.

Funding Statement

This work was supported by Health Data Research UK, which is funded by the UK Medical
Research Council, EPSRC, Economic and Social Research Council, Department of Health and
Social Care (England), Chief Scientist Office of the Scottish Government Health and Social
Care Directorates, Health and Social Care Research and Development Division (Welsh
Government), Public Health Agency (Northern Ireland), British Heart Foundation and the
Wellcome Trust (MC, MJK, and NDM). MJK and NM are affiliated to the National Institute for
Health Research Health Protection Research Units (NIHR HPRUs) in Gastrointestinal
Infections and in Genomics and Enabling Data. The views expressed are those of the
author(s) and not necessarily those of the NIHR, the Department of Health and Social Care
or UKHSA. The funders had no role in study design, data collection and analysis, decision to
publish, or preparation of the manuscript.

Acknowledgements

The authors would like to thank colleagues in microbiology laboratories for their prompt
reporting of iGAS cases and submission of isolates, the UKHSA streptococcal reference
laboratory (Karen Broughton, Roger Daniel, and Chenchal Dhami), and the wider
collaborative group working on iGAS cluster detection, which incorporates Consultants in
Communicable Disease, Field Epidemiology Consultants, Topic Experts, and Public Health
Registrars. We also thank Majel McGranaham and Aoife Barror for valuable discussions.

Software and reproducibility

Analyses were carried out using R ([44], version 3.4.3). t-distributed stochastic neighbour
embedding was performed using the package Rtsne [52] with default parameters. Maps
created with Sf [53] using shapefiles from the GADM database (www.gadm.org) and the
Ordnance Survey Data Hub (osdatahub.os.uk). All software (also including codes for
generating   of   synthetic   data)   is   archived   online   at
https://github.com/mcavallaro/outbreak-detection.

Data cannot be shared publicly as confidential. Data are available from UKHSA and must
abide by the Caldicott principles. Material transfer must be obtained through a Material
Transfer Agreement.

Competing interests

551   The authors declare no competing interests.

Author contribution

Conceptualization MC, NDM, DT, MJK

Data curation, MC, JC, DR, VD, TL, DT

Software, MC

Formal analysis, MC, MJK

Validation, MC

Investigation, MC

Visualization, MC

Methodology, MC, MJK

Funding acquisition NDM, MJK

writing - original draft MC, MJK

writing - review and editing. MC, JC, DR, VD, TL, NDM, DT, MJK

Bibliography

1.   German, R.R., et al., *Updated guidelines for evaluating public health surveillance systems; recommendations from the Guidelines Working Group.* 2001.
2.   Elliot, A.J., et al., *The COVID-19 pandemic: a new challenge for syndromic surveillance.* Epidemiology and Infection, 2020. **148**: p. e122.
3.   Marston, H.D., et al., *Antimicrobial Resistance.* Jama-Journal of the American Medical Association, 2016. **316**(11): p. 1193-1204.
4.   Martinez, R., et al., *Trends in premature avertable mortality from non-communicable diseases for 195 countries and territories, 1990-2017: a population-based study.* Lancet Glob Health, 2020. **8**(4): p. e511-e523.
5.   Brookmeyer, R., et al., *Monitoring the Health of Populations: Statistical Principles and Methods for Public Health Surveillance*. 2004: Oxford University Press, USA.
6.   Harcourt, S.E., et al., *Developing a new syndromic surveillance system for the London 2012 Olympic and Paralympic Games.* Epidemiology and Infection, 2012. **140**(12): p. 2152-2156.
7.   Harcourt, S.E., et al., *Use of a large general practice syndromic surveillance system to monitor the progress of the influenza A(H1N1) pandemic 2009 in the UK.* Epidemiology and Infection, 2012. **140**(1): p. 100-105.
8.   Elliot, A.J., et al., *Establishing an emergency department syndromic surveillance system to support the London 2012 Olympic and Paralympic Games.* Emergency Medicine Journal, 2012. **29**(12): p. 954-960.

586 9. Abat, C., et al., *Traditional and syndromic surveillance of infectious diseases and*
587   *pathogens.* Int J Infect Dis, 2016. **48**: p. 22-8.
588 10. Morbey, R.A., et al., *The application of a novel 'rising activity, multi-level mixed effects,*
589   *indicator emphasis' (RAMMIE) method for syndromic surveillance in England.*
590   Bioinformatics, 2015. **31**(22): p. 3660-3665.
591 11. Smith, G.E., et al., *Novel public health risk assessment process developed to support*
592   *syndromic surveillance for the 2012 Olympic and Paralympic Games.* Journal of Public
593   Health, 2017. **39**(3): p. E111-E117.
594 12. Elliot, A.J., et al., *Syndromic surveillance - a public health legacy of the London 2012*
595   *Olympic and Paralympic Games.* Public Health, 2013. **127**(8): p. 777-781.
596 13. Buckingham-Jeffery, E., et al., *Correcting for day of the week and public holiday*
597   *effects: improving a national daily syndromic surveillance service for detecting public*
598   *health threats.* BMC Public Health, 2017. **17**(1): p. 477.
599 14. Hutwagner, L., et al., *The bioterrorism preparedness and response Early Aberration*
600   *Reporting System (EARS).* Journal of Urban Health-Bulletin of the New York Academy
601   of Medicine, 2003. **80**(2): p. I89-I96.
602 15. Noufaily, A., et al., *An improved algorithm for outbreak detection in multiple*
603   *surveillance systems.* Statistics in Medicine, 2013. **32**(7): p. 1206-1222.
604 16. Glaz, J., J. Naus, and S. Wallenstein, *Scan Statistics.* 2001, New York: Springer-Verlag.
605 17. Kulldorff, M., *A spatial scan statistic.* Communications in Statistics: Theory and
606   Methods, 1997. **26**: p. 1481-1496.
607 18. Kulldorff, M., *Prospective time periodic geographical disease surveillance using a scan*
608   *statistic.* Journal of the Royal Statistical Society: Series A (Statistics in Society), 2001.
609   **164**: p. 61-72.
610 19. Kulldorff, M., et al., *A space-time permutation scan statistic for disease outbreak*
611   *detection.* Plos Medicine, 2005. **2**(3): p. 216-224.
612 20. Kulldorff, M. and N. Nagarwalla, *Spatial disease clusters: detection and inference.*
613   Statistics in Medicine, 1995. **14**: p. 799--810.
614 21. Briggs, A.D.M., et al., *Approaches to the detection of very small, common, and easily*
615   *missed outbreaks that together contribute substantially to human Cryptosporidium*
616   *infection.* Epidemiology and Infection, 2014. **142**: p. 1869-1876.
617 22. Tango, T., K. Takahashi, and K. Kohriyama, *A Space-Time Scan Statistic for Detecting*
618   *Emerging Outbreaks.* Biometrics, 2011. **67**(1): p. 106-115.
619 23. Jiang, W., et al., *Spatiotemporal surveillance methods in the presence of spatial*
620   *correlation.* Statistics in Medicine, 2011. **30**(5): p. 569-583.
621 24. Yao, Z.J., J.M. Tang, and F.B. Zhan, *Detection of arbitrarily-shaped clusters using a*
622   *neighbor-expanding approach: A case study on murine typhus in South Texas.*
623   International Journal of Health Geographics, 2011. **10**.
624 25. Besag, J. and J. Newell, *The Detection of Clusters in Rare Diseases.* Journal of the Royal
625   Statistical Society. Series A: Statistics in Society, 1991. **154**: p. 143-155.
626 26. Oliver, J., et al., *Group A Streptococcus pharyngitis and pharyngeal carriage: A meta-*
627   *analysis.* Plos Neglected Tropical Diseases, 2018. **12**(3).
628 27. Lamagni, T.L., et al., *Epidemiology of severe Streptococcus pyogenes disease in Europe.*
629   Journal of Clinical Microbiology, 2008. **46**(7): p. 2359-2367.
630 28. Gherardi, G., L.A. Vitali, and R. Creti, *Prevalent emm Types among Invasive GAS in*
631   *Europe and North America since Year 2000.* Frontiers in Public Health, 2018. **6**.

29. Barnett, T.C., A.C. Bowen, and J.R. Carapetis, *The fall and rise of Group A*
Streptococcus diseases. Epidemiology and Infection, 2019. **147**: p. 1-6.
30. Bundle, N., et al., *Ongoing outbreak of invasive and non-invasive disease due to group*
A Streptococcus (GAS) type emm66 among homeless and people who inject drugs in
England and Wales, January to December 2016. Eurosurveillance, 2017. **22**(3): p. 2-7.
31. Bosley, H., *An invasive group A streptococcal infection outbreak: a community nursing*
experience. Journal of Infection Prevention, 2020.
32. Anderson, R.M. and R.M. May, *Infectious diseases of humans : dynamics and control.*
Oxford science publications. 1991, Oxford ; New York: Oxford University Press.
33. Held, L., M. Höhle, and M. Hofmann, *A statistical framework for the analysis of*
multivariate infectious disease surveillance counts. Statistical Modelling, 2005. **5**: p.
187-199.
34. Unkel, S., et al., *Statistical methods for the prospective detection of infectious disease*
outbreaks: a review. Journal of the Royal Statistical Society: Series A (Statistics in
Society), 2012. **175**: p. 49-82.
35. Woodall, W.H., *The Use of Control Charts in Health-Care and Public-Health*
Surveillance. Journal of Quality Technology, 2006. **38**(2): p. 89-104.
36. Noufaily, A., et al., *Comparison of statistical algorithms for daily syndromic*
surveillance aberration detection. Bioinformatics, 2019. **35**(17): p. 3110-3118.
37. Cox, D.R. and V. Isham, *Point processes.* Monographs on applied probability and
statistics. 1980, London ; New York: Chapman and Hall.
38. Farrington, P. and N. Andrews, *Outbreak Detection: Application to Infectious Disease*
Surveillance, in *Monitoring the Health of Populations: Statistical Principles and*
Methods for Public Health Surveillance., R. Brookmeyer and D.F. Stroup, Editors.
2004, Oxford University Press: New York, NY.
39. Wilson, E.B., *Probable Inference, the Law of Succession, and Statistical Inference.*
Journal of the American Statistical Association, 1927. **22**(158): p. 209-212.
40. Baddeley, A., E. Rubak, and R. Turner, *Spatial Point Patterns: Methodology and*
Applications with R. Spatial Point Patterns: Methodology and Applications with R,
2016: p. 1-810.
41. Bedubourg, G. and Y. Le Strat, *Evaluation and comparison of statistical methods for*
early temporal detection of outbreaks: A simulation-based study. Plos One, 2017.
12(7).
42. Texier, G., et al., *Building test data from real outbreaks for evaluating detection*
algorithms. Plos One, 2017. **12**(9).
43. Kulldorff, M. and I. Information Management Services, *SaTScanTM v8.0: Software for*
the spatial and space-time scan statistics. 2009.
44. R Core Development Team, *R: A language and environment for statistical computing.*
2018, R Foundation for Statistical Computing: Vienna, Austria.
45. van der Maaten, L. and G. Hinton, *Visualizing Data using t-SNE.* Journal of Machine
Learning Research, 2008. **9**: p. 2579-2605.
46. Merchant, E., et al. 2021.
47. Enki, D.G., et al., *Comparison of Statistical Algorithms for the Detection of Infectious*
Disease Outbreaks in Large Multiple Surveillance Systems. Plos One, 2016. **11**(8).
48. Yang, X.T., et al., *Comparing the similarity and difference of three influenza*
surveillance systems in China. Scientific Reports, 2018. **8**.

49.   Noufaily, A., et al., *Modelling reporting delays for outbreak detection in infectious*
679   *disease data.* Journal of the Royal Statistical Society Series a-Statistics in Society,
680   2015. **178**(1): p. 205-222.
681   50.   Noufaily, A., et al., *Detection of Infectious Disease Outbreaks From Laboratory Data*
682   *With Reporting Delays.* Journal of the American Statistical Association, 2016.
683   **111**(514): p. 488-499.
684   51.   Brookmeyer, R. and J.G. Liao, *The Analysis of Delays in Disease Reporting - Methods*
685   *and Results for the Acquired-Immunodeficiency-Syndrome.* American Journal of
686   Epidemiology, 1990. **132**(2): p. 355-365.
687   52.   Krijthe, J.H., *Rtsne: T-Distributed Stochastic Neighbor Embedding using Barnes-Hut*
688   *Implementation*. 2015: https://github.com/jkrijthe/Rtsne.
689   53.   Pebesma, E., *Simple Features for R: Standardized Support for Spatial Vector Data.* R
690   Journal, 2018. **10**(1): p. 439-446.
691   54.   *Notifications of scarlet fever disease in England and data from invasive Group A*
692   *streptococci (iGAS) surveillance.* 2018; Available from:
693   https://webarchive.nationalarchives.gov.uk/ukgwa/20220202090104/https:/ww
694   w.gov.uk/government/publications/group-a-streptococcal-infections-activity-
695   during-the-2017-to-2018-season.

Supporting information

S1 Text. Supporting information, including: Figs S1, S2, S3, and S4 (PDF).

Figures

*Fig 1. Representation of random neighbourhood covering. **A** Black markers represent point*
events that are generated from an endemic baseline (blue shade). Circles are randomly drawn
to cover the neighbourhoods of the events. The number of events in any circle is consistent
*with the endemic baseline prediction within statistical fluctuations. **B** Three point events (red*
markers) representing an unexpected outbreak are on the region delimited by the dashed line.
Many circles (coloured in red) in this region contain significantly more points than the
baseline prediction.

Fig 2. Simulation experiment. **A** Synthetic baseline point events (black markers) for $1 \leq t \leq 9$.
Additional points representing an outbreak (red markers) are in the areas delimited in red. **B**
Algorithm prediction. Episode markers are coloured by their warning scores, blue ($w = 0$) to
red ($w = 1$). The algorithm assigns high warning scores to the events in the outbreak areas.

Fig 3. Timeliness. As the time τ progresses and new cases and cylinders are added, the
warning-score values of cases previously detected (here at t = 5,6,7, see also Fig. 2) are
updated. In fact, these converge quickly and even the earliest warning values are informative.

*Fig 4. Relative frequency λ_m of selected emm types in England computed retrospectively over the observation times τ. **A** The fraction the total of three most common types (viz., 1.0, 89.0, and 12.0) is either overall declining or stable, yet they still account for more than 38% of all episodes at the end of the study. **B** Selected emm types show concerning relative growth (108.1 and 33.0) or stable pattern (44.0 and 94.0) over recent years.*

735 *Fig 5. Weekly number of iGAS infection cases (black solid line) and fitted temporal baseline*
736 *function λ_{time} (dashed red line, shaded area comprises 95% confidence interval). 2018 was a*
737 *year of heightened transmission across England, as reported in [54]. The overall fitted trend is*
738 *increasing, despite the sharp drop in 2020.*

*Fig 6. Simulation experiment on realistic baseline. **A** The total number of synthetic epidemic cases increases slowly from t=40 until it reaches a peak at week t=48, thus simulating the emergence of an outbreak. **B** The retrospectively-computed warning scores w(x) for the epidemic cases (red markers) are typically larger than those for the endemic cases (blue markers); plotting these vs time highlights the epidemic cluster. **C** Localising the cases with w(x) > 0.95 (orange markers) identifies the outbreak epicentre; in the inset, the true outbreak cases are marked with a red cross and the area detected by SaTScan™ is circled for comparison. Map created with Sf [53] using shapefiles from the GADM database (www.gadm.org) and the Ordnance Survey Data Hub (osdatahub.os.uk). **D** Warning scores of two random replicates with different cylinder volumes are strongly correlated, showing that outbreak detection is robust with respect to the cylinder volume choice. **E** Timeliness: updating a true-case warning score prospectively as new cases are added shows that it increases as the outbreak progresses, thus permitting detection earlier than the peak time t=48 (colour shades are case dates as in A).*

*Fig 7. Data point embedding and cluster analysis for emm type 94.0. A Each iGAS record,
identified by its position in time and space, is embedded in a two-dimensional space of
coordinates C_1 and C_2 according to the t-SNE method. The points are coloured by their
warning scores (0 to 1, blue to red), thus showing the presence of a bright red cluster of points
with high warning scores ($w > 0.95$). Colouring the detected cases by their record time (B)
and region (C) shows that t-SNE preserves the neighbourhoods. D The framed cluster
corresponds to cases localised in the towns of Bournemouth and Weymouth. Map created with
Sf [53] using shapefiles from the GADM database (www.gadm.org) and the Ordnance Survey
Data Hub (osdatahub.os.uk).*

Emm type 108.1

Fig 8. Data point embedding and cluster analysis for emm type `108.0`. *Keys as in Fig. 7. emm*
`108.0` *is a concerning type due to its recent increase in the number of cases. The many cases*
*tagged with high warning scores (**A**) occurred after 2018 (**B**) and are clustered around*
*several geographical locations (**C-D**). Map created with Sf [53] using shapefiles from the*
GADM database (www.gadm.org) and the Ordnance Survey Data Hub (osdatahub.os.uk).

Fig 9. Data point embedding and cluster analysis for emm type `44.0`. *Key as in Fig. 7. The*
analysis detects a compact cluster localised in the East of England in 2019 (rectangle), with
its embedded points mapped to their geographical location by the arrows. Map created with
Sf [53] *using shapefiles from the GADM database (www.gadm.org) and the Ordnance Survey*
Data Hub (osdatahub.os.uk).

Fig 10. Data point embedding and cluster analysis for the rare emm type 33.0. Keys as in Fig
7. Inspection suggests that these cases belong to a single diffused cluster that emerged in
September 2019. Map created with Sf [53] using shapefiles from the GADM database
(www.gadm.org) and the Ordnance Survey Data Hub (osdatahub.os.uk).

Fig 11. Data point embedding for the common emm type 12.0. No detected case is assigned a
warning score higher than 0.95.

Fig 12. Prospective analysis for emm type 44.0. **A** *Warning scores retrospectively computed*
every week as new cases are included; some cases, corresponding to the cluster of March 2019
in the East of England, incur a sharp increase in their warning scores (top right lines, amber
to violet colour indicate serotyping week). **B** *The score of the first cluster case (detected and*
serotyped in the week of 21/03/19) increases rapidly during the first four weeks. **C**
Observation counts (marker) and estimated baselines (lines) in the week of 18/06/20 for both
typed (blue colour) and untyped (grey colour) cases. **D** *Warning scores of selected cluster*
cases. These cases were initial recorded without any serotyping information. Using the
untyped observations, it is possible to obtain early warning scores (grey colour). The sample
are serotyped a week after detection and the warning scores are updated accordingly (amber
to orange colour shade indicate serotype week).

"Cluster detection with random neighbourhood covering:
application to invasive Group A Streptococcal disease"

Massimo Cavallaro[1,2,3]*, Juliana Coelho[3], Derren Ready[3,6], Valerie Decraene[3], Theresa Lamagni[3], Noel D.
McCarthy[1,4,5], Dan Todkill[3,4], Matt J. Keeling[1,2]

[1] The Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research, University of
Warwick, Coventry, CV4 7AL, United Kingdom.

[2] School of Life Sciences and Mathematics Institute, University of Warwick, Coventry, CV4 7AL, United
Kingdom.

[3] National Infection Service, UK Health Security Agency, United Kingdom.

[4] Warwick Medical School, University of Warwick, Coventry, CV4 7AL, United Kingdom.

[5] Institute of Population Health, School of Medicine, Trinity College Dublin, University of Dublin, 2 Dublin,
Ireland.

[6] Health Protection Research Unit in Behavioural Science and Evaluation at the University of Bristol, Bristol,
UK

*m.cavallaro@warwick.ac.uk

*Fig S1. Retrospective assessment of RaNCover performances by means of simulations. **A-B** ROC*
curves (solid black line) of RaNCover predictions over two replicates, obtained using different
cylinder volumes. Solid grey curves are from ROCs obtained by means of bootstrapping. The
dotted red lines intercept the values of sensitivity (horizontal lines, sensitivities equal to 0.54
*and 0.40 for **A** and **B**, respectively) and specificity (vertical lines, specificity equal to 0.94 and*
*0.98 for **A** and **B**, respectively) obtained from setting $1 - \alpha = 0.95$ as a discrimination*

*Fig S2. Illustration of simulation-experiment data aggregated and projected to time (**A-B**) and*
*geography (**C-D**). Cases are simulated from a Poisson point process with endemic intensity*
*function fitted on true iGAS data, thus encoding for seasonality (**A**) and population density*
*(**C**). Other cases representing a small outbreak occurring from t=40 to t=59 in St Albans*
*(Hertfordshire, highlighted areas in **A**,**B**,**C**, and **D**) are included in the synthetic dataset; the*
*overall temporal and geographical patterns are not affected (**B** and **D**), yet the algorithm is*
able to spot the anomaly (Fig 6, main text). Maps created with Sf [53] using shapefiles from
the GADM database (www.gadm.org) and the Ordnance Survey Data Hub (osdatahub.os.uk).
Same data embedded in a two-dimensional plane using t-SNE are illustrated in Fig. S4.

Fig S3. Widths of confidence intervals (CIs) of warning scores in simulation experiment for two choices of total number of random cylinders drew ($N = 10^6$ and $N = 5 \times 10^5$, solid and dashed lines, respectively) and two choices of cylinders' volume (V1 and V2=V1×1.4, blue and grey lines, respectively). The warning scores are proportions of flagged cylinders and appropriate CIs are Wilson's for binomial proportions. Increasing N or increasing the volumes, the numbers at numerator and denominator of the proportion increase and the confidence in its estimate improves.

Simulation experiment

*Fig S4. Illustration of simulation data projected on a two-dimensional plane using t-distributed stochastic neighbour embedding (t-SNE). Each simulated record is identified by a point of coordinates C1 and C2. The points are coloured by their warning scores (**(A)** 0 to 1, blue to red, highlighting the presence of a bright red cluster of points with high warning scores w>0.95) and by their record time **(B)**, showing that t-SNE also preserves temporal proximity.*

858