Objective measures of reward sensitivity and motivation in people with high vs low anhedonia

Chloe Slaney[1,4,5], Adam M. Perkins[2], Robert Davis[3], Ian Penton-Voak[4,6], Marcus R Munafò[4,5,6], Conor J. Houghton[7], Emma S.J. Robinson[1].

[1] School of Physiology, Pharmacology & Neuroscience, Faculty of Life Sciences, University of Bristol, Bristol, BS8 1TD, UK.

[2] Department of Psychological Medicine, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, SE5 8AF, UK.

[3] Department of Psychology, Goldsmiths, University of London, London, SE14 6NW, UK.

[4] School of Psychological Science, University of Bristol, 12a Priory Road, Bristol, BS8 1TU, UK.

[5] MRC Integrative Epidemiology Unit at the University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK.

[6] National Institute for Health Research Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol, Bristol, UK.

[7] Department of Computer Science, Faculty of Engineering, University of Bristol, BS8 1UB, UK.

Author for correspondence: Chloe Slaney (E-mail: chloe.slaney@bristol.ac.uk). Address: School of Physiology, Pharmacology & Neuroscience, Biomedical Sciences Building, University Walk, University of Bristol, Bristol, UK,

Financial support. This research was funded in whole, or in part, by the Wellcome Trust [108899/B/15/Z]. For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. Adam M. Perkins is supported by the National Institute for Health Research Biomedical Research Centre at the South London and Maudsley National Health Service Foundation Trust and King's College London. This report represents independent research funded by the National Institute for Health Research Biomedical

33   **Key words:** anhedonia; reward; motivation; sensitivity; EEfRT; decision-making.

35   **Abstract: 235**

37   **Word count: 4,840**

38   **Abstract**

Background. Anhedonia – a diminished interest or pleasure in activities – is a core self-

reported symptom of depression which is poorly understood and often resistant to conventional

antidepressants. This symptom may occur due to dysfunction in one or more sub-components

of reward processing: motivation, consummatory experience, and/or learning. However, the

precise impairments remain elusive. Dissociating these components (ideally, using cross-

species measures) and relating them to the subjective experience of anhedonia is critical as it

may benefit fundamental biology research and novel drug development.

Methods. Using a battery of behavioural tasks based on rodent assays, we examined reward

motivation (Joystick-Operated Runway Task, JORT; and Effort-Expenditure for Rewards Task,

EEfRT) and reward sensitivity (Sweet Taste Test) in a non-clinical population who scored high

($N = 32$) or low ($N = 34$) on an anhedonia questionnaire (Snaith-Hamilton Pleasure Scale).

Results. Compared to the low anhedonia group, the high anhedonia group displayed marginal

impairments in effort-based decision-making (EEfRT) and reduced reward sensitivity (Sweet

Taste Test). However, we found no evidence of a difference between groups in physical effort

exerted for reward (JORT). Interestingly, whilst the EEfRT and Sweet Taste Test correlated with

anhedonia measures, they did not correlate with each other. This poses the question of whether

there are subgroups within anhedonia; however, further work is required to directly test this

hypothesis.

Conclusions. Our findings suggest that anhedonia is a heterogenous symptom associated with

impairments in reward sensitivity and effort-based decision-making.

63   **Objective measures of reward sensitivity and motivation in people with high vs**

64   **low anhedonia**

Anhedonia, a markedly diminished interest or pleasure in activities (DSM-5; American

Psychiatric Association, 2013), is a core self-reported symptom of Major Depressive Disorder

(MDD) that responds poorly to conventional treatments (Uher et al., 2012). Despite its

importance, the behavioural and neurobiological basis of anhedonia remains poorly understood

(Cooper, Arulpragasam, & Treadway, 2018). This could, in part, be due the over-reliance on

questionnaires to measure this symptom. Objective measures, particularly those that can be

applied in both clinical and preclinical animal research, may help to improve our mechanistic

understanding of anhedonia and aid the development of targeted treatments (Der-Avakian &

Markou, 2012; Pizzagalli, Jahn, & O'Shea, 2005; Thomsen, 2015; Treadway & Zald, 2011).

One way to objectively measure anhedonia is behaviourally. Anhedonia may occur due

to dysfunction in one or more components of reward processing: motivation to obtain rewards

("wanting"), consumption of rewards ("detection" or "liking"), and learning what predicts rewards

("learning") (Berridge & Robinson, 2003). Whilst the taxonomy of reward processing is debated,

preclinical studies suggest that these sub-domains involve partially dissociable neural systems

(Berridge & Robinson, 1998; Berridge et al., 2009). Despite advancements in preclinical

research (Der-Avakian et al., 2013), few studies have tried to assess these different

components in people and relate them to questionnaire measures of anhedonia (McCabe,

2018). Translating behavioural tasks from animals to humans and assessing which components

relate to anhedonia is critical as it may provide an objective biomarker that could be used in the

development and testing of new treatments (Der-Avakian, Barnes, Markou, & Pizzagalli, 2016).

As most preclinical tasks focus on measuring reward motivation and consumption, we also

focus on these two components here.

Reward motivation is the "incentive or desire to act or accomplish goals" (Der-Avakian et

al., 2016). Recently, preclinical reward motivation tasks, such as effort-related choice tasks

(Salamone, Correa, Yang, Rotolo, & Presby, 2018) and the Progressive Ratio Task (Hodos,

1961), have been adapted for human studies (Hershenberg et al., 2016; Treadway, Buckholtz,

Schwartzman, Lambert, & Zald, 2009). For example, the Effort-Expenditure for Rewards Task

(EEfRT; Treadway et al., 2009), requires participants to choose whether to exert more effort for

a high reward or less effort for a low reward over many trials. Interestingly, people with MDD

and people who have higher levels of anhedonia in a non-clinical population choose the high

effort/high reward option less often than healthy controls suggesting impairments in effort-based

decision-making (Treadway, Bossaller, Shelton, & Zald, 2012; Treadway et al., 2009). In the

Progressive Ratio Task, incrementally more effortful responses are required to obtain the same

amount of reward, and motivation is operationalised as the point at which the individual gives up

(i.e., their "breakpoint") (Strauss et al., 2016). In this task, people with MDD give up earlier than

healthy controls (Hershenberg et al., 2016), suggesting a reduced willingness to work for

reward. Whilst there is accumulating evidence showing reduced reward motivation in MDD, few

studies focus on physical effort-expenditure for reward (Halahakoon et al., 2020) (although see

Cléry-Melin et al., 2011). To address this gap, we developed a novel reward motivation task

(Joystick-Operated Runway Task; JORT; Perkins et al., 2009), designed to measure a person's

physical effort exerted for reward.

Reward sensitivity is the consummatory experience of reward. Here, we use the term

reward sensitivity to refer to reward detection, not reward liking which is how it is often used in

the human literature. We focus on reward detection, rather than reward liking, to increase the

back translation of the results to the animal work. The most commonly used reward sensitivity

task in rodents is the sucrose preference test (SPT; Willner et al., 1987), which measures an

animal's ability to detect, and show a preference for, a weak sucrose solution over water.

Researchers have attempted to translate this task into humans using a Sweet Taste Test

(Amsterdam, Settle, Doty, Abelman, & Winokur, 1987; Berlin, Givry-Steiner, Lecrubier, & Puech,

1998; Dichter, Smoski, Kampov-Polevoy, Gallop, & Garbutt, 2010). Although different variations

have been used, they typically require participants to report the intensity and pleasantness of

different sucrose concentrations. Previous studies have reported similar pleasantness ratings of

sucrose in people with MDD compared to healthy controls (Amsterdam et al., 1987; Berlin et al.,

1998; Dichter et al., 2010). However, there is discrepancy across studies when measuring

sucrose intensity: some studies have reported poorer detection of sucrose in people with MDD

compared to healthy controls (Amsterdam et al., 1987; Berlin et al., 1998), whilst others have

failed to find any differences (Dichter et al., 2010). One potential explanation is that point scales,

often used to measure ability to detect sucrose, may not be sensitive enough to reliably detect

impairments (McCabe, 2018).

Despite progress in the development of translational measures (Thomsen, 2015), there

are gaps in the literature. First, most studies examine only one sub-component of reward (e.g.,

motivation) using one assay. However, a battery of tasks designed to probe different reward

components (e.g., motivation, sensitivity) can demonstrate whether different components can be

dissociated and which components are associated with anhedonia (Husain & Roiser, 2018;

Nielson et al., 2020). Second, most studies compare people with MDD to healthy controls, with

few directly measuring anhedonia using anhedonia questionnaires (McCabe, 2018). This is

surprising, as only ~ 37% people with MDD exhibit significant levels of anhedonia (Pelizza &

Ferrari, 2009). In line with the Research Domain Criteria (RDoC), a symptom-based approach

which examines behavioural tasks in relation to symptom-based questionnaires may provide

more rigorous findings (Cuthbert, 2014; Insel et al., 2010).

Here, we compared individuals with high versus low levels of anhedonia in a non-clinical

population (Snaith et al., 1995) on a battery of behavioural tasks designed to measure reward

motivation (EEfRT and JORT) and reward sensitivity (Sweet Taste Test). A novel measure of

reward motivation - the JORT - was designed to assess reward motivation in the absence of

explicit decision-making. These tasks were chosen based on their potential to dissociate reward

components and their potential for cross-species research. Based on previous research, we

predicted that compared to people with low levels of anhedonia, people with high levels of

anhedonia will show reduced reward motivation and reward sensitivity.

145   **Method**

146   This study was pre-registered on the Open Science Framework (https://osf.io/p4zt6) and

147   was approved by the Faculty of Biomedical Sciences Research Ethics Committee at the

148   University of Bristol (Ref: 74082).

150   **Screening**

151   Individuals who scored high (score ≥ 25) or low (score < 18) on the Snaith-Hamilton

Pleasure Scale (SHAPS; see Figure 1)(Snaith et al., 1995) in a non-clinical population were

invited to take part in this study. These individuals scored in the top and bottom 25th percentile

on the SHAPS, which was administered online to 380 volunteers. In the pre-registration, the

original cut-off was a low score of < 17. However, due to initial difficulties with recruitment, this

was increased to < 18 during the study.

Participants

159   Whilst 101 participants (44 low anhedonia, 57 high anhedonia) met the online SHAPS

160   criteria and took part in the study, only 66 (34 low, 32 high) still met their SHAPS criteria when

161   re-tested at the study visit and were included in the primary analysis. For sample size

162   calculations, see supplementary materials.

163   Eligibility criteria were aged ≥ 18 years, fluent in English, not suffering from a mental

health condition or neurological illness (self-report), no current physical injuries, no allergy or

intolerance to sugar, no disorder of taste or smell, not diabetic, not previously participated in a

study using the JORT.

Participants were recruited using advertisements within the University of Bristol and

volunteer databases. Participants were informed that they would receive £20 for their time (~ 2

hours) and an additional performance-based pay (up to £5) based on one of the two reward

motivation tasks (chosen randomly).

Behavioural Measures

Joystick-Operated Runway Task. The JORT was originally developed as a

translational measure of fear and anxiety, here it has been adapted as a measure of reward

motivation. This is a computerised task which uses a force-sensing joystick to measure a

person's physical effort to obtain a reward (Perkins et al., 2009; PH-JS1, Psyal, London). Each

trial begins with a green dot (representing the participant) positioned along an onscreen runway.

A cue in the top left-hand corner of the screen indicates the number of points on offer (0, 10,

100 or 1,000 points; see Figure 2). Following this, a target (black dot) appears and immediately

accelerates away from the participant. Participants are informed that to win the points on offer,

they must push the joystick to chase and catch the onscreen target. The speed of the

participants cursor is linked to the force applied on the joystick (Perkins et al., 2009). Trials vary

in the number of points on offer (0, 10, 100 and 1,000 points; visible to the participant) and the

minimum effort required to win (50, 80, 100 or 120% of their maximum calibrated force; not

visible). If the participant catches the target, they receive feedback that they have won the

number of points on offer. Trials end either upon catching the target or after 7 seconds. The

inter-trial interval varies pseudo-randomly between 3-7 seconds. After each trial, the next trial

automatically begins. Force applied to the joystick is recorded every 15 ms.

Participants complete two blocks of 48 trials (96-total), separated by a short break (~ 5

minutes). Each block contains an equal number of reward-effort combinations ($N = 3$), except

for one combination where two trials were presented due to experimenter error (0 points-80%

effort in second block). All participants complete the same pre-randomised order of trials. For a

proof-of-concept study of the JORT, see supplementary materials.

Prior to the task, participants are given standardised verbal instructions before

completing a calibration phase and four practice trials. In the calibration phase, participants

push the joystick as "hard as they can" five times whenever "GO" is presented on the screen.

Peak force reached during the calibration or practice trials (whichever is highest) is recorded as

a person's max calibrated force. To standardise the amount of effort required across

participants on the main task, effort levels reflect the percentage of a participants max calibrated

force.

Effort-Expenditure for Rewards Task. The EEfRT (Treadway et al., 2009) is a

computerised effort-based decision-making task. On each trial, participants choose between an

"easy" and "hard" option. The easy option requires 30 button presses within 7 seconds (using

the index finger on their dominant hand) for 50p. The hard option requires 98 button presses

(using the little finger on their non-dominant hand) within 21 seconds for 58p - £2. Participants

have 5 seconds to choose, or they are randomly assigned to either option for that trial. Each trial

also contains a visible probability cue (12%, 50%, 88%) which indicates the probability of

winning money if they successfully complete the trial (i.e., win trials). The probability cue applies

to both the easy and hard option, with an equal proportion across the experiment. Participants

are informed that two "win trials" may be paid to them at the end of the experiment

(performance-based pay). Participants complete the same pre-randomised order of trials for 20

minutes. For further details, see supplementary materials.

Sweet Taste Test. We employed a modified version of the Sweet Taste Test (Berlin et

al., 1998), which measures detection threshold of sweet taste. On each trial, participants were

given one sucrose concentration. They were asked to sip one mouthful of the solution, swish it

around their mouth for 5 seconds and then spit it out (Berlin et al., 1998). Participants were

asked if they could detect the presence of sugar in the solution (Berlin et al., 1998). In between

trials, participants sipped one mouthful of water, swished it around their mouth for 5 seconds

and then spat this out. Prior to the task, participants completed two practice trials containing

water to ensure that they were familiar with the procedure and to clean their palette. Seven

concentrations of sucrose were used (0, 0.5, 1, 1.5, 2, 2.5, 5% w/v), each delivered in 15 mL. A

staircase method with five reversals was employed. The first concentration administered was

always the strongest concentration (5% sucrose) to ensure participants were aware of what to

detect. A staircase method was used as it provides a fast determination of threshold. The

detection threshold was calculated as the mean of the five boundaries (i.e., points at which they

changed response from either detect to fail to detect, or vice versa).

Self-Report Measures

The primary measure of anhedonia was the SHAPS (Snaith et al., 1995). This is a 14-

item questionnaire which asks participants to report their ability to experience pleasure in the

last few days. Additional measures of anhedonia, apathy and depression were collected:

Chapman Physical Anhedonia Scale (CPAS; Chapman et al., 1976), Temporal Experience of

Pleasure Scale (TEPS; Gard et al., 2006), Beck Depression Inventory-II (BDI-II; Beck et al.,

1996) and Apathy Evaluation Scale (AES; Marin et al., 1991). Higher scores indicate higher

symptoms on all questionnaires except for the TEPS. For further details, see supplementary

materials.

Procedure

Participants completed the behavioural tasks in the following order: EEfRT, JORT and

the Sweet Taste Test, which took approximately one hour. Participants then completed the

questionnaires: demographic form, SHAPS, CPAS, TEPS, BDI-II, AES.

Data Analysis

Primary Analysis. Data were analysed using SPSS 24 (IBM). Primary analyses were

run on those who met the SHAPS cut-offs at the study visit ($N = 66$). Based on previous

research, *a priori* potential co-variates were examined and their main effects retained in the

model (Engqvist, 2005).

For the JORT, 2 x 4 (group x reward) mixed ANOVAs examined differences between

groups in relative average force, maximum force, and reaction time. The primary outcome was

relative average force. For the EEfRT, 2 x 3 (group x probability) mixed ANCOVAs examined

differences between groups in mean proportion of hard-task choices and reaction time. Sex was

included as a covariate (Treadway et al., 2009). The primary outcome was proportion of hard-

task choices. For the Sweet Taste Test, an ANCOVA examined differences between groups in

detection threshold. Current smoking status (3 levels: daily, less than daily, not at all) was

included as a covariate (Sato, Endo, & Tomita, 2002). All secondary outcomes and data checks

are reported in the supplementary materials. Due to the high and low anhedonia group not

being well-matched regarding sex, sensitivity analyses were run including sex as a covariate to

ensure that any findings were not being driven by sex differences.

Exploratory Analysis. Spearman correlations examined associations between tasks

and self-report measures, and correlations between different tasks. As these correlational

analyses were exploratory, we report them descriptively only.

264   **Results**

Data are available at the University of Bristol data repository, data.bris,

at https://doi.org/10.5523/bris.1wlrhv4jzqs7q2egf8i7ruzta1.

Participant Characteristics

For participant demographics, see Table 1. For distribution of scores on self-report

measures and correlations between measures, see Supplementary Materials.

Joystick-Operated Runway Task

Based on *a priori* criteria, participants who succeeded in over 75% trials ($N = 4$) were

excluded from the analysis. This is because these participants must have achieved at least one

trial designed to be impossible (120% effort trials) and were therefore considered to have not

successfully achieved their maximum force during the calibration. In total, 62 participants (33

low anhedonia, 29 high anhedonia) were included in the analysis.

Average force. There was strong evidence of a main effect of reward, participants

exerted more force for higher reward magnitudes, $F_{(1.45, 87.14)} = 39.41$, $p < .001$, $n_p^2 = .40$, Figure

3. Bonferroni-corrected pairwise comparisons revealed evidence of a difference between all

reward magnitudes (ps ≤ .003). There was no evidence of a main effect of group ($F_{(1,60)} = .004$,

$p = .95$, $n_p^2 < .001$; see Figure 3) or reward x group interaction ($F_{(1.45, 87.14)} = .56$, $p = .52$, n_p^2

= .009). In the sensitivity analyses, where sex was added to the model, there was no evidence

of a main effect of sex ($p = .58$), group ($p = .80$) or reward x group interaction ($p = .66$).

Effort-Expenditure for Rewards Task

Consistent with Treadway (2009), an upper bound of the first 50 trials was used in the

analysis. Trials where participants did not make a choice were not included in the analysis ($M =$

49.20, $SD = 1.90$, range = 39 to 50).

Proportion of hard-task choices. There was strong evidence of a main effect of

probability, participants chose the hard-task choice more often when the probability of winning

was higher, $F_{(2,126)} = 209.12$, $p < .001$, $n_p^2 = .77$, Figure 3. Bonferroni-corrected comparisons

revealed evidence of a difference between all probabilities (ps < .001). There was evidence of a

main effect of group, $F_{(1,63)} = 4.45$, $p = .039$, $n_p^2 = .07$: the high anhedonia group ($M = 28.88$, SE

$= 2.42$) chose the hard-task choice less often than the low anhedonia group ($M = 36.41$, $SE =$

2.58; see Figure 3). There was no evidence of other main effects or interactions (ps ≥ .57).

However, removal of one outlier weakened the evidence of the main effect of group, $F_{(1,62)} =$

2.64, $p =.109$, $n_p^2 = .041$). Independent t-tests indicated modest evidence of a difference

between groups on low probability trials ($p = .016$), with no clear evidence of a difference on

medium or high probability trials (ps ≥ .16).

Button Pressing Rate. There was no evidence of a difference between groups in mean

button pressing rate (button-pressing speed) for the easy or hard-task choices (ps ≥ .11).

Sweet Taste Test

Based on *a priori* criteria, participants who reported detection of sucrose in 0% sucrose

concentrations were excluded. This excluded 5 participants; 61 participants (31 low anhedonia,

30 high anhedonia) were included in the analysis.

The analysis was conducted on log transformed data, which met ANOVA assumptions.

There was evidence of a main effect of group, the high anhedonia group required a higher

concentration of sucrose to detect its presence, $F_{(1,57)} = 4.56$, $p = .037$, $n_p^2 = .07$, see Figure 3.

There was no evidence of a main effect of smoking status, $F_{(2,57)} = .41$, $p = .66$, $n_p^2 = .01$. In the

sensitivity analysis, where sex was added in the model, there was no evidence of a main effect

of sex ($p = .68$), and still evidence of a main effect of group ($p = .039$). Removing one outlier

strengthened the evidence of a main effect of Group ($p = .015$, $n_p^2 = .10$).

Exploratory Analyses

Correlations between tasks and self-report measures. In line with the original EEfRT

study (Treadway et al., 2009), we found evidence of a negative correlation between proportion

of hard-task choices and trait anhedonia (CPAS; r = -.29; 95% CI = -0.47 to -0.10). In the Sweet

Taste Test, we found evidence of positive a correlation between detection threshold and

symptoms of depression (BDI-II), anhedonia (SHAPS and CPAS) and apathy (AES), see Table

2. In the JORT, to facilitate correlational analyses, individual slopes (force exerted across the

four reward magnitudes) were extracted for each participant. There was no clear evidence of

correlations between self-report measures and the JORT, see Table 2.

Correlations between tasks. There was evidence of a positive correlation between

proportion of hard-task choices (EEfRT) and average force slopes (JORT), $r_{(101)}$ = .21, 95% CI =

0.01 to 0.39, participants who were more willing to choose the hard-task option exerted more

force for higher compared to lower reward magnitudes on the JORT. There was no evidence of

a correlation between the EEfRT and Sweet Taste Test ($r_{(92)}$ = -.02, 95% CI = -0.23 to 0.18), or

the JORT and Sweet Taste Test ($r_{(92)}$ = -.07, 95% CI = -0.27 to 0.13).

333   **Discussion**

We found that the JORT was sensitive to reward, with participants exerting more force

for higher reward magnitudes, but we found no evidence of a difference between the high and

low anhedonia group in physical effort exerted for reward. However, compared to the low

anhedonia group, the high anhedonia group displayed altered effort-based decision-making

(EEfRT) and reduced reward sensitivity (Sweet Taste Test). Although the JORT did not find any

effect between groups, there was an association between the EEfRT and JORT, suggesting that

they may measure similar aspects of reward motivation. In this study, the EEfRT was more

sensitive to changes in anhedonia scores, possibly because it involves both physical effort and

cognitive decision-making.

Reward motivation and anhedonia

In the JORT, there was no evidence of a difference between the high and low anhedonia

group, both groups exerted more physical force for higher reward trials. This suggests that

people with higher levels of anhedonia, at least in a non-clinical population, do not display

deficits in their exertion of physical effort for reward on this task. Using a human Progressive

Ratio Task, Hershenberg and colleagues (2016) reported that people with MDD had a lower

breakpoint compared to healthy controls (Hershenberg et al., 2016). However, this task required

cognitive effort and as the rodent Progressive Ratio Task has failed to find a similar effect in

some models of depression (chronic mild stress and maternal separation) (Barr & Phillips, 1998;

Shalev & Kafkafi, 2002), its translational validity has been questioned. Nevertheless, Cléry-

Melin et al. (2011) found that compared to healthy volunteers, people with depression exerted

less physical force to maximize monetary rewards on a handgrip apparatus (Cléry-Melin et al.,

2011). Whilst these studies may seem to conflict the findings presented here, it is important to

note that our study included a non-clinical population, measuring anhedonia as a varying trait,

as opposed to clinical depression. Further studies are required that explore anhedonia and

physical effort for reward using different contexts (e.g., real-life settings), durations (e.g.,

temporary vs sustained effort), and types of effort (e.g., cognitive, social, other types of physical

exertion).

In the EEfRT, the high anhedonia group displayed a reduced proportion of hard-task

choices, compared to the low anhedonia group. However, evidence for this difference was weak

(especially after removing one outlier) and we did not find evidence of a difference between

groups on 50% probability trials, as predicted in our pre-registration. Nevertheless, our

exploratory analysis did replicate the original Treadway (2009) finding that higher trait

anhedonia (measured using the CPAS) was associated with reduced hard-task choices on the

EEfRT (Treadway et al., 2009). Interestingly, both our study and the Treadway study found an

association between the EEfRT and CPAS, but failed to find a robust association between the

EEfRT and SHAPS (Treadway et al., 2009; Yang et al., 2014). One possible reason for this is

that the SHAPS focuses more on consummation of reward ("I would be able to enjoy my

favourite meal") whereas the CPAS also includes questions on interest/desire for future rewards

("I have had very little desire to try new kinds of foods") (Leventhal, Chasson, Tapia, Miller, &

Pettit, 2006; Rizvi, Pizzagalli, Sproule, & Kennedy, 2016), with the latter being more relevant to

the EEfRT (although see Tran et al., 2020). Overall, this finding highlights the value of

employing multiple anhedonia questionnaires.

377   There was a correlation between the JORT and EEfRT suggesting that they may

measure a similar construct of motivation. However, only the EEfRT was related to anhedonia.

The EEfRT examines explicit decision-making (Treadway et al., 2009) (similar to rodent choice-

based tasks) where a participant must weigh up the value of two options, whereas the JORT

(more similar to the Progressive Ratio Task) measures amount of physical effort exerted for

reward. Speculatively, anhedonia may not be related to reduced exertion of physical effort for

reward when engaged in an activity, but rather in the choice to engage in an effortful activity for

reward (decision-making; Halahakoon et al., 2020). In relation to the rodent literature, these

findings suggest that effort-based-choice tasks may provide a more sensitive measure of

motivational impairments relevant to anhedonia than physical effort-for-reward tasks (e.g.,

Progressive Ratio Task). Nevertheless, it is important to consider alternative explanations

including methodological differences between tasks (e.g., presence of probabilistic cues, reward

sizes) and the cognitive demands of the EEfRT (Cooper et al., 2019; Whitton, Merchant, &

Lewandowski, 2020).

Reward sensitivity and anhedonia

In the Sweet Taste Test, the high anhedonia group displayed reduced sensitivity to

sucrose compared to the low anhedonia group. Most previous studies using this task have

compared people with clinical disorders such as MDD to healthy controls, and found conflicting

results (Amsterdam et al., 1987; Berlin et al., 1998; Dichter et al., 2010). In relation to

anhedonia, Bedwell and colleagues (2019) reported that sucrose sensitivity did not relate to

anhedonia measures in the typical population (Bedwell, Spencer, Chirino, & O'donnell, 2019).

One potential explanation for these conflicting results is how sucrose sensitivity is measured.

Most studies have measured sweet taste intensity on a point scale (Bedwell et al., 2019; Dichter

et al., 2010), as opposed to detection threshold (i.e., point at which sucrose can be detected)

(Berlin et al., 1998). It has been suggested that point scales may not be sensitive enough to

reliably detect subtle impairments (McCabe, 2018). This study suggests that anhedonia, at least

in a non-clinical population, is related to reduced reward sensitivity. Future studies should

examine whether this effect is specific to pleasant stimuli (e.g., sucrose) or whether it can also

be observed using negative stimuli (e.g., bitter tastes), indicating a general sensitivity

impairment.

Collectively, using a battery of translational tasks designed to probe different reward

components (motivation, sensitivity), we support and extend preclinical findings by revealing

that these components may also be dissociable in people. This is demonstrated by a lack of

correlations between tasks: individuals who were less willing to choose the hard-task choice on

the EEfRT did not display reduced sensitivity on the Sweet Taste Test. We also found that

anhedonia was related to reduced reward motivation and reward sensitivity. Despite

performance on both tasks being related to anhedonia, there was a lack of correlation between

tasks. Speculatively, anhedonia may be a heterogenous symptom (i.e., there may be sub-

groups of individuals who have dysfunction in different components of reward) (Thomsen, 2015;

Treadway & Zald, 2013). To directly address this question, further studies (with a larger sample

size) should be conducted that: (1) demonstrate that the behavioural tasks used are reliable to

individual effects, and (2) employ cluster analysis on a battery of behavioural tasks (motivation,

sensitivity, and learning) and self-report measures in a clinical population.

Limitations

This study examined anhedonia in a non-clinical, predominantly student, population.

There are benefits of examining non-clinical populations including the reduced likelihood of

cognitive deficits, medication use and comorbidity. However, anhedonia levels in the clinical

population are higher and may be qualitatively different. Therefore, future studies employing a

battery of translational behavioural tasks in a clinical population are needed. Second, there was

a higher proportion of males in the high anhedonia group (53%) compared to the low anhedonia

group (21%), which may have influenced the differences between groups. However, including

sex as a covariate in all analyses did not change the overall findings. Interestingly, few studies

have investigated sex differences in anhedonia, and thus it is unknown whether the higher

proportion of males in the high anhedonia group reflects a real sex difference in anhedonia

(Chan et al., 2012; Xinhua Yang, Wang, Liu, Liu, & Harrison, 2020) or whether this is a result of

sampling. Third, the data did not always meet the ANOVA assumptions. Whilst we attempted to

address this (e.g., transforming the data), this was not always entirely successful. In these

cases, analyses were conducted on the original untransformed data, which should be

considered when interpreting our findings. Nevertheless, ANOVA is relatively robust to minor

deviations from normality if extreme outliers are not the cause of skew (Field, 2005). Fourth, we

did not include pleasantness ratings and therefore we could not assess whether there are

differences in reward liking between groups. We decided not to include pleasantness ratings

because this requires using point scales which may not be sensitive enough to reliably detect

impairments (McCabe, 2018). For example, some studies have reported that adolescences with

symptoms of depression and people with a history of depression display altered neural

responses to hedonic stimuli (chocolate) compared to healthy controls, despite no evidence of a

difference between groups on self-report measures of liking and intensity after tasting chocolate

(McCabe, Cowen, & Harmer, 2009; Rzepa, Fisk, & McCabe, 2017). An objective measure of

reward liking, which does not rely on point scales, in the human literature would be ideal to

address this issue.

To summarize, this study employed a battery of behavioural tasks designed to tap into

different reward components (effort-for-reward, effort-based decision-making, and reward

sensitivity) in humans. Examining these different components in relation to anhedonia, we found

that symptoms of anhedonia in a non-clinical population were related to changes in reward

sensitivity and effort-based decision-making. Speculatively, as performance on the effort-based

decision-making (EEfRT) and reward sensitivity (Sweet Taste Test) tasks did not correlate, this

could suggest heterogeneity within anhedonia (Treadway & Zald, 2013). For example, some

people with anhedonia may display reduced sensitivity to reward, whilst others may display

altered decision-making to engage in effortful activities for reward. Further studies are required

that directly address this question. If this finding is found in clinical populations, it may have

important implications for assessment and treatment of anhedonia.

Acknowledgements. The authors wish to thank Michael T. Treadway for providing us with the

EEfRT.

Conflicts of Interest. R.D. is the owner of Psyal Limited (http://www.psyal.co.uk/), a company

that designs and sells products to the psychology community. This study includes one product

purchased from Psyal Limited: the JORT.

Ethical standards. The authors assert that all procedures contributing to this work comply with

the ethical standards of the relevant national and institutional committees on human

experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*

(5th ed.). American Psychiatric Association.

Amsterdam, J. D., Settle, R. G., Doty, R. L., Abelman, E., & Winokur, A. (1987). Taste and smell

perception in depression. *Biological Psychiatry*, *22*, 1481–1485.

Barr, A. M., & Phillips, A. G. (1998). Chronic mild stress has no effect on responding by rats for

sucrose under a progressive ratio schedule. *Physiology & Behavior*, *64*, 591–597.

Beck, A., Steer, R., & Brown, G. (1996). Manual for the Beck Depression Inventory-II. *San*

Antonio: Psychological Corporation. https://doi.org/10.1037/t00742-000

Bedwell, J. S., Spencer, C. C., Chirino, C. A., & O'donnell, J. P. (2019). The Sweet Taste Test:

Relationships with Anhedonia Subtypes, Personality Traits, and Menstrual Cycle Phases.

Journal of Psychopathology and Behavioral Assessment, *41*, 235–248.

Berlin, I., Givry-Steiner, L., Lecrubier, Y., & Puech, A. J. (1998). Measures of anhedonia and

hedonic responses to sucrose in depressive and schizophrenic patients in comparison with

healthy subjects. *European Psychiatry*, *13*, 303–309.

Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward: hedonic

impact, reward learning, or incentive salience? *Brain Research Reviews*, *28*, 309–369.

Berridge, K. C., & Robinson, T. E. (2003). Parsing reward. *Trends in Neurosciences*, *26*, 507–

513.

Berridge, K. C., Robinson, T. E., & Aldridge, J. W. (2009). Dissecting components of reward:

"liking", "wanting", and learning. *Current Opinion in Pharmacology*, *9*, 65–73.

Chan, R. C. K., Wang, Y., Yan, C., Zhao, Q., McGrath, J., Hsi, X., & Stone, W. S. (2012). A

Study of Trait Anhedonia in Non-Clinical Chinese Samples: Evidence from the Chapman

Scales for Physical and Social Anhedonia. *PLoS One*, *7*.

https://doi.org/10.1371/journal.pone.0034275

Chapman, L. J., Chapman, J. P., & Raulin, M. L. (1976). Scales for physical and social

anhedonia. *Journal of Abnormal Psychology*, *85*, 374–382.

Cléry-Melin, M., Schmidt, L., Lafargue, G., Baup, N., Fossati, P., & Pessiglione, M. (2011). Why

Don't You Try Harder? An Investigation of Effort Production in Major Depression. *PLoS*

ONE, *6*. https://doi.org/10.1371/journal.pone.0023178

Cooper, J. A., Arulpragasam, A. R., & Treadway, M. T. (2018). Anhedonia in depression:

biological mechanisms and computational models. *Current Opinion in Behavioral Sciences*,

22, 128–135.

Cooper, J. A., Barch, D. M., Reddy, L. F., Horan, W. P., Green, M. F., & Treadway, M. T.

(2019). Effortful Goal-Directed Behavior in Schizophrenia: Computational Subtypes and

Associations With Cognition. *Journal of Abnormal Psychology*.

https://doi.org/10.1037/abn0000443

Cuthbert, B. N. (2014). The RDoC framework: facilitating transition from ICD/DSM to

dimensional approaches that integrate neuroscience and psychopathology. *World*

Psychiatry, *13*, 28–35.

Der-Avakian, A., Barnes, S. A., Markou, A., & Pizzagalli, D. A. (2016). Translational assessment

of reward and motivational deficits in psychiatric disorders. *Current Topics in Behavioral*

Neurosciences, *28*, 231–262.

Der-Avakian, A., & Markou, A. (2012). The neurobiology of anhedonia and other reward-related

deficits. *Trends in Neurosciences*, *35*, 68–77.

Der-Avakian, A., Souza, M. D., Pizzagalli, D., & Markou, A. (2013). Assessment of reward

responsiveness in the response bias probabilistic reward task in rats: implications for

cross-species translational research. *Translational Psychiatry*, *3*.

https://doi.org/10.1038/tp.2013.74

Dichter, G. S., Smoski, M. J., Kampov-Polevoy, A. B., Gallop, R., & Garbutt, J. C. (2010).

Unipolar depression does not moderate responses to the sweet taste test. *Depression and*

Anxiety, *27*, 859–863.

Engqvist, L. (2005). The mistreatment of covariate interaction terms in linear model analyses of

behavioural and evolutionary ecology studies. *Animal Behaviour*, *70*, 967–971.

Field, A. P. (2005). *Discovering statistics using IBM SPSS statistics : and sex and drugs and*

rock "n" roll. SAGE.

Gard, D. E., Gard, M. G., Kring, A. M., & John, O. P. (2006). Anticipatory and consummatory

components of the experience of pleasure: A scale development study. *Journal of*

Research in Personality, *40*, 1086–1102.

Halahakoon, D. C., Kieslich, K., O'Driscoll, C., Nair, A., Lewis, G., & Roiser, J. P. (2020).

Reward-Processing Behavior in Depressed Participants Relative to Healthy Volunteers: A

Systematic Review and Meta-analysis. *JAMA Psychiatry*.

https://doi.org/10.1001/jamapsychiatry.2020.2139

Hershenberg, R., Satterthwaite, T. D., Daldal, A., Katchmar, N., Moore, T. M., Kable, J. W., &

Wolf, D. H. (2016). Diminished effort on a progressive ratio task in both unipolar and

bipolar depression. *Journal of Affective Disorders*, *196*, 97–100.

Hodos, W. (1961). Progressive ratio as a measure of reward strength. *Science*, *134*, 943–944.

Husain, M., & Roiser, J. P. (2018). Neuroscience of apathy and anhedonia: A transdiagnostic

approach. *Nature Reviews Neuroscience*, *19*, 470–484.

Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., … Wang, P. (2010).

Research Domain Criteria (RDoC): Toward a New Classification Framework for Research

on Mental Disorders. *American Journal of Psychiatry*, *167*, 748–751.

Leventhal, A. M., Chasson, G. S., Tapia, E., Miller, E. K., & Pettit, J. W. (2006). Measuring

Hedonic Capacity in Depression: A Psychometric Analysis of Three Anhedonia Scales.

Journal of Clinical Psychology, *62*, 1545–1558.

Marin, R. S., Biedrzycki, R. C., & Firinciogullari, S. (1991). Reliability and validity of the Apathy

Evaluation Scale. *Psychiatry Research*, *38*, 143–162.

McCabe, C. (2018). Linking anhedonia symptoms with behavioural and neural reward

responses in adolescent depression. *Current Opinion in Behavioral Sciences*, *22*, 143–151.

McCabe, C., Cowen, P. J., & Harmer, C. J. (2009). Neural representation of reward in recovered

depressed patients. *Psychopharmacology*, *205*, 667–677.

Nielson, D. M., Keren, H., O'Callaghan, G., Jackson, S. M., Douka, I., Vidal-Ribas, P., …

Stringaris, A. (2020). Great Expectations: A Critical Review of and Suggestions for the

Study of Reward Processing as a Cause and Predictor of Depression. *Biological*

Psychiatry. https://doi.org/10.1016/j.biopsych.2020.06.012

Pelizza, L., & Ferrari, A. (2009). Anhedonia in schizophrenia and major depression: state or

trait? *Annals of General Psychiatry*, *8*, 22.

Perkins, A. M., Ettinger, U., Davis, R., Foster, R., Williams, S. C. R., & Corr, P. J. (2009). Effects

of Lorazepam and Citalopram on Human Defensive Reactions: Ethopharmacological

Differentiation of Fear and Anxiety. *Journal of Neuroscience*, *29*, 12617–12624.

Pizzagalli, D. A., Jahn, A. L., & O'Shea, J. P. (2005). Toward an objective characterization of an

anhedonic phenotype: A signal-detection approach. *Biological Psychiatry*, *57*, 319–327.

Rizvi, S. J., Pizzagalli, D. A., Sproule, B. A., & Kennedy, S. H. (2016). Assessing anhedonia in

depression: Potentials and pitfalls. *Neuroscience & Biobehavioral Reviews*, *65*, 21–35.

Rzepa, E., Fisk, J., & McCabe, C. (2017). Blunted neural response to anticipation, effort and

consummation of reward and aversion in adolescents with depression symptomatology.

Journal of Psychopharmacology. https://doi.org/10.1177/0269881116681416

Salamone, J. D., Correa, M., Yang, J.-H., Rotolo, R., & Presby, R. (2018). Dopamine, Effort-

Based Choice, and Behavioral Economics: Basic and Translational Research. *Frontiers in*

Behavioral Neuroscience, *12*, 52.

Sato, K., Endo, S., & Tomita, H. (2002). Sensitivity of Three Loci on the Tongue and Soft Palate

to Four Basic Tastes in Smokers and Non-smokers. *Acta Oto-Laryngologica*, *122*, 74–82.

Shalev, U., & Kafkafi, N. (2002). Repeated maternal separation does not alter sucrose-

reinforced and open-field behaviors. *Pharmacology, Biochemistry, and Behavior*, *73*, 115–

576   122.

577   Snaith, R. P., Hamilton, M., Morley, S., Humayan, A., Hargreaves, D., & Trigwell, P. (1995). A

578   scale for the assessment of hedonic tone the Snaith-Hamilton Pleasure Scale. *The British*

579   *Journal of Psychiatry*, *167*, 99–103.

580   Strauss, G. P., Whearty, K. M., Morra, L. F., Sullivan, S. K., Ossenfort, K. L., & Frost, K. H.

581   (2016). Avolition in schizophrenia is associated with reduced willingness to expend effort

582   for reward on a Progressive Ratio task. *Schizophrenia Research*, *170*, 198–204.

583   Thomsen, K. R. (2015). Measuring anhedonia: impaired ability to pursue, experience, and learn

584   about reward. *Frontiers in Psychology*. https://doi.org/10.3389/fpsyg.2015.01409

585   Tran, T., Hagen, A. E. F., Hollenstein, T., & Bowie, C. R. (2020). Physical- and Cognitive-Effort-

586   Based Decision-Making in Depression: Relationships to Symptoms and Functioning.

587   *Clinical Psychological Science*. https://doi.org/10.1177/2167702620949236

588   Treadway, M. T., Bossaller, N. A., Shelton, R. C., & Zald, D. H. (2012). Effort-Based Decision-

589   Making in Major Depressive Disorder: A Translational Model of Motivational Anhedonia.

590   *Journal of Abnormal Psychology*, *121*, 553–558.

591   Treadway, M. T., Buckholtz, J. W., Schwartzman, A. N., Lambert, W. E., & Zald, D. H. (2009).

592   Worth the 'EEfRT'? The Effort Expenditure for Rewards Task as an Objective Measure of

593   Motivation and Anhedonia. *PLoS ONE*, *4*. https://doi.org/10.1371/journal.pone.0006598

594   Treadway, M. T., & Zald, D. H. (2011). Reconsidering anhedonia in depression: Lessons from

595   translational neuroscience. *Neuroscience and Biobehavioral Reviews*, *35*, 537–555.

596   Treadway, M. T., & Zald, D. H. (2013). Parsing Anhedonia: Translational Models of Reward-

597   Processing Deficits in Psychopathology. *Current Directions in Psychological Science*, *22*,

598   244–249.

599   Uher, R., Perlis, R. H., Henigsberg, N., Zobel, A., Rietschel, M., Mors, O., … McGuffin, P.

600   (2012). Depression symptom dimensions as predictors of antidepressant treatment

601   outcome: replicable evidence for interest-activity symptoms. *Psychological Medicine*, *42*,

967–980.

Whitton, A. E., Merchant, J. T., & Lewandowski, K. E. (2020). Dissociable mechanisms

underpinning effort-cost decision-making across the psychosis spectrum. *Schizophrenia*

Research, *224*, 133–140.

Willner, P., Towell, A., Sampson, D., Sophokleous, S., & Muscat, R. (1987). Reduction of

sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic

antidepressant. *Psychopharmacology*, *93*, 358–364.

Yang, Xin hua, Huang, J., Zhu, C. ying, Wang, Y. fei, Cheung, E. F. C., Chan, R. C. K., & Xie,

G. rong. (2014). Motivational deficits in effort-based decision making in individuals with

subsyndromal depression, first-episode and remitted depression patients. *Psychiatry*

Research, *220*, 874–882.

Yang, Xinhua, Wang, D., Liu, S., Liu, G., & Harrison, P. (2020). Trajectories of state anhedonia

and recent changes in anhedonia in college students: Associations with other psychiatric

syndromes. *Journal of Affective Disorders*, *262*, 337–343.

618   **Tables**

619   **Table 1**

620   *Demographic data and questionnaire scores for participants in the high and low anhedonia*

621   *group.*

	Low Group		High Group		Group Difference		
	Mean	*SD* / %	Mean	*SD* / %	*F*	*p*-value	Post-hoc
Age	24.0	5.1	22.8	8.6	.41	.52	n/a
Males (*N*)	7	21%	17	53%	$X^2 = 7.54$	.006	H > L
Smoke Status	2.7	0.7	2.7	0.6	$X^2 = .23$	.89	n/a
Income	2.3	0.6	2.2	0.6	$X^2 = .64$	.73	n/a
Money Concern	12	36%	16	50%	$X^2 = 1.23$	.27	n/a
SHAPS	15.4	1.0	27.6	2.2	838.34	<.001	H > L
CPAS	9.6	5.7	18.0	7.1	27.67	<.001	H > L
TEPS – A	50.5	4.6	40.3	5.2	71.10	<.001	L > H
TEPS – C	40.6	5.3	30.3	6.2	52.80	<.001	L > H
BDI-II	6.4	5.5	11.4	7.6	9.41	.003	H > L
Apathy Scale	24.6	4.4	32.4	6.2	33.74	<.001	H > L

Note. Current smoke status (1=daily, 2=less than daily, 3=not at all); income (1=high,

2=medium, 3=low); money concern, participants who reported "yes" to finance concerns;

SHAPS, Snaith-Hamilton Pleasure Scale; TEPS, Temporal Experience of Pleasure Scale (A,

anticipatory; C, consummatory); BDI-II, Beck Depression Inventory-II; n/a, not applicable.

Welch's ANOVA and Chi Square to compare groups.

Table 2

Spearman's correlations between behavioural tasks and self-report measures.

Variable		JORT	EEfRT	Sweet Taste Test
SHAPS	*R*	-.09	-.14	.27
	95% CI	-0.28 to 0.11	-0.33 to 0.06	0.07 to 0.46
CPAS	*R*	-.03	-.29	.22
	95% CI	-0.22 to 0.17	-0.47 to -0.10	0.02 to 0.41
TEPS–A	*R*	.001	.16	-.15
	95% CI	-0.20 to 0.20	-0.04 to 0.34	-0.34 to 0.06
TEPS–C	*R*	-.07	.09	-.19
	95% CI	-0.26 to 0.13	-0.11 to 0.28	-0.38 to 0.02
BDI-II	*R*	-.04	-.06	.27
	95% CI	-0.23 to 0.16	-0.25 to 0.14	0.06 to 0.45
Apathy Scale	*R*	.01	-.04	.23
	95% CI	-0.18 to 0.21	-0.24 to 0.15	0.03 to 0.42

Note. Sweet Taste Test - mean detection threshold; EEfRT, Effort-Expenditure for Rewards

Task – overall proportion of hard-task choices; JORT, Joystick-Operated Runway Task – slope

of average force across four reward magnitudes; SHAPS, Snaith-Hamilton Pleasure Scale;

CPAS, Chapman Physical Anhedonia Scale; TEPS, Temporal Experience of Pleasure Scale (A,

Anticipatory; C, Consummatory); BDI-II, Beck Depression Inventory-II; Apathy Scale, Apathy

Evaluation Scale. 95% CI, 95% Confidence Intervals; *N* = 91–101.

Figure 1. CONSORT diagram of study selection process.

Figure 2. Example trial on the JORT. The green dot (representing the participant) is initially displayed along with a cue indicating the number of points on offer. This is followed by the presentation of the target (a black dot). The participant then receives feedback if they catch the target (e.g., "You win 10 points") or no feedback if they fail to catch the target.

Figure 3. Differences between the high and low anhedonia group in the (A) Joystick-Operated Runway Task: relative average force exerted for each reward magnitude (N = 33 low, 29 high) (B) Effort-Expenditure for Rewards Task: mean proportion of hard-task choices across different levels of probability (N = 34 low, 32 high) and (C) Sweet Taste Test (N = 31 low, 30 high). Points represent participants. Error bars represent *SEM*. Data presented with statistical outliers included. * $p < .05$, ** $p < .001$.