The Omega-3 Index is Inversely Associated with the Neutrophil-Lymphocyte Ratio in Adults

Michael I. McBurney[1-3], Nathan L. Tintle[1,4], William S. Harris[1,5]

[1]Fatty Acid Research Institute, Sioux Falls, SD 57106, USA (MIM, NLT, WSH); [2]Department of

Human Health and Nutritional Sciences, University of Guelph, Guelph, ON N1G 2W1, Canada

(MIM), [3]Division of Biochemical and Molecular Biology, Friedman School of Nutrition Science

and Policy, Tufts University, Boston, MA 02111, USA (MIM); [4]Department of Population Health

Nursing Science, College of Nursing, University of Illinois – Chicago, Chicago, IL 60612,

USA(NLT); [5]Sanford School of Medicine, University of South Dakota, Sioux Falls, SD 57105, USA

(WSH).

Authors last names: McBurney, Tintle, Harris

Corresponding author: M.I. McBurney, Fatty Acid Research Institute, Sioux Falls, SD 57106,

USA. E-mail: mim@faresinst.com

Reprint requests: W.S. Harris, Fatty Acid Research Institute, Sioux Falls, SD 57106, USA. E-mail:

wsh@faresinst.com

Sources of Support: This work was supported by the Fatty Acid Research Institute (FARI). FARI is

a non-profit foundation bringing together nutrition scientists and biostatistical experts to

accelerate discovery of the relationships between fatty acids, especially omega-3 fatty acids,

and health. Pending application and approval, data described in the manuscript, code book, and

analytic code will be made available upon request to the Fatty Acid Research Institute

(https://www.faresinst.org/).

Summary

The EPA+DHA content of red blood cell membranes, i.e., the omega-3 index (O3I), reflects omega-3 fatty acid status. EPA and DHA modulate inflammation via effects on cell signaling and the synthesis of pro-resolving lipid mediators. The neutrophil-lymphocyte ratio (NLR) is a measure of innate-adaptive immune system balance and is a biomarker of systemic inflammation. Both a low O3I and an elevated NLR are associated with increased risk for chronic disease and mortality, including cardiovascular diseases and cancer. Our objective was to determine if a relationship existed between O3I and NLR in healthy adults. A healthy cohort of 28,871 non-inflamed individuals [C-reactive protein (CRP) <3 mg/L] was identified from a clinical laboratory dataset. NLR was inversely associated with O3I (p<0.0001) after adjusting for age, sex, body mass index, and CRP. In this healthy population, an O3I >6.6% was associated with a low NLR, reflective of a quiescent, balanced immune system.

40   **Keywords**: neutrophils, lymphocytes, neutrophil-lymphocyte ratio, omega-3 index,

41   eicosapentaenoic acid, docosahexaenoic acid

43   **Abbreviations**: BMI, body mass index; CRP, high-sensitivity C-reactive protein; DHA,

44   docosahexaenoic acid; EPA, eicosapentaenoic acid; NLR, neutrophil-lymphocyte ratio; O3I,

45   omega-3 index; SPM, specialized pro-resolving lipid mediators.

1. Introduction

Cells from both the innate and adaptive immune system circulate in blood. Cells of the former (i.e., neutrophils, phagocytes, dendritic cells, macrophages, eosinophils, basophils, mast cells, and natural killer cells) work together to neutralize pathogens, in part by releasing pro-inflammatory mediators that modulate the latter. Neutrophils are the most abundant leukocytes in circulation. They migrate to sites of tissue injury or damage and are responsible for a non-specific inflammatory response. The adaptive system (acquired immunity) responds to specific pathogens and cytokines by producing antibodies and maintains immunological memory via lymphocytes, i.e., T and B cells. The hallmark of the adaptive immune system is clonal expansion of lymphocytes. Inflammation is generally self-limiting, with its resolution being an active rather than passive process [1]. Chronic or unresolved inflammation can damage host tissues and increase risk for the development of a range of non-communicable diseases, e.g. cardiovascular, cancer, chronic respiratory diseases, diabetes, and autoimmune disorders, e.g. rheumatoid arthritis [2–4].

The neutrophil-lymphocyte ratio (NLR) is an emerging biomarker of systemic inflammation [5–8] and innate-adaptive immune system balance [6,9] that is readily accessible from a complete blood count. It has the benefit of not being influenced by physiological conditions such as dehydration and exercise [6]. NLR has been used to monitor astronaut immune function during long-duration missions [10]. Elevated NLR is also a biomarker of systemic inflammation [5,11–14], cardiovascular disease and events [6,15–18], the severity of covid-19 [19–22], mood disorders [23], cognitive impairment [24], cancer [6,7,25], dry eye [26], periodontitis [27],

rheumatoid arthritis [28], cancer [12,25,29], diabetic kidney disease [30] and total mortality

[6,9,14,17,31,32].

Nutrient deficiencies or inadequacies can impair immune system function and weaken the

immune response [3,33–36]. The proportion of EPA and DHA in immune cell membranes can

modulate inflammation via the synthesis of prostaglandins, leukotrienes, lipoxins and

specialized pro-resolving lipid mediators (SPMs, e.g., resolvins, protectins, and maresins) [1,37].

The EPA and DHA-derived metabolites are less inflammatory than those derived from the

omega-6 precursor, arachidonic acid [1,33,37–42]. SPMs derived from EPA and DHA reduce

neutrophil infiltration and production of reactive oxygen species, regulate the cytokine-

chemokine axis, and temper the inflammatory response without immunosuppression [37,43–

45]. The omega-3 index (O3I) [46] is a stable biomarker of long term EPA and DHA intake

[47,48]. A low O3I [49] and an elevated NLR [14,18,50] are both associated with risk of

cardiovascular events and mortality (**Supplemental Table 1**). Hypothesizing that low blood

EPA+DHA levels may partly contribute to systemic chronic inflammation, we asked if there was

a relationship between the O3I and the NLR in healthy individuals without inflammation.

2. Patients and Methods

This is a cross-sectional analysis of data from blood samples submitted for testing to

Health Diagnostic Laboratory, Inc (HDL, Inc., Richmond VA; now defunct) as part of routine

clinical assessment between 2011-2012. Subjects were adults (≥18y) with data on O3I, NLR,

body mass index (BMI), age, sex, and high-sensitivity C-reactive protein (CRP) (n=44,925).

Individuals with extreme O3I values (i.e., highest and lowest 0.5%) were excluded. Because CRP

> 3mg/L is indicative of inflammation arising from infection, trauma or chronic disease [51–54],

individuals with CRP >3 mg/L were also excluded. Thus, the final study population consisted of

28,871 individuals (**Figure 1**).

2.1 Laboratory Methods

Blood samples were drawn after an overnight fast and shipped with cold packs to HDL, Inc.

for testing. Samples were prepared at each clinical site according to standardized instructions

as previously described [55]. Absolute concentrations of blood cells (neutrophils, lymphocytes,

NLR) were determined with a Beckman-Coulter DxH 800 analyzer (Brea, CA, USA) and

biomarker data (CRP, O3I) were extracted without any linked patient identifiers except age, sex,

and BMI. For fatty acid analysis, RBCs were separated from plasma by centrifugation and

analyzed using gas chromatography as previously described [56]. The University of South

Dakota Institutional Review Board reviewed and approved the analysis of deidentified HLD, Inc.

laboratory data (IRB-21-147).

2.2 Statistical methods

Sample characteristics were summarized using standard statistical methods (e.g., means,

SDs, correlations) with t-tests and adjusted linear models used to compare characteristics of

male and female participants above and below O3I values that were ultimately identified as

primary cut points in the NLR-O3I curves (see below). Splines were fit using 3rd degree

polynomials with knots at each decile in R (version 3.6.2; splines package). Unadjusted models

used a linear model to predict NLR values by splines of O3I. Adjusted models accounted for sex,

age, BMI and CRP values in the linear models accounting for potential non-linear relationships

using splines. To identify significant changes in the NLR-O3I relationship, we used a "sliding O3I

window" approach. The width of each window was three O3I percentage points (e.g., 4% to

7%). By moving the window up by 0.1% increments and repeatedly testing for significant

differences between the mean NLR in the lower vs. the upper half of the window, we sought to

discover O3I cut points where the NLR-O3I relationship appeared to flatten. These would be O3I

values above which the "effect" of an increase in O3I on NLR had little impact. We began with a

window midpoint of O3I =2.6%, and we used adjusted linear models in R to test for upper vs

lower half differences. We identified the first window which did not have a statistically

significant difference in upper and lower mean NLR values. The midpoint of this window was

chosen as the O3I cut point to be used in further analysis.

Statistical interactions with O3I values above and below the subsequently identified

inflection points were tested by placing the interaction term in separate models predicting O3I,

NLR, age, sex, BMI, and CRP. Pearson correlations were used to assess strength and direction of

linear association between covariates and NLR. Statistical significance was set to 0.05 for all

analyses and 95% confidence bands are provided where appropriate.

3. Results

The final dataset consisted of 28,871 healthy adults (**Figure 1**). The average age was

55.1±15.1 years, the average BMI was 27.6±5.4 kg/m^2, and there were slightly more females

(51%) than males. Females were significantly younger and had slightly but significantly higher

O3I and CRP levels and lower NLR, neutrophil count, and BMI than males (**Table 1**). When the

model was also adjusted for age, BMI and CRP, small but significant sex differences remained

with the exception of BMI (**Table 1**).

NLR was significantly (p<0.0001) and inversely associated with O3I (**Figure 2A**). Pearson

correlations (r) of other variables with NLR were r=0.06 (CRP), r=0.14 (age), r=0.01 (BMI), r=0.65

(neutrophil count), and r=-0.24 (lymphocyte count). Adjustment for age, sex, BMI, and CRP

strengthened the relationship (**Figure 2B**). The O3I value where the relationship between O3I

and NLR was no longer changing (i.e., the cut point determined by the sliding window analysis

described in Methods) was 6.6% (**Figure 2B**). That is, below 6.6%, the curve was clearly steeper

than it was above 6.6%. The O3I cut point of 6.6% was explored in more detail. An O3I <6.6%

was associated with significantly higher NLR, neutrophil number, lymphocyte number, CRP,

BMI, and lower age (versus O3I ≥6.6%) (**Table 2**). These O3I-related differences persisted,

except for lymphocyte number, when the model was adjusted for age, sex, BMI and CRP (**Table**

2).

4. Discussion and Conclusions

We report a significant inverse relationship between O3I and NLR. By excluding individuals

with CRP >3 mg/L, we could examine the O3I-NLR relationship in individuals without any

evidence of tissue injury, infection, or systemic acute inflammation [57,58]. NLR values

measured in this cohort (**Table 1**) were similar to those measured in healthy adults globally

[8,24,59–63]. NLR is known to increase with age [9,32,60], obesity [13,60], CRP [64,65], and to

sometimes differ by sex [32,59]. These same relationships were mostly observed here as well

(data not reported), suggesting that the findings from this clinical laboratory cohort are

generally representative.

Inflammation is a normal physiological response to infection and injury, but unrestrained

inflammation can cause tissue damage leading to disease. As noted earlier, EPA and DHA serve

as precursors for the synthesis of prostaglandins, leukotrienes, lipoxins and SPMs that can

mediate immune responses [1,29,33,37,39–41]. SPMs derived from EPA and DHA protect

tissues by limiting acute inflammatory responses and helping achieve homeostasis without

immunosuppression [37,44,45]. A threshold blood EPA+DHA concentration may be required to

fuel sufficient SPM biosynthesis to achieve meaningful resolution of inflammation [37].

Omega-3 supplementation increases the EPA+DHA content of neutrophil membranes in

vivo and inhibits the 5-lipoxygenase pathway and leukotriene B_4-mediated functions of

neutrophils *in vitro* [38]. Omega-3 supplementation significantly increased plasma EPA+DHA

levels (50-100%) and decreased concentrations of IL-6 (10-12% vs 36% increase in placebo

group) and TNFα (0.2 to -2.3% vs 12% increase in placebo group) in healthy, middle-aged and

older individuals [66]. Meta-analyses have confirmed this effect [67]. In a study of 8,237

participants living in Japan without a history of cardiovascular disease, eating fish >4 days per

week was associated with significant lower NLR [68]. High omega-3 dietary intake was

associated with a trend towards lower CRP (p=0.09) and NLR (p=0.17) in men with acute

coronary syndrome [69]. A systematic review and meta-analysis of 18 randomized controlled

trials found marine-derived omega-3 fatty acids lowered pro-inflammatory eicosanoid

concentrations, e.g., in neutrophil leukotriene B_4 [70]. In the long-term, chronically lower levels

of inflammation may have health benefits. For example, a pooled analysis of data from 17

prospective cohort studies involving 15,720 deaths among 43,466 individuals over a median of

16 years of follow-up found blood EPA+DHA levels were inversely associated with risk for death

from all causes, CVD, cancer and other causes [49].

Despite the strength of the NLR-O3I relationship, the most obvious limitation of this study

is that confounding metabolic events, dietary behaviors, smoking habits and/or physical activity

levels associated with O3I may also affect innate-adaptive immune system balance. However, a

nationally representative study of 9,427 individuals living in the US did not find significant

differences in NLR with respect to sex, education, insurance status, or drinking habits [59].

Moreover, this cross-sectional population is similar to 2/3 of the US population who have CRP

<3mg/L [71] and the ~95% of adults living in the USA with circulating O3I percentages below 5%

[72,73].

In this cross-sectional study of healthy individuals, 81% of participants had O3I <6.6%.

Below this threshold, an increase in NLR, reflecting an increasingly imbalanced innate-adaptive

immune system, was observed. NLR continued to decrease with increasing O3I until O3I >8.5%,

similar to the target proposed for cardiovascular health, i.e. O3I >8% [46,74]. Clearly additional

evidence, including intervention studies, is needed to determine whether the NLR-O3I

relationship is causal or coincidental.

In conclusion, red blood cell EPA+DHA levels are significantly and inversely associated with

NLR, a biomarker of inflammation and innate-adaptive immune system balance. We suggest

that an O3I >6.6%, and possibly as high as 8.5%, may be targets for maintaining "optimal" NLR

values reflecting a quiescent, balanced innate-adaptive immune system.

Acknowledgements: The authors wish to thank Steven Varvel, PhD for his help in acquiring and

collating the dataset used in this study.

CRediT authorship contribution statement: M.I. McBurney: Conceptualization & analytical

design. Writing -original draft, review & editing. **W.S. Harris**: Conceptualization & analytical

design. Funding acquisition. Writing- review & editing. **N.L. Tintle**: Formal analysis. Writing-

review & editing.

Disclosures: M.I. McBurney serves on the Board of Directors of the American Society for

Nutrition and has or has held consulting agreements in the past 3 years with Council for

Responsible Nutrition; Church & Dwight; DSM Nutritional Products; International Life Sciences

Institute, North America; McCormick; OmegaQuant Analytics; PepsiCo; and VitaMe

Technologies. W.S. Harris holds an interest in OmegaQuant Analytics, a lab that offers omega-3

blood testing; and is a member of the RB Schiff Science and Innovation Advisory Board. N.L.

Tintle has no conflicts to disclose.

References

[1]   C.N. Serhan, Pro-resolving lipid mediators are leads for resolution physiology, Nature. 510

(2014) 92–101. https://doi.org/10.1038/nature13479.

[2]   P.C. Calder, Marine omega-3 fatty acids and inflammatory processes: Effects, mechanisms

and clinical relevance, Bioch. Biophys. Acta: Mol Cell Biol Lipids. 1851 (2015) 469–484.

https://doi.org/10.1016/j.bbalip.2014.08.010.

[3]   Childs, Calder, Miles, Diet and Immune Function, Nutrients. 11 (2019) 1933.

https://doi.org/10.3390/nu11081933.

[4]   D. Furman, J. Campisi, E. Verdin, P. Carrera-Bastos, S. Targ, C. Franceschi, L. Ferrucci, D.W.

Gilroy, A. Fasano, G.W. Miller, A.H. Miller, A. Mantovani, C.M. Weyand, N. Barzilai, J.J.

Goronzy, T.A. Rando, R.B. Effros, A. Lucia, N. Kleinstreuer, G.M. Slavich, Chronic

inflammation in the etiology of disease across the life span, Nat Med. 25 (2019) 1822–

1832. https://doi.org/10.1038/s41591-019-0675-0.

[5]   F. Imtiaz, K. Shafique, S.S. Mirza, Z. Ayoob, P. Vart, S. Rao, Neutrophil lymphocyte ratio as a

measure of systemic inflammation in prevalent chronic diseases in Asian population,

International Archives of Medicine. 5 (2012) 2. https://doi.org/10.1186/1755-7682-5-2.

[6]   T. Bhat, S. Teli, J. Rijal, H. Bhat, M. Raza, G. Khoueiry, M. Meghani, M. Akhtar, Neutrophil

to lymphocyte ratio and cardiovascular diseases: a review, Expert Rev. Cardiovasc. Ther.

(2013) 6.

[7]   S. Socorro Faria, P.C. Fernandes Jr, M.J. Barbosa Silva, V.C. Lima, W. Fontes, R. Freitas-

Junior, A.K. Eterovic, P. Forget, The neutrophil-to-lymphocyte ratio: a narrative review,

Ecancer. 10 (2016). https://doi.org/10.3332/ecancer.2016.702.

[8] J. Fest, R. Ruiter, M.A. Ikram, T. Voortman, C.H.J. van Eijck, B.H. Stricker, Reference values

for white blood-cell-based inflammatory markers in the Rotterdam Study: a population-

based prospective cohort study, Sci Rep. 8 (2018) 10566. https://doi.org/10.1038/s41598-

018-28646-w.

[9] M. Song, B.I. Graubard, C.S. Rabkin, E.A. Engels, Neutrophil-to-lymphocyte ratio and

mortality in the United States general population, Nature Res. 11 (2021) 464.

https://doi.org/10.1038/s41598-020-79431-7.

[10] A.M. Paul, S.D. Mhatre, E. Cekanaviciute, A.-S. Schreurs, C.G.T. Tahimic, R.K. Globus, S.

Anand, B.E. Crucian, S. Bhattacharya, Neutrophil-to-Lymphocyte Ratio: A Biomarker to

Monitor the Immune Status of Astronauts, Front. Immunol. 11 (2020) 564950.

https://doi.org/10.3389/fimmu.2020.564950.

[11] R. Zahorec, Ratio of neutrophil to lymphocyte counts - rapid and simple parameter of

systemic inflammation and stress in critically ill, Bratisl. Lek. Listy. 102 (2001) 5–14.

[12] L. Venkatraghavan, T.P. Tan, J. Mehta, A. Arekapudi, A. Govindarajulu, E. Siu, Neutrophil

Lymphocyte Ratio as a predictor of systemic inflammation - A cross-sectional study in a

pre-admission setting., F1000Res. 4 (2015) 123.

https://doi.org/10.12688/f1000research.6474.1.

[13] E. Rodríguez-Rodríguez, A.M. López-Sobaler, R.M. Ortega, M.L. Delgado-Losada, A.M.

López-Parra, A. Aparicio, Association between Neutrophil-to-Lymphocyte Ratio with

Abdominal Obesity and Healthy Eating Index in a Representative Older Spanish Population,

Nutrients. 12 (2020) 855. https://doi.org/10.3390/nu12030855.

[14] H. Wada, T. Dohi, K. Miyauchi, R. Nishio, M. Takeuchi, N. Takahashi, H. Endo, M. Ogita, H.

Iwata, T. Kasai, S. Okazaki, K. Isoda, S. Suwa, H. Daida, Neutrophil to Lymphocyte Ratio and

Long-Term Cardiovascular Outcomes in Coronary Artery Disease Patients with Low High-

Sensitivity C-Reactive Protein Level, Int. Heart J. 61 (2020) 447–453.

https://doi.org/10.1536/ihj.19-543.

[15] B. Suh, D.W. Shin, H.-M. Kwon, J.M. Yun, H.-K. Yang, E. Ahn, H. Lee, J.H. Park, B. Cho,

Elevated neutrophil to lymphocyte ratio and ischemic stroke risk in generally healthy

adults, PLoS ONE. 12 (2017) e0183706. https://doi.org/10.1371/journal.pone.0183706.

[16] T. Angkananard, T. Anothaisintawee, M. McEvoy, J. Attia, A. Thakkinstian, Neutrophil

Lymphocyte Ratio and Cardiovascular Disease Risk: A Systematic Review and Meta-

Analysis, BioMed Research International. 2018 (2018) 1–11.

https://doi.org/10.1155/2018/2703518.

[17] N.H. Adamstein, J.G. MacFadyen, L.M. Rose, R.J. Glynn, A.K. Dey, P. Libby, I.A. Tabas, N.N.

Mehta, P.M. Ridker, The neutrophil–lymphocyte ratio and incident atherosclerotic events:

analyses from five contemporary randomized trials, European Heart Journal. 42 (2021)

896–903. https://doi.org/10.1093/eurheartj/ehaa1034.

[18] L. Zhan, Y. Liu, Y. Cheng, W. Guo, J. Yang, Predictive Value of Neutrophil/Lymphocyte Ratio

(NLR) on Cardiovascular Events in Patients with COVID-19, IJGM. Volume 14 (2021) 3899–

3907. https://doi.org/10.2147/IJGM.S317380.

[19] S. Qun, Y. Wang, J. Chen, X. Huang, H. Guo, Z. Lu, J. Wang, C. Zheng, Y. Ma, Y. Zhu, D. Xia,

Y. Wang, H. He, Y. Wang, M. Fei, Y. Yin, M. Zheng, Y. Xu, W. Ge, F. Hu, J. Zhou, Neutrophil-

272  to-Lymphocyte Ratios Are Closely Associated With the Severity and Course of Non-mild

273  COVID-19, Front. Immunol. 11 (2020) 2160. https://doi.org/10.3389/fimmu.2020.02160.

274 [20] S. Sun, X. Cai, H. Wang, G. He, Y. Lin, B. Lu, C. Chen, Y. Pan, X. Hu, Abnormalities of

275  peripheral blood system in patients with COVID-19 in Wenzhou, China, Clinica Chimica

276  Acta. 507 (2020) 174–180. https://doi.org/10.1016/j.cca.2020.04.024.

277 [21] M. Seyit, E. Avci, R. Nar, H. Senol, A. Yilmaz, M. Ozen, A. Oskay, H. Aybek, Neutrophil to

278  lymphocyte ratio, lymphocyte to monocyte ratio and platelet to lymphocyte ratio to

279  predict the severity of COVID-19, The American Journal of Emergency Medicine. 40 (2021)

280  110–114. https://doi.org/10.1016/j.ajem.2020.11.058.

281 [22] E. Vafadar Moradi, A. Teimouri, R. Rezaee, N. Morovatdar, M. Foroughian, P. Layegh, B.

282  Rezvani Kakhki, S.R. Ahmadi Koupaei, V. Ghorani, Increased age, neutrophil-to-lymphocyte

283  ratio (NLR) and white blood cells count are associated with higher COVID-19 mortality, The

284  American Journal of Emergency Medicine. 40 (2021) 11–14.

285  https://doi.org/10.1016/j.ajem.2020.12.003.

286 [23] M.G. Mazza, S. Lucchi, A.G.M. Tringali, A. Rossetti, E.R. Botti, M. Clerici,

287  Neutrophil/lymphocyte ratio and platelet/lymphocyte ratio in mood disorders: A meta-

288  analysis, Progress in Neuro-Psychopharmacology and Biological Psychiatry. 84 (2018) 229–

289  236. https://doi.org/10.1016/j.pnpbp.2018.03.012.

290 [24] J.-H. Liu, Y.-J. Zhang, Q.-H. Ma, H.-P. Sun, Y. Xu, C.-W. Pan, Elevated blood neutrophil to

291  lymphocyte ratio in older adults with cognitive impairment, Archives of Gerontology and

292  Geriatrics. 88 (2020) 104041. https://doi.org/10.1016/j.archger.2020.104041.

293   [25]   A.J. Templeton, M.G. McNamara, B. Šeruga, F.E. Vera-Badillo, P. Aneja, A. Ocaña, R.

294   Leibowitz-Amit, G. Sonpavde, J.J. Knox, B. Tran, I.F. Tannock, E. Amir, Prognostic Role of

295   Neutrophil-to-Lymphocyte Ratio in Solid Tumors: A Systematic Review and Meta-Analysis,

296   JNCI: Journal of the National Cancer Institute. 106 (2014).

297   https://doi.org/10.1093/jnci/dju124.

298   [26]   Y.-F. Meng, Q. Pu, Q. Ma, W. Zhu, X.-Y. Li, Neutrophil/lymphocyte ratio as an inflammatory

299   predictor of dry eye: A case-control study, Therapeutics and Clinical Risk Management. 17

300   (2021) 259–266.

301   [27]   R. Lu, W. Li, X. Wang, D. Shi, H. Meng, Elevated neutrophil-to-lymphocyte ratio but not

302   platelet-to-lymphocyte ratio is associated with generalized aggressive periodontitis in a

303   Chinese population, J Periodontol. 92 (2021) 507–513. https://doi.org/10.1002/JPER.20-

304   0282.

305   [28]   Z. Jin, G. Cai, P. Zhang, X. Li, S. Yao, L. Zhuang, M. Ren, Q. Wang, X. Yu, The value of the

306   neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio as complementary

307   diagnostic tools in the diagnosis of rheumatoid arthritis: A multicenter retrospective study,

308   J Clin Lab Anal. 35 (2021). https://doi.org/10.1002/jcla.23569.

309   [29]   L. Xia, W. Li, J. Zhai, C. Yan, J. Chen, H. Yang, Significance of neutrophil-to-lymphocyte

310   ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio and prognostic

311   nutritional index for predicting clinical outcomes in T1–2 rectal cancer, BMC Cancer. 20

312   (2020) 208. https://doi.org/10.1186/s12885-020-6698-6.

313   [30] L. Winter, L.A. Wong, G. Jerums, J. Seah, M. Clarke, S.M. Tan, M.T. Coughlan, R.J. MacIsaac,

314   E.I. Ekinci, Use of Readily Accessible Inflammatory Markers to Predict Diabetic Kidney

315   Disease, Front. Endocrinol. 9 (2018) 225. https://doi.org/10.3389/fendo.2018.00225.

316   [31] S. Balta, S. Demirkol, M. Unlu, Z. Arslan, T. Celik, Neutrophil to lymphocyte ratio may be

317   predict of mortality in all conditions, Br J Cancer. 109 (2013) 3125–3126.

318   https://doi.org/10.1038/bjc.2013.598.

319   [32] J. Fest, T.R. Ruiter, B. Groot Koerkamp, D. Rizopoulos, M.A. Ikram, C.H.J. van Eijck, B.H.

320   Stricker, The neutrophil-to-lymphocyte ratio is associated with mortality in the general

321   population: The Rotterdam Study, Eur J Epidemiol. 34 (2019) 463–470.

322   https://doi.org/10.1007/s10654-018-0472-y.

323   [33] D. Wu, E.D. Lewis, M. Pae, S.N. Meydani, Nutritional Modulation of Immune Function:

324   Analysis of Evidence, Mechanisms, and Clinical Relevance, Front. Immunol. 9 (2019) 3160.

325   https://doi.org/10.3389/fimmu.2018.03160.

326   [34] P.C. Calder, A.C. Carr, A.F. Gombart, M. Eggersdorfer, Optimal Nutritional Status for a

327   Well-Functioning Immune System Is an Important Factor to Protect against Viral

328   Infections, Nutrients. 12 (2020) 1181. https://doi.org/10.3390/nu12041181.

329   [35] A.F. Gombart, A. Pierre, S. Maggini, A Review of Micronutrients and the Immune System–

330   Working in Harmony to Reduce the Risk of Infection, Nutrients. 12 (2020) 236.

331   https://doi.org/10.3390/nu12010236.

332   [36] C.A. Reider, R.-Y. Chung, P.P. Devarshi, R.W. Grant, S. Hazels Mitmesser, Inadequacy of

333   Immune Health Nutrients: Intakes in US Adults, the 2005–2016 NHANES, Nutrients. 12

334   (2020) 1735. https://doi.org/10.3390/nu12061735.

[37]   P.C. Calder, Eicosapentaenoic and docosahexaenoic acid derived specialised pro-resolving

mediators: Concentrations in humans and the effects of age, sex, disease and increased

omega-3 fatty acid intake, Biochimie. 178 (2020) 105–123.

https://doi.org/10.1016/j.biochi.2020.08.015.

[38]   T.H. Lee, R.L. Hoover, J.D. Williams, R.I. Sperling, J. Ravalese III, B.W. Spur, D.R. Robinson,

E.J. Corey, R.A. Lewis, K.F. Austen, Effect of dietary enrichment with eicosapentaenoic acid

and docosahexaenoic acids on in vitro neutrophil and monocyte leukotriene generation

and neutrophil function, New England Journal of Medicine. 312 (1985) 217–224.

[39]   P. Yaqoob, P.C. Calder, Fatty acids and immune function: new insights into mechanisms, Br

J Nutr. 98 (2007) S41–S45. https://doi.org/10.1017/S0007114507832995.

[40]   E.A. Dennis, P.C. Norris, Eicosanoid storm in infection and inflammation, Nat Rev Immunol.

15 (2015) 511–523. https://doi.org/10.1038/nri3859.

[41]   U. Radzikowska, A.O. Rinaldi, Z. Çelebi Sözener, D. Karaguzel, M. Wojcik, K. Cypryk, M.

Akdis, C.A. Akdis, M. Sokolowska, The Influence of Dietary Fatty Acids on Immune

Responses, Nutrients. 11 (2019) 2990. https://doi.org/10.3390/nu11122990.

[42]   J. McDaniel, Regulation of Neutrophil Function by Marine n-3 Fatty Acids—A Mini Review,

Cell Biochem Biophys. (2021). https://doi.org/10.1007/s12013-021-01001-z.

[43]   C.N. Serhan, N. Chiang, T.E. Van Dyke, Resolving inflammation: dual anti-inflammatory and

pro-resolution lipid mediators, Nat Rev Immunol. 8 (2008) 349–361.

https://doi.org/10.1038/nri2294.

[44] J. Dalli, J.W. Winkler, R.A. Colas, H. Arnardottir, C.-Y.C. Cheng, N. Chiang, N.A. Petasis, C.N.

Serhan, Resolvin D3 and Aspirin-Triggered Resolvin D3 Are Potent Immunoresolvents,

Chemistry & Biology. 20 (2013) 188–201. https://doi.org/10.1016/j.chembiol.2012.11.010.

[45] J.K. Colby, R.-E.E. Abdulnour, H.P. Sham, J. Dalli, R.A. Colas, J.W. Winkler, J. Hellmann, B.

Wong, Y. Cui, S. El-Chemaly, N.A. Petasis, M. Spite, C.N. Serhan, B.D. Levy, Resolvin D3 and

Aspirin-Triggered Resolvin D3 Are Protective for Injured Epithelia, The American Journal of

Pathology. 186 (2016) 1801–1813. https://doi.org/10.1016/j.ajpath.2016.03.011.

[46] W.S. Harris, C. von Schacky, The Omega-3 Index: a new risk factor for death from coronary

heart disease?, Prev. Med. 39 (2004) 212–220.

https://doi.org/10.1016/j.ypmed.2004.02.030.

[47] M.R. Flock, A.C. Skulas-Ray, W.S. Harris, T.D. Etherton, J.A. Fleming, P.M. Kris-Etherton,

Determinants of erythrocyte omega-3 fatty acid content in response to fish oil

supplementation: A dose–response randomized controlled trial, J. Am. Heart Assoc. 2

(2013) e000513. https://doi.org/10.1161/JAHA.113.000513.

[48] R.E. Walker, K.H. Jackson, N.L. Tintle, G.C. Shearer, A. Bernasconi, S. Masson, R. Latini, B.

Heydari, R.Y. Kwong, M. Flock, P.M. Kris-Etherton, A. Hedengran, R.M. Carney, A. Skulas-

Ray, S.S. Gidding, A. Dewell, C.D. Gardner, S.M. Grenon, B. Sarter, J.W. Newman, T.L.

Pedersen, M.K. Larson, W.S. Harris, Predicting the effects of supplemental EPA and DHA

on the omega-3 index, Am. J. Clin. Nutr. 110 (2019) 1034–1040.

https://doi.org/10.1093/ajcn/nqz161.

[49] W.S. Harris, N.L. Tintle, F. Imamura, F. Qian, A.V. Ardisson Korat, M. Marklund, L. Djoussé,

J.K. Bassett, P.-H. Carmichael, Y.-Y. Chen, Y. Hirakawa, L.K. Kupers, M. Lankinen, R.A.

Murphy, C. Samieri, M.K. Senn, P. Shi, J.K. Virtanen, I.A. Brouwer, K.-L. Chien, G.

Eiriksdottir, N.G. Forouhi, J.M. Geleijnse, G.G. Giles, V. Gudnason, C. Helmer, A. Hodge, R.

Jackson, K.-T. Khaw, M. Laakso, H. Lai, D. Laurin, J. Lindsay, R. Micha, J. Mursu, T.

Ninomiya, W. Post, B.M. Psaty, U. Riserus, J.G. Robinson, A.H. Shadyab, L. Snetselaar, A.

Sala-Vila, Y. Sun, L.M. Steffen, M.Y. Tsai, N.J. Wareham, A.C. Wood, J.H.Y. Wu, F. Hu, Q.

Sun, D.S. Siscovick, R.N. Lemaitre, D. Mozaffarian, F. The Fatty Acids and Outcomes

Research Consortium, Blood n-3 fatty acid levels and total and cause-specific mortality

from 17 prospective studies, Nature Commun. 12 (2021) 1–9.

https://doi.org/10.1038/s41467-021-22370-2.

[50] S. Kim, M. Eliot, D.C. Koestler, W.-C. Wu, K.T. Kelsey, Association of Neutrophil-to-

Lymphocyte Ratio With Mortality and Cardiovascular Disease in the Jackson Heart Study

and Modification by the Duffy Antigen Variant, (2018) 8.

[51] M.A. Zimmerman, Diagnostic Implications of C-Reactive Protein, Arch Surg. 138 (2003)

220. https://doi.org/10.1001/archsurg.138.2.220.

[52] A. Villaseñor, S.W. Flatt, C. Marinac, L. Natarajan, J.P. Pierce, R.E. Patterson, Postdiagnosis

C-Reactive Protein and Breast Cancer Survivorship: Findings from the WHEL Study, Cancer

Epidemiol Biomarkers Prev. 23 (2014) 189–199. https://doi.org/10.1158/1055-9965.EPI-

13-0852.

[53] M.L. Knight, The Application of High-Sensitivity C-Reactive Protein in Clinical Practice: A

2015 Update, US Pharmacist. 40 (2015) 50–53.

[54] S. Jafarnejad, V. Boccardi, B. Hosseini, M. Taghizadeh, Z. Hamedifard, A Meta-analysis of

Randomized Control Trials: The Impact of Vitamin C Supplementation on Serum CRP and

Serum hs-CRP Concentrations, (2018).

https://doi.org/10.2174/1381612824666181017101810.

[55] T.D. Dayspring, S.A. Varvel, L. Ghaedi, D.L. Thiselton, J. Bruton, J.P. McConnell, Biomarkers

of cholesterol homeostasis in a clinical laboratory database sample comprising 667,718

patients, Journal of Clinical Lipidology. 9 (2015) 807–816.

https://doi.org/10.1016/j.jacl.2015.08.003.

[56] W.S. Harris, J.V. Pottala, R.S. Vasan, M.G. Larson, S.J. Robins, Changes in erythrocyte

membrane trans and marine fatty acids between 1999 and 2006 in older Americans, J.

Nutr. 142 (2012) 1297–1303. https://doi.org/10.3945/jn.112.158295.

[57] WHO, C-reactive protein concentrations as a marker of inflammation or infection for

interpreting biomarkers of micronutrient status., World Health Organization, Geneva,

Switzerland, 2014.

http://apps.who.int/iris/bitstream/10665/133708/1/WHO_NMH_NHD_EPG_14.7_eng.pdf

?ua=1.

[58] M. Antonelli, I. Kushner, It's time to redefine inflammation, FASEB J. 31 (2017) 1787–1791.

https://doi.org/10.1096/fj.201601326R.

[59] B. Azab, M. Camacho-Rivera, E. Taioli, Average Values and Racial Differences of Neutrophil

Lymphocyte Ratio among a Nationally Representative Sample of United States Subjects,

PLoS ONE. 9 (2014) e112361. https://doi.org/10.1371/journal.pone.0112361.

[60] J. Li, Q. Chen, X. Luo, J. Hong, K. Pan, X. Lin, X. Liu, L. Zhou, H. Wang, Y. Xu, H. Li, C. Duan,

Neutrophil-to-Lymphocyte Ratio Positively Correlates to Age in Healthy Population: NLR

Correlates to Age, J. Clin. Lab. Anal. 29 (2015) 437–443.

https://doi.org/10.1002/jcla.21791.

[61] Y. Chen, Y. Zhang, G. Zhao, C. Chen, P. Yang, S. Ye, X. Tan, Difference in Leukocyte

Composition between Women before and after Menopausal Age, and Distinct Sexual

Dimorphism, PLoS ONE. 11 (2016) e0162953.

https://doi.org/10.1371/journal.pone.0162953.

[62] P. Forget, C. Khalifa, J.-P. Defour, D. Latinne, M.-C. Van Pel, M. De Kock, What is the

normal value of the neutrophil-to-lymphocyte ratio?, BMC Res Notes. 10 (2017) 12.

https://doi.org/10.1186/s13104-016-2335-5.

[63] J.S. Lee, N.Y. Kim, S.H. Na, Y.H. Youn, C.S. Shin, Reference values of neutrophil-lymphocyte

ratio, lymphocyte-monocyte ratio, platelet-lymphocyte ratio, and mean platelet volume in

healthy adults in South Korea, Medicine. 97 (2018) e11138.

https://doi.org/10.1097/MD.0000000000011138.

[64] C.P. de Jager, P.T. van Wijk, R.B. Mathoera, J. de Jongh-Leuvenink, T. van der Poll, P.C.

Wever, Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia

better than conventional infection markers in an emergency care unit, Crit Care. 14 (2010)

R192. https://doi.org/10.1186/cc9309.

[65] J.C. Yombi, P.E. Schwab, E. Thienpont, Neutrophil-to-lymphocyte ratio (NLR) distribution

shows a better kinetic pattern than C-reactive protein distribution for the follow-up of

early inflammation after total knee arthroplasty, Knee Surg Sports Traumatol Arthrosc. 24

(2016) 3287–3292. https://doi.org/10.1007/s00167-015-3921-0.

[66] J.K. Kiecolt-Glaser, M.A. Belury, R. Andridge, W.B. Malarkey, B.S. Hwang, R. Glaser, Omega-

3 supplementation lowers inflammation in healthy middle-aged and older adults: A

randomized controlled trial | Elsevier Enhanced Reader, Brain, Behav. Immun. 26 (2012)

988–995. https://doi.org/10.1016/j.bbi.2012.05.011.

[67] K. Li, T. Huang, J. Zheng, K. Wu, D. Li, Effect of Marine-Derived n-3 Polyunsaturated Fatty

Acids on C-Reactive Protein, Interleukin 6 and Tumor Necrosis Factor α: A Meta-Analysis,

PLoS ONE. 9 (2014) e88103. https://doi.org/10.1371/journal.pone.0088103.

[68] S. Tani, R. Matsuo, W. Atsumi, K. Kawauchi, T. Ashida, T. Yagi, K. Imatake, Y. Suzuki, A.

Takahashi, N. Matsumoto, Y. Okumura, Higher Frequency of Fish Intake May Be Associated

with a Lower Neutrophil/Lymphocyte Ratio: Anti-Atherosclerotic Effects of Fish

Consumption, Ann Nutr Metab. (2021) 1–8. https://doi.org/10.1159/000515915.

[69] A. Sut, K. Chiżyński, M. Różalski, J. Golański, Dietary intake of omega fatty acids and

polyphenols and its relationship with the levels of inflammatory markers in men with

chronic coronary syndrome after percutaneous coronary intervention, Kardiol Pol. 78

(2020) 117–123. https://doi.org/10.33963/KP.15078.

[70] J. Jiang, K. Li, F. Wang, B. Yang, Y. Fu, J. Zheng, D. Li, Effect of Marine-Derived n-3

Polyunsaturated Fatty Acids on Major Eicosanoids: A Systematic Review and Meta-Analysis

from 18 Randomized Controlled Trials, PLoS One. 11 (2016).

https://doi.org/10.1371/journal.pone.0147351.

[71] I. Kushner, D. Rzewnicki, D. Samols, What Does Minor Elevation of C-Reactive Protein

Signify?, The American Journal of Medicine. 119 (2006) 166.e17-166.e28.

https://doi.org/10.1016/j.amjmed.2005.06.057.

[72] R. Murphy, E. Yu, E. Ciappio, S. Mehta, M. McBurney, Suboptimal Plasma Long Chain n-3

Concentrations are Common among Adults in the United States, NHANES 2003–2004,

Nutrients. 7 (2015) 10282–10289. https://doi.org/10.3390/nu7125534.

[73] R.A. Murphy, P.P. Devarshi, S. Ekimura, K. Marshall, S. Hazels Mitmesser, Long-chain

omega-3 fatty acid serum concentrations across life stages in the USA: an analysis of

NHANES 2011–2012, BMJ Open. 11 (2021) e043301. https://doi.org/10.1136/bmjopen-

2020-043301.

[74] W.S. Harris, The Omega-3 Index: Clinical Utility for Therapeutic Intervention, Curr Cardiol

Rep. 12 (2010) 503–508. https://doi.org/10.1007/s11886-010-0141-6.

Table 1. Characteristics of the study population. Mean±SD			
Variable[1]	All (n=28,871)	Males (n=14,211)	Females (n=14,660)	Male vs. Female Adjusted[2] p-value
O3I (%)	4.97±1.84	4.9±1.82	5.05±1.85	0.016
NLR	2.09±1.10	2.13±1.13	2.05±1.07	<0.0001
Neutrophil (10^3/µL)	3.41 ±1.4	3.46±1.4	3.37±1.39	0.0002
Lymphocyte (10^3/µL)	1.81±1.47	1.82±1.77	1.80±1.10	<0.0001
CRP (mg/L)	1.23±0.77	1.21±0.75	1.25±0.79	0.0001
BMI (kg/m^2)	27.6±5.4	28.8±4.9	26.6±5.6	0.54
Age (years)	55.1±15.1	55.3±14.7	54.9±15.5	<0.001
[1] O3I, Omega-3 index; NLR, neutrophil-lymphocyte ratio; CRP, high-sensitivity C-reactive protein; BMI, body mass index. [2] Models were adjusted for age, BMI, and CRP (except when the model was predicting these variables).			

Variable[1]	O3I <6.6% (n=23,510)	O3I ≥6.6% (n=5,361)	Adjusted[2] p-value
Table 2. Stratification by omega-3 index (O3I) classification (n=28,871). Mean±SD			
O3I (%)	4.28±1.11	8.03±1.16	<0.0001
NLR	2.12±1.11	1.96±1.07	<0.0001
Neutrophil (10^3/µL)	3.48±1.40	3.11±1.35	<0.0001
Lymphocyte (10^3/µL)	1.82±1.56	1.76±0.95	0.591
CRP (mg/L)	1.27±0.78	1.07±0.73	<0.0001
BMI (kg/m^2)	28.0±5.4	26.3±4.9	<0.0001
Age (years)	54.1±15.4	59.4±13.1	<0.0001

[1] O3I, Omega-3 index; NLR, neutrophil-lymphocyte ratio; CRP, high-sensitivity C-reactive protein; BMI, body mass index.

[2] Models were adjusted for sex, age, BMI, and CRP (except when the model was predicting these variables).

480   Figure 1. Analytical sample flow. NLR, neutrophil-lymphocyte ratio; O3I, omega-3 index; CRP,

481   high-sensitivity C-reactive protein; BMI, body mass index.

483   Figure 2. The unadjusted (A) and adjusted for sex, age, BMI, and CRP (B) relationship between

484   the neutrophil-lymphocyte ration (NLR) and the omega-3 index (O3I) in 28,871 healthy adults.

485   (Means and 95% confidence bands).