Integrated patient network and genomic plasmid analysis reveal a regional, multi-species outbreak of carbapenemase-producing Enterobacterales carrying both *bla*IMP and *mcr-9* genes

Adhiratha Boonyasiri[1,2*], Ashleigh C. Myall[1,3*] , Yu Wan[1*], Frances Bolt[1,4,5], Alice Ledda[6,7], Siddharth Mookerjee[4], Andrea Y. Weiße[8,9], Jane F. Turton[7], Hala Abbas[1,10], Ruta Prakapaite[11], Akshay Sabnis[11], Alireza Abdolrasouli[4], Kenny Malpartida-Cardenas[1,12], Luca Miglietta[1,12], Hugo Donaldson[10], Mark Gilchrist[1,4], Katie L. Hopkins[1,7], Matthew J Ellington[1,13], Jonathan A. Otter[1], Gerald Larrouy-Maumus[1,14], Andrew M. Edwards[11], Jesus Rodriguez-Manzano[1,5,12], Xavier Didelot[15], Mauricio Barahona[3], Alison H. Holmes[1,4,5], Elita Jauneikaite[1,6#] and Frances Davies[1,4,10#]

1. NIHR Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance, Department of Infectious Disease, Imperial College London, Hammersmith Hospital, London, UK.
2. Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand.
3. Department of Mathematics, Imperial College London, London, UK
4. Imperial College Healthcare NHS Trust, London, UK
5. Centre for Antimicrobial Optimisation, Hammersmith Hospital, Imperial College London, Du Cane Road, London, UK
6. Department of Infectious Disease Epidemiology, School of Public Health, Imperial College London, London, UK
7. HCAI, Fungal, AMR, AMU & Sepsis Division, UK Health Security Agency, London, UK
8. School of Biological Sciences, University of Edinburgh, Scotland, UK
9. School of Informatics, University of Edinburgh, Scotland, UK
10. Department of Microbiology, North West London Pathology, London, UK
11. MRC Centre for Molecular Bacteriology and Infection, Department of Infectious Disease, Faculty of Medicine, Imperial College London, London, United Kingdom
12. Centre for Bio-Inspired Technology, Department of Electrical and Electronic Engineering, Faculty of Engineering, Imperial College London, UK
13. Reference Services Division, UK Health Security Agency, London, UK
14. MRC Centre for Molecular Bacteriology and Infection, Department of Life Sciences, Faculty of Natural Sciences, Imperial College London, London, UK

34   15. School of Life Sciences and Department of Statistics, University of Warwick, UK
35
36
37   * These authors contributed equally; # these senior authors contributed equally.

39   **Keywords:** Carbapenem-resistant Enterobacterales, IMP carbapenemase, antimicrobial
40   resistance, horizontal gene transfer, molecular epidemiology, spatiotemporal network, patient
41   pathways

Abstract

The incidence of carbapenemase-producing Enterobacterales (CPE) is rising globally, yet Imipenemase (IMP) carbapenemases remain relatively rare. This study describes an investigation of the emergence of IMP-encoding CPE amongst diverse Enterobacterales species between 2016 and 2019 in patients across a London regional hospital network.

A network analysis approach to patient pathways, using routinely collected electronic health records, identified previously unrecognised contacts between patients who were IMP CPE positive on screening, implying potential bacterial transmission events. Whole genome sequencing of 85 Enterobacterales isolates from these patients revealed that 86% (73/85) were diverse species (predominantly *Klebsiella* spp, *Enterobacter* spp, *E. coli*) and harboured an IncHI2 plasmid, which carried both bla_{IMP} and the putative mobile colistin resistance gene *mcr-9*. Detailed phylogenetic analysis identified two distinct IncHI2 plasmid lineages, A and B, both of which showed significant association with patient movements between four hospital sites and across medical specialities.

Combined, our patient network and plasmid analyses demonstrate an interspecies, plasmid-mediated outbreak of bla_{IMP}CPE, which remained unidentified during standard microbiology and infection control investigations. With whole genome sequencing (WGS) technologies and large-data incorporation, the outbreak investigation approach proposed here provides a framework for real-time identification of key factors causing pathogen spread. Analysing outbreaks at the plasmid level reveals that resistance may be wider spread than suspected, allowing more targetted interventions to stop the transmission of resistance within hospital networks.

Introduction

Infections by carbapenemase-producing Enterobacterales (CPE) pose a substantial clinical, operational, and financial challenge[1]. These organisms are associated with high morbidity and mortality, and therapeutic options are severely restricted[2,3]. Carbapenemase genes are frequently carried on plasmids, which can easily transfer between bacterial species. CPE outbreaks involving different bacterial species are often unrecognised, as many plasmids are variable in their gene content[4] and have a broad host range[5]. The metallo-β-lactamase – (MBL) Imipenemase gene (IMP)-1, was first described in *Pseudomonas aeruginosa*[6], whilst IMP genes have spread globally, outbreaks in Enterobacterales species are mostly sporadic and often localised to specific geographical locations[7-9]. IMP genes are rarely isolated in the UK, but the number of IMP encoding isolates within Enterobacterales species referred to UK Health Security Agency (UK HSA) has been increasing[10,11].

Therapeutic options are particularly limited for MBL CPEs, with limited efficacy of the newer beta-lactam combination agents[2,3]. Colistin and polymyxin B remain the last-line therapeutic agents in most countries, partly due to lack of access to newer agents; yet resistance is increasing globally, particularly through mobilised colistin resistance (*mcr*) genes on plasmids[12,13]. Since the first discovery of *mcr*-1 in 2016[14], ten *mcr* genes (*mcr*-1 to *mcr*-10) have been described to date, presenting a substantial global healthcare challenge[13,15]. Although *mcr* genes are typically associated with phenotypic polymyxin resistance, *mcr-9*, identified in 2019, does not appear to confer direct colistin resistance[7,16]. It has been detected in clinical isolates co-harbouring different resistance genes in Australia (with $bla_{\text{IMP-4}}$), South Africa (with ESBLs), Egypt and the Netherlands (with bla_{VIM}) and Japan (with $bla_{\text{IMP-1}}$)[7,8,17-20], and is widespread in isolates from human, animal and environmental origins, in a wide range of bacterial species[7,21].

Person-to-person contact is a route of transmission for many infectious diseases. Consequently, understanding the patterns of these contacts, especially in healthcare settings, can offer detailed insight for targeted interventions[22]. Contact patterns, however, become increasingly complex when considering large numbers of interacting individuals over long-time spans, bacterial interactions with the human host and environment, bacterial behaviour under antibiotic and disinfectant pressure, and plasmids that can be shared by multiple bacterial species. Network models provide flexible tools to capture such complex contact patterns,with the availability of

well-developed analytical toolboxes they offer robust and reproducible methodology that has become widespread across disciplines[23]. Network models of infectious diseases[24-26] incorporate both person-to-person transmission through contact networks[27-30] and spatial spread through networks representing physical locations and their relationships[31,32]. So far, studies utlising network models of patient contacts in combination with detailed genomic analysis of bacterial isolates have demonstrated their promise through increasing the detail in outbreak characterisation [33,34].

Here we describe the emergence of bla_{IMP} and *mcr-9* co-harboured by diverse Enterobacterales species that have been isolated from patients across a regional network of London hospitals between 2016 and 2019. We combine in-depth phylogenomic analysis of the IncHI2 plasmid containing *mcr-9* and bla_{IMP} with a network model of patient movements to provide novel insight into the spread and dissemination of bla_{IMP} and *mcr-9* between bacterial species and patients over 3 years, with implications for how CPE clusters are identified and managed in the hospital setting.

Results

Emergence of bla_{IMP}CPE

Following the introduction of enhanced CPE screening in 2015[35], a CPE-carrying bla_{IMP} (bla_{IMP}CPE) was first observed in our hospital group in June 2016 from a routine rectal screening sample. From November 2016, increasing numbers of bla_{IMP}CPE isolates were identified across diverse Enterobacterales species at three of the acute hospital sites (representing 64% of all inpatient beds at our institution) (**Figure 1a**), and simultaneously at two other external hospitals in the regional network, all served by the same microbiology diagnostics laboratory, and closely connected by regular inter-hospital patient transfers. The highest incidence of bla_{IMP}CPE cases occurred in January and April 2019 (**Figure 1b**). In total, bla_{IMP}CPE were isolated from screening or clinical samples from 116 patients who were admitted to our hospital group up until the end of November 2019, when new cases rapidly dropped, despite no targeted changes in infection control interventions. However, only two clusters of cases, (totalling 5 out of 116 cases), fitted the real-time infection control outbreak definition, i.e. two or more cases of the same bacterial species with the same resistance mechanism overlapping in time and space. Pulsed Field Gel Electrophoresis (PFGE) typing

performed at the public health laboratories for those 5 cases had confirmed similarity, suggesting cross transmission events had occured.

To reveal the operational burden of bla_{IMP}CPE, the daily number of occupied beds was computed for all colonised patients. The analysis revealed a continuous burden of patients colonised with bla_{IMP}CPE which was far higher than expected from the new case incidence, particularly for those colonised with *Enterobacter* species (**Figure 1a,b**), with 424 total inpatient bed days per month at the peak (March 2019), across the hospital network (**Figure 1c**).

Figure 1: Characteristics of confirmed bla_{IMP}CPE patient cases and distribution of CPE species. (A) Total number of days patients (rows labelled by CPE isolate identifier) were present in a hospital ward before confirmed to be colonised/infected with bla_{IMP}CPE (*in-hospital infectious period*) and infection prevention and control measures were taken. Additionally, patients with known carriage of bla_{IMP}CPE but without sequenced isolates are shown as unlabelled rows. Patients with paired isolates (IMP 22/24, IMP 25/33, IMP 96/97, IMP 100/101) are on adjacent rows. **(B)** Monthly total number of confirmed bla_{IMP}CPE isolates detected in patients during the study period from 2016 to 2019. **(C)** Weekly cumulative number of bed days of patients with confirmed carriage of bla_{IMP}CPE before they identified as colonised with bla_{IMP}CPE over their infectious in-hospital period. Colours in each of the panels indicate the genus of CPE identified in the study.

Contact network of *bla*IMPCPE *patients*

A detailed patient contact network for all 116 *bla*IMPCPE colonised/infected patients who were
identified in our hospital group, including data from their admissions at the other regional
hospitals, confirmed that 77/116 (66%) patients were in contact with at least one other
*bla*IMPCPE colonised/infected patient (**Figure 2**). The analysis showed that the median number
of contacts was two per patient, however, contact numbers displayed substantial variation. For
example, one patient (Case 85, cluster 1.5) had 10 contacts with 9 other patients which linked
them to 8% of all cases (**Table S1**) and therefore identified the patient as a potential 'super-
spreader' in the outbreak.

Across all *bla*IMPCPE patient contact pairs, 57/96 (59%) pairs differed in bacterial species (8
species identified carrying *bla*IMP), explaining why these contacts were not identified as
outbreak-related using the clinical infection control outbreak definition, which was reliant on
the same bacterial species rather than resistance mechanism. The network of patient contacts
split patients into 12 separate clusters, with interactions occurring across hospitals, as patients
were transferred between wards and sites **(Figure 2)**. The largest cluster (Cluster 1) contained
45 patients and was partitioned into a further seven subclusters, labelled 1.1 to 1.7, that
comprised 13, 12, 2, 6, 5, 5 and 2 patients, respectively (**Figure 2**). The analysis of *bla*IMPCPE
patient contacts, at regional, individual hospital and ward level, suggested that different species

of *bla*~IMP~CPE were involved intransmission events (**Figure 2**). This multi-species involvement

in the ongoing outbreak prompted genomic analysis of the available *bla*~IMP~CPE isolates.

Figure 2. Patient contact network of interactions among *bla*~IMP~CPE positive patients.

Each node of the network represents a patient (coloured according to *bla*~IMP~CPE isolate

species) and each edge represents a contact between two patients. Edges represent recorded

contacts between confirmed *bla*~IMP~CPE cases (patients present on the same ward on the same

day according to their electronic health records). The patient contact network identified 12

distinct clusters (each shaded in a different colour) based on disconnected components of

patient contact. Cluster 1, the largest cluster consisting of 45 patients, was further partitioned

into 7 subclusters using community detection with edges weighted by duration of contact (see

Supplementary methods – Community detection). Six patient contacts re-occurred over

different wards, indicated by additional edges connecting the same patients.

Genomic and phenotypic characterisation of bla*~IMP~*CPE isolates

A total of 85 bacterial isolates from 82/116 *bla*~IMP~CPE colonised patients were sequenced for

genomic analysis (**Table S2**). The *bla*~IMP~ gene was detected in 13 different bacterial species,

most commonly in members of the *Enterobacter cloacae* complex species (55/85), *K.*

pneumoniae (16/85) and *E. coli* (10/85) (**Table S2**). Most patients harbourd only one *bla*~IMP~

encoding species, but there were four patients who were colonised with two different bacterial

species that carried *bla*IMP (**Table S2**).

Genomic analysis identified that two alleles, *bla*IMP-1 (75/85) and *bla*IMP-4 (10/85), co-occurred

with two groups of plasmids: IncHI2 plasmid was identified in 72/85 isolates and IncN3 in

11/85 (**Figure 3**). One isolate of *K. aerogenes* (IMP79) carried both plasmids, and two isolates

(IMP47, IMP76) carried a different plasmid co-occurring with *bla*IMP-1. As expected, all

*bla*IMPCPE isolates had carried multiple β-lactam resistance genes in addition to other

antimicrobial resistance genes (**Table S3**), though only one isolate (IMP89) had another

carbapenem resistance gene, *bla*OXA-48 (**Table S3**).

Figure 3. Phylogenetic tree of 85 Enterobacterales genomes in this study. A neighbour-

joining phylogenetic tree of 85 Enterobacterales genomes was constructed using whole-

genome FastANI distances. The colour at the end of each branch indicates the species identity

for that isolate. The rings are as follows: the innermost ring indicates the type of plasmid

detected; the middle ring indicates the allelic variant of the *bla*IMP gene detected, and the outer

ring indicates the presence or absence of the *mcr-9* gene. NF = not found. The scale bar

indicates the average nucleotide distance (in percent) per site.

The multi-species distribution of the 72 IncHI2 plasmids among contact network linked patients, was highly suggestive of plasmid mobility. The *mcr*-9 gene was detected in 81% (69/85) of all *bla*IMPCPE isolates, all of which harboured the IncHI2 plasmid (94% 69/72 of IncHI2 plasmid carrying isolates; none of IncN3 plasmids had *mcr-9*). The *mcr-9* specific LAMP assay confirmed the presence of *mcr-9* with 100% concordance (**Table S4**) as previously reported[36]. MALDIxin (which detects modifications to the lipid A of the lipopolysaccharide as mediated by *mcr*1-5 and *mcr*-8[37]) did not detect any Lipid A modifications attributable to the *mcr*-9 gene in this study. Altogether, 12/84 isolates tested (all *Enterobacter* species) were resistant to colistin with MICs > 2 µg/ml (range 4 to >64 µg/ml) (**Table S4**), and 5 isolates demonstrated a skipped-well phenomenon[38] suggestive of colistin heteroresistance. However, determining the contribution of *mcr-9* or other mechanisms of resistance to this was beyond the scope of this study, and will be the subject of future investigations.

Population structure of IncHI2 plasmids

Phylogenetic analysis of the 72 IncHI2 plasmids (where reads were mapped to complete genomic sequence of plasmid pKA_P10 (NZ_CP044215.1)) identified two major lineages (A and B) and four sublineages (A.1, A.2, B.1 and B.2, **Figure 4**). While all the isolates were carrying multiple antimicrobial resistance (AMR) genes and multiple plasmids, only plasmids of the B.1 lineage carried *bla*IMP-4 and were present in isolates that also exclusively saw co-occurrence of *tet*(D), *qnrB2, bla*OXA-1 and *mph*(A) (**Figure 4**). Comparison of 233 plasmid core-genome single-nucleotide polymorphisms (cgSNPs) showed some congruency of cgSNPs with the plasmid sequences (**Figure S1**). Furthermore, this analysis grouped the IncHI2 plasmids into 15 polytomic clades (with identical SNP profiles), of which 12 showed identical cgSNPs after masking recombination regions (**Table S5** and **Figure S2**).

Figure 4. Dated phylogenetic tree showing the relationship between 72 IncHI2 plasmids identified in CPE isolates with *bla*IMP. The complete genome of IncHI2 plasmid pKA_P10 was used as the reference for variant calling. The tree was generated and rooted using BactDating. Branch colours denote major plasmid lineages, which are supported by bootstrap values of 100. At the tips of the tree, circles represent plasmids from routine rectal screening samples, and squares represent plasmids from clinical samples. Colours of the tip symbols indicate species. Lineage identifiers are labelled above lineage-specific root branches. The

column panel to the right of the phylogenetic tree indicates presence-absence of mobile antimicrobial resistance genes and plasmid replicons. NA: information was not available.

The root of the plasmid tree was dated in 1949 (95% CI: 1927-1971), with the most recent common ancestors of lineages A and B dated in 1965 (95% CI: 1943–1988) and 1967 (95% CI: 1945–1988), respectively (**Figure S3**). These estimates suggest that not only IncHI2 plasmids have been circulating amongst *Enterobacterales* for many decades, but also that they have undergone extensive evolution in the antibiotic era.

Plasmid lineages and patient cluster comparisons

To investigate if the identified IncHI2 plasmid lineages A and B were responsible for the emergence of *bla*IMPCPE in our hospital network, plasmid genomic data was overlaid onto the

patient contact network (**Figure 5, Figure S4).** IncHI2 plasmid lineage A (3/72 IncHI2

plasmids) was exclusively linked to patients in cluster 5 (**Figure 5**), with contacts confined to

the same hospital site, highly suggestive of a patient-to-patient or common-source transmission

event. The predominant IncHI2 plasmid clade B, representing 69/72 plasmids sequenced and

60% of all cases identified in the study, was distributed widely across the hospital network,

dominated Cluster 1 of patient contact clusters, and showed particularly strong association with

contacts at hospital site 3 (**Figure 5**). The IncN3 plasmid (9.5% of overall cases) displayed

strong association with patients in network Cluster 9, with contacts occurring at three different

hospitals (hospitals 1, 2 and 3) (**Figure 5**). Although 3 of the IncN3 cases in cluster 9 had been

identified as an outbreak (Cases 80, 81 and 82) at the time of initial isolate identification, 5

other linked cases have now been revealed by this combined network and plasmid analysis,

that were previously missed. Similarly, Cases 67 and 71 (Cluster 1.2), identified real-time as

related cases, are revealed by this combined analysis to be linked to multiple (n=34) other cases

across clusters 1.1, 1.2, 1.3 and 1.4, with the IncHI2 plasmid lineage B.2.

Figure 5. Patient contact network of interactions among patients focussed on specific

locations of contact and plasmid information. Each node represents a patient, edges

represent recorded ward contacts between confirmed bla_{IMP}CPE cases. The edges are coloured

according to the hospital site (site 1-4) and the width of the edge is proportional to the duration

of the contact. Cluster nodes are coloured according to the lineage of the IncHI2 plasmid

identified in whole genome sequenced bla_{IMP}CPE isolates; patients with isolates lacking the

IncHI2 plasmid are coloured in white; patients with no whole genome sequenced bla_{IMP}CPE

isolates are displayed as labels without a circle. The presence of an IncN3 plasmid in the

bla_{IMP}CPE isolates is denoted by dark-grey borders surrounding the nodes. Contact network

clusters are highlighted with individual colour shading as in Figure 4.

Bacterial sequence data was missing from some cases that could have clarified links between and within plasmid lineages A and B, as suggested by the contact network analysis. Similarly, data was not available for potentially linked cases identified at other regional sites but not transferred to our hospital network. Despite missing hospital network data, clear associations between plasmid clades and patient pathway network clusters are evident, supporting the hypothesis of plasmid spread throughout the course of this outbreak, particularly at hospital 3. When reviewing the cumulative burden of the plasmid clades over time, it becomes apparent that most of the the cumulative bed days of patients shown in Figure 1 are due to patients colonised with clade B2 (**Figure S5**).

Spatial and temporal analysis

Having identified four plasmid lineages among the 72 bla_{IMP}-CPE IncHI2 plasmids, a network was constructed capturing the structure of patient transfers among hospital wards and used to investigate the spatial distribution of the four major plasmid lineages **(Figure 6).** The smaller plasmid lineages A.1 and A.2 were present in only two and one patients, respectively, and hence, were locally constrained by a smaller set of patient movements. Plasmid lineage A.1 was localised within hospital 3 renal services and had no recorded patient inter-hospital

movement. Plasmid lineage A.2 was localised to a single patient movement within hospital 4,

then an inter-hospital movement to hospital 2. Patients with isolates in plasmid lineage B.1

were recorded on wards from hospital sites 1, 2, 3 and 5. However, inter-hospital transfers of

these patients were only recorded between hospitals 2 and 5, and hospitals 3 and 5. It is thus

likely that some transmission of plasmid lineages B.1 was missed, either through patients who

were not detected on screening, or these were patients whose isolates were not available for

sequencing. Plasmid lineage B.2 was spatially more distributed with recorded multiple bi-

directional inter-hospital transfers among hospital sites 1, 2, 3 and 5. Looking at the association

between plasmid lineage clusters and ward/specialties over the study periodthe most common

associations across all specialities were critical care and renal services (**Table S6**). The only

exception was general internal medicine and general surgery predominated in plasmid lineage

B.1 at hospital site 5, which has more general wards and less specialist services than the other

hospital sites in the network. Despite the predominance of cases being identified in specialties

with high risk for invasive disease, only 4 clinical infections were identified during the study,

and no blood stream infections (**Table S2**)

Figure 6. **Spatial analysis representing the pathways of patient transfers among wards**

and hospitals. Movement of patients carrying bla_{IMP}CPE are represented by arrows between

wards (nodes) and the five hospitals (dotted ovals). For hospital site 5, only hospital-level data

was available, therefore, only intra-hospital movement is shown. Repeated transfers of patients

between wards are aggregated into edges withproportionally greater edge width. Plasmid

clusters, based on plasmid lineage (A.1, A.2, B.1 and B.2) are coloured accordingly. For

patients with multiple CPE isolates, transfers are duplicated to match each isolate uniquely in

the visualisation.

Patient transfer (plasmid cluster)
- A.1
- A.2
- B.1
- B.2
- Non-sequenced

Total patient transfers

Total patient days on ward

Although the first case was identified in 2016 (IMP 51/Case 91), a temporal network analysis

indicated it played only a minor role in onward transmission, due to the small number of

reachable patients over time (**Figure S6**). This may suggest that previous/subsequent patients

were not identified or point to the existence of an environmental reservoir. Considering the

timing and ordering of all contacts, IMP79 could have seeded up to 27 subsequent cases

(restricted to those in Cluster 1 in **Figure 2**), either directly or indirectlty, via possible

transmission routes. The patient carried the isolate that co-harboured both IncN3 and IncHI2

plasmids, potentially suggesting multiple transmission opportunities to, as well as, from, this

patient. There were 14 patients (15 isolates) from the 72 isolates carrying an IncHI2 plasmid,

which had a proportionately larger number of reachable time-ordered contacts, suggesting that

these patients may also have contributed substantially to transmission (**Figure S6**).

These integrated geno-spatio-temporal representations show that the spread of plasmids

containing bla_{IMP} followed patient movements for specialty care around the region, with

possibly multiple introductions and re-introductions of the plasmid at different points during

the patient journeys, and multiple chances for cross transmission events to have occurred.

Discussion

Following the detection of a new mechanism of resistance, investigation of its origin and mode

of transmission is challenging, especially in healthcare settings where investigations usually

focus on single species cross transmissions. With confounding factors such as multiple

bacterial species and spread over different hospital locations, new methods to investigate

potential outbreaks are much needed. The incorporation of plasmid genomics and patient

networks into our analysis changed the way the emergence of bla_{IMP}CPE was visualised and

produced a clearer understanding of the high-risk ward locations and pathways for potential

cross transmission in our regional healthcare system. As patients were found to follow common

routes, with regular re-encounters, this information can provide dynamic risk assessments to

be introduced along those pathways, to prevent future cross transmission events of any

healthcare-associated pathogen from occurring[39]. Detailed genomic analysis of plasmids

enhanced our understanding of the relatedness of different patient isolates to the network

analysis. It moreover revealed concerning information about unsuspected resistance

mechanisms, with potential for antibiotic treatment failures that were missed on conventional

laboratory susceptibility testing. In this study, we characterized IncHI2 plasmids as the main

vehicle in horizontal transfer of the metallo-β-lactamase gene bla_{IMP}. A dated phylogeny of the

IncHI2 plasmid carrying bla_{IMP}, indicated that this plasmid underwent a clonal expansion

around 2008, about 8 years before the onset of the outbreak described in our study. IncHI2

plasmids are common, large plasmids with a wide host range, that has been reported globally[19],

and more recently have been reported to be associated with colistin resistance genes *mcr*-1 and

mcr-3[40,41]. Although in our study we detected *mcr-9* in 81% of the isolates tested, we did not

find evidence of phenotypic expression of this gene, in line with previous observations[7,42]. We

identified the predominant IncHI2 plasmid in multiple different bacterial species across linked

patients, highlighting the need for integration of genomics into routine clinical practice.

Although this study focussed on the emergence of the bla_{IMP} carbapenemase gene in our

hospital network, it supports the concept that plasmid analysis across different resistance

mechanisms as well as among different species should be the standard for investigations in the

future. The small number of clinical infections from this outbreak in comparison to other CPE

outbreaks from our hospital network[43] and other reports of bla_{IMP} CPE[7,19] is noteworthy, and

poses questions about the wider importance of this plasmid and the resistance mechanisms

revealed in this study. This observation reinforces the argument that screening for silent

carriage of CPE in hospitals is key to preventing spread[44-46], and cautious antimicrobial

stewardship is essential to prevent expression of hidden resistance mechanisms[47].The

association of IMP carbapenemases and drainage[48,49], leads to speculation that the acquisition

of this plasmid may provide an environmental survival advantage, enhancing the potential for

spread between human hosts through a poorly maintained plumbing environment[50,51].

We acknowledge some limitations of our study. Firstly, long-read data was available only for

two isolates, pKA_P10 and pEB_P9 (NZ_CP043767.1)[10] , and though both were tested as

references, assumptions had to be made for in-depth genomic analysis of our plasmids. As a

result, our plasmid tree may omit some similarities and differences between identified IncHI2

plasmids. Secondly, full pathway data across the hospital during the three years of the outbreak

was only available for identified positive cases, not for all patients. It was therefore not possible

to fully establish potential missed cases flagging as close contacts but with potential for missed

screening or false negative results. Full pathway movement data for all positive cases identified

within our hospital network was available, yet neither pathway details nor genomic data were

available for other bla_{IMP}CPE-positive cases identified in the two other regional hospitals who

did not visit our institution, thus reducing the understanding in our analysis. Interactions at

other potential hospital locations such as interventional imaging or endoscopy were not

examined in this study, nor was environmental sampling performed, which could inform future

studies on modes of transmission. Nevertheless, this study highlights a previously unidentified

extent of transmission and thus provides valuable new insights into the spread of an emerging

resistance mechanism. Moreover, our novel multi-layered methodology, incorporating plasmid

phylogeny with contact network analysis, provides invaluable tools for outbreak investigation

that can be generalised to a wide range of scenarios.

Materials and Methods

Clinical Setting

This study was carried out using data from a regional network of London hospitals, comprising

of 7 hospital sites with a total of 2000 inpatient beds, with managerial responsibility assigned

to 2 different NHS Trusts. Cases of colonisation/infection with bla_{IMP}CPE were identified at 3

hospital sites from one NHS trust (referred to as our Institution in the results) and 2 hospital

sites from the other. All hospitals in the network are are served by the same centralised

microbiology laboratory and patients are frequently transferred between hospitals in the

regional network for specialist investigations and treatment.

Sampling, microbiology, antimicrobial susceptibility testing and detection of mcr-9

bla_{IMP}-encoding Enterobacterales isolates were cultured from rectal screens or clinical samples

as part of routine diagnostic testing between June 2016 and November 2019. Rectal screens

were cultured on chromogenic media (Colorex™ mSuperCARBA™, E&O Laboratories, UK),

and species identification was performed using Biotyper MALDI-TOF MS (Bruker Daltonics,

Germany) according to the manufacturer's protocol. Antimicrobial susceptibility testing was

performed according to EUCAST guidelines [52], colistin susceptibility testing was performed

by broth microdilution on all viable isolates according to manufacturer's instructions

(MICRONAUT, BioConnections, UK) [52] . The identification and detection of carbapenemase

genes was by PCR (Xpert® Carba-R, Cepheid Inc, USA). After February 2018, all isolates were

tested using the RESIST 3 O.K.N. K.SeT (Coris BioConcept, Belgium), and negative isolates

had the carbapenem inactivation method (CIM) performed, and CIM positive isolates then had

PCR (Xpert® Carba-R, Cepheid Inc, USA) performed as previously described[35]. bla_{IMP} positive

isolates with suspected links were sent to UK Health Security Agency (UK HSA) reference

laboratories for typing by Variable Number Tandem Repeat VNTR (*Klebsiella* sp.) or Pulsed

Field Gel Electrophoresis PFGE (all other species).

Loop-mediated isothermal amplification (LAMP) was used to examine the prevalence of *mcr-*

9 amongst bla_{IMP}CPE isolates as described previously [53]. MALDIxin testing was conducted to

confirm phenotypic colistin resistance, as previously described on isolates IMP 1-58 only[37].

Whole-genome sequencing

Bacterial isolates were grown on Columbia Blood Agar (Oxoid Ltd, UK), incubated aerobically

at 37°C overnight before DNA extraction. Genomic DNA was extracted using GenElute™

Bacterial Genomic DNA Kit (Sigma-Aldrich, USA) following manufacturer instructions.

Multiplexed DNA libraries were generated with Nextera XT (Illumina, USA) and were

sequenced for a minimum of 100-fold average coverage on Illumina HiSeq 4000 systems under

a   150-bp   paired-end   layout.   Seven   isolates   (IMP09/CB_P14_L5_12.18,

IMP29/KP_P12_L5_11.18,   IMP33/EB_P9_L5_03.19,   IMP40/EB_P8_L5_01.19,

IMP41/KQ_P11_L5_04.19, IMP42/KA_P10_L5_03.19, IMP45/ EB_P13_L5_04.19) were

also sent to UK HSA for MinION (Oxford Nanopore, UK) long-read sequencing [10].

Sequence data processing and analyses

Short-read sequence reads were quality assessed using FastQC v0.11.6 [54]. Raw reads were trimmed using Trimmomatic v0.39[55] and were *de novo* assembled using SPAdes v3.11 [56] and annotated with Prokka [57]. Species identification was confirmed with KmerFinder 3.2 [58]. Multilocus sequence typing (MLST) was performed with the MLST v1.8 online server [59] using the Achtman MLST scheme for *E. coli* curated on PubMLST and EnteroBase[60] and the Pasteur MLST scheme for other *Enterobacterales* isolates. AMR genes, and plasmid typing (**Table S7**) were done using ResFinder v3.2 [61], and PlasmidFinder v2.1 [62], respectively. A neighbour-joining tree of CPE isolates used in this study was generated from assemblies using FastANI v1.3 based on whole-genome average nucleotide identities [63].

Phylogenetic analysis of IncHI2 plasmids

A reference complete genome of IncHI2 plasmids was selected from three candidates: pKA_P10 (NZ_CP044215.1; 334 kbp) from *Klebsiella aerogenes*, pEB_P9 (NZ_CP043767.1; 331 kbp) from *Enterobacter hormaechei* isolate[10] and R478 (NC_005211.1; 275 kbp) identified in *Serratia marcescens*[64]. The selection aimed to maximise: (i) overall coverage of each reference genome by reads, (ii) the number of plasmid core-genome SNPs identified from the read alignment, and (iii) bootstrap values supporting major lineages in the plasmid phylogenetic tree reconstructed using the same approach based on each reference. Eventually, plasmid pKA_P10 was determined as the reference for the rest of analysis (**Figures S7**).

Repetitive regions ($\geq$90% nucleotide identity) in the reference plasmid genome were identified using NUCmer of MUMer v3.1[65]. Processed short reads of 73 IMP isolates harbouring IncHI2 plasmids were further filtered for a minimum length of 50 bp using Trimmomatic for improving mapping specificity, given estimated read depths >100 fold per bacterial genome before and after the filtering. Filtered reads were mapped to the reference pKA_P10 using Snippy (github.com/tseemann/snippy) with parameters set to as follows: mapping quality $\geq$ Q60, base quality $\geq$ Q20, base coverage $\geq$10 folds, and variant-call quality $\geq$ Q100. Plasmid pIMP06 was excluded from further steps as its reads covered only 27% of the reference genome. Core-genome SNPs (cgSNPs, sites presented in all genomes) outside of repetitive regions were identified and a whole-sequence alignment of remaining 72 plasmids against the reference was generated using Snippy. A neighbour-joining tree were generated from recombination-free cgSNPs using pairwise SNP counts as distance metrics (namely, SNP distances) (**Figure S8**).

Maximum-likelihood phylogenetic reconstruction from the whole-genome plasmid alignment
(which includes invariant sites, gaps, and masked regions) was performed using IQ-Tree
v1.6.12[66], with the best-fit model "HKY+F+R2" (determined through extended model
selection and full-tree search implemented in IQ-Tree), 10 independent runs (for tree selection),
and 500 bootstrap replicates (for branch supports). The resulting unrooted tree was corrected
for recombination using ClonalFrameML [67] with its default parameters and the whole-genome
alignment of plasmids. Major plasmid lineages were identified from topology of the corrected
tree. Detected recombination regions and reconstructed ancestral sequences were drawn against
the corrected tree using the script cfml_results.R from ClonalFrameML. Sample information
and AMR gene content of IMP isolates harbouring these plasmids were drawn against the
plasmid tree using iTOL v5[68]. The dating of internal nodes of the plasmid tree was inferred by
BactDating using the mixed-Gamma model, root-finding algorithm, 5×10^4 random walks of
Markov Chain Monte Carlo (MCMC), and accounting for recombination identified by
ClonalFrameML. To evaluate the robustness of tree reconstruction, a cgSNP tree was generated
by IQ-Tree (best-fit model: "TIM3e+ASC"; other parameters were the same as reconstructing
the whole-genome tree) and corrected for recombination using ClonalFrameML (with default
parameters). Tree comparisons (**Figures S9–S12**) were performed in R v3.6.3 with function
tanglegram in package dendextend[69].
504
Patient contact network analysis
Ward locations and timings were extracted from the Electronic Health Record (EHR) data for
bla_{IMP}CPE positive patients to establish patient movement history. Movement to other
locations for procedures during an inpatient episode, such as endoscopy, interventional
radiology or operating theatres, was not examined. To reveal potential transmission events
structure, a patient contact network was constructed from patient movement history, which
included information on when and where patients were on the same ward (for more details
please see Suppl. Methods). Time aggregated patient contacts were subsequently clustered to
reveal groups of patients more closely linked together using the Walktrap community detection
algorithm[70]. In this framework, contacts were also weighted by the duration of time spent
together across their hospital episode, capturing that the probability of transmission increases
with the amount of time two patients spend together[71]. Moreover, a temporal analysis of patient
interactions was performed over the time ordered sequence of contacts between patients[72] and
used to assess patient roles and position in transmission (for more details please see Suppl.
Methods).

521 For insights into the spatial distribution of bla_{IMP}CPE, a spatial network of ward/hospital

522 network, connected by the same bla_{IMP}CPE patient movement was generated. From this spatial

523 network, the following was calculated: total number of days patient has spent on each ward

524 over the time before they were known to be colonised with bla_{IMP}CPE and infection prevention

525 and control measures were taken (their infectious in-hospital period); network structure to

526 determine ward/hospital spread of bla_{IMP}CPE; highly visited wards from patients of each

527 specific genomic sub-cluster were also determined (using PageRank[73,74]) and contrasted for

528 evidence of genomic localisation to hospital spaces. All network analysis and visualisation was

529 completed in Python, R, and Cytoscape[75].

531 *Data availability*

532 Illumina reads and draft genome assemblies of 85 isolates were deposited in European

533 Nucleotide Archive (ENA) under the BioProject accession PRJEB38818.

535 *Ethics*

536 This study was carried out in accordance with ethics reference 21/LO/0170 (279677), protocol

537 21HH6538 Investigation of epidemiological and pathogenic factors associated with infectious

538 diseases.

542 **Acknowledgements**

543 We thank the staff of the diagnostic microbiology laboratory of North West London Pathology,

544 for isolate collection and storage. We would also like to acknowledge the support of the

545 Imperial College Healthcare Trust NIHR Biomedical Research Centre (BRC). The

546 Imperial BRC Genomics Facility has provided resources and support that have contributed to

547 the research results reported within this paper. The Imperial BRC Genomics Facility is

548 supported by NIHR funding to the Imperial Biomedical Research Centre.

550 **Funding**

551 This work has been funded in part by faculty of medicine, Siriraj hospital, Mahidol university,

552 Thailand (awarded to AB) and Medical Research Council Clinical Academic Research

553 Fellowship scheme (awarded to FD, grant number MR/T005254/1).

AM is funded in part by a scholarship from the Medical Research Foundation National PhD
Training Programme in Antimicrobial Resistance Research (MRF-145-0004-TPG-AVISO),
the EPSRC Centre for Mathematics of Precision Healthcare (EP/N014529/)and through
an National Institute for Health Research Senior Research Investigator Award. AM and MB
acknowledge funding from EPSRC grant EP/N014529/1 to MB, supporting the EPSRC Centre
for Mathematics of Precision Healthcare. AS is funded by a PhD studentship from the Medical
Research Council Doctoral Training Award to Imperial College London (MR/N014103/1).
AME acknowledges support from the National Institute for Health Research (NIHR) Imperial
Biomedical Research Centre (BRC). EJ is an Imperial College Research Fellow, funded by
Rosetrees Trust and the Stoneygate Trust (M683). HA was supported by the Imperial Health
Charity. XD is supported by the NIHR Health Protection Research Unit in Genomics and
Enabling Data. AL acknowledges funding from the National Institute for Health Research
Health Protection Research Unit (NIHR HPRU) in Modelling Methodology at Imperial
College London (grant HPRU-2012–10080) and the National Institute for Health Research
(NIHR) Health Protection Research Unit in Healthcare Associated Infections and
Antimicrobial Resistance at University of Oxford (NIHR200915) in partnership with UK
Health Security Agency. AH is a National Institute for Health Research (NIHR) Senior
Investigator. AH, FD, EJ, AB, YW, ME are affiliated with the National Institute for Health
Research Health Protection Research Unit (NIHR HPRU) in Healthcare Associated Infections
and Antimicrobial Resistance at Imperial College London in partnership with the UK Health
Security Agency (previously PHE), in collaboration with, Imperial Healthcare Partners,
University of Cambridge and University of Warwick. This report is independent research
funded by the National Institute for Health Research. The views expressed in this publication
are those of the author(s) and not necessarily those of the NHS, the National Institute for Health
Research, the Department of Health and Social Care or the UK Health Security Agency.

Transparency declarations

Author contributions

Conceptualization: FD, EJ, AH, MB, FB, XD, ME. Data curation: YW, BA, EJ, AM, FD, SM,
AW, ME. Formal analysis: AB, AM, YW, FB, FD, EJ, AW, MB, AH, ME. Funding
acquisition: AB, FD, AH, ME. Investigation: AB, YW, AM, HA, RP, AL, AA, JT, LM, KM-

C, AS, GL-M, JR-M. Methodology: FD, AM, AW, SM, YW, EJ, BA, XD, MB, JO, KH, ME, JT, MG. Project administration: FB, AH, FD, EJ. Resources: FD, AH, AE, HD, KH, JT. Supervision: FD, EJ, AH, MB. Validation: KH, JT, ME, HD, AW, SM. Visualization: AB, AM, YW, EJ, FD. Writing – original draft and further drafts: AB, AM, YW, EJ, FD, FB. Writing – review & editing: All authors.

References

Otter, J. A. *et al.* Counting the cost of an outbreak of carbapenemase-producing Enterobacteriaceae: an economic evaluation from a hospital perspective. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* **23**, 188-196, doi:10.1016/j.cmi.2016.10.005 (2017).

Doi, Y. Treatment Options for Carbapenem-resistant Gram-negative Bacterial Infections. *Clinical Infectious Diseases* **69**, S565-S575, doi:10.1093/cid/ciz830 (2019).

Pranita D. Tamma, S. L. A., Robert A. Bonomo, Amy J. Mathers, David van Duin, Cornelius J. Clancy. *Infectious Diseases Society of America Antimicrobial Resistant Treatment Guidance: Gram-Negative Bacterial Infections*, <https://www.idsociety.org/practice-guideline/amr-guidance/> (2020).

Branger, C. *et al.* Extended-spectrum β-lactamase-encoding genes are spreading on a wide range of Escherichia coli plasmids existing prior to the use of third-generation cephalosporins. *Microbial genomics* **4**, doi:10.1099/mgen.0.000203 (2018).

Li, X. *et al.* Diversification of broad host range plasmids correlates with the presence of antibiotic resistance genes. *FEMS microbiology ecology* **92**, doi:10.1093/femsec/fiv151 (2016).

Walsh, T. R., Toleman, M. A., Poirel, L. & Nordmann, P. Metallo-beta-lactamases: the quiet before the storm? *Clinical microbiology reviews* **18**, 306-325, doi:10.1128/cmr.18.2.306-325.2005 (2005).

Kananizadeh, P. *et al.* Emergence of carbapenem-resistant and colistin-susceptible Enterobacter cloacae complex co-harboring blaIMP-1 and mcr-9 in Japan.

Roberts, L. W. *et al.* Genomic analysis of carbapenemase-producing Enterobacteriaceae in Queensland reveals widespread transmission of bla IMP-4 on an IncHI2 plasmid. *Microbial genomics*, doi:10.1099/mgen.0.000321 (2019).

Matsumura, Y. *et al.* Global Molecular Epidemiology of IMP-Producing Enterobacteriaceae. *Antimicrobial agents and chemotherapy* **61**, doi:10.1128/aac.02729-16 (2017).

Turton, J. F. *et al.* IncN3 and IncHI2 plasmids with an In1763 integron carrying bla (IMP-1) in carbapenem-resistant Enterobacterales clinical isolates from the UK. *J Med Microbiol* **69**, 739-747, doi:10.1099/jmm.0.001193 (2020).

England, P. H. Vol. 15 *Health Protection report* 1-14 (Assets publishing service UK government, 2021).

Poirel, L., Jayol, A. & Nordmann, P. Polymyxins: Antibacterial Activity, Susceptibility Testing, and Resistance Mechanisms Encoded by Plasmids or Chromosomes. *Clinical microbiology reviews* **30**, 557-596, doi:10.1128/cmr.00064-16 (2017).

Ling, Z. *et al.* Epidemiology of mobile colistin resistance genes mcr-1 to mcr-9. *Journal of Antimicrobial Chemotherapy* **75**, 3087-3095, doi:10.1093/jac/dkaa205 (2020).

Liu, Y. Y. *et al.* Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* **16**, 161-168, doi:10.1016/s1473-3099(15)00424-7 (2016).

Shen, Y. *et al.* Heterogeneous and Flexible Transmission of mcr-1 in Hospital-Associated Escherichia coli. *mBio* **9**, doi:10.1128/mBio.00943-18 (2018).

Carroll, L. M. *et al.* Identification of Novel Mobilized Colistin Resistance Gene mcr-9 in a Multidrug-Resistant, Colistin-Susceptible Salmonella enterica Serotype Typhimurium Isolate. *mBio* **10**, doi:10.1128/mBio.00853-19 (2019).

Osei Sekyere, J., Maningi, N. E., Modipane, L. & Mbelle, N. M. Emergence of *mcr*-9.1 in Extended-Spectrum-β-Lactamase-Producing Clinical Enterobacteriaceae in Pretoria, South Africa: Global Evolutionary Phylogenomics, Resistome, and Mobilome. *mSystems* **5** (2020).

Soliman, A. M. *et al.* Emergence of a Multidrug-Resistant Enterobacter hormaechei Clinical Isolate from Egypt Co-Harboring mcr-9 and bla(VIM-4). *Microorganisms* **8**, doi:10.3390/microorganisms8040595 (2020).

Macesic, N. *et al.* Silent spread of mobile colistin resistance gene mcr-9.1 on IncHI2 'superplasmids' in clinical carbapenem-resistant Enterobacterales. *Clinical Microbiology and Infection*, doi:https://doi.org/10.1016/j.cmi.2021.04.020 (2021).

Hendrickx, A. P. A. *et al.* A genetic cluster of MDR Enterobacter cloacae complex ST78 harbouring a plasmid containing blaVIM-1 and mcr-9 in the Netherlands. *JAC-Antimicrobial Resistance* **3**, doi:10.1093/jacamr/dlab046 (2021).

Li, Y. *et al.* Characterization of the global distribution and diversified plasmid reservoirs of the colistin resistance gene mcr-9. *Scientific Reports* **10**, 8113, doi:10.1038/s41598-020-65106-w (2020).

Meyers, L. Contact network epidemiology: Bond percolation applied to infectious disease prediction and control. *Bulletin of the American Mathematical Society* **44**, 63-86 (2006).

Newman, M. E. J. The Structure and Function of Complex Networks. *SIAM Review* **45**, 167-256, doi:10.1137/s003614450342480 (2003).

Pastor-Satorras, R., Castellano, C., Van Mieghem, P. & Vespignani, A. Epidemic processes in complex networks. *Reviews of Modern Physics* **87**, 925-979, doi:10.1103/RevModPhys.87.925 (2015).

Klovdahl, A. S. Social networks and the spread of infectious diseases: The AIDS example. *Social Science & Medicine* **21**, 1203-1216, doi:https://doi.org/10.1016/0277-9536(85)90269-2 (1985).

Keeling, M. J., Woolhouse, M. E. J., May, R. M., Davies, G. & Grenfell, B. T. Modelling vaccination strategies against foot-and-mouth disease. *Nature* **421**, 136-142, doi:10.1038/nature01343 (2003).

Salathé, M. *et al.* A high-resolution human contact network for infectious disease transmission. *Proceedings of the National Academy of Sciences* **107**, 22020-22025, doi:10.1073/pnas.1009094108 (2010).

Meyers, L. A., Newman, M. E. J. & Pourbohloul, B. Predicting epidemics on directed contact networks. *Journal of Theoretical Biology* **240**, 400-418, doi:https://doi.org/10.1016/j.jtbi.2005.10.004 (2006).

Ancel Meyers, L., Newman, M. E. J., Martin, M. & Schrag, S. Applying network theory to epidemics: control measures for Mycoplasma pneumoniae outbreaks. *Emerging infectious diseases* **9**, 204-210, doi:10.3201/eid0902.020188 (2003).

Newman, M. E. J. Spread of epidemic disease on networks. *Physical Review E* **66**, 016128, doi:10.1103/PhysRevE.66.016128 (2002).

Brockmann, D. & Helbing, D. The Hidden Geometry of Complex, Network-Driven Contagion Phenomena. *Science* **342**, 1337, doi:10.1126/science.1245200 (2013).

Colizza, V., Barrat, A., Barthélemy, M. & Vespignani, A. The role of the airline transportation network in the prediction and predictability of global epidemics. *Proceedings of the National Academy of Sciences* **103**, 2015-2020, doi:10.1073/pnas.0510525103 (2006).

33 Kwong, J. C. *et al.* Translating genomics into practice for real-time surveillance and response
to carbapenemase-producing Enterobacteriaceae: evidence from a complex multi-
institutional KPC outbreak. *PeerJ* **6**, e4210-e4210, doi:10.7717/peerj.4210 (2018).
34 Gouliouris, T. *et al.* Quantifying acquisition and transmission of Enterococcus faecium using
genomic surveillance. *Nat Microbiol*, doi:10.1038/s41564-020-00806-7 (2020).
35 Otter, J. A. *et al.* Detecting carbapenemase-producing Enterobacterales (CPE): an evaluation
of an enhanced CPE infection control and screening programme in acute care. *Journal of
Antimicrobial Chemotherapy* **75**, 2670-2676, doi:10.1093/jac/dkaa192 (2020).
36 Rodriguez-Manzano, J. *et al.* Rapid Detection of Mobilized Colistin Resistance using a Nucleic
Acid Based Lab-on-a-Chip Diagnostic System. *Scientific reports* **10**, 8448, doi:10.1038/s41598-
020-64612-1 (2020).
37 Furniss, R. C. D. *et al.* Detection of Colistin Resistance in <span class="named-content genus-
species" id="named-content-1">Escherichia coli by Use of the MALDI Biotyper Sirius
Mass Spectrometry System. *Journal of clinical microbiology* **57**, e01427-01419,
doi:10.1128/jcm.01427-19 (2019).
38 Landman, D., Salamera, J. & Quale, J. Irreproducible and uninterpretable Polymyxin B MICs for
Enterobacter cloacae and Enterobacter aerogenes. *Journal of clinical microbiology* **51**, 4106-
4111, doi:10.1128/jcm.02129-13 (2013).
39 Fournier, S. *et al.* Efficiency of different control measures for preventing carbapenemase-
producing enterobacteria and glycopeptide-resistant Enterococcus faecium outbreaks: a 6-
year prospective study in a French multihospital institution, January 2010 to December 2015.
Euro Surveill **23**, 17-00078, doi:10.2807/1560-7917.ES.2018.23.8.17-00078 (2018).
40 Haenni, M. *et al.* Co-occurrence of extended spectrum β lactamase and MCR-1 encoding genes
on plasmids. *Lancet Infect Dis* **16**, 281-282, doi:10.1016/s1473-3099(16)00007-4 (2016).
41 Yin, W. *et al.* Novel Plasmid-Mediated Colistin Resistance Gene mcr-3 in Escherichia coli. *MBio*
8, doi:10.1128/mBio.00543-17 (2017).
42 Kieffer, N. *et al.* mcr-9, an Inducible Gene Encoding an Acquired Phosphoethanolamine
Transferase in Escherichia coli, and Its Origin. *Antimicrobial agents and chemotherapy* **63**,
doi:10.1128/aac.00965-19 (2019).
43 Otter, J. A. *et al.* Emergence and clonal spread of colistin resistance due to multiple mutational
mechanisms in carbapenemase-producing Klebsiella pneumoniae in London. *Scientific reports*
7, 12711, doi:10.1038/s41598-017-12637-4 (2017).
44 Otter, J. A., Mutters, N. T., Tacconelli, E., Gikas, A. & Holmes, A. H. Controversies in guidelines
for the control of multidrug-resistant Gram-negative bacteria in EU countries. *Clinical
microbiology and infection : the official publication of the European Society of Clinical
Microbiology and Infectious Diseases* **21**, 1057-1066, doi:10.1016/j.cmi.2015.09.021 (2015).
45 Jamal, A. J. *et al.* Infection prevention and control practices related to carbapenemase-
producing Enterobacteriaceae (CPE) in acute-care hospitals in Ontario, Canada. *Infection
Control & Hospital Epidemiology* **40**, 1006-1012, doi:10.1017/ice.2019.173 (2019).
46 Harris, A. D., McGregor, J. C. & Furuno, J. P. What Infection Control Interventions Should Be
Undertaken to Control Multidrug-Resistant Gram-Negative Bacteria? *Clinical Infectious
Diseases* **43**, S57-S61, doi:10.1086/504479 (2006).
47 Doron, S. & Davidson, L. E. Antimicrobial stewardship. *Mayo Clin Proc* **86**, 1113-1123,
doi:10.4065/mcp.2011.0358 (2011).
48 Amoureux, L. *et al.* Nosocomial Infections with IMP-19-Producing Pseudomonas aeruginosa
Linked to Contaminated Sinks, France. *Emerging infectious diseases* **23**, 304-307,
doi:10.3201/eid2302.160649 (2017).
49 Mollenkopf, D. F. *et al.* Carbapenemase-Producing Enterobacteriaceae Recovered from the
Environment of a Swine Farrow-to-Finish Operation in the United States. *Antimicrobial agents
and chemotherapy* **61**, e01298-01216, doi:10.1128/AAC.01298-16 (2017).

50   Heireman, L. *et al.* Toilet drain water as a potential source of hospital room-to-room
transmission of carbapenemase-producing Klebsiella pneumoniae. *The Journal of hospital*
infection **106**, 232-239, doi:10.1016/j.jhin.2020.07.017 (2020).
51   Regev-Yochay, G. *et al.* Sink traps as the source of transmission of OXA-48-producing Serratia
marcescens in an intensive care unit. *Infect Control Hosp Epidemiol* **39**, 1307-1315,
doi:10.1017/ice.2018.235 (2018).
52   EUCAST. Breakpoint tables for interpretation of MICs and zone diameters. Version 7.1.
European Committee on Antimicrobial Susceptibility Testing (2017).
53   Rodriguez-Manzano, J. *et al.* Rapid Detection of Mobilized Colistin Resistance using a Nucleic
Acid Based Lab-on-a-Chip Diagnostic System. *Sci Rep* **10**, 8448, doi:10.1038/s41598-020-
64612-1 (2020).
54   Andrews, S. FASTQC. A quality control tool for high throughput sequence data. (2010).
55   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics (Oxford, England)* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170
(2014).
56   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-
Cell Sequencing. *Journal of Computational Biology* **19**, 455-477, doi:10.1089/cmb.2012.0021
(2012).
57   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-2069,
doi:10.1093/bioinformatics/btu153 (2014).
58   Larsen, M. V. *et al.* Benchmarking of methods for genomic taxonomy. *Journal of clinical*
microbiology **52**, 1529-1539, doi:10.1128/jcm.02981-13 (2014).
59   Larsen, M. V. *et al.* Multilocus sequence typing of total-genome-sequenced bacteria. *Journal*
of clinical microbiology **50**, 1355-1361, doi:10.1128/jcm.06094-11 (2012).
60   Jolley, K. A., Bray, J. E. & Maiden, M. C. J. Open-access bacterial population genomics: BIGSdb
software, the PubMLST.org website and their applications. *Wellcome Open Res* **3**, 124,
doi:10.12688/wellcomeopenres.14826.1 (2018).
61   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J Antimicrob*
Chemother **67**, 2640-2644, doi:10.1093/jac/dks261 (2012).
62   Carattoli, A. *et al.* In silico detection and typing of plasmids using PlasmidFinder and plasmid
multilocus sequence typing. *Antimicrobial agents and chemotherapy* **58**, 3895-3903,
doi:10.1128/aac.02412-14 (2014).
63   Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature*
communications **9**, 5114, doi:10.1038/s41467-018-07641-9 (2018).
64   Gilmour, M. W., Thomson, N. R., Sanders, M., Parkhill, J. & Taylor, D. E. The complete
nucleotide sequence of the resistance plasmid R478: defining the backbone components of
incompatibility group H conjugative plasmids through comparative genomics. *Plasmid* **52**,
182-202, doi:https://doi.org/10.1016/j.plasmid.2004.06.006 (2004).
65   Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biology* **5**,
R12, doi:10.1186/gb-2004-5-2-r12 (2004).
66   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
Evolution **32**, 268-274, doi:10.1093/molbev/msu300 (2014).
67   Didelot, X. & Wilson, D. J. ClonalFrameML: efficient inference of recombination in whole
bacterial genomes. *PLoS computational biology* **11**, e1004041,
doi:10.1371/journal.pcbi.1004041 (2015).
68   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display
and annotation. *Bioinformatics (Oxford, England)* **23**, 127-128,
doi:10.1093/bioinformatics/btl529 (2006).

69   Galili, T. dendextend: an R package for visualizing, adjusting and comparing trees of
hierarchical   clustering.   *Bioinformatics   (Oxford,   England)*   **31**,   3718-3720,
doi:10.1093/bioinformatics/btv428 (2015).
70   Pons, P. & Latapy, M.   284-293 (Springer Berlin Heidelberg).
71   Gómez, S., Arenas, A., Borge-Holthoefer, J., Meloni, S. & Moreno, Y. Discrete-time Markov
chain approach to contact-based disease spreading in complex networks. *EPL (Europhysics*
Letters) **89**, 38009, doi:10.1209/0295-5075/89/38009 (2010).
72   Holme,   P.   &   Saramäki,   J.   Temporal   networks.   *Physics   Reports*   **519**,   97-125,
doi:https://doi.org/10.1016/j.physrep.2012.03.001 (2012).
73   Senanayake, U., Piraveenan, M. & Zomaya, A. The Pagerank-Index: Going beyond Citation
Counts in Quantifying Scientific Impact of Researchers. *PloS one* **10**, e0134794-e0134794,
doi:10.1371/journal.pone.0134794 (2015).
74   Page, L., Brin, Sergey, Motwani, Rajeev, Winograd, Terry. The PageRank Citation Ranking:
Bringing Order to the Web. (Stanford University, Stanford InfoLab Publication Server, 1999).
75   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res* **13**, 2498-2504, doi:10.1101/gr.1239303 (2003).