Parents' perceptions and intention to vaccinate their children against COVID-19: Results from a cross-sectional national survey in India

Running title: Children COVID-19 vaccination intention among parents

Bijaya Kumar Padhi[1*], Prakasini Satapathy[2], Vineeth Rajagopal[1*], Neeti Rustagi[3], Jatina Vij[1], Lovely Jain[1], Venkatesan Chakrapani[4], Binod Patro[5], Sitanshu Sekhar Kar[6], Ritesh Singh[7], Star Pala[8], Lalit Sankhe[9], Bhavesh Modi[10], Surya Bali[11], Tanvi Kiran[1], Kapil Goel[1], Arun Kumar Aggarwal[1], Madhu Gupta[1#]

*Contributed equally

#Corresponding author: Madhu Gupta, MD, Professor, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh-160 012, E-mail: madhugupta21@gmail.com

[1]Bijaya Kumar Padhi, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India-160 012

[2]Prakasini Satapathy, Department of Public Health, Utkal University, Odisha, India

[1]Vineeth Rajagopal, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

[3]Neeti Rustagi, Department of Community Medicine & Family Medicine, AIIMS Jodhpur,India

[1]Jatina Vij, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

[1]Lovely Jain, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

[4]Venkatesan Chakrapani, Centre for Sexuality and Health Research and Policy (C-SHaRP), Chennai; India

[5]Binod Patro, Department of Community and Family Medicine, All India Institute of Medical Sciences (AIIMS), Bhubaneswar, Odisha, India

[6]Sitanshu Sekhar Kar, Department of Preventive and Social Medicine, Jawaharlal Institute of Postgraduate Medical Education & Research (JIPMER), Puducherry, India

[7]Ritesh Singh, Department of Community and Family Medicine, All India Institute of Medical Sciences (AIIMS), Kalyani, West Bengal, India

[8]Star Pala, Department of Community Medicine, North Eastern Indira Gandhi Regional Institute of Health & Medical Sciences, (NEIGRIHMS), Mawdiangdiang, Shillong, Meghalaya, India

[9]Lalit Sankhe, Grant Medical College & JJ Hospital, Mumbai, Maharashtra, India

[10]Bhavesh Modi, Community Medicine Department, GMERS Medical College, Gandhinagar, Gujarat, India

[11]Surya Bali, Department of Community and Family Medicine, All India Institute of Medical Sciences(AIIMS), Bhopal, India

[1]Tanvi Kiran, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

[1]Kapi Goel, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

[1]Arun Kumar Aggarwal, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

[1]Madhu Gupta, Department of Community Medicine and School of Public Health, Post Graduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

Abstract

Background: Despite the success of adult vaccination against COVID-19, providing vaccines to children remains a challenge for policymakers globally. As parents are primary decision-makers for their children, we aimed to assess parents' perceptions and intentions regarding COVID-19 vaccination in India.

Methods: A cross-sectional web-based study was designed, parents or caregivers (N=770) were recruited through snowball sampling using Google form. Cross-tabulation was performed by parents' intention to vaccinate their children against COVID-19 virus with sociodemographic characteristics and their risk perception towards COVID-19, trust in the healthcare system, and their history of vaccine hesitancy behavior. Multivariable logistic regression analysis was performed to compute the predictors of child vaccination intention among Indian parents.

Results: 770 parents across the country have completed the survey. Of the 770 participants, 258 (33.5%) have shown intent to vaccinate their children. The stated likelihood of child vaccination was greater among parents who had a bachelor's degree or higher education (aOR: 1.98, 95% CI: 1.15-3.51); as well as among parents who intended to vaccinate themselves (aOR: 2.35, 95% CI: 1.30-4.67). Parental concerns centered around vaccine safety and side effects.

Conclusion: Indian parents reported high knowledge of the COVID-19 virus and were aware of the development of a novel vaccine. However, about one-third of parents intended to vaccinate their children, and about half of them were not sure whether to vaccinate their children or not against the COVID-19 virus. The study highlighted the need for health promotion strategies that promote vaccine uptake among parents.

Keywords: Vaccine acceptance, vaccine hesitancy, Children, Risk perception, Trust

Introduction

The World Health Organization (WHO) has reported more than 242 million confirmed cases of COVID-19 worldwide, including approximately 5 million deaths as of 20 October 2021(1). India reported 34 million confirmed cases and 450 thousand deaths as of 25[th] October 2021(1). An effective strategy to mitigate the morbidity and mortality of the COVID-19 and to ensure population higher levels of immunity is the development of an effective and safe vaccine for all populations, including children. In general, children with COVID-19 present with milder symptoms and are at lower risk of hospitalization and life-threatening complications(2). Numerous public health authorities, including the WHO and United States Centers for Disease Control and Prevention, advocate a vaccine for children. However, acceptance of the COVID-19 vaccine among parents or caregivers remains unexplored.

Since the start of the COVID-19 pandemic, a high number of vaccines is being developed(3). The WHO has declared on the 11[th] of July 2021 that 107 and 184 vaccine candidates are in clinical and pre-clinical stages, respectively, and at least 13 vaccines have been approved and administrated through four platforms(4). As of 25 October 2021, approximately 6 billion vaccine doses have been administrated worldwide(1). In India, as of 25 October 2021, the Ministry of Health and Family Welfare reported over one billion doses to adults(5). To expand the vaccine administration to children, the government of India is planning to implement a COVID-19 vaccine campaign for the age group of 12 to 18 years old(6).

Numerous studies on the COVID-19 vaccine acceptance shows that a lower number of parents were interested in vaccinating their children(7–14). A national survey in the United States reported less than one-half of parents are likely to administer a COVID-19 vaccine to their children(15). In Canada, 63% of parents intend to vaccinate their children against COVID-19(10). In Turkey, only 36.3% of parents were willing to have their children receive the COVID-19 vaccine(13). In China, COVID-19 vaccine hesitancy was found to be 8.44% among reproductive women(8). In a hospital-based study in Saudi Arabia, one-fourth of mothers were hesitant towards childhood immunization(11). To the best of our knowledge, none of the studies conducted in India address the intention and perception of parents' willingness to vaccinate their children in India.

Despite the success of adult vaccination globally, childhood immunization against the COVID-19 vaccine is now a challenge for policymakers. Including an effective and safe vaccine, a high level of vaccine acceptance among the parents or caregivers is also a key factor for global immunization success. Many factors such as fear about the vaccine safety and risk for a child to be infected by the virus impact parents' acceptance of the new COVID-19 vaccination program(13). Therefore, understanding parents' perception and acceptance of the COVID-19 vaccine will help in designing effective strategies for promotion and rolling out COVID-19 vaccination and beyond. To the best of our knowledge, this is the first community-based study in India that investigates parental willingness for COVID-19 vaccination. The study also explored factors associated with vaccine hesitancy among Indian parents or caregivers.

Methods

Study Design and Sample

We conducted a national online cross-sectional survey between November 2020 and January 2021 (before the introduction of the novel COVID-19 vaccine). The overall sample (N = 770) of parents were recruited across major geographical regions in India with an equal weightage to the urban and rural population. Respondents were adults who have one or more children 0–17 years old in their home and have access to the internet or telephone. We estimated a minimal sample size of 768, based on the maximum variability possible in the outcome variable in the population (i.e., a proportion of 0.50), with a margin of error of +/- 5% and 95% confidence intervals (CI). The design effect was kept at 2. Study participants were recruited through snowball sampling. Invitations to participate in the study were distributed to the respondents via email, social media: Twitter, Facebook, and the WhatsApp communication platform of the primary contacts of the study team and requested to transmit further for its maximum reach. Considerations were made to recruit the participants across major geographic regions of India.

To ensure rigor and validity, respondents had unique IDs, and 10% of respondents were contacted by telephone for cross verification of the presence of children 0–17 years old in their home.

Questionnaire Development

A first draft of the questionnaire was prepared based on a review of the relevant literature and reviewed by subject experts. The draft questionnaires were pre-tested and then pilot tested with 50 participants and adjusted for accuracy and clarity. The consistency and stability of the final questionnaires were tested using Cronbach's alpha (0.7). The online survey was administered via Google Forms and was distributed to the parents via Facebook, WhatsApp, and mail groups. Informed consent was obtained from all parents (18 years or older and currently living in India). Detailed responses were recorded further only those who provided informed consent to complete the survey.

Sociodemographic Measures

The survey has detailed information about the sociodemographic characteristics of study participants, including parental age, sex, education level, place of residence, income, occupation, social status in the community.

Parents knowledge of COVID-19 and vaccine intention

The survey tested parents' knowledge of the COVID-19 virus by asking, "Before this interview, were you aware that the COVID-19 virus is currently circulating in the community?". The responses were captured in a 3-point Likert scale (Yes, No, and Don't know). We also asked parents to record their response to vaccine development knowledge by asking, "To the best of your knowledge, is there currently a vaccine being prepared for the pandemic Coronavirus strain

referred to as COVID-19 vaccine?". Parents' own likelihood of getting a COVID-19 vaccine (same 3-point Likert scale: Yes, No, and Not sure) was also recorded.

Intent to vaccinate the child

The survey asked, "Do you intend to vaccinate your child(ren) for COVID-19 once a vaccine is available for children?". The responses were captured in a 3-point Likert scale (Yes, No, and Not sure). We coded "Yes" as "likely to get a COVID-19 vaccine"; all others were labeled as "hesitant."

Statistical Analyses

The key outcome measures of the study were to know the parent's intention to vaccinate their children against the COVID-19 virus. Those who responded "Yes" were labeled as "Intended to vaccinate" vs. "hesitancy." Descriptive statistical analysis was performed by doing cross-tabulation of demographic characteristics with the primary response variable "Intended to vaccinate." We performed both simple and multivariable logistic regression analyses to compute the odds ratio (OR) and a 95% confidence interval (CI). Chi-squared tests were performed for bivariate analysis (cross-tabulation) between the outcome variable and all explanatory variables. A multivariate logistic regression model was constructed by including factors with P-values <.05 at binary comparisons. Inference on significant association was considered with a two-tailed $p \leq 0.05$. STATA 15.0 software (StataCorp LP, Texas, USA) was used for all statistical analyses.

Ethical Considerations

Ethical approval was granted for the study by the institutional Research Ethics Committee of Post Graduate Institute of Medical Education and Research(PGIMER), Chandigarh, India.

Informed assent was taken before participation in the study. Anonymized data was used for analysis, interpretation, and reporting.

Results

Seven hundred and seventy parents participated in this study. Over half of the parents were 30-49 years old (60.0%), 39.6% were female, 28.5% were educated to graduate level or higher, 23.4% employed in government sector, 59.7% had more than six members in their family, 31.8% had a monthly income above 50,000 Indian rupees, 41.3% reside in rural areas, and 48.2% perceived their social status as high in the community (Table 1).

 The rate of parents' willingness to allow their children to receive the COVID-19 vaccine was 33.5%, whereas 40.3% of parents were willing to receive the COVID-19 vaccine. At the time of survey, 27.5% of parents had a history of exposures to a confirmed COVID-19 cases and 62.7% had concerned of getting infected to COVID-19 virus. Most (88.3%) of the parents were aware that COVID-19 virus is currently circulating in the community, and 86.0% of them had knew that a vaccine is being prepared for the pandemic Coronavirus strain referred to as "COVID-19 vaccine". Of the 770 parents, about one-fourth of them had a previous history of vaccine hesitancy, 62.2% had trusted the healthcare system , and 39.1% considered domestic vaccines are good (Table 2).

Table 3 shows the analysis of bivariate and multivariable logistic models, and determined the predictors affecting parents' willingness to allow their children to be given the COVID-19 vaccine. In bivariate model, trust in the healthcare system, higher risk perception, domestic vaccine, female respondent, employed in government sector, and being a rural residents were found to be associated with parent's intention to vaccinate their children. After adjusting for confounding variables, parent's own willingness to receive the COVID-19 vaccine (aOR: 2.35,

95% CI: 1.30-4.67), and parent's who had a bachelor's degree or higher education (aOR: 1.98, 95% CI: 1.15-3.51) were found to be associated with parents' willingness to receive the COVID-19 vaccine for their children (Table 3).

Figure 1, shows the concerns of parents' in accepting the novel COVID-19 vaccine. The key reasons for reluctance to allow their children to receive the COVID-19 vaccine included safety and effectiveness (86.4%), side effects (78.2%), doses not known (63.7%), lack of sufficient scientific data (53.2%), childrens are not affected from COVID-19 (52.8%), and trust in vaccine (43.4%).

Discussions

In this study, we aimed to determine the acceptability of the COVID-19 vaccine among parents in India and the associated factors. To the best of our knowledge, this is the first study in India that investigates parents' willingness to administer COVID-19 vaccine to their children. Understanding the factors associated with the intention will help in implementation of vaccination program in India and elsewhere. We found that nearly 33.5% of parents in India would be willing to administer a COVID-19 vaccine to their child. Sociodemographic factors, such as higher education level and parents who intended to vaccinate themselves, were found to be associated with childhood vaccination. The main concerns raised by parents in our study were around the safety and effectiveness of the novel COVID-19 vaccine.

Our findings contribute to an emerging literature suggesting that there is a low demand for childhood COVID-19 vaccine among parents(7–9,11,16). A study conducted in Texas, USA reported that about 38.3% of mothers had no intention to consider COVID-19 vaccine for their children(16). Proportion of willing parents in India is considerably lower as compared to other

countries as per studies from England (89%)(7); New Zealand (80%)(17); China (73%)(18); USA (65%)(19); Canada (63%)(10); and Turkey (42%)(20). A large proportion of participants in our study were aware about COVID 19 virus(88.3%) ; developmental status of COVID 19 vaccine (86%) and were exposed to COVID 19 (72.5%). Inspite of that, less than half of partcipants were willing to get themselves readily vaccinated (40.3%). This supports the findings that parent's intention to themselves get vaccinated is the most influential factor determining their intent towards children vaccination. Factors such as newness of vaccine , rapid development and unknown long term side effects influence parent's perception of vaccine safety and their intention to get themselves and their children vaccinated(13). As reported by numerous studies, vaccine safety is fundamental to maintaining the public's trust in vaccines(12,21,22). Thus future studies are required to understand long term change in perception of parents and factors influencing their intention to get themselves and their children vaccinated. Health care providers play a key role in influencing decision of parents towards vaccine acceptance and uptake. It is essential to thus engage various level of health care providers and strategize vaccine messages to parents especially from lower education status

Limitations of this study that should be considered include the sampling strategy adopted by the study which may not be representative of all parents in the country, hence limits the generalizability of our finding. The cross-sectional design of the study should be interpretated carefully when accessing overall prevalence of vaccine hesitancy among parents in India. Our sample represented individuals with internet access and having mobile literacy. This does not necessarily reflect the country wide perception of parents towards vaccine and their intention to get children vaccinated.

Despite the above limitations, our study assessed nationally representative sample of parents regarding their perceptions and intentions to vaccinate their children at a critical time, just as adult COVID-19 vaccination programs was initiated in India. Our findings reminiscent of recent research suggesting that there is a low demand for COVID-19 vaccine among parents, and highlights the need for longitudinal studies to measure the acceptability of a COVID-19 vaccine at different intervals. Future studies are thus required to supplement our current findings to enhance vaccine uptake among children in India. Nevertheless, our study establishes evidence regarding parent's hesitancy towards getting their children vaccinated and their concerns in coming months must be addressed through targeted communication stratgeies.

Conclusion

This study depicts that Indian parents have good knowledge of COVID-19 and vaccine developments. However, the demand for COVID-19 vaccine among parents was found to be low, with many parents unsure about their acceptance of COVID-19. The study warents implementation of strategies to improve parents' knowledge about the safety, efficacy and side effects of COVID-19 vaccine.

Abbreviations

COVID-19: Coronavirus disease-2019; WHO: World Health organization

Acknowledgements

The authors would like to acknowledge the study participants for their time and contributions to the study.

Authors' contributions

BKP, MG and AKG conceptualized the study, designed the tools; BKP, LJ, JV, PS, VC, BP, SSK, RS, SP, SB, NR, VR, TK, KG, BM, LS, MG, AKG conducted the study at national level, and collected the data. All authors reviewed drafts, provided edits, and approved the final submission.

Funding

None.

Availability of data and materials

De-identified data is available upon request.

Declarations

Ethics approval and consent to participate

The study protocol was reviewed and approved by the Institutional Ethical Committee of PostGraduate Institute of Medical Education and Research (PGIMER), Chandigarh, India. Informed consent was obtained from all participants. Anonymized data was used for interpretation and reporting.

Consent for publication

Not applicable.

Competing interests

The authors declare they have no competing interests.

References

1.   WHO. WHO Coronavirus (COVID-19) Dashboard. (2021) Available at:

https://covid19.who.int/

2.   Hoang A, Chorath K, Moreira A, Evans M, Burmeister-Morton F, Burmeister F, Naqvi R,

Petershack M, Moreira A. COVID-19 in 7780 pediatric patients: A systematic review.

EClinicalMedicine (2020) **24**:100433. doi:10.1016/j.eclinm.2020.100433

3.   Lurie N, Saville M, Hatchett R, Halton J. Developing Covid-19 Vaccines at Pandemic

Speed. *N Engl J Med* (2020) **382**:1969–1973. doi:10.1056/NEJMp2005630

4.   WHO. WHO COVID-19 - Landscape of novel coronavirus candidate vaccine development

worldwide. *COVID-19 vaccine tracker Landsc* (2021) Available at:

https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-

vaccines

5.   MOHFW. Ministry of Health and Family Welfare, Government of India. COVID-19

dashboard. (2021) Available at: https://mohfw.gov.in/

6.   MOHFW. Ministry of Health and Family Welfare, Government of India. COVID-19

dashboard. (2021)

7.   Bell S, Clarke R, Mounier-Jack S, Walker JL, Paterson P. Parents' and guardians' views on

the acceptability of a future COVID-19 vaccine: A multi-methods study in England.

Vaccine (2020) **38**:7789–7798. doi:10.1016/j.vaccine.2020.10.027

8.   Du M, Tao L, Liu J. The Association Between Risk Perception and COVID-19 Vaccine

Hesitancy for Children Among Reproductive Women in China: An Online Survey. *Front*

Med (2021) **8**: doi:10.3389/fmed.2021.741298

9.   Aggarwal S, Madaan P, Sharma M. Vaccine Hesitancy Among Parents of Children With

Neurodevelopmental Disabilities: A Possible Threat to COVID-19 Vaccine Coverage. *J*

Child Neurol (2021)088307382110421. doi:10.1177/08830738211042133

10.   Humble RM, Sell H, Dubé E, MacDonald NE, Robinson J, Driedger SM, Sadarangani M,

Meyer SB, Wilson S, Benzies KM, et al. Canadian parents' perceptions of COVID-19

vaccination and intention to vaccinate their children: Results from a cross-sectional

national survey. *Vaccine* (2021) doi:10.1016/j.vaccine.2021.10.002

11.   Aldakhil H, Albedah N, Alturaiki N, Alajlan R, Abusalih H. Vaccine hesitancy towards

childhood immunizations as a predictor of mothers' intention to vaccinate their children

against COVID-19 in Saudi Arabia. *J Infect Public Health* (2021) **14**:1497–1504.

doi:10.1016/j.jiph.2021.08.028

12.   Goldman RD, Krupik D, Ali S, Mater A, Hall JE, Bone JN, Thompson GC, Yen K, Griffiths

MA, Klein A, et al. Caregiver Willingness to Vaccinate Their Children against COVID-19

after Adult Vaccine Approval. *Int J Environ Res Public Health* (2021) **18**:10224.

doi:10.3390/ijerph181910224

13.   Yılmaz M, Sahin MK. Parents' willingness and attitudes concerning the COVID-19 vaccine:

A cross-sectional study. *Int J Clin Pract* (2021) **75**: doi:10.1111/ijcp.14364

14.   Catma S, Reindl D. Parents' willingness to pay for a COVID-19 vaccine for themselves and

their children in the United States. *Hum Vaccin Immunother* (2021) Available at:

https://www.tandfonline.com/doi/abs/10.1080/21645515.2021.1919453?casa_token=q

8XSKKpOy-YAAAAA:KMDHO8I8ygwOa4NEWYDkkSIXz8FgKfGsxGflf4bE9tiDBVRUKW3g9-

Kvv2Q3aEUdMJ6WLtjYmf8evQ

15. Szilagyi PG, Shah MD, Delgado JR, Thomas K, Vizueta N, Cui Y, Vangala S, Shetgiri R,

Kapteyn A. Parents' Intentions and Perceptions About COVID-19 Vaccination for Their

Children: Results From a National Survey. *Pediatrics* (2021) **148**:e2021052335.

doi:10.1542/peds.2021-052335

16. Zakeri M, Li J, Sadeghi SD, Essien EJ, Sansgiry SS. Strategies to decrease COVID-19 vaccine

hesitancy for children. *J Pharm Heal Serv Res* (2021) doi:10.1093/jphsr/rmab060

17. Jeffs E, Lucas N, Walls T. <scp>CoVID</scp> -19: Parent and caregiver concerns about

reopening New Zealand schools. *J Paediatr Child Health* (2021) **57**:403–408.

doi:10.1111/jpc.15234

18. Zhang KC, Fang Y, Cao H, Chen H, Hu T, Chen YQ, Zhou X, Wang Z. Parental Acceptability

of COVID-19 Vaccination for Children Under the Age of 18 Years: Cross-Sectional Online

Survey. *JMIR Pediatr Parent* (2020) **3**:e24827. doi:10.2196/24827

19. Goldman RD, Yan TD, Seiler M, Parra Cotanda C, Brown JC, Klein EJ, Hoeffe J, Gelernter R,

Hall JE, Davis AL, et al. Caregiver willingness to vaccinate their children against COVID-19:

Cross sectional survey. *Vaccine* (2020) **38**:7668–7673. doi:10.1016/j.vaccine.2020.09.084

20. Akarsu B, Canbay Özdemir D, Ayhan Baser D, Aksoy H, Fidancı İ, Cankurtaran M. While

studies on COVID-19 vaccine is ongoing, the public's thoughts and attitudes to the future

COVID-19 vaccine. *Int J Clin Pract* (2021) **75**: doi:10.1111/ijcp.13891

21. Opel DJ, Diekema DS, Ross LF. Should we mandate a COVID-19 vaccine for children?

JAMA Pediatr (2021) Available at:

https://jamanetwork.com/journals/jamapediatrics/article-abstract/2770123

22.   Kumar J, Meena J, Kasi SG, Shah A. COVID-19 Vaccine in Children: Where Do We Stand?

Indian Pediatr (2021) Available at: https://www.researchgate.net/profile/Jogender-

Kumar-3/publication/349119982_COVID-

19_Vaccine_in_Children_Where_Do_We_Stand/links/60217a2e299bf1cc26b23cca/COVI

D-19-Vaccine-in-Children-Where-Do-We-Stand.pdf

Table 1: Sociodemographic characteristics of the study respondents (N=770)

Variables	n (%)
Parental age	
18-29	120 (15.6)
30-49	462 (60.0)
above 50	188 (24.4)
Parental sex	
Male	465 (60.4)
Female	305 (39.6)
Parental education	
No Formal Education	26 (3.4)
Primary School	193 (25.1)
Diploma/High School	262 (34.0)
Undergraduate	130 (16.9)
Postgraduate and above	89 (11.6)
Parent's employment status	
Government sector	180 (23.4)
Private sector	282 (36.6)
Self-employed	230 (29.9)
Unemployed	78 (10.1)
Family size	
Five and below	310 (40.3)
Six and above	460 (59.7)
Family income (monthly, INR)	
below 10000	52 (6.8)
11000-20000	291 (37.8)
21000-50000	182 (23.6)
above 50000	245 (31.8)
Social status	
Low	108 (14.0)
Medium	291 (37.8)
High	371 (48.2)
Region of residence	
Eastern	192 (24.9)
Western	91 (11.8)
Northern	210 (27.3)
Southern	108 (14.0)
Central	86 (11.2)
North-east	83 (10.8)
Area of residence	
Urban	452 (58.7)
Rural	318 (41.3)
Social caste	
Other Backward Caste	312 (40.5)
Unreserved	215 (27.9)
Scheduled Caste	150 (19.5)
Scheduled Tribe	93 (12.1)
Religion	
Hindu	367 (47.7)
Muslim	173 (22.5)
Christian	85 (11.0)
Sikhs	105 (13.6)
Other	40 (5.2)

Table 2: Willingness to accept COVID-19 vaccine, contact history with COVID-19 patients, risk perception, and vaccine preferences among parents (N=770)

Variables	n(%)
Intended to vaccinate the child	
Yes	258 (33.5)
No	164 (21.3)
Not sure	348 (45.2)
Intended to vaccinate themselves	
Yes	310 (40.3)
No	270 (35.1)
Not sure	190 (24.7)
Exposed to COVID-19 cases	
No	212 (27.5)
Yes	558 (72.5)
Knowledge about the COVID-19 virus	
No	90 (11.7)
Yes	680 (88.3)
Knowledge about the development of the COVID19 vaccine	
No	108 (14.0)
Yes	662 (86.0)
History of vaccine hesitancy	
Yes	210 (27.3)
No	560 (72.7)
High-risk perception	
Yes	483 (62.7)
No	287 (37.3)
Trust in the healthcare system	
No	291 (37.8)
Yes	479 (62.2)
Trust in domestic vaccines	
Better	301 (39.1)
Neutral	282 (36.6)
Worse/Not Sure	187 (24.3)

Table 3: Unadjusted and adjusted odds ratios for the association between parent's intention to vaccinate their children with sociodemographic and other COVID-19 behaviors (N=770)

Variable	"Intended to vaccinate children"		
	OR [95% CI]	aOR [95% CI]	P-value
Parents' COVID-19 vaccination intention for themselves			
No	Ref	Ref	
Yes	2.88 [1.21-4.06]	2.35 [1.30-4.67]	0.001
Exposed to COVID-19 cases			
No	Ref	Ref	
Yes	1.71[0.81-2.61]	1.28 [0.46-2.25]	0.311
Trust in the healthcare system			
No	Ref	Ref	
Yes	2.42 [1.41-3.55]	2.11 [0.98-3.17]	0.068
History of vaccine hesitancy			
Yes	Ref	Ref	
No	1.48 [0.81-3.66]	1.45 [0.27- 2.74]	0.241
Higher risk perception			
No	Ref	Ref	
Yes	2.46 [1.19-3.80]	1.87 [0.99-3.17]	0.067
Opinion in domestic vaccines			
Worse	Ref	Ref	
Better	2.72 [1.02-5.08]	1.67 [0.71-3.97]	0.165
Neutral	2.30 [1.04-2.87]	1.40 [0.26-2.83]	0.151
Age			
18 – 29	Ref	Ref	
30-49	1.93 [0.85-2.23]	1.13 [0.48-2.77]	0.126
above 50	1.78 [0.91-2.14]	1.52 [0.35-2.17]	0.132
Sex			
Male	Ref	Ref	
Female	1.97 [1.07-2.20]	1.35 [0.97-2.44]	0.081
Employment status			
Unemployed/self-employed	Ref	Ref	
Private sector	1.51 [0.65-2.20]	1.18 [0.31-2.70]	0.142
Government sector	1.91 [1.07- 3.31]	1.30 [0.71-2.67]	0.095
Education			
Primary or below	Ref	Ref	
Diploma/High School	2.18 [1.19-5.82]	1.86 [0.95-4.61]	0.087
Undergraduate and higher	3.41[1.32-6.11]	1.98 [1.15-3.51]	0.001
Social status in the community			
Low	Ref	Ref	
Medium	1.80 [0.59-2.72]	1.57 [0.92-2.67]	0.085
High	1.83 [0.58-2.88]	1.16 [0.75-2.24]	0.116
Place of residence			
Urban	Ref	Ref	
Rural	1.92 [1.07-3.12]	1.25 [0.44-3.05]	0.134

Figure 1: Parents' concerns for not willing to vaccinate of their children for COVID-19 (n = 770)