

NIBS in Dystonia: A Meta-Analysis

Effects of non-invasive brain stimulation in dystonia: a systematic review and meta-analysis

Authors: Jordan Morrison-Ham¹, Gillian M. Clark¹, Elizabeth G. Ellis¹, Andris Cerins¹, Juho Joutsa^{2,3}, Peter G. Enticott¹, Daniel T. Corp^{1,4}.

1. Cognitive Neuroscience Unit, School of Psychology, Deakin University, Geelong, Australia
2. Turku Brain and Mind Center, Clinical Neurosciences, University of Turku, Turku, Finland
3. Turku PET Centre, Neurocenter, Turku University Hospital, Turku, Finland
4. Center for Brain Circuit Therapeutics, Brigham and Women's Hospital, Boston, MA, USA

Corresponding authors: Jordan Morrison-Ham (jmorriso@deakin.edu.au) & Daniel T. Corp (daniel.corp@deakin.edu.au)

Keywords. Transcranial magnetic stimulation. Transcranial direct current stimulation. Dystonia. Meta-Analysis.

Funding/Conflicts of interest: JMH, EGE & AC are funded by an Australian Government Research Training Program Scholarship. JJ is funded by the Instrumentarium Research Foundation, the Finnish Foundation for Alcohol Studies, University of Turku (private donation, Sigrid Juselius Foundation) and Turku University Hospital (ERVA funds), and has received a lecturer honorarium from Lundbeck. PGE is funded by a Future Fellowship from the Australian Research Council (FT160100077).

NIBS in Dystonia: A Meta-Analysis

Abstract

Background. Deep brain stimulation is a highly effective treatment of dystonia, but is invasive and associated with risks, such as intraoperative bleeding and infections. Previous research has used non-invasive brain stimulation (NIBS) in an attempt to alleviate symptoms of dystonia. The results of these studies, however, have been variable, leaving efficacy unclear. This study aimed to evaluate the effects of NIBS on symptoms of dystonia and determine whether methodological characteristics are associated with variability in effect size.

Methods. Embase and MEDLINE Complete databases were searched for articles using any type of NIBS as an intervention in dystonia patients, with changes in dystonia symptoms the primary outcome of interest.

Results. Meta-analysis of 26 studies demonstrated a small effect size for NIBS in reducing symptoms of dystonia (random-effects Hedges' $g = 0.21$, $p = .002$). Differences in the type of NIBS, type of dystonia, and brain region stimulated had a significant effect on dystonia symptoms. Meta-regression revealed that 10 sessions of active stimulation, and the application of concurrent motor training programs resulted in significantly larger mean effect sizes.

Conclusion. NIBS has yielded small improvements to dystonic symptoms, but effect sizes depended on methodological characteristics, with more sessions of stimulation producing a larger response. Future research should further investigate the application of NIBS parallel to motor training, in addition to providing a greater quantity of sessions, to help define optimal parameters for NIBS protocols in dystonia.

Registration. PROSPERO 2020, CRD42020175944.

Acronyms. ACC = anterior cingulate cortex; CD = cervical dystonia; dPM = dorsal premotor cortex; FHD = focal hand dystonia; M1 = primary motor cortex; NIBS = non-invasive brain stimulation; rTMS = repetitive transcranial magnetic stimulation; S1 = primary somatosensory cortex; TBS = theta-burst stimulation; tDCS = transcranial direct current stimulation; TMS = transcranial magnetic stimulation.

NIBS in Dystonia: A Meta-Analysis

Introduction

Dystonia is a chronic neurological disorder characterized by involuntary muscle contractions and postures¹. Dystonia can involve any body region, and is one of the most common movement disorders, with prevalence estimates of approximately 16.43 cases per 100,000 people². Dystonia can be idiopathic or secondary to other brain pathologies, such as focal brain lesions.

Invasive neuromodulation is highly effective in the treatment of dystonias³. Deep brain stimulation (DBS) to the globus pallidus interna (GPi) is the most widely used neuromodulation treatment for dystonia, and the subthalamic nucleus has also shown success^{3,4}. The mechanism of action for DBS in dystonia is not yet fully understood, but it is considered to modulate the function of the sensorimotor network, regions of which are often functionally abnormal in dystonia patients^{5,6}. Nevertheless, DBS is invasive and only considered in more severe cases that do not respond to botulinum toxin injections and oral pharmacotherapy⁷.

As a result, non-invasive brain stimulation (NIBS) has been suggested as a potential therapeutic treatment for dystonia symptoms, due to its ability to non-invasively modulate the functioning of abnormal neural networks^{8,9}. NIBS techniques, such as transcranial magnetic stimulation (TMS) and transcranial electrical stimulation (tES), have been effective in the treatment of other neurological and neuropsychiatric disorders, such as depression¹⁰, migraine¹¹, and obsessive compulsive disorder¹². Applied cortically, repetitive TMS (rTMS) and tES induce a plasticity-like response and can upregulate or downregulate neuronal activity at local and regional levels¹³.

Given that DBS in dystonia affects a large brain network, it is likely that there are multiple nodes that could be modulated via NIBS for therapeutic benefit. Several studies have demonstrated loss of inhibition, increased excitability, or abnormal plasticity in dystonia, in cortical regions associated with sensorimotor function including the somatosensory cortex (S1), primary motor cortex (M1), dorsal premotor cortex (dPM), and cerebellum¹⁴⁻¹⁹. However, NIBS to these cortical areas has returned variable results. Whilst previous research has suggested that rTMS and transcranial direct current stimulation (tDCS) provide some relief from symptoms of dystonia²⁰, other studies suggest little to no effect on dystonia symptoms in comparison to sham stimulation^{21,22}. Given these conflicting results, it is not yet known whether NIBS is effective in dystonia, nor whether specific NIBS methods or brain regions may enhance therapeutic effects.

NIBS in Dystonia: A Meta-Analysis

Therefore, the aim of this systematic review and meta-analysis is to pool all studies that have used NIBS in dystonia to comprehensively evaluate the effect of NIBS methods on dystonia symptoms, and to better understand which protocols may be most effective.

Methods

Study Selection

Systematic Search

A search of Embase and MEDLINE Complete was conducted during March and April 2020, using a combination of synonyms of the following terms: dystonia; transcranial magnetic stimulation (TMS); theta-burst stimulation (TBS); transcranial alternating current stimulations (tACS); transcranial direct current stimulation (tDCS); transcranial electrical stimulation (tES); transcranial random noise stimulation (tRNS); and non-invasive brain stimulation (NIBS). Exact search syntax is provided in supplementary file 1. No publication status or year limiters were applied, however only studies reported in English were considered. The reference lists of all included articles were searched for studies missed in the initial search.

Inclusion & Exclusion Criteria

Studies were screened using inclusion and exclusion criteria based on the PICO (participants, intervention, control, outcome) framework²³. Studies were first selected for *qualitative review* based on the following criteria: (P) participants who had a clinical diagnosis of dystonia (any type), with a study sample size of 1 or more; (I) non-invasive brain stimulation (any type) used as an intervention intended to reduce dystonia symptom severity; (C) no comparison group or randomization necessary for qualitative review; and (O) an outcome measure that assessed changes in clinical symptoms of dystonia (e.g., Burke-Fahn-Marsden Dystonia Rating Scale).

Studies were selected for *quantitative review* (i.e., meta-analysis) based on the following criteria: (P) participants who had a clinical diagnosis of dystonia (any type), with a study sample size of at least 3; (I) as above; (C) a comparison group of dystonia controls who received sham stimulation (parallel trials), or a design where dystonia patients received both sham and real stimulation (crossover trials); (O) as above. Studies that examined dystonia participants who were actively receiving DBS were excluded, as DBS can influence the response to NIBS, even where the DBS stimulator is switched off^{24, 25}.

NIBS in Dystonia: A Meta-Analysis

Screening & Data Extraction

Literature search results were exported to EndNote (version X9) and Rayyan²⁶. Two reviewers independently screened titles and abstracts obtained from the literature search against the inclusion and exclusion criteria. Full text articles were then assessed against inclusion criteria, with disagreements resolved through discussion, and where necessary by a third member of the study team (D.C.).

Following the screening and inclusion of full text articles, data were extracted from individual studies into custom Microsoft Excel spreadsheets, including participant demographics, clinical information, trial characteristics, NIBS protocols, and symptom scores. The primary outcome was changes in dystonia symptoms, post-intervention. In this review, we analysed dystonia symptoms measured by clinically validated rating scales (e.g., the Toronto Western Spasmodic Torticollis Rating Scale [TWSTRS]); subjective patient symptom scales created specifically for the empirical study; and changes in motor performance in the affected limb post-intervention. The potential influence of outcome measures on effect sizes was later analysed using meta-regression.

Effect Size Calculations

Due to the small sample sizes of the included articles, a Hedges' *g* effect size was calculated to correct for potential overestimation of the population standardized mean difference (SMD)²⁷. For all studies, Hedges' *g* was calculated so that positive values indicated NIBS improved dystonia symptoms, and negative values indicated NIBS worsened dystonia symptoms. Hedges' *g* was calculated to compare the change in dystonia symptoms from baseline (either clinical or task-based) between the NIBS and sham conditions. This effect size was calculated from pre- and post-stimulation mean scores (or change from baseline scores) and standard deviations (SDs) for both NIBS and sham groups, using Comprehensive Meta-Analysis (CMA; version 3.3.070) software. In studies where the means and SDs were reported in graphs or images, Plot Digitizer (version 2.6.8; <http://plotdigitizer.sourceforge.net/>) software was used to extract values. If standard errors (SE) or confidence intervals (CI) were reported for mean scores, they were converted to SDs using the equations:

$$SD = SE * \sqrt{N} \text{ (for SEs) and (for CIs): } SD = \sqrt{N} * \frac{\text{upper limit} - \text{lower limit}}{3.92}$$

NIBS in Dystonia: A Meta-Analysis

Where N is the total number of participants²⁸. All formulas for effect size calculations are provided in Supplementary 2.

Pooling of Effect Sizes

For studies that used more than one outcome measure to assess symptoms of dystonia (e.g., a task-based measure along with a clinically validated rating scale)²⁹⁻³⁴, effect sizes and variances for each outcome were averaged within studies, to obtain one overall effect size for each study. All effect sizes were then pooled using a random effects model in CMA software. Both study level and the overall pooled effect size were considered significant if $p < .05$.

Meta-Analysis

All meta-analysis forest plots and sensitivity analyses were conducted in Stata/SE (version 15.1). A leave-one-out sensitivity analysis was performed to detect the presence of any outliers, using the ‘metainf’ command³⁵. In order to obtain an effect size estimate for each level within categorical variables, individual meta-analyses were run separating studies by NIBS type (e.g., tDCS, rTMS), brain region stimulated, type of dystonia, and outcome measures: clinically validated rating scales, unvalidated rating scales (i.e., rating scales devised for the study), and task-based outcomes (e.g., timed handwriting tests). Separate meta-analyses were conducted for each of the aforementioned variables (rather than comparing levels of the variable with a technique such as meta-regression) as there were a high number of levels per variable (e.g., high and low frequency rTMS, intermittent and continuous TBS, and tDCS for the variable NIBS type) and few study effect sizes per level, therefore insufficient statistical power to utilise a number of these variables within a meta-regression³⁶.

Between-study heterogeneity in effect sizes was quantified using the I^2 statistic³⁷. As per Higgins³⁷ the effect of heterogeneity was considered low, moderate, or high for I^2 values of 25%, 50%, and 75%, respectively.

Meta-Regression

Meta-regression analyses were conducted in Stata/SE (version 15.1) to determine the influence of mean age, gender ratio, number of active sessions of stimulation, etiology of dystonia, and concurrent motor training on NIBS outcomes. The ‘metareg’³⁸ function was used for continuous variables (mean age and gender ratio), and the ‘maanova’³⁹ function on the categorical variables (number of active sessions of stimulation, dystonia etiology and concurrent

NIBS in Dystonia: A Meta-Analysis

motor training). Prior to conducting the regression analysis, data were checked visually for normality and collinearity using histograms and scatterplots. Levels of independent variables were omitted from the regression analysis if they did not comprise at least three studies, ensuring that there were enough data for each level to provide a reliable regression estimate³⁶.

Evaluation of Bias

The methodological quality of each study was assessed using the Cochrane Collaboration's Risk of Bias (RoB) checklists⁴⁰. For parallel trials, the revised Cochrane Risk of Bias tool for randomized trials (RoB 2)⁴⁰ was used, while a modified version of the RoB 2 for repeated measures designs was utilised for crossover trials. The RoB 2 checklist assesses studies on the domain's randomization, blinding of participants and personnel, outcome measurement and assessor blinding, incomplete outcome data, and selective outcome reporting. Each domain was judged to be of low, unclear, or high risk of bias, with an overall judgement given for each study. For crossover trials, bias arising from period or carryover effects was also assessed.

The presence of publication bias across studies was assessed using funnel plots where effect sizes for each study were plotted against their standard error⁴¹. In the absence of publication bias, symmetrical distribution of effect sizes around the overall effect size is observed. The symmetry of the funnel plot was assessed both visually, and statistically using Egger's test⁴¹.

Results

Study Selection

In total, 1124 records were identified across the two databases. After duplicate removal, and title and abstract screening, 121 full-text articles were assessed for eligibility. 46 studies were included for qualitative synthesis, with 26 studies (11 parallel and 15 crossover trials) meeting inclusion criteria for the meta-analysis (Figure 1).

NIBS in Dystonia: A Meta-Analysis




Figure 1. PRISMA flowchart of search method and screening process.

Study Characteristics

A total 513 participants were included across 46 studies, with ages ranging from 7-70 years ($M = 45.93$, $SD = 11.93$). Three studies included patients with acquired dystonia, associated with Wilson's disease⁴³ or cerebral palsy^{44, 45}. Thirteen studies included participants who were not on oral medications (e.g., benzodiazepines), and 18 studies examined participants who had their last botulinum toxin injection more than 4 weeks prior to stimulation. Low frequency rTMS (1Hz or 0.2Hz; 21 studies) was the most utilised form of NIBS, followed by tDCS (anodal or cathodal; 17 studies). A single study applied tACS⁵⁰. Dystonia-specific motor training or biofeedback was

NIBS in Dystonia: A Meta-Analysis

employed concurrently with NIBS in seven studies^{29, 30, 32, 33, 46-48}. All study designs, participant demographics and characteristics are provided in Table 1.

NIBS in Dystonia: A Meta-Analysis

Table 1. Included articles' participant demographics and characteristics, study designs and outcomes.

Study	Design	Participants	Dystonia Cause	Intervention	Stimulation Target	Outcome Measure	Symptom change after NIBS
<i>Allam et al. (2007)</i> ⁴⁹	Case study	1 CD/WC patient (37 years)	Idiopathic	5 sessions rTMS (1Hz, 1200 stimuli @ 90% RMT – 20 mins)	Left dPM	BFMDRS	CD ↓ for 4 months WC no change
<i>Angelakis et al. (2013)</i> ⁵⁰	Sham-controlled case study	1 CD patient (54 years)	Idiopathic	a. 5 real, 5 sham sessions tDCS (1.5mA – 15 mins)	a. Cathode C4, anode P3	TWSTRS, patient rating scale	↓
				b. 7 sessions tACS (15Hz, 5Hz, 15Hz – 6/3/6 mins)	b. SMC		TWSTRS ↓ 29.5pts
<i>Benninger et al. (2011)</i> ²¹	RCT	12 WC patients ($M = 57.1$)	Idiopathic	3 sessions real or sham cathodal tDCS (2mA – 20 mins)	Contralateral M1	ADDS, WCRS, handwriting kinematics	↑
<i>Betti et al. (2018)</i> ⁵¹	Case study	1 MD patient (55 years) and 1 HC (50 years)	Idiopathic	5 sessions rTMS (1Hz, 1800 stimuli @ 90% RMT – 30 mins)	Left M1	Motor tasks	↓
<i>Bhanpuri et al. (2015)</i> ⁴⁴	Crossover RCT	a. 7 HD patients ($M = 15.7$)	a. 1 DYT1, 6 acquired (4 cerebral palsy, 1 vitamin E deficient, 1 TBI)	a. 5 real, 5 sham sessions cathodal tDCS (2mA – 9 mins)	a. Contralateral M1	BADs, EMG tracking task	↓
		b. 6 HD patients ($M = 14.8$)	b. 6 acquired (1 TBI, 5 cerebral palsy)	b. 5 real, 5 sham sessions anodal tDCS (2mA – 9 mins)	b. Contralateral M1		↑
<i>Bologna et al. (2016)</i> ⁵²	Crossover RCT	13 FHD ($M = 48.5$), 13 CD ($M = 46.7$) and 13 HC ($M = 49.9$)	Idiopathic	1 real, 1 sham session cTBS (600 stimuli – 40 secs)	Cerebellum (ipsilateral to affected hand in FHD)	TWSTRS, WCRS, cortical excitability	No change
<i>Borich et al. (2009)</i> ⁵³	Crossover RCT	6 FHD patients ($M = 46.5$) and 9 HC ($M = 33$)	Idiopathic	5 sessions rTMS (1Hz, 900 stimuli @ 90% RMT), 3 patients received sham before crossing over to the intervention	Contralateral dPM	Handwriting kinematics, cortical excitability	Handwriting (pen velocity) ↓*

NIBS in Dystonia: A Meta-Analysis

Study	Design	Participants	Dystonia Cause	Intervention	Stimulation Target	Outcome Measure	Symptom change after NIBS
<i>Bradnam et al. (2014)</i> ⁵⁴	Case study	1 CD patient (47 years)	Idiopathic	20 sessions anodal tDCS (2mA – 2 x 15 min blocks, separated by 5 mins)	Right/left cerebellum (5 sessions each), right M1/left cerebellum (10 sessions)	TWSTRS, CDQ-24, CDIP-58	TWSTRS, CDQ-24, CDIP-58 ↓ 18+ pts
Bradnam et al. (2015) ⁵⁵	Crossover RCT	8 FHD patients ($M = 59$) and 8 HC ($M = 61.3$)	Idiopathic	1 session each anodal, cathodal, and sham tDCS (2mA – 20mins)	Cerebellum	Handwriting kinematics, cortical excitability	Anodal tDCS handwriting ↓*
Bradnam et al. (2016) ⁸	RCT	16 CD patients ($M = 51.9$)	Idiopathic	10 sessions real or sham iTBS (600 stimuli each hemisphere – 8 mins)	Bilateral cerebellum	TWSTRS, CDQ-24, hand dexterity, cortical excitability	TWSTRS, CDQ-24, hand dexterity ↓*
Buttkus et al. (2010a) ⁵⁶	Crossover RCT	10 MD patients ($M = 48.8$)	Idiopathic	1 real, 1 sham session cathodal tDCS (2mA – 20 mins)	Left M1	ADDS, FAM, assessment of fine motor control	No change
<i>Buttkus et al. (2010b)</i> ⁴⁷	Case study	1 MD patient (43 years)	Idiopathic	5 cathodal, 5 anodal, 5 sham sessions tDCS + training (2mA – 20 mins)	Left M1	Assessment of fine motor control	Anodal, cathodal tDCS ↓*
Buttkus et al. (2011) ²⁹	Crossover RCT	9 MD patients ($M = 44$)	Idiopathic	1 cathodal, 1 anodal, 1 sham session tDCS + training (2mA – 20mins)	Left M1	Assessment of fine motor control	No change
<i>Conte et al. (2014)</i> ⁵⁷	Crossover RCT	12 WC patients ($M = 43$) and 12 HC ($M = 42$)	Idiopathic	1 iTBS, 1 cTBS, 1 sham session (iTBS = 600 stimuli @ 80% AMT, cTBS = 600 stimuli @ 80% AMT – 40 secs)	Contralateral S1	Writing task, STDT	No change
Furuya et al. (2014) ³⁰	Crossover RCT	10 MD patients ($M = 39.6$) and 10 HC ($M = 27.9$)	Idiopathic	1 cathodal, 1 anodal, 1 anodal over unaffected hemisphere, 1 sham session (all + training), and 1 cathodal session (no training), 5 sessions total (2mA – 24 mins)	Contralateral M1, except for 1 anodal session over ipsilateral M1	Assessment of fine motor control	↓ for cathodal (+training) only*
Havrankova et al. (2010) ⁵⁸	Crossover RCT	11 WC patients ($M = 40.3$)	Idiopathic	5 real, 5 sham sessions rTMS (1Hz, 1800 stimuli @ 90% AMT – 30 mins)	Contralateral S1	BFMDRS, handwriting, 2-minute writing test	Handwriting ↓*

NIBS in Dystonia: A Meta-Analysis

Study	Design	Participants	Dystonia Cause	Intervention	Stimulation Target	Outcome Measure	Symptom change after NIBS
Huang <i>et al.</i> (2010) ⁵⁹	RCT	7 patients ($M = 44.9$) and 9 HC ($M = 42.7$)	Idiopathic	1 session cTBS (600 stimuli @ 80% AMT – 40 secs)	Contralateral dPM	Handwriting, cortical excitability	Handwriting ↓*
Huang <i>et al.</i> (2012) ⁶⁰	RCT	18 WC patients ($M = 41.4$)	Idiopathic	5 sessions real or sham cTBS (600 stimuli @ 80% AMT – 40 secs)	Contralateral dPM	Handwriting, cortical excitability	No change
Kimberley <i>et al.</i> (2013) ⁶¹	RCT	17 FHD patients ($M = 46.5$)	Idiopathic	5 sessions real or sham rTMS (1Hz, 1800 stimuli @ 90% RMT – 30 mins)	Contralateral dPM	Handwriting kinematics, cortical excitability	Handwriting (pen pressure) ↓*
Kimberley <i>et al.</i> (2015a) ⁴⁸	Crossover RCT	9 FHD patients ($M = 46$)	Idiopathic	5 sessions rTMS + training, 5 sessions rTMS only (1Hz, 1200 stimuli @ 80% RMT – 20 mins)	Contralateral dPM	ADDS, GROC, WCERS, handwriting kinematics, cortical excitability	ADDS ↓ for both conditions*
Kimberley <i>et al.</i> (2015b) ⁶²	Case study	2 FHD patients ($M = 43.5$)	Idiopathic	6 sessions rTMS (1Hz, 1200 stimuli @ 90% RMT – 20 mins)	Right dPM	Handwriting kinematics, cortical excitability, patient rating scale	Handwriting ↓*
Koch <i>et al.</i> (2014) ⁹	RCT	20 CD patients ($M = 54$) and 10 HC ($M = 51.2$)	Idiopathic	10 sessions real or sham cTBS (600 stimuli per hemisphere @ 80% AMT – 4 mins)	Bilateral cerebellum	TWSTRS, BFMDRS, cortical excitability	TWSTRS ↓*
Kranz <i>et al.</i> (2009) ⁶³	Randomised crossover study	7 BEB patients ($M = 62.6$)	Idiopathic	a. 4 sessions rTMS (0.2Hz, 180 stimuli @ 90% RMT – 15 mins)	a. M1, dPM, SMA, ACC	Physician/patient rating scales, BRR	Rating scales ↓ for all targets*
				b. 4 sessions cTBS (600 stimuli @ 80% AMT – 40 secs)	b. M1, dPM, SMA, ACC		No change
				c. 2 sessions tDCS (1mA – 20 mins)	c. M1/dPM, SMA/ACC		No change
Kranz <i>et al.</i> (2010) ⁶⁴	Crossover RCT	12 BEB patients ($M = 61.4$)	Idiopathic	2 rTMS (H-coil and C-coil), 1 sham session (0.2Hz, 180 stimuli @ 100% AMT – 15 mins)	ACC	Physician/patient rating scales, BRR	Rating scales, BRR ↓ for both coils*

NIBS in Dystonia: A Meta-Analysis

Study	Design	Participants	Dystonia Cause	Intervention	Stimulation Target	Outcome Measure	Symptom change after NIBS
<i>Lefaucher et al. (2004)</i> ⁶⁵	Case series	3 CD/lower limb dystonia patients ($M = 38.3$)	Acquired (1 viral encephalitis, 1 neonatal TBI, 1 disulfiram intoxication)	5 sessions rTMS (1Hz, 1200 stimuli @ 90% RMT – 20 mins)	Left dPM	BFMDRS, spasm number/intensity	Spasm number/intensity ↓
<i>Linssen et al. (2015)</i> ⁶⁶	Crossover RCT	10 WC patients and 10 HC [†]	Not provided	1 real, 1 sham cTBS session (80% AMT)	Ipsilateral cerebellum	Writing tasks	No change
<i>Lozeron et al. (2017)</i> ⁴³	Crossover RCT	13 FHD patients ($M = 42.3$)	Inherited (Wilson's Disease)	1 real, 1 sham session rTMS (1Hz, 1200 stimuli @ 80% RMT – 20 mins)	Contralateral S1	WCRS, FAR, UWDRS	No change
<i>Marceglia et al. (2017)</i> ⁶⁷	Sham-controlled crossover study	2 MD patients ($M = 41$)	Idiopathic	5 cathodal, 5 anodal, 5 sham sessions tDCS (2mA – 20mins)	Bilateral M1/dPM	SSS, FSS, TC, writing tasks	Cathodal tDCS ↓
<i>Murase et al. (2005)</i> ⁶⁸	Crossover RCT	9 WC patients ($M = 38$) and 7 HC [†] ($M = 36$)	Idiopathic	3 real, 1 sham session rTMS (0.2Hz, 180 stimuli @ 85% RMT/AMT for SMA)	Contralateral M1, dPM, SMA (1 session each)	Handwriting, cortical excitability	↓ for dPM target*
<i>Naro et al. (2019)</i> ⁶⁹	Case study	1 WC patient (25 years)	Idiopathic	75 sessions rTMS (1Hz, 1200 stimuli @ 90% AMT)	Left dPM	WCRS, 1 minute writing test	↓ for 12 months
<i>Odorfer et al. (2019)</i> ⁷⁰	RCT	8 CD patients and 8 HC	Idiopathic	2 real, 1 sham session cTBS (600 stimuli @ 80% AMT – 40 secs)	Left dPM, bilateral cerebellum (1 session each)	TWSTRS, cortical excitability	No change
<i>Okada et al. (2018)</i> ⁴⁹	Case study	1 WC patient (47 years)	Idiopathic	6 sessions tDCS (2mA – 30 mins) +EMG biofeedback	Bilateral M1	WCRS	↓
<i>Pirio Richardson et al. (2015)</i> ³¹	Crossover RCT	9 CD patients ($M = 53$)	Idiopathic	4 real, 1 sham session rTMS (0.2Hz, 180 stimuli @ 85% RMT – 15 mins)	Left ACC, dPM, M1, SMA (1 session each)	TWSTRS, cortical excitability	↓ dPM, SMA, M1 targets
<i>Rossett-Llobet et al. (2014)</i> ³²	RCT	4 FHD patients ($M = 35.5$) and 30 FHD controls ($M = 34.4$)	Idiopathic	10 sessions cathodal tDCS +training/10 sessions training only, or 20 sessions cathodal tDCS +training (2mA – 20 mins)	Contralateral M1	Motor tasks, patient rating scale	20 sessions cathodal tDCS +training ↓*

NIBS in Dystonia: A Meta-Analysis

Study	Design	Participants	Dystonia Cause	Intervention	Stimulation Target	Outcome Measure	Symptom change after NIBS
Rossett-Llobet et al. (2015) ³³	RCT	26 MD patients ($M = 35$)	Idiopathic	10 session real or sham cathodal tDCS + training (2mA – 20 mins)	Left M1	Dystonia severity rating, therapist rating	↓*
Sadnicka et al. (2014) ²²	Crossover RCT	10 WC patients ($M = 52.8$)	Idiopathic	1 real, 1 sham session anodal tDCS (2mA – 15 mins)	Cerebellum	WCERS, cortical excitability	No change
<i>Salatino et al. (2019)</i> ⁷¹	Sham-controlled case study	1 FHD patient (41 years) and 1 HC [†] (32 years)	Idiopathic	a. 1 real, 1 sham rTMS sessions (1Hz, 900 stimuli @ 90% RMT)	Left dPM	Handwriting, therapist rating	Handwriting ↓
				b. 6 sessions rTMS (1Hz, 900 stimuli @ 90% RMT)			Handwriting ↓
<i>Sharma et al. (2019)</i> ⁷²	Case study	1 lower limb dystonia patient (65 years)	Idiopathic	10 sessions rTMS (1Hz, 2600 stimuli 90% AMT – 44 mins)	Right M1	PHQ-9, CGI	No change
<i>Shin et al. (2019)</i> ⁷³	Case study	1 lower limb dystonia patient (57 years)	Acquired (cerebellar lesion)	5 sessions rTMS (1Hz, 600 stimuli @ 90% RMT)	Right cerebellum	BFMDRS	↓
Siebner et al. (1999) ³⁴	Crossover RCT	16 WC patients ($M = 47$) and 11 HC ($M = 40$)	Idiopathic	1 session rTMS (1Hz, 1800 stimuli @ 90% RMT – 30 mins), 10 WC patients received sham before crossing over to the intervention	Left M1	Handwriting kinematics, cortical excitability	Handwriting pressure ↓*
Siebner et al. (2003) ⁷⁴	Crossover RCT	7 FHD patients ($M = 48$) and 7 HC ($M = 48$)	Idiopathic	1 real, 1 sham session rTMS (1Hz, 1800 stimuli @ 90% RMT – 2x15 min trains)	Left dPM	WCERS, physician rating scale, handwriting	No change
<i>Trebossen et al. (2017)</i> ⁷⁵	Case study	1 BEB patient (70 years)	Not provided	10 sessions tDCS (2mA – 30 mins)	Left dPFC	Eye blink rate, depressive symptom report	↓ for 10 days
<i>Veugen et al. (2013)</i> ⁷⁶	RCT	15 WC patients ($M = 58$) and 10 HC ($M = 58$)	Idiopathic	1 session cTBS (600 stimuli @ 70% RMT)	Contralateral dPM	Writing tasks, cortical excitability	Handwriting (spiral maze) ↓*

NIBS in Dystonia: A Meta-Analysis

Study	Design	Participants	Dystonia Cause	Intervention	Stimulation Target	Outcome Measure	Symptom change after NIBS
<i>Vucurovic et al. (2017)</i> ⁷⁷	Case study	1 left multifocal dystonia patient (41 years)	Acquired (Schizencephaly-related left hemiparesis with spastic dystonia)	3 sessions rTMS (1Hz, 1200 stimuli @ 100% RMT)	Left M1	UDRS	↓ for 1 month
Wagle-Shukla et al. (2018) ⁷⁸	RCT	12 BEB patients ($M = 69.1$)	Not provided	10 sessions real or sham rTMS (0.2Hz, 180 stimuli @ 100% AMT – 15 mins)	ACC	JRS, CDQ-24, BRR	JRS, CDQ-24 ↓ for 2 weeks*
Young et al. (2014) ⁴⁵	Crossover RCT	14 HD patients ($M = 12.6$)	2 idiopathic, 12 acquired (11 cerebral palsy, 1 TBI)	1 real, 1 sham session cathodal tDCS (1mA – 2x9 mins, with 20 min break in between)	Contralateral M1	BADS, EMG tracking task	EMG task (overflow) ↓*

Note. Where boxes are left blank, the information was not provided. Where authors are italicised, studies were included in qualitative analysis only. WC = writer's cramp; rTMS = repetitive transcranial magnetic stimulation; RMT = resting motor threshold; BFMDRS = Burke-Fahn-Marsden Dystonia Rating Scale; SMC = sensorimotor cortex; TWSTRS = Toronto Western Spasmodic Torticollis Rating Scale; RCT = randomised controlled trial; ADDS = Arm Dystonia Disability Scale; WCRS = Writer's Cramp Rating Scale; MD = Musician's dystonia; HC = healthy control; HD = hand dystonia; TBI = traumatic brain injury; BADS = Barry-Albright Dystonia Rating Scale; EMG = electromyogram; cTBS = continuous theta-burst stimulation; CDQ-24 = Craniocervical Dystonia Questionnaire; CDIP-58 = Cervical Dystonia Impact Profile; iTBS = intermittent theta-burst stimulation; FAM = frequency of abnormal movements scale; AMT = active motor threshold; STDT = sensory-temporal discrimination task; GROC = Global Rating of Change; BEB = Benign Essential Blepharospasm; SMA = supplementary motor area; BRR = blink reflex recovery; FAR = flow, accuracy and rhythmicity evaluation; UWDRS = Unified Wilson's Disease Rating Scale; SSS = Symptom Severity Scale; FSS = Functional Status Scale; TC = Tubiana and Champagne scale; PHQ-9 = Patient Health Questionnaire; CGI = Clinical Global Impression; dPFC = dorsolateral prefrontal cortex; UDRS = Unified Dystonia Rating Scale; JRS = Jankovic Rating Scale.

* denotes statistically significant results in comparison to baseline or sham condition ($p < .05$). † refers to participants used for electrophysiological data comparison only. ↓ refers to a decrease in symptoms after NIBS on outcome measures. ↑ refers to an increase in symptoms after NIBS on outcome measure.

NIBS in Dystonia: A Meta-Analysis

Qualitative Analysis

20 studies met criteria for qualitative analysis only, encompassing 73 participants with dystonia and 32 healthy control subjects (Table 1 – see italicized author studies). Overall, 19 of the 20 studies reported some reduction in dystonia symptoms after the application of NIBS, however many did not report whether this was statistically significant. Two studies^{72, 73} applied rTMS to patients with lower limb dystonia, and one study applied rTMS in a patient with left side multifocal dystonia, which affected the upper and lower limbs⁷⁷. The average number of stimulation sessions was 10 ($SD = 15.1$), with a maximum of 75 sessions⁶⁹.

Meta-Analysis


Meta-analysis was performed on 26 studies, totaling 352 participants with dystonia (hand dystonias, inclusive of task-specific FHD, musician's dystonia and writer's cramp, 19 studies; CD, 5 studies; blepharospasm, 2 studies). Included studies were either parallel ($n = 11$; where participants were randomly assigned to sham or intervention groups) or crossover ($n = 15$; where participants completed both sham and intervention conditions) group designs. One crossover group study⁷⁴ only provided post-stimulation data, and thus was treated as a parallel group design. Participant mean age was 44.6 years ($SD = 13.7$). The mean number of sessions of stimulation was 4.61 ($SD = 3.57$), inclusive of sham stimulation sessions in crossover trials. Of the 26 studies, 15 showed a reduction in dystonia symptoms after the application of NIBS.

Prior to conducting the meta-analysis, a leave-one-out sensitivity analysis was performed, demonstrating the presence of an outlier⁶⁴ (Supplementary 3). This study was therefore removed from all subsequent analyses. Nevertheless, meta-analysis conducted with this study⁶⁴ included was still significant (Supplementary 4).

Overall meta-analysis demonstrated a small effect size favoring active stimulation over sham stimulation for a reduction in dystonia symptoms, random-effects Hedges' $g = 0.21$, 95% CI [0.08, 0.35], $p = .002$ (Figure 2). Between-study heterogeneity was significant ($I^2 = 45.04\%$, $p = .012$), therefore meta-regressions were conducted to find moderators of the effect.

Meta-analyses were then run separating studies by selected variables (Table 2). These analyses demonstrate significance for rTMS overall ($p = .002$), 0.2Hz rTMS ($p < .001$), cathodal tDCS ($p = .04$), brain regions ACC ($p = .02$), M1 ($p = .03$) and dPM ($p = .001$), and blepharospasm ($p = .02$), task-specific FHD ($p = .002$), musician's dystonia ($p = .01$), and writer's cramp ($p = 0.007$). All forest plots are available in the supplementary materials.

NIBS in Dystonia: A Meta-Analysis



NOTE: Weights are from random effects analysis

NIBS in Dystonia: A Meta-Analysis

Figure 2. Forest plot of the random effects meta-analysis, demonstrating a small, significant effect for NIBS in decreasing dystonia symptoms. Where protocol states “Anodal+Cathodal”, participants received both anodal and cathodal tDCS. Separate effect sizes were calculated for each protocol and then combined into one overall study effect size.

NIBS in Dystonia: A Meta-Analysis

Table 2. Effect sizes for separate meta-analyses on categorical variables.

Variable	<i>n</i>	Hedges' <i>g</i>	95% CI	I ²
NIBS type				
rTMS	9	0.36*	[0.1, -0.61]	36.4%
1Hz	6	0.12	[-0.12, 0.35]	0%
0.2Hz	3	0.64*	[0.36, 0.95]	0%
TBS	7	0.04	[-0.14, 0.23]	0%
<i>iTBS</i>	1	0.74	[-0.23, 1.71]	-
<i>cTBS</i>	6	0.02	[-0.17, 0.24]	0%
tDCS	11	0.22	[-0.03, 0.47]	59.8%*
<i>Cathodal</i>	7	0.38*	[0.02, 0.74]	59.4%*
<i>Anodal</i>	1	-0.41	[-0.90, 0.08]	-
<i>Anodal+Cathodal</i>	3	0.17	[-0.13, 0.47]	31.5%*
Brain region				
ACC	1	1.37*	[0.19, 2.55]	-
CRB	7	0.02	[-0.16, 0.19]	0%
M1	11	0.29*	[0.04, 0.55]	57.3%*
dPM	6	0.46*	[0.19, 0.73]	0%
S1	2	0.06	[-0.25, 0.38]	32.8%
Dystonia type				
BEB	1	1.37*	[0.19, 2.55]	-
CD	5	0.22	[-0.03, 0.47]	25.2%
HD	6	-0.01	[-0.23, 0.21]	35.5%
Task-specific FHD	15	0.32*	[0.11, 0.52]	35.3%
<i>MD</i>	5	0.6*	[0.13, 1.07]	61.9%*
<i>WC</i>	7	0.28*	[0.08, 0.49]	0%
Outcome type				
Validated scale	13	0.17	[-0.05, 0.38]	54.8%*
Unvalidated scale	2	1.06	[-1.05, 3.17]	87.3%*
Task-based measure	20	0.13	[-0.01, 0.26]	29.5%
Overall	27	0.21*	[0.08, 0.35]	42.3%*

Note. I² statistics were not calculated for several variables due to there only being one study. Three studies that assessed task-specific FHD could not be separated further into musician's dystonia and writer's cramp, as the studies included both participants. CRB = cerebellum; BEB = blepharospasm; HD = hand dystonia; MD = musician's dystonia; WC = writer's cramp. *denotes significance at the $p < .05$ level

NIBS in Dystonia: A Meta-Analysis

Meta-Regression

Meta-regression conducted on the number of active sessions of stimulation demonstrated a significant difference between the three groups, $Q(2) = 10.97, p = .004$. Pairwise comparisons revealed 10 sessions of active stimulation resulted in significantly larger mean effect sizes for NIBS reducing dystonia symptoms (one session $g = 0.2, p = .01$, five sessions $g = 0.04, p = .77$, 10 sessions $g = 0.92, p < .001$; Figure 3). Two and three sessions of stimulation were removed as they did not meet the number of studies to be included in the analysis ($n \geq 3$).




Figure 3. Meta-regression and pairwise comparisons conducted on the number of active sessions of stimulation. Significant differences were found between 1 and 10 sessions, and 5 and 10 sessions of stimulation. **denotes significance at the $p < .05$ level.

There were no significant differences between idiopathic and acquired dystonia study effect sizes ($Q(1) = 2.12, p = .13$), although idiopathic dystonia studies displayed a significant mean effect (idiopathic $g = 0.26, p < .001$), whereas acquired studies did not ($g = 0.02, p = 0.89$;

NIBS in Dystonia: A Meta-Analysis

Figure 4). The lack of significant difference in pairwise comparisons is likely due to the small number of acquired dystonia studies.




Figure 4. Meta-regression conducted on dystonia etiology. No significant differences were found between idiopathic and acquired dystonia, however, idiopathic dystonia effect sizes were significant. ** denotes significance and the $p < .05$ level.

Effect sizes for studies which utilised motor training concurrently with NIBS were significantly larger than studies which applied NIBS alone, $Q(1) = 4.43, p = .04$. Overall mean effect sizes for both groups were significant, NIBS and motor training $g = 0.55, p = .001$ and NIBS alone $g = 0.15, p = 0.03$ (Figure 5).

NIBS in Dystonia: A Meta-Analysis




Figure 5. Meta-regression conducted on NIBS with and without concurrent motor training. No significant difference was found between NIBS and training and NIBS only, however, individual effects were significant. ** denotes significance and the $p < .05$ level.

Meta-regressions on mean age and gender ratio of participants were not significant: mean age $b = 0.002$, $SE = 0.005$, $p = 0.66$ and gender ratio $b = -0.03$, $SE = 0.03$, $p = 0.28$ (supplementary 11 and 12). Additionally, these moderators showed no significant effect when the outlier⁶⁴ study was included (supplementary file 13 and 14).

Evaluation of Bias

Methodological quality of studies, as assessed by the RoB2, is presented in Figure 6. An overall judgement of high risk of bias was given where studies had a high risk of bias in at least one domain. Three studies were considered to be at high risk of bias. Borich et al.⁵³ was considered to be at high risk of bias due to missing outcome data. Bradnam et al.⁵⁵ and Rossett-Llobet et al.³³ indicated that participant allocation to sham or active NIBS group was not concealed, and thus were judged at high risk of bias for the domain of random sequence generation. Most studies were judged to be at an unclear risk of bias in the domain of random sequence generation due to a lack of reporting how participants were randomized, and whether the allocation sequence was concealed. Furthermore, the domain of selective outcome reporting

NIBS in Dystonia: A Meta-Analysis

was judged to be at an unclear risk of bias for most studies, due to insufficient information available to permit a judgement of low risk (e.g., trial protocols)⁴⁰. Overall, the literature was characterized by an unclear risk of bias.

NIBS in Dystonia: A Meta-Analysis

	Benninger et al. (2011)	Bhanpuri et al. (2015)	Bologna et al. (2016)	Borich et al. (2009)	Bradnam et al. (2015)	Bradnam et al. (2016)	Buttkus et al. (2010a)	Buttkus et al. (2011)	Furuya et al. (2014)	Havrankova et al. (2010)	Huang et al. (2012)	Kimberley et al. (2013)	Koch et al. (2014)	Kranz et al. (2010)	Linssen et al. (2015)	Lozeron et al. (2017)	Murase et al. (2005)	Odorfer et al. (2019)	Pirio Richardson et al. (2015)	Rossett-Llobet et al. (2014)	Rossett-Llobet et al. (2015)	Sadnicka et al. (2014)	Siebner et al. (1999)	Siebner et al. (2003)	Wagle Shukla et al. (2018)	Young et al. (2014)
Randomisation	?	?	?	?	-	?	?	?	?	?	?	?	?	?	?	+	?	?	?	?	-	?	?	?	?	?
Blinding of participants/personnel	+	+	+	?	+	+	+	+	+	+	+	+	+	?	?	+	?	?	+	?	+	+	?	+	?	+
Incomplete outcome data	+	+	?	-	+	+	+	+	?	+	+	+	+	+	+	+	+	+	+	+	+	?	+	?	+	+
Blinding/measurement of outcomes	+	+	+	+	+	+	+	+	+	+	+	+	+	?	+	+	+	+	+	?	+	+	+	+	+	+
Selective outcome reporting	+	?	?	?	+	?	?	?	?	?	?	?	?	?	?	?	?	?	+	?	?	?	?	?	?	?
Overall	?	?	?	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	-	?	?	?	?	?

Figure 6. Risk of bias assessment for individual studies. Green boxes (+) = low risk of bias; orange boxes (?) = unclear risk of bias; red boxes (-) = high risk of bias.

NIBS in Dystonia: A Meta-Analysis

The funnel plot analysis revealed two studies outside the boundaries of the funnel^{32, 44} (supplementary 15). Egger's test trended towards significance ($t(26) = 1.95$, $p = 0.06$). Whilst this is indicative of symmetry within the funnel plot, suggesting that publication bias may not have affected this meta-analysis, results should be interpreted with caution given that an outlier study⁶⁴ was not included in this analysis.

Discussion

The primary aim of this systematic review and meta-analysis was to evaluate the efficacy of NIBS on dystonia symptoms. Overall meta-analysis of 27 studies demonstrated a small, yet significant effect for NIBS decreasing symptoms of dystonia. Further meta-analyses were then conducted separating studies by the different types of NIBS, dystonias, brain regions stimulated, and outcome measures. These analyses showed significantly reduced dystonia symptoms for 0.2Hz rTMS and cathodal tDCS, blepharospasm and task-specific FHD (including writer's and musician's dystonias individually), and the ACC, M1 and dPM. Lastly, meta-regression analyses suggested that 10 sessions of active stimulation, or NIBS applied concurrently with motor training had a significant effect on study effect size.

Brain region stimulated and type of non-invasive brain stimulation

Studies stimulating the M1, dPM and ACC demonstrated significantly reduced dystonia symptoms. However, the ACC effect was only contributed to by one study, therefore this result should be interpreted with caution. Further, two inhibitory protocols were found increase the effect of NIBS – specifically, 0.2Hz rTMS and cathodal tDCS. That stimulation of the M1 and dPM, and use of inhibitory NIBS protocols significantly predicted an effect of NIBS on dystonia symptoms is in line with prior research, demonstrating increased excitability in sensorimotor areas including the motor, premotor and somatosensory cortices¹⁵⁻¹⁷. This can be seen through the excessive contraction of both agonist and antagonist muscles in dystonia, leading to unwanted muscle spasms and motor overflow⁸⁰. Thus, the application of inhibitory NIBS protocols to these cortical areas may downregulate cortical and network activity, leading to a reduction in symptoms.

Type of dystonia

When separating meta-analysis by type of dystonia, NIBS significantly reduced symptoms in blepharospasm and task-specific FHD, inclusive of musician's dystonia and writer's cramp. However, the effect for blepharospasm should be interpreted with caution, as only one study was

NIBS in Dystonia: A Meta-Analysis

included in this analysis⁷⁸. While task-specific FHDs significantly benefitted from the application of NIBS, hand dystonia did not reach significance. Hand dystonia NIBS targets were spread over several brain regions, including the cerebellum and sensorimotor areas. Further, both inhibitory and excitatory NIBS protocols were used, with cTBS, anodal and cathodal tDCS, and 1Hz rTMS all trialed. The variability in protocol and targets in hand dystonia, along with the lack of contributing studies, is likely to have contributed to the non-significant finding. Conversely, task-specific FHD studies mainly targeted the M1 and dPM, with the most common NIBS protocol cathodal tDCS (or anodal and cathodal protocols combined in the same study) to the M1. Future trials in task-specific FHD should consider utilizing inhibitory protocols targeting the M1 and dPM, in order to maximize the therapeutic effects of NIBS in this cohort.

Number of non-invasive brain stimulation sessions

Studies ranged from a single session of NIBS to several sessions over multiple weeks. 16 of the 20 studies included in the qualitative review applied multiple sessions of stimulation, all reporting a reduction in dystonia symptoms upon completion of the NIBS sessions – however, statistical significance for many studies was not reported. Meta-regression analysis demonstrated that 10 sessions of active stimulation was more effective for improving dystonia symptoms than one or five sessions of stimulation. The finding of 10 sessions of active stimulation having a larger mean effect than one session is consistent with previous research that suggests consecutive sessions of NIBS, such as rTMS, are more effective in inducing longer-lasting plastic changes within cortical regions such as the M1⁸¹. It is also consistent with clinical protocols for NIBS treatments in neuropsychiatric disorders where rTMS is applied over a number of sessions, for example, depression (30 sessions over 4-6 weeks)⁸² and obsessive-compulsive disorder (29 sessions)⁸³. Nonetheless, optimal parameters for both NIBS protocols and session quantity and timing for dystonia are yet to be established. Future clinical trials should include at least 10 sessions of NIBS to increase therapeutic efficacy, and further examine cumulative effects of NIBS paradigms within dystonia patients.

Concurrent non-invasive brain stimulation and motor training

There was a significant difference in effect sizes between studies which implemented concurrent NIBS and motor training and those where only NIBS was applied, with studies which applied concurrent NIBS and motor training having a larger overall effect on dystonia symptoms. All studies included in the meta-regression which implemented concurrent NIBS and motor training did so in musician's dystonia patients, using tDCS to the M1. Studies utilised motor training programs such as sensory-motor retuning^{32, 33}, a type of therapy commonly used in

NIBS in Dystonia: A Meta-Analysis

musician's dystonia that facilitates proprioceptive changes in the affected limb, and helps to modify abnormal cortical organization of sensory areas⁸⁴. Research in stroke patients indicates that utilizing tDCS over the sensorimotor areas in conjunction with motor training can improve motor function and produce functional changes in sensorimotor areas beyond that of training alone⁸⁵⁻⁸⁷. The use of tDCS may assist with improvement of motor functioning by modulating cortical excitability and increasing plasticity within the targeted cortical area, allowing for optimal conditions in which to consolidate the effects of motor training or therapy⁸⁸. Thus, future research should further examine the promising therapeutic effects of combined tDCS and motor training programs, such as sensory-motor retuning, in other types of dystonia beyond musician's dystonia.

Idiopathic versus acquired dystonia

Meta-regression demonstrated that, although there was no significant difference between idiopathic and acquired dystonia study effect sizes, idiopathic dystonia studies had a significant mean effect. Of the studies that utilised acquired dystonia patients in the overall meta-analysis, two studies recruited participants with cerebral palsy^{44, 45} and one with Wilson's disease⁴³. Given that the basal ganglia are thought to be involved in dystonia as part of the sensorimotor network, the atrophy or lesioning of this brain region, as is often seen in cerebral palsy and Wilson's disease patients, may result in different NIBS outcomes for those with acquired dystonia in comparison to those with idiopathic dystonia. Previous research in idiopathic writer's cramp patients has demonstrated reduced functional connectivity in comparison to healthy controls, in areas such as the bilateral thalamus, putamen and globus pallidus, and left dPM⁸⁹. However, a single session of rTMS induced a significant increase in connectivity in basal ganglia regions, specifically the bilateral thalamus and putamen⁸⁹. This suggests that although NIBS is applied to the cortex, effects extend to the basal ganglia and other subcortical structures, highlighting the need for an unaltered pathway between basal ganglia and stimulated cortex in dystonia patients to optimize NIBS outcomes⁴⁵.

Limitations

A limitation of this meta-analysis was only reviewing dystonia outcomes at the first time-point of assessment after the NIBS intervention. Several studies examined the effects of the NIBS at multiple timepoints (e.g., mid-intervention or four weeks post-intervention), and thus only estimating the effect of NIBS at the immediate end point of the intervention may have led to an overestimation of the true intervention effect, and may not accurately inform how effective the use of NIBS on symptoms of dystonia is long-term.

NIBS in Dystonia: A Meta-Analysis

A moderate level of between-study heterogeneity was found in this meta-analysis. Whilst secondary analyses were conducted to find moderators of the effect, other methodological differences between studies may have contributed to the significant level of heterogeneity – for example, the number of pulses applied in rTMS protocols. The overall methodological quality of the evidence was mixed, with Figure 6 demonstrating the uncertainty in whether randomization and selective outcome reporting influenced individual study results, and thus overall effect size. Notably, the inability to judge the domain of selective outcome reporting as low risk may suggest that the study-level effect sizes were, to a degree, overestimated. Although Egger’s test was non-significant, suggesting that the research field may not suffer from publication bias, meta-analysis results should be considered bearing in mind the standard of reporting.

Conclusions

The present systematic review and meta-analysis found a small effect size in favor of NIBS reducing symptoms of dystonia. The use of ‘inhibitory’ NIBS protocols (i.e., 0.2Hz rTMS and cathodal tDCS), stimulation of the M1 and dPM, protocols employing a greater number of sessions, and concurrent motor training protocols demonstrated the highest treatment effects for NIBS. Future research should apply 10 sessions or more of NIBS and further investigate the use of motor training concurrently with NIBS, to yield the high-quality evidence needed to translate this promising therapeutic technique to clinical use.

NIBS in Dystonia: A Meta-Analysis

Author roles: Jordan Morrison-Ham 1abc, 2ab, 3ab.

Gillian M. Clark 2abc, 3b.

Elizabeth G. Ellis 1c, 3b.

Andris Cerins 1c, 3b.

Juho Joutsa 1ab, 3c.

Peter G. Enticott 1ab, 3c.

Daniel T. Corp 1abc, 2abc, 3b.

- 1. Research Project.** a. Conception, b. organization, c. execution.
- 2. Statistical Analysis.** a. design, b. execution, c. review/critique.
- 3. Manuscript.** a. writing of the first draft, b. review/critique.

NIBS in Dystonia: A Meta-Analysis

References

1. Tarsy D, Simon, DK. Dystonia. *N Engl J Med*. 2006;355(8):818-29.
2. Steeves TD, Day L, Dykeman J, Jette N, Pringsheim T. The Prevalence of Primary Dystonia: A Systematic Review and Meta-analysis. *Mov Disord*. 2012;27(14):1789-96.
3. Albanese A, Romito LM, Calandrella D. Therapeutic advances in dystonia. *Mov Disord*. 2015;30(11):1547-56.
4. Wu YS, Ni LH, Fan RM, Yao MY. Meta-Regression Analysis of the Long-Term Effects of Pallidal and Subthalamic Deep Brain Stimulation for the Treatment of Isolated Dystonia. *World Neurosurg*. 2019;129:e409-16.
5. Okromeijdze L, Tsuboi T, Eisinger R, et al. Functional and Structural Connectivity Patterns Associated with Clinical Outcomes in Deep Brain Stimulation of the Globus Pallidus Internus for Generalized Dystonia. *AJNR Am J Neuroradiol*. 2020;41(3):508-14.
6. Breakefield XO, Blood AJ, Li Y, Hallett M, Hanson PI, Standaert DG. The pathophysiological basis of dystonias. *Nat Rev Neurosci*. 2008;9:222-34.
7. Albanese A, Asmus F, Bhatia KP, et al. EFNS guidelines on diagnosis and treatment of primary dystonias. *Eur J Neurol*. 2011;18(1):5-18.
8. Bradnam LV, McDonnell MN, Ridding MC. Cerebellar Intermittent Theta-Burst Stimulation and Motor Control Training in Individuals with Cervical Dystonia. *Brain Sci*. 2016;6(4):56.
9. Koch G, Porcacchia P, Ponzo, V, et al. Effects of two weeks of cerebellar theta burst stimulation in cervical dystonia patients. *Brain Stimul*. 2014;7(4): 64-72.
10. Mutz J, Vipulanathan V, Carter B, Hurlmann R, Fu CHY, Young AH. Comparative efficacy and acceptability of non-invasive surgical brain stimulation for the acute treatment of major depressive episodes in adults: systematic review and network meta-analysis. *BMJ* 2019;364:l1079.

NIBS in Dystonia: A Meta-Analysis

11. Feng Y, Zhang B, Zhang J, Yin Y. Effects of Non-invasive Brain Stimulation on Headache Intensity and Frequency of Headache Attacks in Patients With Migraine: A Systematic Review and Meta-Analysis. *Headache* 2019;59(9):1436-47.
12. Carmi L, Alyagon U, Barnea-Ygael N, Zohar J, Dar R, Zangen A. Clinical and electrophysiological outcomes of deep TMS over the medial prefrontal and anterior cingulate cortices in OCD patients. *Brain Stimul.* 2018;11(1):158-65.
13. Klomjai W, Katz R, Lackmy-Vallee A. Basic principles of transcranial magnetic stimulation (TMS) and repetitive TMS (rTMS). *Ann Phys Rehabil Med.* 105;58(4):209-13.
14. Quartarone A, Hallett M. Emerging concepts in the physiological basis of dystonia. *Mov Disord.* 2013;28(7):958-67.
15. Beck S, Pirio Richardson S, Shamim EA, Dang N, Schubert M, Hallett M. Short intracortical and surround inhibition are selectively reduced during movement initiation in focal hand dystonia. *J Neurosci.* 2008;28(41):10363-9.
16. Corp DT, Joutsa J, Darby RR, et al. Network localization of cervical dystonia based on causal brain lesions. *Brain* 2019;142(6):1660-74.
17. Delnooz CCS, Helmich RC, Menendorp WP, van de Warrenburg BPC, Toni I. Writer's cramp: increased dorsal premotor activity during intended writing. *Hum Brain Mapp.* 2013;34(3):613-25.
18. Porcacchia P, de Toledo PA, Rodríguez-Baena A, et al. Abnormal cerebellar connectivity and plasticity in isolated cervical dystonia. *PLoS One* 2019;14(1):e0211367.
19. Tamura Y, Ueki Y, Lin P, et al. Disordered plasticity in the primary somatosensory cortex in focal hand dystonia. *Brain* 2009;132(3):749-55.
20. Cho HJ, Hallett M. Non-Invasive Brain Stimulation for Treatment of Focal Hand Dystonia: Update and Future Directions. *Mov Disord.* 2016;9(2):55-62.

NIBS in Dystonia: A Meta-Analysis

21. Benninger DH, Lomarev M, Lopez G, Pal N, Luckenbaugh DA, Hallett M. Transcranial direct current stimulation for the treatment of focal hand dystonia. *Mov Disord.* 2011;26(9):1698-702.
22. Sadnicka A, Hamada M, Bhatia KP, et al. Cerebellar Stimulation Fails to Modulate Motor Cortex Plasticity in Writing Dystonia. *Mov Disord.* 2013;29(10):1304-7.
23. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
24. Hidding U, Baurmer T, Siebner HR, et al. MEP latency shift after implantation of deep brain stimulation systems in the subthalamic nucleus in patients with advanced Parkinson's disease. *Mov Disord.* 2006 21(9):1471-6.
25. Kuhn AA, Trottenberg T, Kupsch A, Meyer BU. Pseudo-bilateral hand motor responses evoked by transcranial magnetic stimulation in patients with deep brain stimulation. *Clin Neurophysiol.* 2002;113(3):341-5.
26. Ouzanni M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan – a web and mobile app for systematic reviews. *Syst Rev.* 2016;5(1):210.
27. Hedges GV, Olkin I. *Statistical methods for meta-analysis.* Orlando: Academic Press; 1985.
28. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.2 [updated February 2021]. Cochrane, 2021. Available from www.training.cochrane.org/handbook
29. Buttkus F, Baur V, Jabusch HC, et al. Single-session tDCS-supported retraining does not improve fine motor control in musician's dystonia. *Restor Neurol Neurosci.* 2011;29(2):85-90.

NIBS in Dystonia: A Meta-Analysis

30. Furuya S, Nitsche MA, Paulus W, Altenmüller E. Surmounting retraining limits in musicians' dystonia by transcranial stimulation. *Ann Neurol*. 2014;75(5):700-7.
31. Pirio Richardson S, Tinaz S, Chen R. Repetitive transcranial magnetic stimulation in cervical dystonia: effect of site and repetition in a randomized pilot trial. *PLoS One* 2015;10(4):e0124937.
32. Rosset-Llobet, J., Fàbregas-Molas, S., & Pascual-Leone, Á. (2014). Transcranial direct current stimulation improves neurorehabilitation of task-specific dystonia: A Pilot Study. *Med Probl Perform Art*. 2014;29(1):16-8.
33. Rosset-Llobet J, Fàbregas-Molas S, Pascual-Leone A. Effect of transcranial direct current stimulation on neurorehabilitation of task-specific dystonia: A double-blind, randomized clinical trial. *Med Probl Perform Art*. 2015;30(3):178-84.
34. Siebner HR, Tormos JM, Ceballos-Baumann AO, et al. Low-frequency repetitive transcranial magnetic stimulation of the motor cortex in writer's cramp. *Neurology* 1999;52(3):529-37.
35. Steichen T. METANINF: Stata module to evaluate influence of a single study in meta-analysis estimation [STATA command]. *Statistical Software Components S419201*. Boston College Department of Economics; 2001. [cited July 2021]. Available from: <http://fmwww.bc.edu/RePEc/bocode/m/>
36. Corp DT, Bereznicki HGK, Clark GM, et al. Large-scale analysis of interindividual variability in theta-burst stimulation data: Results from the 'Big TMS Data Collaboration'. *Brain Stimul*. 2020;13(5):1476-88.
37. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*, 2003;327(7414):557-60.
38. Harbord R, Higgins JPT. METAREG: Stata module to perform meta-analysis regression [STATA command]. *Statistical Software Components S446201*. Boston College

NIBS in Dystonia: A Meta-Analysis

Department of Economics; 2009. [cited July 2021]. Available from:

<http://fmwww.bc.edu/RePEc/bocode/m>

39. Wilson DB. META_ANALYSIS: Stata module to perform subgroup and regression-type fixed- and random-effects meta-analyses [STATA command]. Statistical Software Components S458944. Boston College Department of Economics; 2021. [cited July 2021]. Available from: <http://fmwww.bc.edu/RePEc/bocode/m>
40. Sterne JAC, Savovic J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:I4898.
41. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to Meta-Analysis*. West Sussex: John Wiley & Sons, Ltd; 2009.
42. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34.
43. Lozeron, P, Poujois A, Meppiel E, et al. Inhibitory rTMS applied on somatosensory cortex in Wilson's disease patients with hand dystonia. *J Neural Transm (Vienna)* 2017;124(10):1161-70.
44. Bhanpuri NH, Bertucco M, Young SJ, Lee AA, Sanger TD. Multiday transcranial direct current stimulation causes clinically insignificant changes in childhood dystonia: A pilot study. *J Child Neurol*. 2015;30(12):1604-15.
45. Young SJ, Bertucco M, Sanger TD. Cathodal transcranial direct current stimulation in children with dystonia: A sham-controlled study. *J Child Neurol*. 2014;29(2):232-9.
46. Buttkus F, Baur V, Jabusch H-C, Paulus W, Nitsche MA, Altenmüller E. Retraining and transcranial direct current stimulation in musician's dystonia - a case report. *Mov Disord*. 2010;25(11):1758-60.

NIBS in Dystonia: A Meta-Analysis

47. Kimberley TJ, Schmidt RLS, Chen M, Dykstra DD, Buetefisch CM. Mixed effectiveness of rTMS and retraining in the treatment of focal hand dystonia. *Front Hum Neurosci*. 2015;9:385.
48. Okada Y, Shibamoto C, Osumi Y, et al. Transcranial Direct Current Stimulation Combined with Action Observation and Electromyographic Biofeedback Training in a Patient with Writer's Cramp. *J Mov Disord*. 2018;11(2):82-6.
49. Allam N, Brasil-Neto JP, Brandão P, Weiler F, de Barros Filho J, Tomaz C. Relief of primary cervical dystonia symptom by low frequency transcranial magnetic stimulation of the premotor cortex. *Arq Neuropsiquiatr*. 2007;65(3A):697-9.
50. Angelakis E, Liouta E, Andreadis N, et al. Transcranial alternating current stimulation reduces symptoms in intractable idiopathic cervical dystonia: A case study. *Neurosci Lett*. 2013;533:39-43.
51. Betti S, Spoto A, Castiello U, Sartori L. Testing rTMS-induced neuroplasticity: A single case study of focal hand dystonia. *Neural Plast*. 2018;2018:6464896.
52. Bologna M, Paparella G, Fabbrini A, et al. Effects of cerebellar theta-burst stimulation on arm and neck movement kinematics in patients with focal dystonia. *Clinical Neurophysiol*. 2016;127(11):3472-9.
53. Borich M, Arora S, Kimberley TJ. Lasting effects of repeated rTMS application in focal hand dystonia. *Restor Neurol Neurosci*. 2009;27(1):55-65.
54. Bradnam LV, Frasca J, Kimberley TJ. Direct Current Stimulation of Primary Motor Cortex and Cerebellum and Botulinum Toxin in a Person With Cervical Dystonia. *Brain Stimul*. 2014;7(6):909-11.
55. Bradnam LV, Graetz LJ, McDonnell MN, Ridding MC. Anodal transcranial direct current stimulation to the cerebellum improves handwriting and cyclic drawing kinematics in focal hand dystonia. *Front Hum Neurosci*. 2015;9:286.

NIBS in Dystonia: A Meta-Analysis

56. Buttkus F, Weidenmüller M, Schneider S, et al. Failure of cathodal direct current stimulation to improve fine motor control in musician's dystonia. *Mov Disord.* 2010;25(3):389-94.
57. Conte A, Rocchi L, Ferrazzano G, et al. Primary somatosensory cortical plasticity and tactile temporal discrimination in focal hand dystonia. *Clin Neurophysiol.* 2014;125(3):537-43.
58. Havrankova P, Jech R, Walker ND, et al. Repetitive TMS of the somatosensory cortex improves writer's cramp and enhances cortical activity. *Neuro Endocrinol Lett.* 2010;31(1):73-86.
59. Huang YZ, Rothwell JC, Lu CS, Wang J, Chen RS. Restoration of motor inhibition through an abnormal premotor-motor connection in dystonia. *Mov Disord.* 2010;25(6):696-703.
60. Huang Y-Z, Lu C-S, Rothwell JC, et al. Modulation of the disturbed motor network in dystonia by multisession suppression of premotor cortex. *PLoS One* 2012;7(10):e47574.
61. Kimberley TJ, Borich MR, Arora S, Siebner HR. Multiple sessions of low-frequency repetitive transcranial magnetic stimulation in focal hand dystonia: Clinical and physiological effects. *Restor Neurol Neurosci.* 2013;31(5):533-42.
62. Kimberley TJ, Borich MR, Schmidt RL, Carey JR, Gillick B. Focal hand dystonia: Individualized intervention with repeated application of repetitive transcranial magnetic stimulation. *Arch Phys Med Rehabil.* 2015;96(4 Suppl):S122-8.
63. Kranz G, Shamim EA, Lin PT, Kranz GS, Voller B, Hallett M. Blepharospasm and the modulation of cortical excitability in primary and secondary motor areas. *Neurology* 2009; 73(23):2031-6.

NIBS in Dystonia: A Meta-Analysis

64. Kranz G, Shamim EA, Lin PT, Kranz GS, Hallett M. Transcranial magnetic brain stimulation modulates blepharospasm: A randomized controlled study. *Neurology* 2010;75(16):1465-71.
65. Lefaucheur J-P, Fénelon G, Ménard-Lefaucheur I, Wendling S, Nguyen JP. Low-frequency repetitive TMS of premotor cortex can reduce painful axial spasms in generalized secondary dystonia: a pilot study of three patients. *Neurophysiol Clin*. 2004;34(3-4):141-5.
66. Linssen MW, van Gaalen J, Munneke MAM, Hoffl BS, Hulstijn W, van de Warrenburg BPC. A single session of cerebellar theta burst stimulation does not alter writing performance in writer's cramp. *Brain* 2015;136(Pt 6):e355.
67. Marceglia S, Mrakic-Sposta S, Fumagalli M, et al. Cathodal transcranial direct current stimulation improves focal hand dystonia in musicians: A two-case study. *Front Neurosci*. 2017;11:508.
68. Murase N, Rothwell JC, Kaji R, et al. Subthreshold low-frequency repetitive transcranial magnetic stimulation over the premotor cortex modulates writer's cramp. *Brain* 2005;128(1):104-15.
69. Naro A, Billeri L, Portaro S, Bramanti P, Calabrò RS. Lasting Effects of Low-Frequency Repetitive Transcranial Magnetic Stimulation in Writer's Cramp: A Case Report. *Front Hum Neurosci*. 2019;13:314.
70. Odorfer TM, Homola GA, Reich MM, Volkmann J, Zeller D. Increased Finger-Tapping Related Cerebellar Activation in Cervical Dystonia, Enhanced by Transcranial Stimulation: An Indicator of Compensation? *Front Neurol*. 2019;10:231.
71. Salatino A, Boccia G, Dardanello D, et al. Acute and cumulative effects of rTMS on behavioural and EMG parameters in Focal Hand Dystonia. *Heliyon* 2019;5(11):e02770.

NIBS in Dystonia: A Meta-Analysis

72. Sharma K, Cucca A, Lee A, Agarwal S, Frucht SJ, Biagioni MC. Transcranial magnetic stimulation therapy for focal leg dystonia: A case report. *J Clin Mov Disord.* 2019;6:1.
73. Shin, H.-W., Youn, Y. C., & Hallett, M. (2019). Focal Leg Dystonia Associated with Cerebellar Infarction and Application of Low-Frequency Cerebellar Transcranial Magnetic Stimulation: Evidence of Topographically Specific Cerebellar Contribution to Dystonia Development. *Cerebellum* 2019;18(6):1147-50.
74. Siebner HR, Filipovic SR, Rowe JB, et al. Patients with focal arm dystonia have increased sensitivity to slow-frequency repetitive TMS of the dorsal premotor cortex. *Brain* 2003;126(12):2710-25.
75. Trebossen V, Bouaziz N, Benadhira R, Januel D. Transcranial direct current stimulation for patients with benign essential blepharospasm: a case report. *Neurol Sci.* 2017;38(1):201-2
76. Veugen LC, Hoffl BS, Stegeman DF, Van De Warrenburg BP. Inhibition of the dorsal premotor cortex does not repair surround inhibition in writer's cramp patients. *Exp Brain Res.* 2013;221(1):85-92.
77. Vucurovic K, Emeriau S, Coulon JM, Peruzzi P, Litré CF, Kaladjian A. Functional brain neuroimaging-guided repetitive transcranial magnetic stimulation in neurodevelopmental disorders: The case of a schizencephaly-related spastic dystonia. *J Neurol Sci.* 2017;378:167-9.
78. Wagle Shukla A, Hu W, Legacy J, Deeb, W, Hallet, M. Combined effects of rTMS and botulinum toxin therapy in benign essential blepharospasm. *Brain Stimul.* 2018;11(3):645-7.
79. Hallett M. Neurophysiology of dystonia: The role of inhibition. *Neurobiol Dis.* 2011;42(2):177-84.

NIBS in Dystonia: A Meta-Analysis

80. Phukan J, Albanese A, Gasser T, Warner T. Primary dystonia and dystonia-plus syndromes: clinical characteristics, diagnosis, and pathogenesis. *Lancet Neurol*. 2011;10(12):1274-85.
81. Baumer T, Lange R, Liepert J, et al. Repeated premotor rTMS leads to cumulative plastic changes of motor cortex excitability in humans. *Neuroimage* 2003;20(1):550-60.
82. McClintok SM, Reti IM, Carpenter LL, et al. Consensus recommendations for the clinical application of repetitive transcranial magnetic stimulation (rTMS) in the treatment of depression. *J Clin Psychiatry* 2018;79(1):16cs10905.
83. Roth Y, Tendler A, Arikan MK, et al. Real-world efficacy of deep TMS for obsessive-compulsive disorder: Post-marketing data collected from twenty-two clinical sites. *J Psychiatr Res* 2021; 137; 667-72.
84. Candia V, Rosset-Llobet J, Elbert T, Pascual-Leone A. Changing the brain through therapy for musicians' hand dystonia. *Ann N Y Acad Sci*. 2005;1060:335-42.
85. Lindenberg R, Renga V, Zhu LL, Nair D, Schlaug G. Bihemispheric brain stimulation facilitates motor recovery in chronic stroke patients. *Neurology* 2010;75(24):2176-84.
86. Nair DG, Renga V, Lindenberg R, Zhu L, Schlaug G. Optimizing recovery potential through simultaneous occupational therapy and non-invasive brain stimulation using tDCS. *Restor Neurol Neurosci*. 2011;29(6):411-20.
87. Alisar SC, Ozen S, Sozay S. Effects of Bihemispheric Transcranial Direct Current Stimulation on Upper Extremity Function in Stroke Patients: A randomized Double-Blind Sham-Controlled Study. *J Stroke Cerebrovasc Dis*. 2020;29(1):104454.
88. Bolognini N, Pascual-Leone A, Fregni F. Using non-invasive brain stimulation to augment motor training-induced plasticity. *J Neuroeng Rehabil*. 2009;6:8.

NIBS in Dystonia: A Meta-Analysis

89. Bharath RD, Biswal BB, Bhaskar MV, et al. Repetitive transcranial magnetic stimulation induced modulations of resting state motor connectivity in writer's cramp. *Eur J Neurol.* 2015;22(5):795-805.