IS MODULAR CONTROL RELATED TO FUNCTIONAL OUTCOMES IN INDIVIDUALS WITH KNEE OSTEOARTHRITIS AND FOLLOWING TOTAL KNEE ARTHROPLASTY?

Rebekah R. Koehn[1], Sarah A. Roelker[2], Xueliang Pan[3], Laura C. Schmitt[4-6], Ajit M.W. Chaudhari[1,4-8], Robert A. Siston[1,4,5,7,8]

[1]Department of Mechanical and Aerospace Engineering, The Ohio State University, Columbus, OH, USA
[2]Department of Kinesiology, University of Massachusetts Amherst, Amherst, MA, USA
[3]Center for Biostatistics and Bioinformatics, The Ohio State University, Columbus, OH, USA
[4]School of Health and Rehabilitation Sciences, The Ohio State University, Columbus, OH, USA
[5]Sports Medicine Research Institute, The Ohio State University Wexner Medical Center, Columbus, OH, USA
[6]Division of Physical Therapy, School of Health and Rehabilitation Sciences, The Ohio State University, Columbus, OH, USA
[7]Department of Biomedical Engineering, The Ohio State University, Columbus, OH, USA
[8]Department of Orthopaedics, The Ohio State University, Columbus, OH, USA

Author Contributions:
RK: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Visualization, Writing – Original Draft Preparation
SR: Conceptualization, Data Curation, Formal Analysis, Methodology, Software, Writing – Review & Editing
XP: Conceptualization, Formal Analysis, Funding Acquisition, Software, Writing – Review & Editing
LS: Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing
AC: Conceptualization, Funding Acquisition, Methodology, Resources, Writing – Review & Editing
RS: Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Writing – Review & Editing

Word count: 6704

Short Title: Modular Control and Function Before and After TKA

*Address Correspondence to:
Robert A. Siston
Department of Mechanical and Aerospace Engineering
Scott Laboratory
201 W 19th Ave
Columbus, OH 43210 USA
Phone: 614-247-2721
Email: siston.1@osu.edu

ABSTRACT

BACKGROUND: Individuals who undergo total knee arthroplasty (TKA) for treatment

of knee osteoarthritis often experience suboptimal outcomes. Investigation of neuromuscular

control strategies in these individuals may reveal factors that contribute to these functional deficits.

The purpose of this pilot study was to determine the relationship between patient function and

modular control during gait before and after TKA. **METHODS:** Electromyography data from 36

participants (38 knees) were collected from 8 lower extremity muscles on the TKA-involved limb

during ≥ 5 over-ground walking trials before (n=30), 6-months after (n=26), and 24-months after

(n=13) surgery. Muscle modules were estimated using non-negative matrix factorization. The

number of modules was determined from 500 resampled trials. **RESULTS:** A higher number of

modules was related to better performance-based and patient-reported function before and 6-

months after surgery. Participants with organization similar to healthy, age-matched controls

trended toward better function 24-months after surgery, though these results were not statistically

significant. We also observed plasticity in the participants' modular control strategies, with 100%

of participants who were present before and 24-months after surgery (10/10) demonstrating

changes in the number of modules and/or organization of at least 1 module. **CONCLUSIONS:**

This pilot work suggests that functional improvements following TKA may initially present as

increases in the number of modules recruited during gait. Once individuals can produce discrete

neuromuscular control patterns, further improvements may present as improved module

organization. **NOTEWORTHY:** This work is the first to characterize motor modules in TKA both

before and after surgery and to demonstrate changes in the number and organization of modules

over the time course of recovery, which may be related to changes in patient function. The

plasticity of modular control following TKA is a key finding which has not been previously

documented and may be useful in predicting or improving surgical outcomes through novel

rehabilitation protocols.

INTRODUCTION

Total knee arthroplasty (TKA) is the definitive end-stage treatment for knee osteoarthritis (KOA), with over 750,000 primary TKAs occurring in 2014 and over 1.25 million primary TKAs expected in the U.S. annually by 2030 (1). While TKA improves function in most patients, more than 20% of adults demonstrate functional deficits 2-years post-TKA (2). These functional deficits include slower walking speeds (3), stiff-knee gait (4), and difficulty in stair descent (5) and sit-to-stand transfer (6). Several studies have suggested that these deficits are influenced by a variety of factors including implant design (7), surgical technique (8), ligament laxity and soft tissue balance (9), quadriceps strength (10), strength of the non-operated limb (11), and preoperative conditions (12). However, the primary factor or combination of factors which reliably predict patient outcomes has yet to be determined.

Several studies have also examined neuromuscular control as a factor which may explain the functional deficits seen in this population. One study found that TKA patients, both before and 1-month after surgery, demonstrated higher co-activation of the quadriceps and hamstrings during stand-to-sit transfer and performed the five-times-sit-to-stand task more slowly than healthy adults (13). Another study determined that TKA patients demonstrated quadriceps muscle activation deficits after surgery, but these deficits were not associated with quadriceps strength 3-months after TKA (14). Some work has also been done to investigate changes in patient function and neural control in this population from the preoperative to postoperative condition. Hubley-Kozey et al. (15) studied patient function and muscle activation patterns in patients before and 1-year following TKA and found that while functional improvements were accompanied by shifts in activation patterns toward the patterns of asymptomatic individuals, the patterns were still statistically different from the asymptomatic individuals. While these studies provide valuable

insight into the influence of activation patterns on function in KOA and TKA, it is still unknown

how these electromyography (EMG)-measured factors may change over the time course of

recovery after surgery. Further, it remains unclear if changes in co-activations and muscle

activation patterns are indicative of changes in the underlying neuromuscular control strategies.

Since individual EMG waveforms alone do not provide insight into the underlying mechanisms of

control, other tools are necessary in order to determine the influence of neuromuscular control on

patient function.

The theory of modular muscle control has gained popularity as a tool to explore

neuromuscular control (16). The theory describes a simplified neural control strategy in which a

group of muscles is activated synergistically by a common neural command, reducing the set of

activation profiles used to complete a motion (17). The number of these commands, or "motor

primitives" (18), and the organization of muscles within the modules, which are selected out of an

individual's library of possible commands in order to complete a task, represent varying

complexity of control patterns (19-21). A higher number of modules used to complete a task, such

as walking, may indicate a more complex neuromuscular control strategy and greater flexibility in

control (21). Muscles that are organized into a specific module activate as a group according to

the module's timing profile, and recent literature has suggested that populations with neurologic

movement disorders, including Parkinson's disease, post-stroke hemiparesis, and incomplete

spinal cord injury, demonstrate altered module organization in comparison to healthy controls (22-

25). To our knowledge, there has been only one study examining modular control in the TKA

population. Ardestani et al. (21) found that 1-year following a cruciate retaining (CR) TKA, a low-

functioning group of participants demonstrated 2-3 motor modules while a high-functioning group

demonstrated 4-5 modules, which more closely resembled a healthy control group that

demonstrated 5-6 modules. The results of that study suggest that the complexity of neural control

strategy may be an underlying mechanism that contributes to the large range of post-operative

functional outcomes observed in this population. However, the relationship between module

organization and function has yet to be characterized in the TKA population. Additionally, it has

been suggested that rehabilitation programs may influence the complexity and organization of

modular control in individuals with spinal cord injuries or neurological disorders (22, 26).

However, it is still unknown whether changes in modular control exist in individuals with

orthopaedic disorders and whether such changes may be related to the time course of functional

improvements following surgical interventions, like TKA.

Therefore, the purpose of this pilot study was to determine the relationship between patient

function and modular control during gait before and at two timepoints after surgery and to

investigate the plasticity of neuromuscular control in the KOA and TKA populations. We

hypothesized that better patient function before surgery, 6-months after surgery, and 24-months

after surgery would be associated with (I) a higher number of modules and (II) module

organization more similar to that of healthy controls. Additionally, we aimed to examine the

relationship between changes in function and changes in the number and organization of modules

over the time course of recovery from TKA.

MATERIALS & METHODS

Data Collection

Prior to enrolling in the original longitudinal study (27), 36 individuals (38 knees) with

medial compartment KOA (19/19 R/L) provided written informed consent, in accordance with The

Ohio State University's Institutional Review Board. The cohort in the current study is a superset

of that reported in Freisinger et al. (27), which examined gait biomechanics before surgery, and

Chaudhari et al. (28), which examined muscle strength before and after surgery. Three orthopaedic

surgeons (JFG, AHG, MDB; see acknowledgements) identified potential participants based on

consultation for a TKA at The Ohio State Wexner Medical Center. Participants were required to

have a body-mass index (BMI) less than 45, the ability to walk 20 meters without an assistive

device, and no history of previous TKA or osteotomy. Recruitment began in April 2012 and

continued on a rolling basis, and data was collected from May 2012 to May 2017. All participants

were scheduled to undergo a primary posterior-stabilizing (PS) TKA (Zimmer NexGen LPS Flex

Knee) prior to study recruitment. By consensus, two fellowship-trained musculoskeletal

radiologists (JP, AR; see acknowledgements) determined the Kellgren-Lawrence classification

(KL-grade) of each participant's right and left knees (29).

The participants were tested in the Clinical, Functional, and Performance Biomechanics

Laboratory at The Ohio State University approximately 1-month before, 6-months after, and 24-

months after undergoing surgery. Each participant performed a minimum of 5 over-ground

walking trials. Surface EMG data were collected at 1500 Hz (Telemyo DTS System, Noraxon,

Scottsdale, AZ) from 16 Ag/AgCl dual-electrodes (Peripheral Nerve Stimulation Dual Element

Electrodes, rectangular, 1.625 in x 3.25 in, 0.42 in sensor diameter, 1.625 in inter-electrode

distance, Vermed, Buffalo, NY) affixed over the bellies of 8 lower extremity muscles, bilaterally:

rectus femoris, vastus lateralis, vastus medialis, biceps femoris, medial hamstrings, lateral

gastrocnemius, medial gastrocnemius, and soleus. Prior to affixing the electrodes, the skin over

each muscle was shaved to remove any hair and then cleansed and lightly abraded with alcohol

wipes.

In addition to gait trials, several performance-based and patient-report measures of function were recorded. Each participant completed three clinical performance-based assessments: the timed stair-climbing test (SCT) (30), the timed up-and-go test (TUG) (31), and the six-minute walk test (6MW) (32). Each participant also provided self-reported functional data using four of the Knee Injury and Osteoarthritis Outcome Score (KOOS) (33) survey subscales: pain, symptoms, activities of daily living (ADL), and quality of life (QOL). For each question, the participants' answers were scored on a scale from 0 to 4, with a higher score indicating better self-reported function. The scores were totaled within each subscale and normalized such that a maximum score of 100 represented high function and a minimum score of 0 represented poor function.

Due to high BMI (e.g. 33.9 ± 5.1 kg/m^2 before TKA), a common trait in individuals with osteoarthritis, soft tissue motion artifact was present in the EMG data in many of the trials. Due to this motion artifact, several participants were excluded from the study. Additionally, all 36 participants did not appear at all three data collection sessions. We were able to include data for 30 participants before surgery, 26 participants at 6-months post-TKA, and 13 participants at 24-months post-TKA (Table 1) in this pilot study.

Identification of Muscle Modules

We processed the EMG data for the TKA-involved limb from each gait trial to prepare for module extraction. The EMG data were demeaned and bandpass filtered to frequencies between 50 Hz and 300 Hz (Butterworth, 6th order) due to higher levels of motion artifact associated with high BMI values in this population, as recommended by Kieliba et al. (34) and Santuz et al. (35). The EMG data were then full-wave rectified and smoothed using a 6 Hz lowpass filter (Butterworth, 6th order). Trials with excessive motion artifact were excluded. We extracted the

maximum number of available gait cycles from each participant, and those with fewer than 5

available gait cycles were excluded. Linear envelopes were formed by discretizing the data such

that each trial contained 201 data points, accounting for every 0.5% of the gait cycle, concatenating

all available gait cycles, and normalizing each muscle first to the maximum value across all gait

cycles and then to unit variance (36).

Muscle modules were calculated using non-negative matrix factorization (NMF) (37).

Given that neuron firing rates and strengths cannot be negative, NMF provides insight into an

individual's neural control strategy that can be interpreted physiologically (16). First, the linear

envelopes for each subject were organized into *m x t* matrices representing the original EMG

(EMG$_o$), where *m* is the number of muscles (i.e. *m* = 8) and *t* is the number of data points (i.e. *t* =

201 x the number of gait cycles available). The NMFs were performed using the Statistics and

Machine Learning Toolbox in Matlab R2017a (The MathWorks Inc.; Natick, MA) which

populated an *n x t* matrix (Pattern Matrix) and an *m x n* matrix (Weighting Matrix), where *n* is the

number of modules. The Pattern Matrix represents the temporal waveform of each module while

the Weighting Matrix represents the weight of each muscle within each of the modules, such that

the product of these matrices (EMG$_r$) reconstructs EMG$_o$ (Fig. 1). The sum of the squared errors

between EMG$_r$ and EMG$_o$ was quantified with the percent of variability accounted for (VAF). The

VAF was calculated for each muscle (mVAF) of each solution using:

$$mVAF = \left(1 - \frac{\left(EMG_{o,m} - EMG_{r,m}\right)^2}{EMG_{o,m}^2}\right) \times 100\%$$ Equation 1

and the total VAF (tVAF) for each solution was calculated using:

$$tVAF = \left(1 - \frac{\sum_{m=1}^{8}(EMG_{o,m} - EMG_{r,m})^2}{\sum_{m=1}^{8} EMG_{o,m}^2}\right) \times 100\%$$ Equation 2

215
216
217

218
Fig. 1: Example of EMG Reconstruction using Non-Negative Matrix Factorization. Muscle
modules were calculated using non-negative matrix factorization (NMF) in which an *m x n*
Weighting Matrix is multiplied by an *n x t* Pattern Matrix to reconstruct the EMG patterns in an
m x t matrix, where *m* is the number of muscles (8 in this study), *n* is the number of modules (3
in this example), and *t* is the number of data points in the trial (201 in this study). The Pattern
Matrix represents the timing profile of each module. The Weighting Matrix represents how
strongly each muscle is represented in the module. This process is iterated, increasing *n* from 1 to
m until the error between the reconstructed EMG (black dotted line) and the original EMG (solid
gray line) is reduced to some acceptable level determined by the variability accounted for (VAF)
criteria.

In order to find the number of modules required to reconstruct EMG_o, the number of

modules was increased from 1 to *m* (i.e., *m* = 8; the number of muscles recorded) in separate NMFs

with a maximum of 10,000 iterations until the following criteria were satisfied: the lower bound

of the 95% confidence interval (CI) of tVAF was greater than 90% for all muscles and (a) the

minimum lower bound of the 95% CI of all mVAFs was greater than 75%, or (b) the minimum

lower bound of the 95% CI for all mVAFs was greater than 75% and the addition of another

module did not raise the lower bound of the 95% CI for that mVAF by more than 5% of its value

in the previous NMF (24). The 95% CIs of the VAFs were found using a bootstrapping technique

adopted from Allen et al. (22), wherein EMG_o was resampled 500 times with replacement. The

95% CI was assembled from the VAFs of all 500 bootstrapped samples. Once the appropriate

number of modules was determined, we ran a final NMF on the original EMG using a

multiplicative update algorithm with 50 replicates. To allow for comparison of modules between

participants and timepoints, the Weighting Matrix was normalized to the maximum muscle weight

within that module. Additionally, we evaluated tVAF for all possible module solutions (i.e. 1

module, 2 modules, …, 8 modules) as a measure of the complexity of the control strategy. A higher

tVAF value for any given module solution represents a better fit between EMG_r and EMG_o.

Therefore, a low tVAF value indicates that the solution is not capable of fully capturing the

complexity of the actual control strategy.

To characterize modular organization, we adapted definitions of W_{musc} and W_{sum} from

Hayes et al. (25). The number of significantly active muscles in each module was defined as W_{musc},

and the sum of the weights of the significantly active muscles in each module was defined as W_{sum}.

In Hayes et al. (25) a muscle was considered significantly active if the 95% CI of weights for the

500 bootstrapped gait cycles did not contain 0. However, participants in the current study often

had confidence intervals with a lower bound greater than zero but a very small upper bound. Under

the original definition from Hayes et al. (25), all modules for all participants in the TKA cohort

had a W_{musc} of 7 or 8, meaning all muscles were significantly active in all modules. To uncover

more subtle differences in W_{musc} and W_{sum} between participants in our cohort, we adapted the

definition of "significantly active," such that a muscle was considered significantly active if the

95% CI of weights did not contain 0 and the upper bound was greater than 0.25. Modules from

each of the 500 samples were sorted using a k-means algorithm to ensure like-modules were

grouped prior to determining the 95% CIs for the weights (38).

Healthy Control Group

Motor modules from 10 older adults with no known lower extremity osteoarthritis, a subset

of the cohort used in Roelker et al. (38), were used as healthy, age-matched controls to compare

to the KOA and TKA data. The processes for data collection and identification of muscle modules

are consistent with this study, with the exception of the bandpass filter used to process the EMG

data. Roelker et al. (38) used a 30-300 Hz bandpass filter while we used a 50-300 Hz bandpass

filter in the current study due to the previously mentioned high BMI and resulting motion artifact

observed in the TKA cohort. We determined W_{musc} and W_{sum} for each module for each healthy

participant based on our new definition for "significantly active" muscles.

Comparison of Module Organization

To determine similarities in module organization between populations, we compared the

module weightings of the study participants to those of the healthy controls. The muscle weights

of each module for each participant at each timepoint were compared to the averaged weights of

healthy age-matched controls with the same number of modules using Pearson correlations. A

critical correlation coefficient of $\rho \geq 0.834$ was selected based on the number of muscles (8) and

a *p*-value of 0.01 (22, 23, 25). A module was considered to have proper organization if its weights

were correlated with the healthy group and its W_{musc} and W_{sum} values were within the 95% CIs of

the healthy group. Participants were considered to have proper organization if all of their modules

were organized like healthy controls and poor organization if at least one module did not have

healthy organization.

Minimum Detectable Changes in Functional Performance

Differences in performance-based and self-reported measures were evaluated based on

minimum detectable changes (MDCs), defined by previously reported 90% CIs from

representative populations. Improvements were indicated by decreases in the time to complete

SCT (≥ 1.9 seconds; (39)) and TUG (≥ 2.49 seconds; (32)), increases in the distance walked during

6MW (≥ 61.34 meters; (32)), and increases in all KOOS subscales (≥ 10 points; (33)).

Statistics

To address our hypotheses, we investigated the relationship between functional

performance measures and modules in terms of their number (Hypothesis I) and organization

(Hypothesis II). At each time point, the functional measures were summarized by the number or

organization of modules. Generalized linear mixed models (GLMM) for repeated measures were

used to estimate the effects of number of modules and module organization (proper or poor) on all

functional measures at each time point. Time point, module number/organization, and their

interaction were included as fixed effects and participants were included as random effects in the

GLMMs. GLMMs allowed us to include all participants, even with missing data, assuming data

was missing at random in the analysis.

We examined differences in demographics, function, and modules between time points for

the TKA cohort and also examined differences in demographics and number of modules between

the TKA and healthy cohorts. Separate GLMMs were used to estimate the changes over time in

demographics data (height, mass, BMI, self-selected walking speed, stride length normalized to

leg length), performance-based function (SCT, TUG, 6MW), KOOS-subscale scores (Pain,

Symptoms, ADL, QOL), number of modules, and module organization (proper or poor). Separate

unpaired, two-tailed t-tests (all data was normally distributed; Anderson-Darling test for normality,

all $p \geq 0.052$) compared demographics data between the TKA group at 24-months after surgery

and the healthy control group. A Chi-square test compared number of modules between the healthy

control group and the group tested at 24-months post-TKA. As an additional examination of

module complexity, we used GLMMs to estimate differences in tVAF for all possible module

solutions (i.e. 1 to 8 modules) between time points.

All GLMMs were performed using Statistical Analysis Software (SAS Institute Inc.; Cary,

NC). Main effects were estimated from the fixed effects parameters or their corresponding contract

of the parameters from the GLMM, and, if significant, pairwise differences were assessed based

on the least squares means difference. All other statistical analysis was performed using Minitab

(Minitab, LLC; State College, PA). A significance level of $\alpha<0.05$ was established *a priori* for all

tests.

RESULTS

Population Demographics and Modules

324   The TKA group had statistically significant changes over time in self-selected walking

325   speed ($p = 0.001$) and BMI ($p = 0.043$; Table 1). Self-selected walking speed increased over time

and was slower before surgery than at 6- (pairwise $p = 0.007$) and 24-months (pairwise $p < 0.001$) after surgery. BMI also increased over time and was higher BMI 24-months after surgery than before (pairwise $p = 0.018$) or 6-months (pairwise $p = 0.049$) after surgery. Between the TKA cohort (24-months after surgery) and the healthy age-matched cohort, there were no significant differences in height, self-selected walking speed, or normalized stride length (all $p \geq 0.145$), but the TKA group had significantly higher mass ($p = 0.001$) and BMI ($p < 0.001$) than the healthy group (Table 1).

Table 1: Demographics, Modules, and Function.

		Patient Population; n = 38					Healthy Controls	Between Populations (t-test or Chi-square test)	
		Pre-TKA	6-months Post-TKA	24-months Post-TKA	Between Time Points (GLMM)				
		n = 30	n = 26	n = 13	p	F	n = 10	p	t/X^2
Demographics	Sex	13M, 17F	9M, 17F	5M, 8F	-	-	5M, 5F	-	-
	Height (m)	1.70 ± 0.10	1.67 ± 0.10	1.69 ± 0.11	0.611	$F_{2,66} = 0.50$	1.69 ± 0.08	0.906	$t_{21} = -0.12$
	Age (y)	59.7 ± 7.8	60.3 ± 7.0	61.9 ± 6.9	-	-	63.5 ± 3.4	0.227	$t_{21} = 1.24$
	Mass (kg)	96.3 ± 18.7	96.8 ± 20.3	100.5 ± 20.4	0.124	$F_{2,13} = 2.46$	71.5 ± 13.5	**0.001**	$\mathbf{t_{21} = -3.72}$
	BMI (kg/m^2)	33.7 ± 5.1	34.0 ± 5.7	35.5 ± 6.4	**0.043^{bc}**	$\mathbf{F_{2,12} = 4.04}$	24.8 ± 2.6	**<0.001**	$\mathbf{t_{21} = -4.56}$
	Walking Speed (m/s)	0.97 ± 0.26	1.11 ± 0.22	1.22 ± 0.13	**0.001^{ab}**	$\mathbf{F_{2,24} = 9.34}$	1.12 ± 0.18	0.145	$t_{21} = -1.51$
	Normalized Stride Length (m/m)	1.42 ± 0.25	1.53 ± 0.24	1.57 ± 0.22	0.056	$F_{2,12} = 3.36$	1.51 ± 0.16	0.248	$t_{21} = -1.19$
Mods	Number of Modules	2.5 ± 0.6	2.5 ± 0.6	2.5 ± 0.5	0.844	$F_{2,18} = 0.17$	2.7 ± 0.5	0.673	$X^2_1 = 0.178$
	Proper Module Organization	6 of 30	11 of 26	6 of 13	0.143	$F_{2,19} = 2.15$	-	-	-
Function Metrics	SCT (s)	25.7 ± 14.6	19.5 ± 8.8*	18.3 ± 6.8	**0.049^{ab}**	$\mathbf{F_{2,16} = 3.65}$	-	-	-
	TUG (s)	11.5 ± 4.4	10.0 ± 1.9	8.8 ± 1.8	**0.016^{ab}**	$\mathbf{F_{2,15} = 5.46}$	-	-	-
	6MW (m)	401.1 ± 106.9	457.9 ± 89.0	495.5 ± 78.2	**0.001^{abc}**	$\mathbf{F_{2,17} = 11.28}$	-	-	-
	KOOS-Pain (pts)	48.7 ± 20.4	73.7 ± 22.2*	79.6 ± 15.5	**<0.001^{ab}**	$\mathbf{F_{2,14} = 21.99}$	-	-	-
	KOOS-Symptoms (pts)	46.4 ± 21.2	63.6 ± 18.7*	72.3 ± 15.9	**0.006^{ab}**	$\mathbf{F_{2,12} = 7.91}$	-	-	-
	KOOS-ADL (pts)	55.8 ± 20.8	78.2 ± 21.4*	80.2 ± 14.1	**<0.001^{ab}**	$\mathbf{F_{2,14} = 18.61}$	-	-	-
	KOOS-QOL (pts)	24.7 ± 21.0	54.3 ± 22.2*	64.1 ± 24.2	**<0.001^{ab}**	$\mathbf{F_{2,10} = 27.51}$	-	-	-

Symbols indicate differences between: [a]Pre-TKA and 6-months Post-TKA, [b]Pre-TKA and 24-months Post-TKA, [c]6- and 24- months Post-TKA
*clinically meaningful improvement in function compared to the previous time point; M = male; F = female
Average demographics, module metrics, and function metrics grouped by time point and population (± 1 standard deviation). The healthy control group was compared to the group of participants tested at 24-months post-TKA.

Of the 10 healthy individuals, 3 demonstrated 2 modules and 7 demonstrated 3 modules (2.7 ± 0.5 modules; Table 1). Those with 2 modules had 1 module dominated by the plantarflexors and 1 module dominated by the quadriceps and hamstrings. Those with 3 modules had modules

dominated by the plantarflexors, quadriceps, and hamstrings, separately. Before surgery, 16

individuals with TKA demonstrated 2 modules and 14 demonstrated 3 modules, and 6 of the 30

(20.0%) participants had proper organization. By 6-months after surgery, 12 demonstrated 2

modules and 14 demonstrated 3 modules, and 11 of the 26 (42.3%) participants had proper module

organization. By 24-months after surgery, 5 participants demonstrated 2 modules and 8

demonstrated 3 modules, and 6 of the 13 (46.2%) participants had proper organization. There was

no difference in number of modules ($p = 0.844$) or module organization ($p = 0.143$) between time

points (Table 1).

There were differences in mean tVAF values between time points for solutions with 1, 2,

6, and 7 modules (Fig. 2). For the 1-module solution, the group tested before surgery ($79.8 \pm 6.3\%$)

had a statistically higher mean tVAF value than those at 6-months ($75.7 \pm 6.3\%$; pairwise $p =$

0.001) and 24-months ($72.9 \pm 8.2\%$; pairwise $p = 0.003$) after surgery. For the 2-module solution,

the group tested before surgery ($91.2 \pm 2.6\%$) had a statistically higher mean tVAF than those at

24-months after surgery ($89.3 \pm 2.7\%$; pairwise $p = 0.033$). For the 6-module solution, the group

tested before surgery ($99.3 \pm 0.3\%$) had a statistically higher mean tVAF than those at 24-months

after surgery ($99.1 \pm 0.3\%$; pairwise $p = 0.016$). For the 7-module solution, the group tested before

surgery ($99.7 \pm 0.1\%$) had a statistically higher mean tVAF than those at 6-months ($99.7 \pm 0.1\%$;

pairwise $p = 0.040$) and 24-months ($99.6 \pm 0.2\%$; pairwise $p = 0.004$) after surgery.

Fig. 2: tVAF for All Module Solutions. Estimated mean tVAF by time point for number of modules varying from 1 to 8. Error bars represent +1 standard deviation. * indicates statistically significant differences between time points.

The worst averaged functional scores were observed before surgery and the best averaged functional scores were observed 24-months after surgery (Table 1). All measures were statistically different between timepoints (all $p \leq 0.049$). From before to 6-months after surgery, all measures had statistically significant improvements (all pairwise $p \leq 0.031$), and these improvements were also clinically meaningful for the SCT and all KOOS subscales. From before to 24-months after surgery, all measures had statistically significant (all pairwise $p \leq 0.020$) and clinically meaningful improvements. Average 6MW distance also had a statistically significantly improvement from 6- to 24-months after surgery (pairwise $p = 0.036$), but this improvement was not clinically meaningful.

Hypothesis I: Function and Number of Modules

A higher number of modules was associated with better performance in a few performance-

based and patient-reported functional measures before and 6-months after surgery but with poorer

performance in a few functional measures 24-months after surgery (Figs. 3 & 4). Before surgery,

participants with 3 modules demonstrated better scores on average than those with 2 modules in

all measures, and this difference was clinically meaningful in all measures except the 6MW

distance (Table 2). These results were statistically significant for the TUG ($p = 0.002$) and KOOS-

Pain ($p = 0.005$) scores and approached significance for the SCT ($p = 0.060$), KOOS-ADL ($p = $

0.057), and KOOS-QOL ($p = 0.065$) scores. By 6-months after surgery, participants with 3

modules demonstrated better scores on average than those with 2 modules in all measures except

the KOOS-QOL subscale (Table 2). These differences were clinically meaningful for only the

SCT, which also approached statistical significance ($p = 0.052$) and were statistically significant

but not clinically meaningful for the TUG test ($p = 0.028$). By 24-months after surgery, participants

with 2 modules had better scores on average than those with 3 modules in SCT, 6MW, KOOS-

Symptoms scores (Table 2). These differences were clinically meaningful for the SCT and for the

6MW, which was also statistically significant ($p = 0.006$).

Fig. 3: Performance-Based Function. Performance-based functional scores (SCT, TUG, 6MW) for each participant at each time point. Participants who were present for consecutive testing time points are connected. Filled symbols indicate participants with proper module organization. Thick black lines indicate the group estimated means and 95% confidence intervals at each time point, determined by the GLMM.

*/† = statistically significant (*) and clinically meaningful (†) differences in function between time points.

@/# = statistically significant (@) and clinically meaningful (#) differences in function between groups demonstrating the same number of modules at each time point.

& = clinically meaningful differences in function between groups demonstrating proper or poor module organization at each time point. There were no statistically significant differences.

Fig. 4: Patient-Reported Function. KOOS survey subscale scores (Pain, Symptoms, ADL, QOL) for each participant at each time point. Participants who were present for consecutive testing time points are connected. Filled symbols indicate participants with proper module organization. Thick black lines indicate the group estimated means and 95% confidence intervals at each time point, determined by the GLMM.

*/† = statistically significant (*) and clinically meaningful (†) differences in function between time points.

@/# = statistically significant (@) and clinically meaningful (#) differences in function between groups demonstrating the same number of modules at each time point.

& = clinically meaningful differences in function between groups demonstrating proper or poor module organization at each time point. There were no statistically significant differences.

Table 2: Functional Measures within Module Number Groups.

		Two Modules	Three Modules	p	Estimate	95% CI
Number of Subjects	Pre-TKA	n = 16	n = 14	-	-	-
	6-months Post-TKA	n = 12	n = 14	-	-	-
	24-months Post-TKA	n = 5	n = 8	-	-	-
SCT (s)	Pre-TKA	30.2 ± 15.3	21.0 ± 11.2	0.060[†]	9.2	[-0.42, 18.90]
	6-months Post-TKA	22.5 ± 11.5	16.4 ± 4.2	0.052[†]	6.1	[-0.05, 12.19]
	24-months Post-TKA	16.1 ± 3.6	19.5 ± 7.6	0.434[†]	-3.4	[-12.44, 5.70]
TUG (s)	Pre-TKA	13.1 ± 5.3	9.6 ± 2.3	**0.002[†]**	**3.5**	**[1.63, 5.35]**
	6-months Post-TKA	10.8 ± 2.5	9.4 ± 1.4	**0.028**	**1.4**	**[0.17, 2.60]**
	24-months Post-TKA	9.2 ± 1.9	8.9 ± 1.6	0.803	0.3	[-2.51, 3.16]
6MW (m)	Pre-TKA	379.7 ± 123.8	425.9 ± 76.8	0.128	-46.7	[-107.79, 14.49]
	6-months Post-TKA	453.8 ± 116.3	467.9 ± 53.7	0.621	-13.1	[-67.74, 41.44]
	24-months Post-TKA	546.1 ± 55.7	468.4 ± 48.4	**0.006[†]**	**78.5**	**[29.27, 127.72]**
KOOS-Pain (points)	Pre-TKA	39.2 ± 16.6	58.8 ± 19.8	**0.005[†]**	**-19.6**	**[-32.58, -6.57]**
	6-months Post-TKA	69.8 ± 24.6	78.3 ± 12.4	0.186	-8.6	[-21.82, 4.67]
	24-months Post-TKA	80.6 ± 13.5	80.7 ± 17.0	0.987	-0.1	[-19.98, 19.68]
KOOS-Symptom (points)	Pre-TKA	40.3 ± 18.8	52.4 ± 22.5	0.152[†]	-12.1	[-28.94, 4.72]
	6-months Post-TKA	60.5 ± 19.8	67.1 ± 17.3	0.393	-6.6	[-22.92, 9.71]
	24-months Post-TKA	79.0 ± 13.8	69.2 ± 16.5	0.418	9.7	[-15.79, 35.26]
KOOS-ADL (points)	Pre-TKA	49.1 ± 17.9	63.0 ± 22.3	0.057[†]	-13.9	[-28.17, 0.46]
	6-months Post-TKA	76.4 ± 25.6	80.0 ± 9.5	0.482	-3.6	[-15.35, 8.08]
	24-months Post-TKA	77.8 ± 12.0	83.2 ± 15.9	0.520	-5.4	[-23.85, 12.98]
KOOS-QOL (points)	Pre-TKA	17.7 ± 15.0	32.3 ± 24.1	0.065[†]	-14.7	[-30.49, 1.16]
	6-months Post-TKA	57.2 ± 20.0	53.1 ± 20.1	0.515	4.1	[-9.36, 17.55]
	24-months Post-TKA	61.7 ± 28.4	64.8 ± 22.5	0.832	-3.1	[-34.98, 28.69]

Average functional measures within groups of participants demonstrating the same number of modules at each timepoint (estimated mean ± 1 standard deviation). [†] indicates a clinically meaningful difference in function between the Two Modules and Three Modules groups.

Hypothesis II: Function and Module Organization

Better module organization was not statistically associated with better performance-based function (SCT, TUG, or 6MW) or patient-reported function (KOOS subscales) before or after surgery (Figs. 3 & 4). However, the largest differences in function between the proper and poor organization groups occurred 24-months after surgery, with the proper organization group demonstrating better function on average in all performance-based and patient-reported measures (Table 3). These results approached statistical significance for the KOOS-Pain subscale ($p = 0.061$) and were clinically meaningful for the SCT, KOOS-Pain, and KOOS-Symptoms scores.

437
438 **Table 3: Functional Measures within Module Organization Groups.**

		Poor	Proper	p-Value	Estimate	95% CI
Number of Subjects	Pre-TKA	n = 24	n = 6	-	-	-
	6-months Post-TKA	n = 15	n = 11	-	-	-
	24-months Post-TKA	n = 7	n = 6	-	-	-
SCT (s)	Pre-TKA	26.1 ± 14.2	24.2 ± 17.3	0.750[†]	2.0	[-10.64, 14.60]
	6-months Post-TKA	19.9 ± 10.7	19.1 ± 5.4	0.824	0.7	[-6.39, 7.88]
	24-months Post-TKA	19.6 ± 9.3	16.2 ± 2.6	0.487[†]	3.4	[-7.06, 13.93]
TUG (s)	Pre-TKA	11.3 ± 4.8	12.0 ± 2.5	0.695	-0.6	[-4.04, 2.78]
	6-months Post-TKA	10.2 ± 1.9	9.8 ± 2.0	0.679	0.3	[-1.25, 1.88]
	24-months Post-TKA	8.5 ± 2.3	9.2 ± 0.8	0.540	-0.7	[-3.11, 1.72]
6MW (m)	Pre-TKA	401.6 ± 110.3	399.9 ± 97.4	0.963	1.7	[-75.03, 78.46]
	6-months Post-TKA	454.4 ± 96.2	462.2 ± 79.6	0.762	-7.7	[-60.33, 44.85]
	24-months Post-TKA	489.3 ± 89.8	502.0 ± 69.3	0.727	-12.7	[-92.51, 67.11]
KOOS-Pain (points)	Pre-TKA	49.5 ± 19.2	46.4 ± 26.8	0.729	3.2	[-15.41, 21.74]
	6-months Post-TKA	77.3 ± 19.1	68.1 ± 25.7	0.116	9.2	[-2.75, 21.19]
	24-months Post-TKA	75.2 ± 14.6	86.1 ± 12.8	0.061[†]	-11.0	[-22.53, 0.60]
KOOS-Symptom (points)	Pre-TKA	46.8 ± 19.6	45.7 ± 29.1	0.920	1.1	[-20.41, 22.54]
	6-months Post-TKA	63.4 ± 15.7	63.4 ± 22.5	0.999	0.0	[-18.11, 18.09]
	24-months Post-TKA	67.3 ± 17.3	79.1 ± 13.8	0.315[†]	-11.8	[-36.93, 13.40]
KOOS-ADL (points)	Pre-TKA	56.1 ± 20.4	55.9 ± 24.2	0.984	0.2	[-18.51, 18.88]
	6-months Post-TKA	79.2 ± 17.1	76.6 ± 26.8	0.641	2.6	[-10.72, 15.97]
	24-months Post-TKA	78.7 ± 14.5	83.9 ± 10.6	0.478	-5.2	[-21.13, 10.70]
KOOS-QOL (points)	Pre-TKA	24.4 ± 17.7	27.5 ± 33.2	0.756	-3.0	[-23.10, 17.01]
	6-months Post-TKA	56.8 ± 22.4	50.5 ± 20.8	0.355	6.3	[-9.43, 21.97]
	24-months Post-TKA	63.2 ± 26.3	67.7 ± 21.3	0.759	-4.5	[-36.12, 27.19]

439
440 Average functional measures within groups of participants with modules that were (Proper) or
441 were not (Poor) organized like healthy controls at each timepoint (estimated mean ± 1 standard
442 deviation). [†] indicates a clinically meaningful difference in function between the Proper and Poor
443 organization groups.
444

Secondary Analysis: Changes in Modules

While there were no significant differences in modules between participants grouped by

time point, we observed changes in module number and organization in individual participants

throughout the time course of recovery (Table 4). From before to 6-months after surgery, 63.2%

of participants present at both time points (12/19) demonstrated changes in module number and/or

organization. From 6- to 24-months after surgery, 80.0% of participants present at both time points

(8/10) demonstrated changes in module number and/or organization. From before to 24-months

after surgery, 100% of participants present at both time points (10/10) demonstrated changes in

module number and/or the organization of at least 1 module (Table 4).

Table 4: Count of Participants with Changes in Modules.

		Pre-TKA to 6-m Post-TKA n = 19	6-m to 24-m Post-TKA n = 10	Pre-TKA to 24-m Post-TKA n = 10
Number of Modules	Increase	3 (15.8%)	1 (10.0%)	4 (40.0%)
	No Change	12 (63.2%)	7 (70.0%)	4 (40.0%)
	Decrease	4 (21.1%)	2 (20.0%)	2 (20.0%)
Proper vs. Poor Organization	Improved	5 (26.3%)	2 (20.0%)	4 (40.0%)
	No Change	12 (63.2%)	6 (60.0%)	5 (50.0%)
	Worsened	2 (10.5%)	2 (20.0%)	1 (10.0%)
Any Module Characteristic	Change	12 (63.2%)	8 (80.0%)	10 (100.0%)
	No Change	7 (36.8%)	2 (20.0%)	0 (0.0%)

457
Number of participants (%) that were present at multiple time points and demonstrated changes
in module characteristics. Proper organization describes participants whose organization matched
those of healthy controls for all modules. Poor organization describes participants with at least 1
module that was not organized like healthy controls. Participants who are listed as having a
change in "Any Module Characteristic" had a change in the number of modules and/or the
organization of at least 1 module compared to healthy controls.
464

DISCUSSION

In an effort to identify neuromuscular control strategy as a factor related to functional

deficits in individuals after TKA, the purpose of this study was to determine the relationship

between modular control strategy and function in individuals before and at two time points after

TKA. Our findings partially confirmed our first hypothesis that better function would be related to

a higher number of modules. We found that participants with better performance-based function

before and 6-months after surgery and better patient-reported function before surgery had a higher

number of modules. However, at 24-months after surgery, a higher number of modules were

related to worse 6MW performance, which contradicts our first hypothesis. Our findings did not

strongly confirm our second hypothesis that better function would be related to module

organization that more closely resembled that of healthy controls. However, by 24-months after

surgery, participants with proper module organization had better function on average than those

with poor organization in all performance-based and patient-reported measures, with some

differences being clinically meaningful. Considerable participant drop-out by the 24-month time

point may have prevented us from detecting statistically significant differences in function

between groups demonstrating proper and poor module organization. While we did not find

differences in population-averaged modular control between timepoints, we did observe changes

in both module number and organization in individuals and in population-mean complexity, as

measured by tVAF. While the differences in tVAF between testing time points were statistically

significant for the 6- and 7-module solutions, they were likely not clinically meaningful, though a

clinically meaningful difference in tVAF values has not been established in the literature. Larger

differences in tVAF between time points were observed for the 1- and 2-module solutions, which

may indicate that module complexity is malleable in this population. Together with our

observations on module organization, these results indicate that motor modules are plastic and

suggest that neural control strategy is influenced by surgical intervention and rehabilitation in this

population.

The association we observed between number of modules and function partially confirms

the findings of Ardestani et al. (21), who found that a higher number of modules was associated

with better patient-reported function 1-year following TKA. While we found that better function

was associated with a higher number of modules in individuals with KOA prior to TKA, we did

not find this association by 24-months after surgery. Several methodological differences exist

between the current study and Ardestani et al. (21) which may explain the differences in results.

Ardestani et al. (21) measured patient-reported function using the Knee Functional Survey (KFS)

(40), which assesses general patient satisfaction and symptoms, ability to complete activities of

daily living, and ability to participate in sports and exercise. The KFS may most closely relate to

the KOOS-Symptoms, KOOS-ADL, and KOOS-QOL subscales, which were not statistically

related to number of modules in the current study. However, our small sample sizes, particularly

by 24-months after surgery, may have prevented us from detecting associations between patient-

reported function and number of modules after surgery. Additionally, the methods for module

extraction used by Ardestani et al. (21), including a strict error criterion of 90% accuracy between

the experimental and reconstructed EMG, resulted in a range of 2-5 modules, whereas we saw only

2-3 modules. We believe our approach for calculating modules using 500 bootstrapped samples

(41) and a previously developed, conservative VAF criteria (24) is robust. Lastly, participants in

Ardestani et al. (21) received cruciate-retaining implants while participants in the current study

received posterior-stabilizing implants. The comparison of our results with those of Ardestani et

al. (21) may suggest that proprioceptive inputs, which are lost as a result of sacrificing the cruciate

ligaments, influence neuromuscular control strategy. However, the lack of consistency in

methodologies across studies warrants further investigation into the influence of retaining the

posterior cruciate ligament on post-surgical neuromuscular control. Further, while Ardestani et al.

(21) indicated an association between a higher number of modules and better patient-reported

function at 1-year following TKA, our results indicate that the relationship between number of

modules and function changes over the time course of recovery.

The observed relationship between a higher number of modules and better function in some

measures before TKA reversed in some measures by 24-months after TKA. The group with 2

modules 24-months after surgery demonstrated 6MW distances that were on average better,

statistically and clinically, than those with 3 modules. Further analysis revealed that there were 2

participants with 2 modules that had significantly higher 6MW distances than the other participants

at this time point which drove this result. Both participants demonstrated module organization that

was different from the healthy controls, with one module that was dominated by the quadriceps

and hamstrings and one module that was dominated by the plantarflexors and hamstrings such that

the hamstrings were constantly active throughout the gait cycle. These participants also

demonstrated better-than-average SCT and TUG performance at 24-months but had unremarkable

KOOS scores and demographics measures. When examining all participants at 24-months post-

TKA, we found that 3 of the 5 participants (60.0%) with 2 modules had proper organization while

only 3 of the 8 participants (37.5%) with 3 modules had proper organization. These results indicate

that module organization, rather than number, may be a factor which is related to function in the

long-term post-operative time frame (24-months after TKA).

Though there were no statistically significant differences in function between the proper

and poor organization groups at any time point, we observed trends toward better function in all

functional measures, except TUG, for the group with proper module organization at 24-months

after TKA. While the number of modules has frequently been used as a metric for motor control,

recent studies suggest that module analysis which examines the composition of modules across

multiple trials may provide further insight into functional performance in tasks like walking (22).

Furthermore, relationships between module organization and function have been observed in

populations with neurologic movement disorders. Brough et al. (42) found that individuals with

post-stroke hemiparesis without an independent plantarflexors module demonstrated poorer

locomotor function compared to patients and controls with an independent plantarflexors module.

Hayes et al. (25) also observed altered module composition and slower walking speeds in

individuals with chronic incomplete spinal cord injury compared to able-bodied individuals. While

there was not a significant relationship in the current study between function and module

organization, as it relates to the organization of healthy individuals, the observed trends toward

better function in those with module organization similar to healthy controls, with several

measures presenting with clinically meaningful differences between groups, indicates that there

may be a relationship between module composition and function which should be explored further.

Investigation of the number and organization of modules over time revealed that

neuromuscular control was plastic in individual participants, and this neuroplasticity may be

related to functional outcomes. Previously, Hubley-Kozey et al. (15) found that shifts in activation

patterns of some muscles from before surgery to 1-year after TKA were accompanied by

improvements in knee flexor and extensor strength and patient-reported outcomes (via WOMAC

Index (43)) but were still statistically different from the patterns of asymptomatic individuals. In

the present study, we observed statistically significant and clinically meaningful improvements in

all functional measures from before to after surgery which accompanied population-mean

decreases in several tVAF measures. While small sample sizes (n < 10 present at all time points)

prevented us from using statistical analysis to assess the association between changes in function

and changes in module characteristics between time points, these results suggest that there may be

an underlying relationship between improvements in function and increased module complexity.

Further, when comparing function at 6- to 24-months, there was a larger improvement in all

functional measures for groups with proper organization than those with poor organization, with

the exception of TUG (Table 3). This observation may indicate that individuals with proper module

organization may have a greater capacity to improve function than those with poor organization.

Though we were not able to characterize it in the present study, we believe these observations

suggest that there is a relationship between changes in function and changes in modular control

which should be investigated in future studies.

Our evidence that the neuromuscular control strategies adopted before surgery are not

permanent suggests that surgical intervention and rehabilitation have an influence on modular

control strategy. These results contrast Shuman et al. (44) who found no change in the distribution

of muscle weightings within modules after treatment in children with cerebral palsy. However,

while the present study focuses on an orthopaedic condition rather than a neurological disorder,

our results support the concepts presented by Ting et al. (26), who suggested that neuromuscular

control patterns can be influenced by rehabilitation in individuals with spinal cord injury, stroke,

and Parkinson's disease. Our results also support the findings of Roelker et al. (38) who found

differences in motor modules between individuals with KOA and healthy, age-matched controls,

indicating that the onset of KOA may alter neuromuscular control strategy. It remains unclear

whether the changes in neural control observed in the present study resulted from changes in

neuroanatomical structures (i.e. muscles, motor units, or even spinal/supraspinal structures) or are

products of the neuromechanical interactions (26) caused by changes in the mechanical

environment of the knee during surgery. While we did not collect detailed data on the rehabilitation

plans of the participants in the present pilot study, our results suggest that changes in

neuromuscular control strategies, explored through motor modules, occur throughout the treatment

and rehabilitation of an orthopaedic condition.

The plasticity of control in this population and the association between modular control

and patient function at individual time points may provide a basis for modifying rehabilitation

programs to improve functional outcomes after TKA. The association between better function and

a higher number of modules before and 6-months after surgery and trends toward better average

function in those with proper organization 24-months after surgery suggests that the relationships

between function and module number or organization may be uncoupled. Our results before and

6-months after TKA support the findings of others, who have suggested that higher modular

complexity indicates a higher number of independent motor strategies that can be used to complete

a task (21, 45), thus improving adaptability and function. Our work suggests that improvements in

function in TKA patients may initially present as increases in the number of modules recruited

during gait and that once individuals are capable of producing complex and discrete neuromuscular

control patterns, further functional improvements may present as improvements in module

organization. Further investigation is needed to determine the cause of the observed changes in

modular control. The findings of this future work will help to determine if such motor control

changes may be driven through rehabilitation programs in order to improve patient function.

This pilot study was based on a secondary analysis of EMG data collected in a separate

study, and as such, small sample sizes may have prevented us from observing greater associations

between modular control characteristics and function. The high BMI values, which are

representative of the general TKA population, resulted in increased motion artifact during EMG

collection and also limited the number of available trials for analysis. Further investigation is

necessary to further characterize the relationship between changes in modular control

characteristics and changes in function at the population level. A larger study may provide the

statistical power needed to detect potential associations between modular complexity and

functional outcomes.

CONCLUSIONS

Some performance-based and patient-reported functional measures were related to modular complexity and organization. The number and organization of modules in individuals changed between the three testing time points in our study. We believe this new approach to evaluating motor control in individuals with TKA provides valuable insight into the plasticity of neural control strategies in patients and may help inform rehabilitation programs for patients demonstrating unsatisfactory outcomes. Future work will build on the findings of this pilot study and explore the influence of rehabilitation techniques, which target re-training neuromuscular control, on functional outcomes.

ACKNOWLEDGEMENTS

The authors thank Dr. Jacqueline M. Lewis, Dr. Gregory M. Freisinger, Dr. Elena J. Caruthers, Rachel K. Hall, and Elizabeth M. Leszcz for their contributions to data collection and processing. We also thank Drs. Jeffrey F. Granger, Andrew H. Glassman, and Matthew D. Beal, who performed the surgeries in this study, and Drs. Jason Payne and Alan Rogers, who performed the radiographic grading of OA severity.

GRANTS

This research was supported by the National Science Foundation Graduate Research Fellowship Program under Grant Nos. DGE-1343012 (SAR) and DGE-0822215 (EJC), a fellowship from The Ohio State University (RRK), and by Grant Number R01AR056700 from the National Institute of Arthritis and Musculoskeletal and Skin Diseases. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institute of Arthritis

and Musculoskeletal and Skin Diseases nor the National Institutes of Health.

CONFLICT OF INTEREST STATEMENT

None of the authors have conflicts of interest regarding the contents of this manuscript.

REFERENCES

1.   Sloan M, Premkumar A, Sheth NP. Projected Volume of Primary Total Joint
Arthroplasty in the U.S., 2014 to 2030. J Bone Joint Surg Am. 2018;100(17):1455-60.
2.   Singh JA, O'Byrne M, Harmsen S, Lewallen D. Predictors of moderate-severe functional
limitation after primary Total Knee Arthroplasty (TKA): 4701 TKAs at 2-years and 2935 TKAs
at 5-years. Osteoarthritis and cartilage. 2010;18(4):515-21.
3.   Berman AT, Zarro VJ, Bosacco SJ, Israelite C. Quantitative gait analysis after unilateral
or bilateral total knee replacement. J Bone Joint Surg Am. 1987;69(9):1340-5.
4.   Benedetti MG, Catani F, Bilotta TW, Marcacci M, Mariani E, Giannini S. Muscle
activation pattern and gait biomechanics after total knee replacement. Clinical biomechanics
(Bristol, Avon). 2003;18(9):871-6.
5.   Wilson SA, McCann PD, Gotlin RS, Ramakrishnan HK, Wootten ME, Insall JN.
Comprehensive gait analysis in posterior-stabilized knee arthroplasty. J Arthroplasty.
1996;11(4):359-67.
6.   Su FC, Lai KA, Hong WH. Rising from chair after total knee arthroplasty. Clin Biomech
(Bristol, Avon). 1998;13(3):176-81.
7.   Yercan HS, Ait Si Selmi T, Sugun TS, Neyret P. Tibiofemoral instability in primary total
knee replacement: a review, Part 1: Basic principles and classification. The Knee.
2005;12(4):257-66.
8.   Stulberg SD, Loan P, Sarin V. Computer-assisted navigation in total knee replacement:
results of an initial experience in thirty-five patients. J Bone Joint Surg Am. 2002;84-A Suppl
2:90-8.
9.   Jeffery RS, Morris RW, Denham RA. Coronal alignment after total knee replacement.
The Journal of bone and joint surgery British volume. 1991;73(5):709-14.
10.   Bade MJ, Kohrt WM, Stevens-Lapsley JE. Outcomes before and after total knee
arthroplasty compared to healthy adults. The Journal of orthopaedic and sports physical therapy.
2010;40(9):559-67.
11.   Zeni JA, Jr., Snyder-Mackler L. Early postoperative measures predict 1- and 2-year
outcomes after unilateral total knee arthroplasty: importance of contralateral limb strength.
Physical therapy. 2010;90(1):43-54.
12.   Smith AJ, Lloyd DG, Wood DJ. Pre-surgery knee joint loading patterns during walking
predict the presence and severity of anterior knee pain after total knee arthroplasty. Journal of

orthopaedic research : official publication of the Orthopaedic Research Society. 2004;22(2):260-
6.
13.   Davidson BS, Judd DL, Thomas AC, Mizner RL, Eckhoff DG, Stevens-Lapsley JE.
Muscle activation and coactivation during five-time-sit-to-stand movement in patients
undergoing total knee arthroplasty. J Electromyogr Kinesiol. 2013;23(6):1485-93.
14.   Marmon AR, Snyder-Mackler L. Activation deficits do not limit quadriceps strength
training gains in patients after total knee arthroplasty. Int J Sports Phys Ther. 2014;9(3):329-37.
15.   Hubley-Kozey CL, Hatfield GL, Astephen Wilson JL, Dunbar MJ. Alterations in
neuromuscular patterns between pre and one-year post-total knee arthroplasty. Clin Biomech
(Bristol, Avon). 2010;25(10):995-1002.
16.   Ting LH, McKay JL. Neuromechanics of muscle synergies for posture and movement.
Curr Opin Neurobiol. 2007;17(6):622-8.
17.   Lee WA. Neuromotor synergies as a basis for coordinated intentional action. J Mot
Behav. 1984;16(2):135-70.
18.   Mussa-Ivaldi FA, Giszter SF, Bizzi E. Linear combinations of primitives in vertebrate
motor control. Proc Natl Acad Sci U S A. 1994;91(16):7534-8.
19.   Olree KS, Vaughan CL. Fundamental patterns of bilateral muscle activity in human
locomotion. Biol Cybern. 1995;73(5):409-14.
20.   Ivanenko YP, Poppele RE, Lacquaniti F. Five basic muscle activation patterns account
for muscle activity during human locomotion. J Physiol. 2004;556(Pt 1):267-82.
21.   Ardestani MM, Malloy P, Nam D, Rosenberg AG, Wimmer MA. TKA patients with
unsatisfying knee function show changes in neuromotor synergy pattern but not joint
biomechanics. Journal of electromyography and kinesiology : official journal of the International
Society of Electrophysiological Kinesiology. 2017;37:90-100.
22.   Allen JL, McKay JL, Sawers A, Hackney ME, Ting LH. Increased neuromuscular
consistency in gait and balance after partnered, dance-based rehabilitation in Parkinson's disease.
J Neurophysiol. 2017;118(1):363-73.
23.   Allen JL, Kesar TM, Ting LH. Motor module generalization across balance and walking
is impaired after stroke. J Neurophysiol. 2019;122(1):277-89.
24.   Clark DJ, Ting LH, Zajac FE, Neptune RR, Kautz SA. Merging of healthy motor
modules predicts reduced locomotor performance and muscle coordination complexity post-
stroke. J Neurophysiol. 2010;103(2):844-57.
25.   Hayes HB, Chvatal SA, French MA, Ting LH, Trumbower RD. Neuromuscular
constraints on muscle coordination during overground walking in persons with chronic
incomplete spinal cord injury. Clin Neurophysiol. 2014;125(10):2024-35.
26.   Ting LH, Chiel HJ, Trumbower RD, Allen JL, McKay JL, Hackney ME, et al.
Neuromechanical principles underlying movement modularity and their implications for
rehabilitation. Neuron. 2015;86(1):38-54.
27.   Freisinger GM, Hutter EE, Lewis J, Granger JF, Glassman AH, Beal MD, et al.
Relationships between varus-valgus laxity of the severely osteoarthritic knee and gait, instability,
clinical performance, and function. J Orthop Res. 2017;35(8):1644-52.
28.   Chaudhari AMW, Schmitt LC, Freisinger GM, Lewis JM, Hutter EE, Pan X, et al.
Perceived Instability Is Associated With Strength and Pain, Not Frontal Knee Laxity, in Patients
With Advanced Knee Osteoarthritis. J Orthop Sports Phys Ther. 2019;49(7):513-7.
29.   Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthrosis. Ann Rheum Dis.
1957;16(4):494-502.

30.   Rejeski WJ, Ettinger WH, Jr., Schumaker S, James P, Burns R, Elam JT. Assessing
performance-related disability in patients with knee osteoarthritis. Osteoarthritis Cartilage.
1995;3(3):157-67.
31.   Dobson F, Hinman RS, Roos EM, Abbott JH, Stratford P, Davis AM, et al. OARSI
recommended performance-based tests to assess physical function in people diagnosed with hip
or knee osteoarthritis. Osteoarthritis Cartilage. 2013;21(8):1042-52.
32.   Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and
change of four performance measures: a longitudinal study evaluating outcome following total
hip and knee arthroplasty. BMC Musculoskelet Disord. 2005;6:3.
33.   Roos EM, Lohmander LS. The Knee injury and Osteoarthritis Outcome Score (KOOS):
from joint injury to osteoarthritis. Health Qual Life Outcomes. 2003;1:64.
34.   Kieliba P, Tropea P, Pirondini E, Coscia M, Micera S, Artoni F. How are Muscle
Synergies Affected by Electromyography Pre-Processing? IEEE Trans Neural Syst Rehabil Eng.
2018;26(4):882-93.
35.   Santuz A, Ekizos A, Janshen L, Baltzopoulos V, Arampatzis A. On the Methodological
Implications of Extracting Muscle Synergies from Human Locomotion. Int J Neural Syst.
2017;27(5):1750007.
36.   Torres-Oviedo G, Ting LH. Muscle synergies characterizing human postural responses. J
Neurophysiol. 2007;98(4):2144-56.
37.   Lee DD, Seung HS. Learning the parts of objects by non-negative matrix factorization.
Nature. 1999;401(6755):788-91.
38.   Roelker SA, Koehn RR, Caruthers EJ, Schmitt LC, Chaudhari AMW, Siston RA. Effects
of Age and Knee Osteoarthritis on the Modular Control of Walking: A Pilot Study. medRxiv.
2021:2020.05.22.20110536.
39.   Almeida GJ, Schroeder CA, Gil AB, Fitzgerald GK, Piva SR. Interrater reliability and
validity of the stair ascend/descend test in subjects with total knee arthroplasty. Arch Phys Med
Rehabil. 2010;91(6):932-8.
40.   Weiss JM, Noble PC, Conditt MA, Kohl HW, Roberts S, Cook KF, et al. What functional
activities are important to patients with knee replacements? Clin Orthop Relat Res.
2002(404):172-88.
41.   Sawers A, Allen JL, Ting LH. Long-term training modifies the modular structure and
organization of walking balance control. J Neurophysiol. 2015;114(6):3359-73.
42.   Brough LG, Kautz SA, Bowden MG, Gregory CM, Neptune RR. Merged plantarflexor
muscle activity is predictive of poor walking performance in post-stroke hemiparetic subjects. J
Biomech. 2019;82:361-7.
43.   Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of
WOMAC: a health status instrument for measuring clinically important patient relevant
outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. J
Rheumatol. 1988;15(12):1833-40.
44.   Shuman BR, Goudriaan M, Desloovere K, Schwartz MH, Steele KM. Muscle synergies
demonstrate only minimal changes after treatment in cerebral palsy. J Neuroeng Rehabil.
2019;16(1):46.
45.   Safavynia SA, Torres-Oviedo G, Ting LH. Muscle Synergies: Implications for Clinical
Evaluation and Rehabilitation of Movement. Top Spinal Cord Inj Rehabil. 2011;17(1):16-24.
761