1   **Title: Association of preference and frequency of teleworking with work functioning**
2   **impairment: a nationwide cross-sectional study of Japanese full-time employees**
3

4   Satoshi Yamashita[1,2], M.D., Tomohiro Ishimaru[1], M.D., M.P.H., Ph.D., Tomohisa Nagata[2],
5   M.D., Ph.D., Seiichiro Tateishi[3], M.D., Ph.D., Ayako Hino[4], M.D., Ph.D., Mayumi Tsuji[5],
6   M.D., Ph.D., Kazunori Ikegami[6], M.D., Ph.D., Keiji Muramatsu[7], M.D., Ph.D., and Yoshihisa
7   Fujino[1], M.D., M.P.H., Ph.D., for the CORoNaWork Project
8

[1] Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences, University of
Occupational and Environmental Health, Japan
[2] Department of Occupational Health Practice and Management, Institute of Industrial Ecological Sciences,
University of Occupational and Environmental Health, Japan
[3] Department of Occupational Medicine, School of Medicine, University of Occupational and
Environmental Health, Japan
[4] Department of Mental Health, Institute of Industrial Ecological Sciences, University of Occupational and
Environmental Health, Japan
[5] Department of Environmental Health, School of Medicine, University of Occupational and Environmental
Health, Japan
[6] Department of Work Systems and Health, Institute of Industrial Ecological Sciences, University of
Occupational and Environmental Health, Japan
[7] Department of Preventive Medicine and Community Health, School of Medicine, University of
Occupational and Environmental Health, Japan

Abstract

Objective: We examined whether teleworking preference and frequency were associated with

work functioning impairment.

Methods: This online cross-sectional study was conducted using a self-administered

questionnaire among 27,036 full-time Japanese workers. The Work Functioning Impairment

Scale was used to measure work functioning impairment, and we performed multilevel

logistic regression analysis.

Results: Higher odds ratios for work functioning impairment were observed among

employees who preferred to telework compared with those who preferred working in the

workplace. A similar trend was observed among employees who teleworked 4 or more days a

week compared with those who almost never teleworked. When teleworking preference and

frequency were adjusted, only teleworking preference was associated with work functioning

impairment.

Conclusions: A preference for teleworking was associated with work functioning impairment

and one factor that increased the teleworking frequency.

Keywords: COVID-19; Japan; occupational health; presenteeism; remote working; telework;

work functioning impairment

Introduction

Teleworking has increased among companies worldwide during the COVID-19 pandemic to prevent contact among workers and limit the spread of infection. [1,2] Telework involves working at a location other than a central office or workplace, such as at home. Individuals work separately from other employees during working time and use information and communications technology to interact with coworkers, as needed.[3] A recent survey in Japan revealed that 26.7% of employees were teleworking as of March 2020, the beginning of the COVID-19 pandemic, and this percentage increased to 52.7% as of May 2020 during the first the state of emergency declaration by the Japanese government.[4] A previous survey in Japan showed that the percentage of teleworkers peaked in May 2020 at more than 50%; however, after the first state of emergency declaration period, the percentage of teleworkers had decreased to approximately 30% by June 2020.[5] To not only prevent infection but also maintain business continuity, some workers combine teleworking with going to their office or workplace.[6]

Teleworking is considered to have both positive and negative effects on workers' health. Teleworking is characterized by the absence of commuting time and flexibility regarding working location and time. However, teleworking also involves a lack of supervision by superiors, sitting and working at home for long periods, and little communication with colleagues.[3,7–9] The positive effects of teleworking on workers include

the reduction in commuting time, which allows for more sleep, fewer interruptions from

colleagues at work, and greater flexibility in how employees use their time to improve their

work–life balance.[3,7,8,10] However, the negative effects of teleworking on workers include

musculoskeletal disorders owing to an inappropriate work environment, excessive weight

gain because of physical inactivity, increased risk of cardiovascular disease and diabetes, and

adverse effects on mental health owing to limited opportunities to interact with colleagues and

superiors.[3,9] Teleworking affects the ease of work and performance through its impact on

workers' health.

The negative effects of teleworking may cause work functioning impairment. Work

functioning impairment is loss of the ability to perform a work task as a result of presenteeism,

which is defined as "reduced performance at work, besides illness."[11,12] A teleworking system

not only permits employees to work flexibly but it also enables them to work, even when they

have a health problem, such as fatigue or depression.[8,13] Previous studies have shown that

teleworking increases sedentary behavior and that sedentary behavior leads to

presenteeism.[8,14] The introduction of teleworking may promote presenteeism and working

with some degree of work functioning impairment.

It is not known how teleworking affects work functioning impairment. In particular,

the rapid introduction of teleworking during the early part of the COVID-19 pandemic may

have created a mismatch between workers' preferences and teleworking practices. This rapid

introduction of teleworking differed from teleworking before the COVID-19 pandemic

because workers could not choose to work remotely at will. The option of teleworking was

previously selected according to workers' preferences; however, teleworking during the

COVID-19 pandemic was imposed owing to safety and social requirements, regardless of

workers' preferences. Imposing shift work on workers who do not prefer shift work is

associated with distress and burnout.[15] However, the impact of a preference for teleworking

among workers is unclear. In this study, we examined how teleworking preferences and

frequency are associated with work functioning impairment.

Methods

Study design and participants

This was a cross-sectional study using online questionnaire survey data from the

Collaborative Online Research on Novel-coronavirus and Work study (CORoNaWork study).

The questionnaires were sent by e-mail to target populations who were preregistered with

Cross Marketing Inc. (Tokyo, Japan) and monitors who responded so as to participate in the

survey. The data collection period was from December 22 to 26, 2020. To recruit participants,

605,381 people were emailed a request to participate in the survey; 55,045 people were

enrolled as monitors, and 33,087 people matched the survey criteria. The survey was

completed by 33,087 Japanese full-time workers between the ages of 20 to 65 years. After

excluding 6,051 surveys with invalid responses, 27,036 people were included in the analysis

(Figure 1). Invalid responses were those from surveys completed in less than 6 minutes;

participants with weight less than 30 kg or height less than 140 cm; inconsistent responses;

and wrong answers to the following question: Select the third largest number out of the

following five numbers (231, 245, 323, 252, 312, and the correct answer is 252). Stratified

sampling was conducted by region (five regions according to the regional COVID-19

incidence level), sex (male/female), and job type (office worker/non-office worker). National

statistics were used to obtain the regional COVID-19 incidence level during the study

period.[16] We classified this into five levels according to the cumulative incidence of

COVID-19 in each prefecture. Thus, respondents were categorized into 20 units sampled by

region, sex, and job type. Each unit had an equal number of 1650 respondents. Recruitment

continued until the required number was reached in each unit. Details of the sample selection

have been described elsewhere.[17] This study was conducted with the approval of the Ethics

Committee of the University of Occupational and Environmental Health, Japan (approval no.

R2-079). Informed consent was obtained via a form on the survey website.

Outcome

The Work Functioning Impairment Scale (WFun) was used as a measure of work

functioning impairment. The WFun is a self-administered measure of work functioning

impairment validated and developed according to the Consensus-based Standards for the

Selection of Health Measurement Instruments and based on the Rasch model.[18] The

instrument comprises seven items, with five response options. The total of the seven items is

the overall score, which ranges from 7 to 35 points, with higher scores indicating higher work

functioning impairment. We used the 6-item version of the WFun. This simplified version is

also based on the Rasch model and is converted to the same score as in the original

measurement instrument by multiplying by 1.17 (7–35 points).[19,20] We defined work

functioning impairment as a WFun score of 21 or higher, in line with previous studies.[21]

Explanatory variables

Teleworking frequency and teleworking preferences were included as explanatory variables.

Participants responded to the following question about teleworking preferences: "How do you

feel about teleworking?" with the following options: a) I would like to telework as much as

possible; b) I prefer teleworking; c) I don't mind either; d) I prefer regular work; and e) I

prefer regular work as much as possible. Participants responded to the following question

regarding teleworking frequency: "Do you work from home?" with the following response

options: a) 4 or more days a week; b) 2 or 3 days a week; c) 1 day a week; d) more than once

a month and less than once a week; and e) almost never.

Covariates

On the basis of relevant studies, the following socioeconomic and lifestyle factors

were selected as covariates: sex,[22,23] age,[22–24] marital status[22] (married, divorced/widowed,

and unmarried), annual equivalent household income,[24] education[23] (junior high school, high

school, vocational school, junior college, university, and graduate school), smoking,[22] alcohol

consumption,[22] job type[14,23,25] (mainly desk work, mainly involving interpersonal

communication, mainly physical work), number of employees,[4] and COVID-19 incidence rate

in the month prior to the survey.[4]

Statistical analysis

A WFun score of 21 was used as the cutoff value. Scores above 21 were dichotomized

as work functioning impairment and scores below 21 as no work functioning impairment. For

teleworking frequency, "once a week" was included in "more than once a month and less than

once a week" owing to the small number of respondents. Multilevel logistic regression

analysis was performed to assess the relationship of teleworking preference and frequency

with WFun score. The hierarchical structure nested in the prefecture of residence according to

the cumulative incidence of COVID-19 in each prefecture was adjusted to account for the

effects of intraclass correlation. The associations between teleworking and work functioning

impairment were evaluated using three models: an age- and sex-adjusted model, Model 1, and

Model 2. In Model 1, we adjusted for sex, age, marital status, annual equivalent household

income, education, smoking, alcohol consumption, job type, number of employees, and

COVID-19 incidence rate. Model 2 was adjusted for teleworking preferences and teleworking

frequency, in addition to those factors adjusted in Model 1. All statistical analyses were

conducted using Stata/SE 16.1 (StataCorp LLC, College Station, TX, USA). A p-value less

than 0.05 was considered statistically significant.

Results

The characteristics of respondents are shown in Table 1. Of 27,036 respondents,

approximately half were men (51.1%) and office workers (49.8%). Approximately

three-quarters (72.9%) had a technical college, junior college, university, or graduate degree.

Most respondents (78.7%) reported nearly no teleworking; 10.3% of respondents reported

very frequent (4 or more days a week) teleworking. Respondents who reported nearly no

teleworking had the lowest percentage (22.1%) of high WFun scores. There were no missing

data because the survey system was designed to ensure that all questions were answered.

Table 2 shows the association of work functioning impairment with teleworking

preference and frequency. In the age- and sex-adjusted model, strong preference for

teleworking was associated with work functioning impairment (odds ratio [OR]: 2.08, 95%

confidence interval [CI]: 1.90–2.28). The same trend was observed for Model 1, which was

adjusted for socioeconomic and lifestyle factors (OR: 2.20, 95% CI: 2.01–2.42) and Model 2,

which was adjusted for socioeconomic and lifestyle factors and teleworking frequency (OR:

2.27, 95% CI: 2.05–2.51), with participants who prefer teleworking as much as possible

associated with work functioning impairment. In the age- and sex-adjusted model, the OR of

work functioning impairment was significantly higher among participants who teleworked 4

or more days a week than workers who almost never teleworked (OR: 1.18, 95% CI:

1.07–1.3). In Model 1, adjusted for socioeconomic and lifestyle factors, the OR of work

functioning impairment was significantly higher among employees who teleworked 4 or more

days a week (OR: 1.32, 95% CI: 1.18–1.46) and those who teleworked 2 or 3 days a week

(OR: 1.28, 95% CI: 1.12–1.46) compared with those who almost never teleworked. In Model

2, adjusted for teleworking preference in addition to socioeconomic and lifestyle factors, the

ORs of work functioning impairment were no longer significantly different among workers

who teleworked 4 or more days a week (OR: 0.90, 95% CI: 0.80–1.01) and those who

teleworked 2 or 3 days a week (OR: 1.01, 95% CI: 0.88–1.16).

Discussion

In this study, we found an association between teleworking frequency and increased

work functioning impairment. Additionally, workers who preferred teleworking experienced

increased work functioning impairment. When teleworking preference and frequency were

adjusted for, only teleworking preference was associated with work functioning impairment.

A remarkable finding of our study was that workers who preferred to telework had

greater work functioning impairment. The preference for teleworking could be based on

difficulties with working in the workplace owing to ill health or work–family conflicts.

Previous studies have reported that workers who prefer telework choose teleworking when

feeling unwell because this allows them to work at their own pace and freely adjust their

break time.[26,27] Other previous studies state that female employees with children choose to

telework but are not highly satisfied with telework.[28,29] Attention to the background related to

teleworking preferences is important to support teleworkers with severe work functioning

impairment.

The current study showed that frequent teleworking was associated with work

functioning impairment. One important mechanism for this association could be that

individuals with health problems are more likely to choose to telework. [27,30] In this study,

when teleworking preference and teleworking frequency were adjusted, a teleworking

preference was associated with work functioning impairment whereas the teleworking

frequency was no longer associated with work functioning impairment. Some studies have

suggested that teleworking is associated with poor health conditions, such as depression and

musculoskeletal pain.[31,32] Key to detecting signs of work functioning impairment would be to

follow up on the health conditions of employees who frequently telework.

This study has several public health implications. First, governments should create a

system that enables teleworkers to continue working with reduced work function. Early

detection of work functioning impairment is more difficult in telework than in the workplace

where there is face-to-face interaction. Strategies to check work function are necessary, even

for teleworkers. For example, a checklist to verify the work function of teleworkers should be

developed and promoted for regular use. Second, companies should pay attention to

background factors that reduce the work function of teleworkers. Identifying needed support

for teleworkers will prevent impaired work function. Third, companies should assess the risk

of absence from work for individual workers and provide appropriate work accommodations

in cooperation with occupational health staff. Occupational health staff should have online

meetings with teleworkers to check for any health problems and provide information on

health management when teleworking.

Our study has several limitations: First, the self-reported data used in our study may

be inaccurate. Self-reporting may lead to systematic errors, which may reduce or increase the

significance of the differences. Second, this was a cross-sectional study, and it is not possible

to determine causality. For example, we cannot determine whether there is a causal

relationship between teleworking frequency and teleworking preferences.

In conclusion, the current study findings suggest that a teleworking preference is

associated with work functioning impairment and one factor that increases the teleworking

frequency. This result may be explained by workers who have difficulties with working in the

workplace owing to ill health or work–family conflicts prefer to telework. Employees who

telework frequently should be carefully monitored to determine whether the ability to perform

their job is impaired. Investigating the background regarding the choice to telework could

provide clues to understanding the support needed by teleworkers, which will help to prevent

health problems and lead to increased productivity among these workers.

References

1. Belzunegui-Eraso A, Erro-Garcés A. Teleworking in the context of the Covid-19 crisis.
Sustain Sci Pract Policy. 2020;12(9):3662.

2. Bick A, Blandin A, Mertens K. Work from Home after the Covid-19 Outbreak. Published
online July 1, 2020. Accessed February 9, 2021.
https://papers.ssrn.com/abstract=3650114

3. Allen TD, Golden TD, Shockley KM. How Effective Is Telecommuting? Assessing the
Status of Our Scientific Findings. *Psychol Sci Public Interest*. 2015;16(2):40-68.

4. Sasaki N, Imamura K, Kataoka M, et al. COVID-19 measurements at the workplace in
various industries and company sizes: a 2-month follow-up cohort study of full-time
employees in Japan. *Environmental and Occupational Health Practice*. 2021;3(1).
doi:10.1539/eohp.2020-0017-OA

5. Goto M, Hamano W. Trends in surveys about home teleworking during the COVID-19
pandemic in Japan. *J Inst Nucl Safe Syst*. 2020;27:252-274.

6. Shao Y, Fang Y, Wang M, Chang CHD, Wang L. Making daily decisions to work from
home or to work in the office: The impacts of daily work- and COVID-related stressors
on next-day work location. *J Appl Psychol*. 2021;106(6):825-838.

7. Ishimaru T, Fujino Y. Association between work style and presenteeism in the Japanese
service sector. *J Occup Health*. 2021;63(1):e12211.

8. Barone Gibbs B, Kline CE, Huber KA, Paley JL, Perera S. COVID-19 shelter-at-home and work, lifestyle and well-being in desk workers. *Occup Med* . Published online February 18, 2021. doi:10.1093/occmed/kqab011

9. van der Lippe T, Lippényi Z. Co-workers working from home and individual and team performance. *New Technol Work Employ*. 2020;35(1):60-79.

10. Bailey DE, Kurland NB. A review of telework research: Findings, new directions, and lessons for the study of modern work. *J Organ Behav*. Published online 2002. https://onlinelibrary.wiley.com/doi/abs/10.1002/job.144?casa_token=8bJsEQHv2msAAA AA:q3feFnhk7EQdyHkhJDwuDv5_ZmLEHZUVwkaifKaPKewkQymf7xdWBA40oaPhi qAKwSbtZnduMggWzcY

11. Fujino Y, Shazuki S, Izumi H, et al. Prospective Cohort Study of Work Functioning Impairment and Subsequent Absenteeism Among Japanese Workers. *J Occup Environ Med*. 2016;58(7):e264-7.

12. Lerner DJ, Amick BC 3rd, Malspeis S, Rogers WH. A national survey of health-related work limitations among employed persons in the United States. *Disabil Rehabil*. 2000;22(5):225-232.

13. Chong S, Huang Y, Chang CHD. Supporting interdependent telework employees: A moderated-mediation model linking daily COVID-19 task setbacks to next-day work withdrawal. *J Appl Psychol*. 2020;105(12):1408-1422.

14. Brown HE, Ryde GC, Gilson ND, Burton NW, Brown WJ. Objectively measured sedentary behavior and physical activity in office employees: relationships with presenteeism. *J Occup Environ Med*. 2013;55(8):945-953.

15. Hulsegge G, van Mechelen W, Proper KI, Paagman H, Anema JR. Shift work, and burnout and distress among 7798 blue-collar workers. *Int Arch Occup Environ Health*. 2020;93(8):955-963.

16. Ministry of Health Labour and Welfare. About the situation by which prefectures is medical offer organizations (the situation of the load and the infection by which it's for a medical offer organization). Accessed January 6, 2022. https://www.mhlw.go.jp/stf/seisakunitsuite/newpage_00035.html?SLANG=ja&TLANG= en&XMODE=0&XCHARSET=utf-8&XJSID=0.

17. Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a nationwide Internet-based health survey in workers during the COVID-19 pandemic in 2020. *medRxiv*. Published online 2021. https://www.medrxiv.org/content/10.1101/2021.02.02.21249309v1.abstract

18. Fujino Y, Uehara M, Izumi H, et al. Development and validity of a work functioning
impairment scale based on the Rasch model among Japanese workers. *J Occup Health*.
2015;57(6):521-531.

19. Igarashi Y, Tateishi S, Harada A, et al. Relationship Between Support for Workers
with Illness and Work Functioning Impairment in Japan During the COVID-19 Pandemic.
J Occup Environ Med. Published online January 25, 2022.
doi:10.1097/JOM.0000000000002500

20. Okawara M, Ishimaru T, Tateishi S, et al. Association Between the Physical Work
Environment and Work Functioning Impairment While Working From Home Under the
COVID-19 Pandemic in Japanese Workers. *J Occup Environ Med*.
2021;63(9):e565-e570.

21. Nagata T, Fujino Y, Saito K, et al. Diagnostic Accuracy of the Work Functioning
Impairment Scale (WFun): A Method to Detect Workers Who Have Health Problems
Affecting their Work and to Evaluate Fitness for Work. *J Occup Environ Med*.
2017;59(6):557-562.

22. Yu J, Wang S, Yu X. Health risk factors associated with presenteeism in a Chinese
enterprise. *Occup Med* . 2015;65(9):732-738.

23. Cho YS, Park JB, Lee KJ, Min KB, Baek CI. The association between Korean
workers' presenteeism and psychosocial factors within workplaces. *Ann Occup Environ
Med*. 2016;28(1):41.

24. Navarro A, Salas-Nicás S, Moncada S, Llorens C, Molinero-Ruiz E. Prevalence,
associated factors and reasons for sickness presenteeism: a cross-sectional nationally
representative study of salaried workers in Spain, 2016. *BMJ Open*. 2018;8(7):e021212.

25. Hendriksen IJM, Bernaards CM, Steijn WMP, Hildebrandt VH. Longitudinal
Relationship Between Sitting Time on a Working Day and Vitality, Work Performance,
Presenteeism, and Sickness Absence. *J Occup Environ Med*. 2016;58(8):784-789.

26. Ruhle SA, Schmoll R. COVID-19, Telecommuting, and (Virtual) Sickness
Presenteeism: Working From Home While Ill During a Pandemic. *Front Psychol*.
2021;12:734106.

27. Vargas-Llave O, Mandl I, Weber T, Wilkens M. Telework and ICT-based mobile
work: Flexible working in the digital age. *Luxembourg: Publications Office of the
European Union*. Published online 2020. https://www.voced.edu.au/content/ngv:86087

28.   Sousa-Uva M, Sousa-Uva A, E Sampayo MM, Serranheira F. Telework during the
COVID-19 epidemic in Portugal and determinants of job satisfaction: a cross-sectional
study. *BMC Public Health*. 2021;21(1):2217.

29.   Zhang S, Moeckel R, Moreno AT, Shuai B, Gao J. A work-life conflict perspective on
telework. *Transp Res Part A: Policy Pract*. 2020;141:51-68.

30.   Steidelmüller C, Meyer SC, Müller G. Home-Based Telework and Presenteeism
Across Europe. *J Occup Environ Med*. 2020;62(12):998-1005.

31.   Radulović AH, Žaja R, Milošević M, Radulović B, Luketić I, Božić T. Work from
home and musculoskeletal pain in telecommunications workers during COVID-19
pandemic: a pilot study. *Arh Hig Rada Toksikol*. 2021;72(3):232-239.

32.   Henke RM, Benevent R, Schulte P, Rinehart C, Crighton KA, Corcoran M. The
Effects of Telecommuting Intensity on Employee Health. *Am J Health Promot*.
2016;30(8):604-612.

Figure legends

Figure 1. Flow diagram of the study population

```
Internet panel
(n= 605,581)                    ──────▶  No response
                                         (n= 550,336)

Enrolled as monitors             ──────▶  Did not match the survey criteria
(n= 55,045)                               (n= 21,978)

Complete responses               ──────▶  Invalid responses (n= 6,051)
(n= 33,067)                               A) Surveys completed in less than 6
                                             minutes
                                          B) Weight less than 30 kg
                                          C) Height less than 140 cm
Valid responses                           D) Inconsistent responses
(n= 27,016)                               E) Wrong answer to a screening
                                             question
```

Table 1. Participant characteristics

	4 or more days a week (n=2,790)		2 or 3 days a week (n=1,477)		More than once a month and less than once a week (n=1,493)		Almost never (n=21,276)	
	n	%	n	%	n	%	n	%
Age, mean (standard deviation)	49.6	(10.0)	48.7	(10.3)	47.6	(10.4)	46.5	(10.6)
Sex, male	1,576	56.5	860	58.2	925	62.0	10,453	49.1
Marital status								
Married	1,406	50.4	915	61.9	959	64.2	11,749	55.2
Divorced/Widowed	284	10.2	114	7.7	106	7.1	2,339	11.0
Unmarried	1,100	39.4	448	30.3	428	28.7	7,188	33.8
Job type								
Mainly desk work	2,042	73.2	1,058	71.6	952	63.8	9,416	44.3
Mainly involving interpersonal communication	495	17.7	333	22.5	391	26.2	5,708	26.8
Mainly physical work	253	9.1	86	5.8	150	10.0	6,152	28.9
Annual equivalent household income (JPY)								
< 2,500,000	751	26.9	207	14.0	189	12.7	4,563	21.4
2,500,000–3,750,000	643	23.0	297	20.1	305	20.4	6,305	29.6
3,750,000–5,000,000	576	20.6	362	24.5	371	24.8	5,316	25.0
> 5,000,000	820	29.4	611	41.4	628	42.1	5,092	23.9
Education								
Junior high school	31	1.1	13	0.9	6	0.4	318	1.5
High school	507	18.2	198	13.4	238	15.9	6,010	28.2
Vocational school, junior college, university, or graduate school	2,252	80.7	1,266	85.7	1,249	83.7	14,948	70.3
Current smoker	693	24.8	406	27.5	441	29.5	5,464	25.7
Alcohol consumption								
6 or more days a week	658	23.6	360	24.4	341	22.8	4,315	20.3
4 or 5 days a week	234	8.4	165	11.2	163	10.9	1,515	7.1
2 or 3 days a week	316	11.3	195	13.2	241	16.1	2,514	11.8
Within 1 day a week	445	15.9	262	17.7	281	18.8	3,559	16.7
Almost never	1,137	40.8	495	33.5	467	31.3	9,373	44.1
Number of employees								
1–9	1,670	59.9	343	23.2	275	18.4	3,877	18.2
10–99	228	8.2	230	15.6	245	16.4	6,237	29.3
100–999	321	11.5	335	22.7	375	25.1	6,122	28.8
>1000	571	20.5	569	38.5	598	40.1	5,040	23.7
Work functioning impairment scale (Wfun) score								
7–20	2,163	77.5	1,134	76.8	1,154	77.3	16,794	78.9
21–35	627	22.5	343	23.2	339	22.7	4,482	21.1
Teleworking preference								
I would like to telework as much as possible	1,789	64.1	463	31.3	259	17.3	2,196	10.3
I prefer teleworking	409	14.7	440	29.8	376	25.2	2,666	12.5
I don't mind either	436	15.6	376	25.5	483	32.4	6,637	31.2
I prefer to work in the workplace	98	3.5	154	10.4	272	18.2	2,845	13.4
I prefer to work in the workplace as much as possible	58	2.1	44	3.0	103	6.9	6,932	32.6

Table 2. Odds ratio of work functioning impairment associated with teleworking preference and frequency

	Age- and sex-adjusted			Model 1			Model 2		
	OR	95% CI		*p*	OR	95% CI		*p*	OR	95% CI		*p*
Teleworking preference												
I would like to telework as much as possible	2.08	1.90	2.28	<0.001	2.20	2.01	2.42	<0.001	2.27	2.05	2.51	<0.001
I prefer teleworking	1.74	1.58	1.92	<0.001	1.78	1.62	1.97	<0.001	1.79	1.62	1.98	<0.001
I don't mind either	1.24	1.14	1.35	<0.001	1.25	1.14	1.36	<0.001	1.25	1.14	1.36	<0.001
I prefer to work in the workplace	1.21	1.09	1.35	<0.001	1.22	1.09	1.36	<0.001	1.22	1.09	1.36	<0.001
I prefer to work in the workplace as much as possible	reference				reference				reference			
Teleworking frequency												
4 or more days a week	1.18	1.07	1.30	0.001	1.32	1.18	1.46	<0.001	0.90	0.80	1.01	0.068
2 or 3 days a week	1.21	1.06	1.37	0.004	1.28	1.12	1.46	<0.001	1.01	0.88	1.16	0.863
More than once a month and less than once a week	1.13	1.00	1.28	0.057	1.18	1.04	1.35	0.011	1.04	0.91	1.19	0.545
Almost never	reference				reference				reference			

OR: odds ratio, CI: confidence interval.

Model 1: adjusted for sex, age, marital status, equivalent household income, education, smoking, alcohol consumption, job type, number of employees, and COVID-19 incidence rate in the month before the survey.

Model 2: adjusted for teleworking preference and teleworking frequency in addition to the factors adjusted in Model 1.