Title: Changing molecular epidemiology and characteristics of methicillin-resistant

Staphylococcus aureus isolated from bloodstream infections: nationwide surveillance in Japan in

2019

Running title: Surveillance of MRSA isolated from BSI

Authors: Norihito KAKU[1,2,3*], Daisuke SASAKI[2], Kenji OTA[1,2], Taiga MIYAZAKI[4], Katsunori

YANAGIHARA[1,2]

Affiliations**:**

[1]Department of Laboratory Medicine, Nagasaki University Graduate School of Biomedical

Sciences, Nagasaki City, Nagasaki, Japan

[2]Department of Laboratory Medicine, Nagasaki University Hospital, Nagasaki City, Nagasaki,

Japan

[3]Division of Pulmonary and Critical Care Medicine, University of Michigan, Ann Arbor,

Michigan, United States

[4]Division of Respirology, Rheumatology, Infectious Diseases, and Neurology, Department of

Internal Medicine, Faculty of Medicine, University of Miyazaki, Miyazaki City, Miyazaki, Japan

[*]**Corresponding Author:** Norihito Kaku

1-7-1 Sakamoto, Nagasaki City, Nagasaki, Japan 852-8501

Email: norihitk@gmail.com

Tel: +81-95-819-7574 Fax: +81-95-819-7422

Synopsis

Objectives: Some single-centre studies have reported that MRSA carrying the staphylococcal

cassette chromosome mec (SCC*mec*) type IV has been increasing in bloodstream infections

(BSIs) in Japan. Therefore, we conducted nationwide surveillance for MRSA BSI to investigate

the extent of such change across Japan.

Methods: We recruited 51 Japanese hospitals from the Japanese Association for Infectious

Diseases. MRSA isolates detected in two or more sets of blood cultures were collected between

January and September 2019 and subjected to antimicrobial susceptibility testing. Whole-

genome sequencing was also performed to determine SCC*mec* typing and multilocus typing and

detect drug-resistance and virulence genes.

Results: Two hundred seventy MRSA isolates were collected from 45 hospitals. The major

combination types were ST8 with SCC*mec* type IV (ST8-IV) (30.7%), ST1-IV (29.6%),

ST2725-IV (9.5%), ST764-II (8.1%), and ST5-II (7.8%). However, there were regional

differences among the most major types. The most common types in western, eastern, and

northern Japan were ST1-IV, ST8-IV, and ST5-II, respectively. ST8-IV, ST1-IV, and ST2725-

IV exhibited greater susceptibility to clindamycin and minocycline than ST764-II and ST5-II,

but *erm*(A) was detected in 93.8% and 100% of ST1-IV and ST2725-IV, respectively. Based on

drug-resistance and virulence genes, characteristics of ST8-IV were different from those of ST1-

IV and ST2725-IV. In addition, there were two major ST8-IV types with different characteristics.

Conclusions: This study revealed that SCC*mec* type IV replaced SCC*mec* type II in MRSA BSI.

In addition, SCC*mec* type IV was divided into several types with different characteristics.

Introduction

MRSA is an important drug-resistant pathogen in bloodstream infections (BSIs) because *S. aureus,* including MRSA, is the second-most frequently encountered pathogen in both community- and hospital-onset BSI.[1] In the United States, the percentage of patients with BSIs involving MRSA increased from 22% in 1995 to 57% in 2001,[2] but the incidence of MRSA BSIs has been decreasing since the middle of the 2000s through various infection control efforts.[3] However, a new threat involves the spread of community-associated MRSA (CA-MRSA), represented by the USA300 clone. CA-MRSA strains carrying staphylococcal cassette chromosome *mec* (SCC*mec*) types IV or V have different characteristics than healthcare-associated MRSA (HA-MRSA) strains carrying SCC*mec* types I, II, or III. CA-MRSA strains are more susceptible to antimicrobial agents such as fluoroquinolones, macrolides, lincosamides, and aminoglycosides; moreover, they sometimes produce Panton-Valentine leucocidin (PVL), which causes skin and soft-tissue infections and necrotising pneumonia.[4] In 2009, a mathematical model predicted that CA-MRSA would become the dominant MRSA strain in hospitals and healthcare facilities,[5] and in fact, there were some reports that CA-MRSA has replaced HA-MRSA.[6–8] In addition, the clonal diversity of CA-MRSA is increasing, and it is no longer possible to identify CA-MRSA only by SCC*mec* type IV or production of PVL.[9]

In Japan, sequence type (ST) 5 carrying SCC*mec* type II, represented by the New York/Japan clone, has been the most common clone in MRSA BSIs.[10] However, the proportion of SCC*mec* type II has been decreasing, while that of SCC*mec* type IV has been increasing in MRSA BSIs.[11,12] On the other hand, most SCC*mec* type IV strains in Japan lack PVL.[11–13] However, since these data were obtained from a single institution, the molecular epidemiology and characteristics of MRSA isolated from BSIs in Japan are unclear. In this study, we collected

MRSA isolates from patients with BSIs throughout Japan in 2019 and performed antimicrobial

susceptibility testing and whole-genome sequencing to reveal the characteristics of these MRSA

isolates.

Materials and Methods

Study design

We recruited participating medical institutions through a mailing list obtained from the

Japanese Association for Infectious Diseases in December 2018; 51 Japanese hospitals were

recruited. MRSA isolates detected in two or more blood samples obtained at the same time were

collected at each hospital between January 22 and September 30, 2019. Only the first isolate per

patient was included in this study. No MRSA isolates met the criteria in the six hospitals.

Eventually, 274 isolates collected from 45 hospitals were analysed at Nagasaki University

Hospital. Four isolates were excluded from the analysis because they were identified as

methicillin-susceptible *Staphylococcus aureus* at Nagasaki University Hospital (Supplementary

Figure 1A). Finally, the remaining 270 isolates were analysed (Supplementary Figure 1B). They

were stored in Microbank (Iwaki & Co., Ltd., Tokyo, Japan) at -80°C and transferred to

Nagasaki University Hospital. The following data regarding MRSA isolates were collected from

the participating medical institutions: blood culture collection location (outpatient or inpatient)

and day following hospitalization on which blood cultures were collected.

Antimicrobial susceptibility testing

The MICs for ampicillin, oxacillin, cefoxitin, cefazolin, imipenem, meropenem, levofloxacin,

erythromycin, clindamycin, minocycline, trimethoprim/sulfamethoxazole, vancomycin,

teicoplanin, linezolid, daptomycin, and arbekacin were determined by broth microdilution testing

using Dry Plate Eiken (Eiken, Tokyo, Japan) according to the manufacturer's instructions. MIC_{50}

and MIC_{90} values were calculated as previously reported.[14] Antimicrobial susceptibility was

measured according to CLSI guidelines (M100-Ed31) and EUCAST v.11.0.[15,16]

WGS and molecular characteristics

All procedures were performed according to the manufacturers' instructions. DNA was

extracted from MRSA isolates using a Quick-DNA Fungal/Bacterial Kit (Zymo Research, Irvine,

CA, USA). DNA libraries for sequencing were generated using the Invitrogen Collibri ES DNA

Library Prep Kit for Illumina (A38607096, ThermoFisher Scientific, Waltham, MA, USA), and

sequencing was performed on a MiSeq system (Illumina, San Diego, CA, USA) using MiSeq

Reagent kit v. 3 (600 cycles) (Illumina). Genome assemblies with de Bruijn graphs, MLST, and

detection of antimicrobial resistance genes in ResFinder were performed using the CLC

Genomics Workbench and Microbial Genomics Modules (QIAGEN N. V., Venlo, Netherlands).

SCC*mec* types were determined using SCC*mec*Finder v.1.2

(https://cge.cbs.dtu.dk/services/SCCmecFinder/).[17] The subtypes of SCC*mec* type IV were also

detected using SCC*mec*Finder v.1.2. If two or more subtypes were detected, it is classified as

non-subtypeable. Virulence genes were detected using VirulenceFinder v.2.0.3

(https://cge.cbs.dtu.dk/services/VirulenceFinder/, software v.2.0.3, database v.2020-05-29).[18,19]

Core genome MLST (cg MLST) and staphylococcal protein A (spa) typing were performed

using Ridom SeqSphere+ ver.8.3.1(Ridom GmbH, Münster, Germany) and genome assemblies

for cgMLST and spa typing were performed with SKESA ver.2.3.0.[20] A Minimum Spanning

Tree (MST) was created based on MLST, cgMLST, and *S. aureus* Accessory using Ridom

114 SeqSphere+. Samples with more than 10% missing values of the items for distance calculation

115 were excluded in MST.

117 *Statistical analysis*

118  All statistical analyses were performed using R software (v. 4.0.3; R Foundation for Statistical

119 Computing, Vienna, Austria). Fisher's exact test with Bonferroni correction was used to compare

120 categorical variables. The statistical significance level was set at $P < 0.05$. P values are listed in

121 Table S1.

123 *Ethics*

124  The Ethics Committee approved this study of Nagasaki University Hospital (19012123).

125 MRSA isolates were anonymized and individually numbered when they were isolated from

126 blood cultures. All data and samples were fully anonymized before being sent to Nagasaki

127 University Hospital. The Ethics Committee of Nagasaki University Hospital waived

128 requirements for informed consent.

130 *Data availability*

131 Raw data were generated at Nagasaki University Hospital. The derived data supporting the

132 findings of this study are presented in this paper. WGS data reported in this study are available in

133 the DDBJ Sequence Read Archive (https://www.ddbj.nig.ac.jp/dra/index-e.html) under accession

134 number DRA013058.

Results

Sequence and SCCmec typing

Among 270 BSI isolates of MRSA, the percentages of SCC*mec* types I, II, IV, V, and IX were 1.9%, 18.5%, 77.4%, 1.9%, and 0.4%, respectively (Table S2). The subtypes of SCC*mec* type IV were IVa (118, 56.5%), IVc (11, 5.3%), IVg (4, 1.9%), IVh (4, 1.9%), and non-subtypeable (72, 34.4%). The major combination types in Japan can be seen in Figure 1A. However, there were regional differences among major types. The most common types in western Japan, eastern Japan, and northern Japan were ST1-IV, ST8-IV, ST5-II, and ST764-II, respectively (Figure 1B). The characteristics of the major types are listed in Table 1. The percentages of hospital-acquired BSIs detected from samples obtained 48 h after hospitalization, involving ST1-IV, ST2725-IV, ST8-IV, ST5-II, and ST764 were 62.5%, 61.5%, 60.2%, 71.4%, 72.7%, respectively. There were no significant differences in patient backgrounds among the types (Table S1). The most detected subtype of SCC*mec* IV in ST8-IV, ST1-IV, and ST2725-IV were non-typeable (80.7%), IVa (100%), and IVa (92.3%) (Table 1).

Spa type and cgMLST

The most common spa type was t784 for ST1-IV and ST2725-IV, and t002 for ST764-II and ST5-II (Table 1). There were two main spa types in ST8-IV: t1767 and t5071. In cgMLST, 203 isolates were classified into 191 complex types. There were nine complex types containing two or three isolates, and isolates of the same complex type were all detected in the same hospital. The number of cgMLST types containing two or more isolates in ST8-IV, ST1-IV, ST2725-IV, ST764-II, and ST5-II was 2, 2, 3, 1, and 1, respectively. MST for all isolates indicated that ST1 and ST2725 were in the same cluster (Figure 2).

Antimicrobial susceptibility

Resistance rates calculated using CLSI/EUCAST criteria are shown in Table 1. The resistance rates for anti-MRSA agents were very low in all types. However, for some antimicrobial agents, there were significant differences among the types. The drug resistance rate for levofloxacin in both CLSI and EUCAST criteria in ST8-IV was significantly lower than that of other types ($p < 0.001$). The resistance rates for clindamycin for ST8-IV, ST1-IV, and ST2725-IV were significantly lower than ST5-II and ST764-II ($p < 0.05$). In addition, the drug resistance rate for clindamycin in ST1-IV and ST2725-IV was significantly lower than that for ST8-IV ($p < 0.001$). The resistance rates for minocycline in EUCAST criteria in ST8-IV, ST1-IV, and ST2725-IV were significantly lower than those of ST5-II and ST764-II ($p < 0.05$).

There were differences among the major types in the MICs for β-lactams (Table 2). Both MIC_{50} and MIC_{90} for cefazoline, imipenem, and meropenem were lower in ST8-IV, ST1-IV, and ST2725-IV than in ST5-II and ST764-II. These differences were particularly pronounced for imipenem. The MIC_{90} for imipenem was ≤ 0.25, ≤ 0.25, and 0.5, respectively, in ST8-IV, ST1-IV, and ST2725-IV, respectively, whereas it was ≥ 32 in both ST5-II and ST764-II.

Drug-resistance genes

Table 3 shows the positive rate for drug-resistance genes in each type. Almost all major types carried aminoglycoside-resistance genes. However, specific aminoglycoside-resistance genes differed among the major types; for example, the positive rate for *ant(9)-Ia* for ST8-IV was significantly lower than for other types ($p < 0.001$). The positive rate for the macrolide-resistance gene *erm*(A) for ST8-IV was also significantly lower than for other types ($p < 0.001$). In ST764-

II, the positive rate for *blaZ* was significantly lower than for other types (p <0.001), whereas that

for the fosfomycin-resistance gene was significantly higher than for other types (p <0.05). The

positive rate for the tetracycline-resistance gene was significantly higher for ST5-II and ST764-II

than for ST8-IV, ST1-IV, and ST2725-IV (p < 0.05).

Virulence genes

Table 4 shows the positive rate for virulence genes in each type. Almost all isolates carried

exoenzyme genes, such as *aur*, *splA*, *and splB*. Although > 90% of the isolates in ST1-IV and

ST2725-IV also carried *splE*, no isolate in ST5-II and ST764-II carried it. In terms of toxins,

almost all isolates carried *hlgA, hlgB, hlgC, LukD,* and *LukE*. Only five (6.0%) isolates in ST8-

IV carried *LukF-PV*. CC1-IV, composed of ST1-IV and ST2725-IV, whereas CC5-II, composed

of ST5-II and ST764-II, had unique virulence genes. CC1-IV carried *sea*, *seh*, *sek*, and *seq*,

whereas CC5-II carried *sec, sei, sem, sen, seo,* and *seu*. However, there were some differences

between ST5-II and ST764-II; ST5-II carried *sec, sel,* and *tst* more frequently than ST764-II. For

other virulence genes, the arginine catabolic mobile element (ACME) was detected in 1.3%,

6.0%, and 31.8% for ST1-IV, ST8-IV, and ST765-II, respectively.

Diversity in ST8-IV

Based on positive rate for drug-resistance and virulence genes, ST8-IV isolates were classified

into three groups: USA300 carrying both *LukF-PV* and ACME (4, 4.8%), *LukF*-PV-negative

isolates carrying *sec* and *tst* known as CA-MRSA/J (36, 43.4%), and others (43, 51.8%) (Table

5). Of the isolates classified as others, nineteen isolates were spa type t5071. MST for ST8-IV

isolates indicated that others with spa type t5071 were classified to a different cluster than

USA300 and CA-MRSA/J. The characteristics of others with spa type t5071 differed from CA-

MRSA/J in that they showed a high rates of resistance to levofloxacin (100%), erythromycin

(94.7%), clindamycin (76.7%), minocycline (78.9%) in EUCAST criteria. Other isolates of spa

type t5071 had higher rates of resistance genes, such as *ant(9)-Ia* (100%), *erm(A)* (100%), *tet(M)*

(68.4%), and *tet(S/M)* (63.2%); they also showed the higher positive rate for *splE* (84.2%) than

CA-MRSA/J. Of other isolates with spa type t5071, 42.1% were detected in the Kinki region and

73.7% were isolated from in-patients after 48 h of hospitalization, although there were only two

isolates of the same complex type in cgMLST.

Discussion

The most common MRSA type isolated from BSI in Japan in 2019 was ST8-IV, closely

followed by ST1-IV. In contrast, ST5-II, which had been the most prevalent in Japan,[10] ranked

fifth, representing only 7.8%. A previous Japanese study, which analysed 151 MRSA isolates

detected in blood cultures from 53 medical institutions in 2011, reported that the percentage of

SCC*mec* type IV was only 19.9%, whereas that of SCC*mec* type II was 75.6%.[21] However, a

previous study from Tokyo in the Kanto region reported that the relative abundances of SCC*mec*

types II and IV had reversed between 2015 and 2017, with SCC*mec* type IV the most prevalent,

comprising 73.5% of MRSA BSI cases.[12] Therefore, although participating medical institutions

and analytic methods differed between earlier surveillance programs and this study,[21] our data

indicated that ST8-IV and ST1-IV have replaced ST5-II in MRSA BSIs in Japan. However, there

were regional differences among the major types. The most common types in Northern, Eastern,

and Western Japan were ST5-II and ST764-II, ST1-IV, and ST8-IV, respectively. The most

common type in northern Japan, in the Hokkaido region, was very different from other regions,

and the percentage of SCC*mec* type II was 71.4%. This is similar to the surveillance results

conducted in the Hokkaido region between 2017 and 2019,[22] although the number of medical

institutions that participated in the Hokkaido region in this study was only two. Thus, there may

be regional differences in the spread of ST8-IV and ST1-IV. SCC*mec* type IV has been

associated with community-acquired infection,[23] but in this study, more than 60% of ST8-IV,

ST1-IV and ST2725-IV cases were hospital-acquired BSI. Some cgMLST types contained two

or more isolates of ST8-IV, ST1-IV, and ST2725-IV, but their numbers were small. Therefore,

although there had been cases of SCC*mec* type IV spreading in outbreaks in hospitals, the rate is

not considered high. On the other hand, the rate of hospital-acquired BSI in SCC*mec* type IV is

about 10% lower than SCC*mec* type II in this study. Considering the very high rate of hospital-

acquired BSI (82.6%) in Japan for MRSA,[24] ST8-IV, ST1-IV and ST2725-IV spread in hospitals

and communities.

We observed differences among the major types in terms of drug resistance. In antimicrobial

susceptibility testing, SCC*mec* type IV (ST8-IV, ST1-IV, and ST2725-IV) exhibited a

significantly lower resistance rate for clindamycin and minocycline than SCC*mec* type II of ST5-

II and ST764-II. The resistance rate for clindamycin was almost the same as the *erm*(A) gene

retention rate in ST8-IV, whereas the resistance rate for clindamycin was much lower than the

erm(A) gene retention rate in ST1-IV and ST2725-IV. Since the *erm*(A) gene is related to

inducible clindamycin resistance,[25] these results suggested that D-zone test should be performed

to detect inducible clindamycin resistance in areas where ST1-IV and ST2825-IV are spreading.

In contrast, the drug-resistance rate for minocycline in EUCAST criteria was almost the same as

the positive rate for the tetracycline resistance gene. SCC*mec* type IV also showed lower MIC_{50}

and MIC$_{90}$ values for cefazolin, imipenem, and meropenem than SCC*mec* type II. This result was

similar to a previous study.[21] Since β-lactams exhibit a synergic effect in combination with anti-

MRSA agents in several studies,[26,27] β-lactams might be considered to treat MRSA BSI caused

by SCC*mec* type IV in combination with anti-MRSA agents.

Almost all isolates involving all the major types carried *aur*, *splA, splB*, *hlgA, hlgB, hlgC,*

LukD, LukE, sak, and *scn*. ST1-IV and ST2725-IV, both of which are CC1-IV, had a similar

pattern of virulence genes. CC1-IV carried *splE, sea, seh, sek,* and *seq* more frequently than the

other types. In addition, MST for all isolates indicated that ST1 and ST2725 were in the same

cluster. Therefore, there might be no need to distinguish between ST1-IV and ST2725-IV in

Japan. At the clonal complex level, CC1-IV was the most abundant type (39.3%). However,

since no isolate carried *LukF-PV* and only one isolate carried ACME in CC1-IV, they differed

from USA400.[28] Several Japanese studies have reported the same characteristics for CC1-IV.[29,30]

In addition, *LukF-PV*-negative CC1-IV has been reported in Europe and Australia.[31,32] The

characteristics of CC1-IV in this study and previous Japanese studies differ from those of

European *LukF-PV*-negative CC1-IV regarding virulence genes,[32] whereas Japanese CC1-IV

have a similar pattern to *LukF-PV*-negative CC1-IV in Australia.[31] However, there exists a

difference in macrolide-resistance genes between Japan and Australia. In Australia, *erm*(C) was

detected in CC1-IV,[31] but *erm*(A) was detected in CC1-IV in this study. These results indicate

that CC1-IV has evolved and spread independently in Japan. ST5-II and ST764-II, both of which

are CC5-II, also had a similar pattern of virulence genes, but there were significant differences

between them. ST5-II carried *sec, sel,* and *tst* more frequently than ST764-II. They also had

different drug resistance genes, such as *aaD*, *fosD*, *and tet(S*/M*)*. Since the spread of ST764-II

among elderly Japanese in long-term care facilities in Japan,[33] the change in the ratio of CC5-II

should be carefully monitored.

ST8-IV was the most frequently detected type in this study, but it carried diverse virulence

genes. The positive rate for *sec* and *tst* was 44.6%, with one isolate carrying both *LukF-PV* and

ACME. The *LukF*-PV-negative ST8 clone CA-MRSA/J, characterized by carrying *sec* and *tst*,

has emerged and spread in Western Japan since 2003;[34] therefore, it is noteworthy that 43.4% of

ST8-IV in this study were estimated as CA-MRSA/J. Four isolates carried both *LukF-PV* and

ACME in *sec* and *tst*-negative ST8-IV, and three isolates with spa type identified were all t008.[35]

Based on these results, 4.8% of ST8-IV estimated as USA300. The characteristics of ST8-IV

isolates classified as others differed from CA-MRSA/J and USA300 isolates. Focusing on the

spa type t5071, the most common spa type in the others, they were classified to a different

cluster than USA300 and CA-MRSA/J. The others with spa type t5071 showed higher resistance

to several antimicrobial agents, such as fluoroquinolones, macrolides, lincosamides, and

aminoglycosides, than CA-MRSA/J. Both CA-MRSA/J and others with spa type t5071 were

detected mainly in Western Japan, but others with spa type t5071 were prevalent in the Kinki

region. In addition, others with spa type t5071 tended to be detected in inpatients after 48 h of

hospitalization. However, there were only two isolates of the same complex type in cgMLST.

These findings suggest the spread of two major types of ST8-IV, CA-MRSA/J and spa type

t5071 ST8-IV, mainly in Western Japan.

This study had several limitations. First, the impact of changes in SCC*mec* type on the

clinical course is unknown because we did not collect the information on patient backgrounds,

such as age, underlying diseases, as well as severity and prognosis of BSI. Second, we evaluated

the diversity of each MRSA type based on patterns of drug-resistance, virulence genes, spa type,

and cgMLST. However, complex type in cgMLST and spa type were not identified in 24.8% and

19.6% of the isolates in Ridom SeqSphere+. We had optimized sequencing in the MiSeq system

to determine MLST and detect drug-resistance genes in the CLC workbench. Even though the

CLC workbench was able to identify ST of MLST, Ridom SeqSphere+ did not identify ST in 59

(21.9%) of the isolates. The algorithms for genome assembly are different, which may have

affected the results. These suggest that the results of WGS may differ depending on the software,

algorithm, and database. Therefore, it is necessary to validate the best analysis tool for MRSA

WGS in the future. Third, our analyses differed from those used in previous Japanese

surveillance programs, as this is the first Japanese surveillance using WGS. In addition, the

backgrounds of participating medical institutions in this surveillance are different from those of

previous Japanese programs. The participating medical institutions in this study have in-house

microbiology laboratories and are considered secondary or tertiary hospitals. On the other hand,

the participating medical institutions in the previous study are considered primary hospitals or

long-stay hospitals because they outsourced microbiological testing.[21] It is necessary to continue

this surveillance to validate changes in MRSA because the patient populations in the two studies

should be different. Fourth, we focused on MRSA BSI. Since BSIs occur secondary to other

infections, such as pneumonia and skin and soft-tissue infections, or medical procedure, the

major types we found in this study may be spread around medical institutions. Therefore, it is

necessary to confirm whether those types are really throughout the community in Japan by

conducting nationwide active epidemiological surveillance, including a healthy population.

Conclusion

This study revealed that SCC*mec* type IV replaced SCC*mec* type II in MRSA BSI. However, the

characteristics of the major types in Japanese SCC*mec* type IV were different from USA300 and

USA400, with a low prevalence of *LukF-PV* and ACME. This study also indicated that the three

main types of SCC*mec* type IV, CC1-IV, CA-MRSA/J, and spa type t5071 ST8-IV, have spread

throughout Japan, although there were regional differences.

Transparency declarations

The authors have no conflicts of interest directly relevant to the content of this article.

Funding

This work was supported by the Japanese Association for Infectious Diseases, Grant for Clinical

Research Promotion [the 1st (2018)]. Taiga Miyazaki, a member of the Committee for Clinical

Research Promotion of the Japanese Association for Infectious Diseases, advised on study design,

sample collection, and manuscript preparation. The funders had no role in the analysis or the

decision to publish.

Acknowledgements

We thank Professor Kazuyoshi Kawakami (Medical Microbiology, Mycology and Immunology,

Tohoku University Graduate School of Medicine) for advice on study design and sample

collection as chair of the Committee for Clinical Research Promotion of the Japanese

Association for Infectious Diseases, Shuji Miyazaki (Department of Laboratory Medicine,

Nagasaki University of Hospital) for support of the analysis of bacteria, and Editage

(www.editage.com) for English language editing.

We also thank all staff members cooperated in this study at the Japanese Association for

Infectious Diseases, Furano Hospital, Sapporo Medical University Hospital, Aomori Prefectural

Central Hospital, Odate Municipal General Hospital, Fukushima Medical University, Saitama

Medical University International Medical Center, Jikei University Katsusika Medical Center,

Yokohama City University Hospital, Yokohama Municipal Citizen's Hospital, Yokohama City

University Medical Center, Tokyo Bay Urayasu Ichikawa Medical Center, Showa University

Hospital, Tokyo Saiseikai Central Hospital, Hamamatsu University Hospital, Ishikawa

Prefectural Central Hospital, Chuno Kosei Hospital, Asanogawa General Hospital, Nagaoka Red

Cross Hospital, University of Fukui Hospital, Toyama University Hospital, Nagoya University

Hospital, Nagoya City University West Medical Center, Niigata University Medical & Dental

Hospital, Japanese Red Cross Society Suwa Hospital, Daiyukai General Hospital, Osaka Medical

and Pharmaceutical University Hospital, Osaka City General Hospital, Kobe City Nishi-Kobe

Medical Center, Osaka Univerisity Hospital, Kindai University Hospital, University Hospital

Kyoto Prefectural University of Medicine, Nara Prefectural General Medical Center, Shimane

University Hospital, Tottori University Hospital, Ehime University Hospital, Chikamori Hospital,

Uwajima City Hospital, Oita Prefectural Hospital, Saga-Ken Medical Centre Koseikan,

University of The Ryukyus Hospital, Fukuoka University Hospital, National Hospital

Organization Nagasaki Medical Center, Fukuoka Children's Hospital, Nagasaki Goto Chuoh

Hospital, and Nagasaki University Hospital for collection of MRSA isolates.

Access to Data

The derived data supporting the findings of this study are presented in this paper. WGS data

reported in this study are available in the DDBJ Sequence Read Archive

(https://www.ddbj.nig.ac.jp/dra/index-e.html) under accession number DRA013058.

Contributions

All authors were involved in the study design, acquisition, and interpretation of the data. NK, DS,

KO, and KY were involved in data analysis. NK wrote the original manuscript, and all authors

revised the manuscript and have approved the manuscript for publication.

References

1. Kern W V., Rieg S. Burden of bacterial bloodstream infection—a brief update on

epidemiology and significance of multidrug-resistant pathogens. *Clin Microbiol Infect* 2020; **26**:

151–7.

2. Wisplinghoff H, Bischoff T, Tallent SM, *et al.* Nosocomial Bloodstream Infections in US

Hospitals: Analysis of 24,179 Cases from a Prospective Nationwide Surveillance Study. *Clin*

Infect Dis 2004; **39**: 309–17.

3. Kourtis AP, Hatfield K, Baggs J, *et al.* Vital Signs: Epidemiology and Recent Trends in

Methicillin-Resistant and in Methicillin-Susceptible Staphylococcus aureus Bloodstream

Infections - United States. *MMWR Morb Mortal Wkly Rep* 2019; **68**: 214–9.

4. Naimi TS, LeDell KH, Como-Sabetti K, *et al.* Comparison of Community- and Health Care -

Associated Methicillin-Resistant Staphylococcus aureus Infection. *J Am Med Assoc* 2003; **290**:

2976–84.

5. D'Agata EMC, Webb GF, Horn MA, *et al.* Modeling the invasion of community-acquired

methicillin-resistant Staphylococcus aureus into hospitals. *Clin Infect Dis* 2009; **48**: 274–84.

6. David MZ, Cadilla A, Boyle-Vavra S, Daum RS. Replacement of HA-MRSA by CA-MRSA

infections at an academic medical center in the midwestern United States, 2004-5 to 2008. *PLoS*

One 2014; **9**: e92760.

7. Nichol KA, Adam HJ, Golding GR, *et al.* Characterization of MRSA in Canada from 2007 to

2016. *J Antimicrob Chemother* 2019; **74**: iv55–63.

8. Coombs GW, Daley DA, Mowlaboccus S, *et al*. Australian Group on Antimicrobial

Resistance (AGAR) Australian Staphylococcus aureus Sepsis Outcome Programme (ASSOP)

Annual Report 2019. *Commun Dis Intell* 2020; **44**.

9. Henderson A, Nimmo GR. Control of healthcare- and community-associated MRSA: recent

progress and persisting challenges. *Br Med Bull* 2018; **125**: 25–41.

10. Yamada K, Yanagihara K, Hara Y, *et al.* Clinical Features of Bacteremia Caused by

Methicillin-Resistant Staphylococcus aureus in a Tertiary Hospital. *Tohoku J Exp Med* 2011;

224: 61–7.

11. Kaku N, Yanagihara K, Morinaga Y, *et al.* Influence of antimicrobial regimen on decreased

in-hospital mortality of patients with MRSA bacteremia. *J Infect Chemother* 2014; **20**: 350–5.

12. Hamada M, Yamaguchi T, Sato A, *et al.* Increased Incidence and Plasma-Biofilm Formation

Ability of SCCmec Type IV Methicillin-Resistant Staphylococcus aureus (MRSA) Isolated From

Patients With Bacteremia. *Front Cell Infect Microbiol* 2021; **11**: 602833.

13. Kimura Y, Morinaga Y, Akamatsu N, *et al.* Antimicrobial susceptibility and molecular

characteristics of methicillin-resistant Staphylococcus aureus in a Japanese secondary care

facility. *J Infect Chemother* 2016; **22**: 14–8.

14. Schwarz S, Silley P, Simjee S, *et al.* Editorial: Assessing the antimicrobial susceptibility of

bacteria obtained from animals. *J Antimicrob Chemother* 2010; **65**: 601–4.

15. CLSI. Performance *Standards for Antimicrobial Susceptibility Testing-Thirty-First Edition:*

M100. 2021.

16. The European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for

interpretation of MICs and zone diameters, version 11.0, 2021.

17. Kaya H, Hasman H, Larsen J, *et al.* SCC mec Finder, a Web-Based Tool for Typing of

Staphylococcal Cassette Chromosome mec in Staphylococcus aureus Using Whole-Genome

Sequence Data. *mSphere* 2018; **3**: e00612-17.

18. Tetzschner AMM, Johnson JR, Johnston BD, *et al.* In Silico Genotyping of Escherichia coli

Isolates for Extraintestinal Virulence Genes by Use of Whole-Genome Sequencing Data. *J Clin*

Microbiol 2020; **58**: e01269-20.

19. Joensen KG, Scheutz F, Lund O, *et al.* Real-time whole-genome sequencing for routine

typing, surveillance, and outbreak detection of verotoxigenic Escherichia coli. *J Clin Microbiol*

2014; **52**: 1501–10.

20. Souvorov A, Agarwala R, Lipman DJ. SKESA: strategic k-mer extension for scrupulous

assemblies. *Genome Biol* 2018; **19**: 153.

21. Miura Y, Yamaguchi T, Nakamura I, *et al.* Epidemiological Trends Observed from

Molecular Characterization of Methicillin-Resistant Staphylococcus aureus Isolates from Blood

Cultures at a Japanese University Hospital, 2012–2015. *Microb Drug Resist* 2018; **24**: 70–5.

22. Aung MS, Urushibara N, Kawaguchiya M, *et al.* Clonal diversity of methicillin-resistant

Staphylococcus aureus (MRSA) from bloodstream infections in northern Japan: Identification of

spermidine N-acetyltransferase gene (speG) in staphylococcal cassette chromosomes (SCCs)

associated with type II and IV. *J Glob Antimicrob Resist* 2021; **24**: 207–14.

23. DeLeo FR, Otto M, Kreiswirth BN, *et al.* Community-associated meticillin-resistant

Staphylococcus aureus. *Lancet* 2010; **375**: 1557–68.

24. Takeshita N, Kawamura I, Kurai H, *et al.* Unique characteristics of community-onset

healthcare- associated bloodstream infections: a multi-centre prospective surveillance study of

bloodstream infections in Japan. *J Hosp Infect* 2017; **96**: 29–34.

25. Leclercq R. Mechanisms of resistance to macrolides and lincosamides: Nature of the

resistance elements and their clinical implications. *Clin Infect Dis* 2002; **34**: 482–92.

26. Ono D, Yamaguchi T, Hamada M, *et al.* Analysis of synergy between beta-lactams and anti-

methicillin-resistant Staphylococcus aureus agents from the standpoint of strain characteristics

and binding action. *J Infect Chemother* 2019; **25**: 273–80.

27. Davis JS, Hal S van, Tong SYC. Combination antibiotic treatment of serious methicillin-

resistant staphylococcus aureus infections. *Semin Respir Crit Care Med* 2015; **36**: 3–16.

28. Mulvey MR, MacDougall L, Cholin B, *et al.* Community-associated Methicillin-resistant

Staphylococcus aureus, Canada. *Emerg Infect Dis* 2005; **11**: 844-850.

29. Nakaminami H, Takadama S, Ito A, *et al.* Characterization of sccmec type IV methicillin-

resistant staphylococcus aureus clones increased in Japanese hospitals. *J Med Microbiol* 2018;

67: 769–74.

30. Osaka S, Okuzumi K, Koide S, *et al.* Genetic shifts in methicillin-resistant Staphylococcus

aureus epidemic clones and toxin gene profiles in Japan: Comparative analysis among pre-

epidemic, epidemic and post-epidemic phases. *J Med Microbiol* 2018; **67**: 392–9.

31. Coombs GW, Monecke S, Pearson JC, *et al.* Evolution and diversity of community-

associated methicillin-resistant Staphylococcus aureus in a geographical region. *BMC Microbiol*

2011; **11**: 215.

32. Earls MR, Shore AC, Brennan GI, *et al.* A novel multidrug-resistant PVL-negative CC1-

MRSA-IV clone emerging in Ireland and Germany likely originated in South-Eastern Europe.

Infect Genet Evol 2019; **69**: 117–26.

33. Kawamura K, Kitaoka K, Kimura K, *et al.* Spread of seb-Positive Methicillin-Resistant

Staphylococcus aureus SCCmec Type II-ST764 Among Elderly Japanese in Nonacute Care

Settings. *https://home.liebertpub.com/mdr* 2019; **25**: 915–24.

34. Iwao Y, Ishii R, Tomita Y, *et al.* The emerging ST8 methicillin-resistant Staphylococcus

aureus clone in the community in Japan: associated infections, genetic diversity, and comparative

genomics. *J Infect Chemother* 2012; **18**: 228–40.

35. O'Hara FP, Suaya JA, Ray GT, *et al.* spa Typing and Multilocus Sequence Typing Show

Comparable Performance in a Macroepidemiologic Study of Staphylococcus aureus in the

United States. *Microb Drug Resist* 2016; **22**: 88-96.

Figure Legends

Figure 1. The major types in Japan

The major types in Japan (A) and each region (B) were determined by a combination of sequence

type (ST) and SCC*mec* type.

Figure 2. Minimum Spanning Tree for all isolates.

One hundred twenty-two samples with more than 10% missing values of the items for distance

calculation were excluded from MST. MST for 149 samples was created based on MLST (8),

cgMLST (1861), and *S. aureus* Accessory (706) using Ridom SeqSphere+. The number in the

node is a complex type determined by cgMLST.

MST, Minimum Spanning Tree; cgMLST, core genome MLST.

Figure 3. Minimum Spanning Tree for ST8-IV isolates

Twenty-two samples with more than 10% missing values of the items for distance calculation

were excluded from MST. MST for 51 samples was created based on MLST (7), cgMLST

(1861), and *S. aureus* Accessory (706) using Ridom SeqSphere+. The number in the node is a

complex type determined by cgMLST.

MST, Minimum Spanning Tree; cgMLST, core genome MLST.

Table 1. Characteristics of the major types

	ST8-IV	ST1-IV	ST2725-IV	ST764-II	ST5-II
	(n = 83)	(n = 80)	(n = 26)	(n = 22)	(n = 21)
Patient background					
Outpatient	21.7%	23.8%	19.2%	13.6%	-
Inpatient within 48 h of hospitalization	16.9%	13.8%	19.2%	13.6%	28.6%
Inpatient after 48 h of hospitalization	60.2%	62.5%	61.5%	72.7%	71.4%
Unknown	1.2%	-	-	-	-
Subtypes of SCC*mec* IV					
IVa	8.4%	100%	92.3%	-	-
IVc	4.8%	-	-	-	-
IVg	1.2%	-	-	-	-
IVh	4.8%	-	-	-	-
Non-subtypeable	80.7%	-	7.7%	-	-
Spa type					
t002	-	-	-	59.1%	61.9%
t045	-	-	-	4.5%	-
t088	-	-	-	4.5%	-
t458	-	-	-	-	4.8%
t568	-	-	-	-	4.8%
t693	1.3%	-	-	-	-
t1767	34.9%	-	-	4.5%	4.8%
t1779	1.2%	-	-	-	-
t1784	1.2%	72.5%	84.6%	-	-
t4133	2.4%	-	-	-	-
t4407	1.2%	-	-	-	-
t4494	-	2.5%	-	-	-

t5071	24.1%	-	-	-	-
t7083	-	-	-	-	9.5%
t7744	-	5.0%	3.8%	-	-
t17749	1.2%	-	-	-	-
t18300	1.2%	-	-	-	-
t18648	1.2%	-	-	-	-
Unknown	3.6%	3.8%	-	4.5%	-
Unable to analyze	15.7%	15.0%	11.5%	22.7%	14.3%
Drug resistance rate in CLSI/EUCAST					
Oxacillin	96.4 / 96.4	88.8 / 88.8	88.5 / 88.5	100 / 100	100 / 100
Cefoxitin	96.4 / 96.4	97.5 / 97.5	92.3 / 92.3	100 / 100	100 / 100
Levofloxacin	55.4 / 56.6	100 / 100	100 / 100	100 / 100	100 / 100
Erythromycin	59.0 / 61.4	93.8 / 93.8	100 / 100	100 / 100	100 / 100
Clindamycin	44.6 / 45.8	3.8 / 5.0	3.8 / 3.8	100 / 100	85.7 / 85.7
Minocycline	0.0 / 30.1	0.0 / 6.3	0.0 / 3.8	4.5 / 100	4.8 / 71.4
Trimethoprim / Sulfamethoxazole	0.0 / 0.0	0.0 / 0.0	0.0 / 0.0	0.0 / 0.0	0.0 / 0.0
Vancomycin	1.2 / 2.4	0.0 / 1.3	0.0 / 3.8	0.0 / 0.0	0.0 / 0.0
Teicoplanin	0.0 / 1.2	1.3 / 1.3	0.0 / 0.0	0.0 / 0.0	0.0 / 0.0
Linezolid	1.2 / 1.2	1.3 / 1.3	3.8 / 3.8	4.5/ 4.5	0.0 / 0.0

SCC*mec*, staphylococcal cassette chromosome *mec*

488
489

Table 2. Antimicrobial susceptibility of MRSA isolates

MIC_{50} / MIC_{90}	ST8-IV (n = 83)	ST1-IV (n = 80)	ST2725-IV (n = 26)	ST764-II (n = 22)	ST5-II (n = 21)
Ampicillin	$\geqq$32 / $\geqq$32	$\geqq$32 / $\geqq$32	$\geqq$32 / $\geqq$32	8 / 16	$\geqq$32 / $\geqq$32
Oxacillin	$\geqq$8 / $\geqq$8	$\geqq$8 / $\geqq$8	$\geqq$8 / $\geqq$8	$\geqq$8 / $\geqq$8	$\geqq$8 / $\geqq$8
Cefoxitin	8 / 8	8 / 16	8 / 16	$\geqq$32 / $\geqq$32	$\geqq$32 / $\geqq$32
Cefazolin	4 / 16	4 / 8	2 / 16	$\geqq$32 / $\geqq$32	$\geqq$32 / $\geqq$32
Imipenem	$\leqq$0.25 / $\leqq$0.25	$\leqq$0.25 / $\leqq$0.25	0.25 / 0.5	$\geqq$32 / $\geqq$32	$\geqq$32 / $\geqq$32
Meropenem	1 / 4	$\leqq$0.25 / 1	1 / 1	16 / $\geqq$32	16 / $\geqq$32
Levofloxacin	4 / 8	$\geqq$32 / $\geqq$32	16 / $\geqq$32	$\geqq$32 / $\geqq$32	$\geqq$32 / $\geqq$32
Erythromycin	$\geqq$16 / $\geqq$16	$\geqq$16 / $\geqq$16	$\geqq$16 / $\geqq$16	$\geqq$16 / $\geqq$16	$\geqq$16 / $\geqq$16
Clindamycin	$\leqq$0.125 / $\geqq$16	$\leqq$0.125 /	$\leqq$0.125 /	$\geqq$16 / $\geqq$16	$\geqq$16 / $\geqq$16

		≦0.125	≦0.125		
Minocycline	≦0.25 / 8	≦0.25 / ≦0.25	≦0.25 / ≦0.25	8 / 8	8 / 8
Trimethoprim / Sulfamethoxazole	≦0.5/9.5 / ≦0.5/9.5	≦0.5/9.5 / ≦0.5/9.5	≦0.5/9.5 / ≦0.5/9.5	≦0.5/9.5 / ≦0.5/9.5	≦0.5/9.5 / ≦0.5/9.5
Vancomycin	≦0.25 / 1	≦0.25 / 1	≦0.25 / 2	1 / 1	1 / 2
Teicoplanin	≦0.25 / ≦0.25	≦0.25 / 1	both ≦0.25	≦0.25 / ≦0.25	≦0.25 / 1
Linezolid	2/ 2	2/ 2	2/ 2	1 / 2	2 / 2
Daptomycin	≦0.25 / 1	≦0.25 / 1	≦0.25 / 0.5	≦0.25 / 1	≦0.25 / 2
Arbekacin	4 / ≧16	2 / 8	2 / 8	4 / 8	4 / ≧16

490
491

Table 3. The positive rate for drug-resistance genes

	ST8-IV	ST1-IV	ST2725-IV	ST764-II	ST5-II
	(n = 83)	(n = 80)	(n = 26)	(n = 22)	(n = 21)
Aminoglycoside-resistance gene	92.8%	95.0%	96.2%	100%	100%
aac(6')-aph(2'')	56.6%	8.8%	11.5%	81.8%	42.9%
aaD	27.7%	1.3%	-	40.9%	85.7%
ant(6)-Ia	2.4%	-	-	4.5%	-
ant(9)-Ia	45.8%	95.0%	96.2%	100%	100%
aph(3')-III	2.4%	-	-	-	-
β-lactamase gene *blaZ*	97.6%	98.8%	92.3%	-	71.4%
Chloramphenicol-resistance gene, *cat(pC221)*	-	-	-	4.5%	-
Fosfomycin-resistance gene	6.0%	-	-	40.9%	4.8%
fosB4	-	-	-	4.5%	4.8%
fosD	6.0%	-	-	40.9%	-
Macrolide-resistance gene	56.6%	93.8%	100%	100%	100%
erm(A)	47.0%	93.8%	96.2%	100%	100%
erm(B)	-	1.3%	-	4.5%	-
erm(C)	8.4%	1.3%	4.4%	4.5%	-
mph(C)	2.4%	-	-	4.5%	-
msr(A)	2.4%	-	-	4.5%	-
Methicillin-resistance gene, *mecA*	100%	100%	100%	100%	100%

	ST8-IV	ST1-IV	ST2725-IV	ST764-II	ST5-II
Tetracycline-resistance gene	28.9%	-	3.8%	100%	76.2%
tet(K)	2.4%	-	-	4.5%	-
tet(M)	26.5%	-	3.8%	100%	76.2%
tet(S/M)	24.1%	-	-	100%	57.1%
Trimethoprim-resistance gene, dfrG	1.2%	-	-	-	-

492
493

Table 4. The positive rate for virulence genes

	ST8-IV	ST1-IV	ST2725-IV	ST764-II	ST5-II
	(n = 83)	(n = 80)	(n = 26)	(n = 22)	(n = 21)
Exoenzyme genes					
aur	100%	100%	100%	100%	100%
splA	96.4%	98.8%	96.2%	100%	100%
splB	94.0%	98.8%	96.2%	100%	100%
splE	47.0%	91.3%	96.2%	-	-
Toxin genes					
edinA	18.1%	-	-	-	-
hlgA	98.8%	100%	96.2%	100%	95.2%
hlgB	100%	100%	100%	100%	100%
hlgC	86.7%	100%	100%	100%	100%
LukD	97.6%	100%	100%	100%	95.2%
LukE	98.8%	100%	100%	100%	95.2%
LukF-PV	6.0%	-	-	-	-
sea	4.8%	88.8%	80.8%	-	4.8%
seb	1.2%	-	3.8%	86.4%	9.5%
sec	44.6%	-	3.8%	13.6%	61.9%
seg	-	-	-	95.5%	81.0%
seh	-	100%	96.2%	-	4.8%
sei	-	-	-	95.5%	100%
sek	6.0%	85.0%	80.8%	13.6%	-

sel	39.8%	-	-	13.6%	71.4%
sem	-	-	-	100%	100%
sen	-	-	-	100%	90.5%
seo	-	-	-	100%	95.2%
sep	67.5%	-	-	9.1%	28.6%
seq	7.2%	85.0%	80.8%	13.6%	4.8%
seu	-	-	-	100%	85.7%
tst	44.6%	3.8%	3.8%	13.6%	66.7%
Other					
ACME	6.0%	1.3%	-	31.8%	-
sak	94.0%	98.8%	100%	72.7%	85.7%
scn	96.4%	98.8%	100%	86.4%	85.7%

Table 5. Characteristics of CA-MRSA/J, USA300, and other isolates in ST8-IV

	USA300	CA-MRSA/J	Others	
			All	spa type t5071
	(n = 4)	(n = 36)	(n = 43)	(n = 19)
Subtypes of SCC*mec* type IV				
IVa	100%	5.6%	2.3%	-
Ivc	-	5.6%	4.7%	-
Ivg	-	-	2.3%	-
Ivh	-	-	9.3%	10.5%
Non-subtypeable	-	88.9%	81.4%	89.5%
Spa type				
t008	75.0%	8.3%	7.0%	-
t211	-	-	2.3%	-
t1767	-	61.1%	16.3%	-
t1779	-	-	2.3%	-
t1784	-	-	2.3%	-
t4133	-	2.8%	2.3%	-
t4407	-	-	2.3%	-

t5071	-	2.8%	44.2%	100.0%
t17749	-	-	2.3%	-
t18300	-	2.8%	-	-
t18658	-	2.8%	-	-
unknown	-	2.8%	4.7%	-
unable to analyze	25.0%	16.7%	14.0%	-
Region				
Hokkaido	-	-	-	-
Tohoku	-	2.8%	-	-
Kanto	25.0%	22.2%	9.3%	10.5%
Chubu	0.0%	25.0%	14.0%	15.8%
Kinki	25.0%	16.7%	41.9%	42.1%
Chugoku	0.0%	2.8%	7.0%	5.3%
Shikoku	25.0%	8.3%	4.7%	5.3%
Kyusyu	25.0%	22.2%	23.3%	21.1%
Patient background				
Outpatient	50.0%	19.4%	20.9%	21.1%
Inpatient within 48 h of hospitalization	0.0%	22.2%	14.0%	5.3%
Inpatient after 48 h of hospitalization	25.0%	58.3%	65.1%	73.7%
Unknown	25.0%	-	-	-
Drug resistance rate				
Oxacillin	100/100	97.2/97.2	95.3/95.3	89.5/89.5
Cefoxitin	75.0/75.0	94.4/94.4	100/100	100/100
Levofloxacin	100/100	11.1/13.9	88.4/88.4	100/100
Erythromycin	75.0/75.0	27.8/30.6	83.7/86.0	94.7/94.7
Clindamycin	0.0/0.0	11.1/13.9	76.7/76.7	76.7/76.7
Minocycline	0.0/0.0	0.0/5.6	0.0/53.5	0.0/78.9
Trimethoprim / Sulfamethoxazole	0.0/0.0	0.0 / 0.0	0.0/2.3	0.0/0.0
Vancomycin	0.0/0.0	2.8/2.8	0.0/0.0	0.0/0.0
Teicoplanin	0.0/0.0	0.0/2.8	0.0/0.0	0.0/0.0
Linezolid	0.0/0.0	2.8/2.8	0.0/0.0	0.0/0.0
Aminoglycoside-resistace gene	100.0%	88.9%	95.3%	100.0%
aac(6')-aph(2'')	75.0%	72.2%	41.9%	26.3%
aaD	0.0%	50.0%	11.6%	0.0%
ant(6)-Ia	50.0%	0.0%	0.0%	0.0%
ant(9)-Ia	0.0%	5.6%	83.7%	100.0%
aph(3')-III	50.0%	0.0%	0.0%	0.0%

β-lactamase gene, *blaZ*	75.0%	100.0%	97.7%	100.0%
Fosfomycin-resistance gene, *fosD*	0.0%	0.0%	11.6%	15.8%
Macrolide-resistance gene	50.0%	22.2%	86.0%	100.0%
erm(A)	0.0%	8.3%	83.7%	100.0%
erm(C)	0.0%	13.9%	4.7%	5.3%
mph(C)	20.0%	0.0%	0.0%	0.0%
msr(A)	20.0%	0.0%	0.0%	0.0%
Tetracycline-resistance gene	0.0%	8.3%	48.8%	68.4%
tet(K)	0.0%	5.6%	0.0%	0.0%
tet(M)	0.0%	2.8%	48.8%	68.4%
tet(S/M)	0.0%	0.0%	46.5%	63.2%
Trimethoprim-resistance gene, *dfrG*	0.0%	0.0%	2.3%	0.0%
Exoenzyme genes				
aur	100.0%	100.0%	100.0%	100.0%
splA	75.0%	94.4%	100.0%	100.0%
splB	75.0%	94.4%	95.3%	94.7%
splE	75.0%	2.8%	81.4%	84.2%
Toxin genes				
edinA	0.0%	30.6%	9.3%	0.0%
hlgA	100.0%	100.0%	97.7%	94.7%
hlgB	100.0%	100.0%	100.0%	100.0%
hlgC	100.0%	86.1%	86.0%	94.7%
LukD	100.0%	97.2%	97.7%	100.0%
LukE	100.0%	97.2%	100.0%	100.0%
LukF-PV	100.0%	0.0%	2.3%	0.0%
sea	0.0%	11.1%	0.0%	0.0%
seb	0.0%	2.8%	0.0%	0.0%
sec	0.0%	100.0%	2.3%	0.0%
sek	100.0%	2.8%	0.0%	0.0%
sel	0.0%	88.9%	2.3%	0.0%
sep	0.0%	55.6%	83.7%	94.7%
seq	100.0%	2.8%	2.3%	0.0%
tst	0.0%	100.0%	23.0%	0.0%
Other virulence genes				
ACME	100.0%	0.0%	2.3%	0.0%
sak	75.0%	91.4%	97.7%	94.7%
scn	100.0%	94.4%	97.7%	94.7%

SCC*mec*, staphylococcal cassette chromosome *mec*

(A)

Breakdown of others

ST	SCC*mec*	%
CC1 other than ST1 and ST2725	IV	1.9%
ST5	IV	1.5%
CC5 other than ST5 and ST764	II	1.1%
	IV	0.7%
ST8	I	0.9%
	II	1.5%
	IX	0.4%
CC8 other than ST8	IV	1.9%
Other STs	IV	1.9%
	V	1.5%

(B)

Figure 1. Combination of Sequence type and SCC*mec* type
A percentage of ST and SCC*mec* type in Japan (A) and each region (B).
MRSA, methicillin-resistant *Staphylococcus aureus*; ST, sequence type; CC, clonal complex

Figure 2. Minimum Spanning Tree for all isolates.
One hundred twenty-two samples with more than 10% missing values of the items for distance calculation were excluded from MST. MST for 149 samples was created based on MLST (8), cgMLST (1861), and *S. aureus* Accessory (706) using Ridom SeqSphere+. The number in the node is a complex type determined by cgMLST.
MST, Minimum Spanning Tree; cgMLST, core genome MLST.

Figure 3. Minimum Spanning Tree for ST8-IV isolates

Twenty-two samples with more than 10% missing values of the items for distance calculation were excluded from MST. MST for 51 samples was created based on MLST (7), cgMLST (1861), and *S. aureus* Accessory (706) using Ridom SeqSphere+. The number in the node is a complex type determined by cgMLST. MST, Minimum Spanning Tree; cgMLST, core genome MLST.

(A)

Fifty-one Japanese hospitals participated in surveillance

No MRSA isolate that meet the criteria in 6 hospitals

Two hundred seventy-four MRSA isolates isolated from 45 hospitals were analyzed at Nagasaki University

Four isolates were excluded from the final analysis because they were identified as MSSA.

Two hundred seventy MRSA isolates were included in the final analysis

(B)

Hokkaido
14 strains from 2 hospitals
(5.2%)

Kinki region
51 strains from
7 hospitals (18.9%)

Chugoku region
8 strains from
2 hospitals (3.0%)

Tohoku region
10 strains from
3 hospitals (3.7%)

Kanto region
56 strains from
8 hospitals (20.7%)

Shikoku region
16 strains from
3 hospitals (5.9%)

Chubu region
67 strains from
12 hospitals (24.8%)

Kyushu region
48 strains from
8 hospitals (17.8%)

Supplementary Figure 1. Selection flow diagram and participating hospitals
MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin-susceptible *Staphylococcus aureus*