Early and rapid identification of COVID-19 patients with neutralizing type I-interferon

auto-antibodies by an easily implementable algorithm

Bengisu Akbil[1,2]*, Tim Meyer[3]*, Paula Stubbemann[4]*, Charlotte Thibeault[4]*, Olga

Staudacher[3,5], Daniela Niemeyer[1,6], Jenny Jansen[1,2], Barbara Mühlemann[1,6], Jan Doehn[4],

Christoph Tabeling[2,4], Christian Nusshag[7], Cédric Hirzel[8], David Sökler Sanchez[9,10],

Alexandra Nieters[11,] Achim Lother[12], Daniel Duerschmied[12], Nils Schallner[13,14], Jan Nikolaus

Lieberum[13,14], Dietrich August[15], Siegbert Rieg[15], Valeria Falcone[16], Hartmut Hengel[16], Uwe

Kölsch[3], Nadine Unterwalder[3], Ralf-Harto Hübner[2,4], Terry C. Jones[1,6,17], Norbert Suttorp[4],

Christian Drosten[1,6], Klaus Warnatz[9,10], Thibaud Spinetti[18], Joerg C. Schefold[18], Thomas

Dörner[19], Leif Sander[4], Victor M. Corman[1,6,20], Uta Merle[21], Pa-COVID study Group[22], Florian

Kurth[4,23#], Horst von Bernuth[2,3,5,24#], Christian Meisel[3,25#], Christine Goffinet[1,2#]

[1]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and

Humboldt-Universität zu Berlin, Institute of Virology, 10117 Berlin, Germany

[2]Berlin Institute of Health at Charité - Universitätsmedizin Berlin, Berlin, Germany

[3]Labor Berlin GmbH, Department of Immunology, Berlin, Germany

[4]Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and

Humboldt-Universität zu Berlin, Department of Infectious Diseases and Respiratory

Medicine, Berlin, Germany

[5]Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin,

Humboldt-Universität zu Berlin, and Berlin Institute of Health, Department of Pediatric

Respiratory Medicine, Immunology and Critical Care Medicine, Berlin, Germany

[6]DZIF German Centre for Infection Research (DZIF), partner site Charité, 10117 Berlin,

Germany

[7]Department of Nephrology, Heidelberg University Hospital, Heidelberg, Germany

[8]Department of Infectious Diseases, Inselspital, Bern University Hospital, University of Bern,

Bern, Switzerland

[9]Department of Rheumatology and Clinical Immunology, Medical Center – University of

Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

[10]Center for Chronic Immunodeficiency (CCI), Medical Center – University of Freiburg,

Faculty of Medicine, University of Freiburg, Freiburg, Germany

[11]University Medical Center Freiburg and Faculty of Medicine, University of Freiburg, Center

for Biobanking, FREEZE-Biobank, Freiburg, Germany

[12]Cardiology and Medical Intensive Care, Heart Center, Faculty of Medicine, University of

Freiburg

[13]Department of Anesthesiology and Critical Care, Medical Center - University of Freiburg,

Germany

[14]Faculty of Medicine, University of Freiburg, Germany

[15]Division of Infectious Diseases, Department of Medicine II, Medical Centre – University of

Freiburg, Faculty of Medicine, University of Freiburg, 79106 Freiburg, Germany

[16]Institute of Virology, Freiburg University Medical Center, Faculty of Medicine, Albert-

Ludwigs-University of Freiburg, Freiburg, Germany.

[17]Centre for Pathogen Evolution, Department of Zoology, University of Cambridge, Downing

St., Cambridge, CB2 3EJ, U.K.

[18]Department of Intensive Care Medicine, Inselspital, Bern University Hospital, University of

Bern, Freiburgstrasse, CH-3010 Bern, Switzerland

[19]Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and

Humboldt-Universität zu Berlin, Department of Medicine/Rheumatology and Clinical

Immunology , Berlin, Germany and DRFZ, Berlin, Germany

[20]Labor Berlin GmbH, Berlin, Germany

[21] Department of Gastroenterology, Heidelberg University Hospital, Heidelberg, Germany

[22]Collaborators listed at the end of the article

[23]Department of Tropical Medicine, Bernhard Nocht Institute for Tropical Medicine and

Department of Medicine I, University Medical Centre Hamburg-Eppendorf, Hamburg,

Germany

[24]Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin,

Humboldt-Universität zu Berlin, and Berlin Institute of Health (BIH), Berlin-Brandenburg

Center for Regenerative Therapies (BCRT), Berlin, Germany

[25]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and

Humboldt-Universität zu Berlin, Institute of Medical Immunology, Berlin, Germany

*These authors contributed equally

#These authors contributed equally, co-corresponding authors

Corresponding authors:

Florian Kurth; Department of Infectious Diseases and Respiratory Medicine, Charité -

Universitätsmedizin Berlin, Augustenburger Platz 1, D-13353 Berlin, Germany,

florian.kurth@charite.de, +49 30 450 665023

Horst von Bernuth; Department of Pediatric Respiratory Medicine, Immunology and Critical

Care Medicine, Charité - Universitätsmedizin Berlin, Augustenburger Platz 1, D-13353

Berlin, Germany, horst.von-bernuth@charite.de, +49 30 450 566693

Christian Meisel, Labor Berlin, Charité - Universitätsmedizin Berlin, Sylter Str. 2, 13353

Berlin, Germany, christian.meisel@laborberlin.com, +49 30 405026471

Christine Goffinet, Institute for Virology, Charité – Universitätsmedizin Berlin, Charitéplatz 1,

10117 Berlin, Germany, christine.goffinet@charite.de, +49 30 450525489

ABSTRACT

Purpose

Six-19% of critically ill COVID-19 patients display circulating auto-antibodies against type I

interferons (IFN-AABs). Here, we establish a clinically applicable strategy for early

identification of IFN-AAB-positive patients for potential subsequent clinical interventions.

Methods

We analysed sera of 430 COVID-19 patients with severe and critical disease from four

hospitals for presence of IFN-AABs by ELISA. Binding specificity and neutralizing activity

were evaluated via competition assay and virus-infection-based neutralization assay. We

defined clinical parameters associated with IFN-AAB positivity. In a subgroup of critically ill

patients, we analyzed effects of therapeutic plasma exchange (TPE) on the levels of IFN-

AABs, SARS-CoV-2 antibodies and clinical outcome.

Results

The prevalence of neutralizing AABs to IFN-α and IFN-ω in COVID-19 patients was 4.2%

(18/430), while being undetectable in an uninfected control cohort. Neutralizing IFN-AABs

were detectable exclusively in critically affected, predominantly male (83%) patients (7.6%

IFN-α and 4.6% IFN-ω in 207 patients with critical COVID-19). IFN-AABs were present early

post-symptom onset and at the peak of disease. Fever and oxygen requirement at hospital

admission co-presented with neutralizing IFN-AAB positivity. IFN-AABs were associated with

higher mortality (92.3% versus 19.1 % in patients without IFN-AABs). TPE reduced levels of

IFN-AABs in three of five patients and may increase survival of IFN-AAB-positive patients

compared to those not undergoing TPE.

Conclusion

IFN-AABs may serve as early biomarker for development of severe COVID-19. We propose

to implement routine screening of hospitalized COVID-19 patients according to our algorithm

for rapid identification of patients with IFN-AABs who most likely benefit from specific

therapies.

3-5 KEYWORDS

COVID-19, SARS-CoV-2, type I interferon, autoantibodies

INTRODUCTION

Since its first detection in Wuhan, China, in 2019, Severe-acute-respiratory syndrome

Coronavirus 2 (SARS-CoV-2) has placed an unprecedented burden on health care systems

worldwide. The clinical spectrum of the associated disease, COVID-19, ranges from

asymptomatic infection to severe disease with hypoxemia, acute respiratory distress

syndrome (ARDS), multiorgan failure, and death (Pfortmueller et al. 2021). Approximately

35% of patients remain asymptomatic, 55% develop upper respiratory tract infections,

whereas 15% develop severe pneumonia (defined as SpO2 < 90% at room air) and 5%

critical pneumonia (defined as Acute Respiratory Distress Syndrome (ARDS), requiring

mechanical ventilation or extra-corporeal membrane (ECMO) (Sah et al. 2021).

Scores containing clinical and laboratory parameters support risk stratification and

resource allocation in clinical practice worldwide (Knight et al. 2020; "Information on COVID-

19 Treatment, Prevention and Research" n.d.). Demographic and clinical risk factors for a

severe disease course include advanced age, male sex, and pre-existing comorbidities

(Williamson et al. 2020). Moreover, genetic polymorphisms are associated with progression

to severe disease (COVID-19 Host Genetics Initiative 2021). Cell-intrinsic innate viral

sensors and antiviral cytokines, including type I and type III interferons (IFNs), orchestrate

the control of SARS-CoV-2 infection (Kim and Shin 2021). Inherited mutations of genes

involved in IFN induction and signalling and circulating auto-antibodies (AABs) that

neutralize type I IFNs have been found to predispose infected individuals to severe COVID-

19 (Bastard et al. 2020; Zhang et al. 2020), presumably by contributing to an ineffective

immune response with delayed or abolished type I IFN signaling. Neutralizing type I IFN-

AABs are present in 6-17% of hospitalized COVID-19 patients with severe pneumonia

(Bastard et al. 2020; Koning et al. 2021; Troya et al. 2021) and 11-19% in critically ill

COVID-19 patients (Bastard et al. 2020; van der Wijst et al. 2021; Goncalves et al. 2021),

greatly exceeding estimated prevalences of around 0.33% (Bastard et al. 2020) in uninfected

individuals. Intriguingly, while neutralizing IFN-AABs in patients with autoimmune

polyendocrine syndrome type 1 (APS-1) can associate with a severe course of SARS-CoV-2

infection (Bastard et al. 2021; Beccuti et al. 2020; Carpino et al. 2021; Ferré et al. 2021;

Lemarquis et al. 2021), their mere presence does not inevitably lead to severe disease

(Meisel et al. 2021). A recent global multi-cohort study reports prevalence of neutralizing

IFN-AABs in 4% of uninfected individuals over 70 years of age, suggesting that IFN-AABs

may pre-exist in some individuals that develop a critical course of COVID-19 (Bastard et al.

2021). Thus, we reasoned that IFN-AABs may serve as biomarkers that could, in conjunction

with other clinical parameters, help to predict risk for developing severe COVID-19 and to

stratify patients for specific therapies.

Specific therapies may comprise the administration of recombinant IFN-β or

therapeutic plasma exchange (TPE). However, the clinical benefit of TPE and other

approaches remains to be defined and requires studies involving large numbers of patients.

With IFN-AABs present in up to 18% of deceased COVID-19 patients (Bastard et al. 2021)

and given the limited therapeutic options for severely affected COVID-19 patients, testing

specific therapeutic approaches is of high urgency, yet clinically implementable strategies for

rapid   and   early   identification   of   IFN-AAB-positive   patients   upfront   are   missing.

METHODS

Study cohorts and data collection

Patients were recruited and data and sample collection was performed within one of four

prospective observational studies conducted at Charité - Universitätsmedizin Berlin,

Germany (Cohort A, (Kurth et al. 2020)), Inselspital Universitätsspital Bern, Switzerland

(Cohort B), Universitätsklinikum Freiburg, Germany (Cohort C) and Universitätsklinikum

Heidelberg, Germany (Cohort D). For this analysis, all patients with a maximum WHO score

of 3-8 were included from Cohorts A-C (henceforth summarized as cross-sectional cohorts,

CSC). For cohort D (therapeutic plasma exchange cohort, TPEC) only patients who

underwent therapeutic plasma exchange for treatment of COVID-19-associated

hyperinflammatory syndrome at the Department of Internal Medicine IV of Heidelberg

University Hospital, Germany (Nusshag et al. 2021) and supplementary methods) were

retrospectively selected. All TPE procedures were performed in accordance with the German

Medical Devices Act ("Medizinproduktegesetz"). Healthy controls were recruited from a study

on SARS-CoV-2 exposition in health care workers (HC cohort). Samples from APS-1

patients were obtained from a published study (Meisel et al. 2021) and published values are

shown here for reference. All studies were conducted according to the Declaration of

Helsinki and Good Clinical Practice principles (ICH 1996).

Detection of IFN-AABs by reverse ELISA

IFN-AABs were detected using an electrochemiluminescence immunoassay (ECLIA)-

platform (MSD, Rockville, U.S.), as described recently (Meisel et al. 2021). Briefly, MSD

GOLD 96-well small spot streptavidin SECTOR Plates (MSD) were washed with wash buffer

(MSD), and blocked with 150 µl blocking buffer (Thermo Fisher, Waltham, U.S.) per well at

4°C overnight. All further incubations were performed for 60 minutes at room temperature.

After blocking, plates were incubated with IFN-α2 (Merck Sharp & Dohme, Kenilworth, USA)

or IFN-ω (Peprotech, Rocky Hill, USA) linked to biotin (Thermo Scientific, Waltham, USA).

Next, plates were incubated with patients' sera following dilution at 1:100 in blocking buffer.

Cytokine AABs were detected using a monoclonal mouse antibody to human IgG (D20JL-6,

MSD). After incubation and washing, 150 μl of read buffer (ReadBufferT (4x), MSD) were

added, incubated for ten minutes at room temperature, and plates were analyzed using the

MESO QuickPlex SQ 120 analyzer (MSD). Data are shown as light signal counts (LSC).

Competition assays

All sera whose IFN-α2-AABs and/or IFN-ω-AABs levels exceeded the 97.5th percentile of

AAB levels of the analysed HCW sera and 27 samples that scored close to, but below this

cut-off were assessed by competition assay using unbiotinylated IFN-α2 or IFN-ω. The sera

of interest were diluted 1:100 with blocking buffer and incubated overnight at 4°C with

2.5mg/ml, 0.025mg/ml, and 0.00025mg/ml unbiotinylated IFN-α2 or IFN-ω. After incubation,

reverse ELISA was performed, as described above. IFN-AABs in a given serum scored

specific when preincubation with the highest concentration of IFN-α2 or IFN-ω resulted in an

at least four-fold reduction of LSC in comparison to analysis of the identical serum without

IFN-α2 or IFN-ω pre-incubation.

Virus infection-based neutralization assays

Calu-3 cells were pre-incubated with 1% human serum in the presence or absence of 200-

400 IU/ml IFN-α2a (Roferon®-A, Roche) or 20-50 ng/ml IFN-ω (PeproTech). After 24 hours,

IFN and serum were removed and cells were infected with SARS-CoV-2 at MOI 0.01. Virus

inoculum was removed after one hour, cells were washed with PBS and 100 μl medium was

added per well. 24 hours post-infection, cell culture supernatant was collected for viral RNA

quantification by RT-PCR and infectious titer determination by plaque assay.

Cytokine and chemokine measurements

Cytokines and chemokines from a subset of patients from cohort A were analyzed using

Quanterix' single molecule array technology or multiplex ECLIA.

RESULTS

We analyzed 430 serum samples collected within four independent observational clinical

studies on COVID-19 for IFN-AAB positivity (Table 1). Median age of patients in the CSC

was 61 years (IQR 52-71) and 72.2% (291/403) were male. Median Charlson Comorbidity

Index (CCI) was 3 (IQR 1-4). 27 patients with critical disease course (median max. WHO

score 7 (IQR 7-8)) who underwent TPE as compassionate use were selected retrospectively

from center D (TPEC). Median age of patients in the TPEC was 65 years (IQR 56-72), 74.1

% (20/27) were male and median CCI was 4 (IQR 3-5). All patients from the TPEC required

invasive mechanical ventilation (IMV), 77.8% (21/27) hemodialysis, one patient was treated

with extracorporeal membrane oxygenation (ECMO), and 13 out of 27 (48.2%) patients died

despite maximum care. 667 serum samples from a healthy cohort (HC) consisting of health-

care workers (Table S1) were screened for the presence of neutralizing IFN-AABs to set the

cut-off for IFN-AAB positivity.

Prevalence of AABs against IFN-α2 and IFN-ω in patients with COVID-19

We first aimed to establish a sensitive screening assay for type I IFN-AABs. To this end, we

first screened samples of our HC for prevalence of AABs against IFN-α and/or IFN-ω by

ELISA. Samples were considered positive when the respective LSC value exceeded the

97.5th percentile of AAB levels of the analyzed sera from the HC (cut-off for IFN-α = 1980

LSC, IFN-ω = 1961 LSC). We then screened sera obtained at the peak of the disease from

patients of cohorts A-C (CSC) and cohort D (TPEC). The proportion of ELISA-positive

patients in the CSC was 5.0% (20/403) for IFN-☐ AABs and 4.2% (17/403) for IFN-ω AABs.

It was significantly higher in cohort D (TPEC) (IFN-α AABs 18.5%, 5/27, p = 0.0035 and IFN-

ω AABs 14.8%, 4/27, p = 0.0132), as expected (Fig. 1a, Fig. S1). Some sera displayed

values approaching or equaling those detected in sera from patients with APS-1, a genetic

disease involving the generation of high titer neutralizing type I IFN-AABs (Fig. 1a, (Meisel et

al. 2021)).

Nonspecific binding is a common phenomenon in immunoassays for the detection of

AABs, and high levels of inflammatory parameters such as CRP correlate with non-specific

binding of (auto-)antibodies (Güven et al. 2014). Therefore, we probed the specificity of all

samples exceeding the 97.5th percentile of the IFN-AAB ELISA and 27 samples that scored

right below the cut-off in a competition assay (Fig. 1b). Based on the assumption that sera

scoring below the cut-off did not contain IFN-AABs, we established a prevalence of specific

IFN-□-AAB of 3.2% (13/403) and of specific IFN-ω-AAB of 1.7% (7/403) in the CSC (Fig. 1a-

b). Cohort D (TPEC) showed 18.5% of sera specifically binding IFN-□ (5/27) and 14.8% for

IFN-ω (4/27) or both (14.8%, 4/27) (Fig. 1a-b). As expected, all tested samples scoring below

the 97.5th percentile in the ELISA scored negative in the specificity assay with the exception

of one patient (B044) who scored slightly below the cut-off in the IFN-ω-AAB assay.

Importantly, none of the tested sera from the HC displayed antibodies that specifically bound

IFN-□ or IFN-□.

IFN-AABs neutralize exogenous IFN in a virus infection-based assay

We next analyzed whether the presence of detectable and specifically IFN-binding AABs

corresponded to a functional neutralization of IFN during infection. To this end, we applied a

previously established assay of IFN-based inhibition of SARS-CoV-2 infection of the

immortalized lung cell line Calu-3 (Meisel et al. 2021), which we consider the gold standard

for analysis of IFN neutralization. We tested the extent to which sera neutralize the antiviral

activity of type I IFNs, resulting in efficient infection despite presence of IFNs. We tested all

ELISA-positive sera as well as 102 ELISA-negative sera as a reference. 13 (3.2 %) and 8

(2.0 %) of the 403 sera of the CSC specifically neutralized exogenous IFN-□ and IFN-□,

respectively, as judged by PCR-based quantification of SARS-CoV-2 genomic RNA in the

supernatant and plaque assays that quantify infectivity of virus progeny (Fig. 2a-b, Fig. S1).

In cohort D (TPEC), 5 (18.5%) and 3 (11.1%) of 27 sera neutralized IFN-□ and IFN-□

activity, respectively (Fig. 2a-b, Fig. S1).

Strikingly, among competition assay-positive samples, 18 (100%) of 18, and 9 (90%)

of 10 sera displayed IFN-□ and IFN-□-neutralizing activity, respectively, indicating that a

positive result in the competition assay associates with neutralization activity with a high

likelihood (Fig. 1c-d). Conversely, a negative result in the competition assay was predictive

of absence of neutralization ability. Specifically, among ELISA-positive, but competition

assay-negative sera, 0 (0%) of 23 and 0 (0%) of 29 sera were able to neutralize IFN-□ and

IFN-□, respectively. Among IFN-α AAB-ELISA-negative sera, we observed no IFN-

neutralizing activity. Interestingly, among IFN-ω AAB-ELISA-negative sera, we identified two

samples that neutralized IFN-ω despite scoring negative in the IFN-ω-AAB ELISA (i.e.,

having LSC counts below the 97.5th percentile cut-off) (Fig. 2b, d). The pronounced ability of

these exact two sera to neutralize IFN-α (Fig. 2a, c) was the reason why we included them in

the IFN-ω test, and suggests a potential cross-reactivity of IFN-α-AAB with IFN-ω.

Merging results from all three assays (Fig. 2c-d) revealed that an LSC value in the

screening ELISA of >35.639 (IFN-□) and >12.603 (IFN-□) predicted specific binding in the

competition assay and neutralization ability in the functional assay. Finally, the prevalence of

ten individual antiphospholipid-ABs did not differ between patients with and without

neutralizing IFN-AABs from cohort A (Fig. S2), suggesting that the presence of AABs is not

generally increased in IFN-AAB-positive patients.

Clinical phenotype of COVID-19 patients displaying type-I IFN-AABs

We next aimed to characterize the clinical phenotype of IFN-neutralizing AAB-positive

COVID-19 patients in the CSC at hospital admission, and to identify discriminatory markers

that may serve as pre-selection criteria for their early identification and stratification.

Interestingly, there were no statistically significant differences regarding clinical baseline

characteristics, including demographic criteria and pre-existing comorbidities between

patients with and without IFN-neutralizing AABs in the CSC using univariate analyses (Table

1). Yet, of all patients with available symptom records (cohorts A and C), the proportion of

patients who reported fever and required supplemental oxygen therapy within 72 hours from

admission was higher in patients with neutralizing IFN-AABs than in those without (fever:

100%, 12/12 versus 62.6% (234/374), p=0.0079 and oxygen: 100% (12/12) versus 76.0%

(250/329), p=0.0528).

Furthermore, patients with IFN-AABs for which respective data were available

displayed higher median values of C-reactive protein (CRP), procalcitonin, lactate

dehydrogenase (LDH), ferritin, total leukocyte and neutrophil count, as well as neutrophil-to-

lymphocyte ratio within the first three days of hospital admission compared to patients

without IFN-AABs (Fig. 3, Fig. S3). In addition, patients with neutralizing IFN-AABs showed

low levels of CD169/Siglec-1 expression on monocytes, a well-known type I IFN-response

marker (Fig. S4). Interestingly, there was a tendency towards a negative correlation between

CRP levels and CD169/Siglec-1 within 72 hours from hospital admission.

In IFN-AAB-positive patients, high quantities of neutralizing IFN-α2-AABs were

present both soon post-symptom onset and at the peak of disease

Next, we evaluated the temporal dynamics of IFN-AAB levels in sera from COVID-19

patients soon after symptom onset as compared to the peak of the disease. Available

samples obtained in cohort A, and in cohort D (TPEC) prior to TPE were analyzed (Fig. 4,

Fig. S5). In all patient sera with detectable neutralizing IFN-AABs at the peak of the disease,

early sera corresponding to ten (min. 4 - max. 20) days post-symptom onset (PSO)

contained abundant and neutralizing (Fig. 4, Fig. S5) IFN-AABs, suggesting that IFN-AABs

either existed prior to the infection or were generated very early PSO.

Cytokine and humoral responses to SARS-CoV-2 infection in IFN-AAB-positive

patients

We next aimed to identify potential quantitative and/or qualitative differences in cytokine

responses, viral load and seroconversion kinetics in IFN-AAB-positive as opposed to IFN-

AAB-negative patients. We analyzed serum cytokine levels in a subset of critical patients

(WHO max. 6-8) from cohort A. Patients with neutralizing IFN-AABs demonstrated

significantly higher levels of IFN-γ, and IFN-γ-induced protein 10 (IP-10) one to two weeks

PSO while monocyte chemoattractant protein-1 (MCP-1) and TNF-α concentrations were

similar compared to sera from patients without neutralizing IFN-AABs (Fig. S6). However,

levels equalized among the two groups at three to four weeks PSO. As expected, patients

with IFN-AABs had undetectable serum IFN-□ levels. Of note, by comparing upper-

respiratory tract swabs and sera from patients with and without IFN-AABs from all infected

cohorts, we failed to identify detectable differences in viral load level or decay over time (Fig.

S7a) and we found no evidence for a difference in duration until seroconversion PSO (Fig.

S7b). In conclusion, some cytokine responses were aberrantly elevated in patients with IFN-

AABs within the first two weeks PSO. However, they normalized at weeks three and four,

and viral RNA production and time to seroconversion remained indistinguishable from

patients without IFN-AABs.

Clinical Outcome of COVID-19 Patients with Neutralizing IFN-AABs

Neutralizing IFN-AAB-positive patients developed significantly higher max. WHO scores than

patients without neutralizing IFN-AABs (median max. WHO score 8 (IQR 8-8) vs 6 (IQR 4-7),

respectively; p<0.0001, Fig. 5a). All patients with neutralizing IFN-AABs in the CSC required

IMV (13/13, 100%), compared to 44.5% (166/372) in patients without IFN-AABs (p<0.0001,

Fig. 5b). Similarly, the proportion of neutralizing IFN-AAB-positive patients requiring

haemodialysis and/or ECMO was markedly higher than in those without IFN-AABs

(haemodialysis: 69.2%, 9/13 versus 26.6%, 98/369, p = 0.0008, ECMO: 46.2%, 6/13 versus

15.6%, 58/372, p = 0.0036, Table 1). Twelve out of thirteen neutralizing IFN-AAB-positive

patients (92.3%) died in hospital compared to 19.1% (71/372) of patients without IFN-AABs

(p<0.0001) in the CSC (Table 1). Median survival of patients with neutralizing IFN-AABs was

28 days (IQR 22-65 days). The probability of surviving to 150 days PSO is 81.3% (300/369)

for the patients from the non-neutralizing group, as opposed to 7.7% (1/13) for the patients

of the neutralizing IFN-AAB-positive group (Fig. 5c). Conclusively, IFN-AAB positivity was

associated with severe disease trajectories of COVID-19 and a worse clinical outcome in our

cohorts.

Inter-individual effect of therapeutic plasma exchange on IFN-AABs and SARS-CoV-2

antibodies.

Cohort D (TPEC) allowed us to compare trajectories of IFN-AAB-positive patients

undergoing TPE to those not undergoing TPE. Criteria for initiation of TPE were: presence of

ARDS requiring IMV and/or vasopressor-dependent circulatory shock, clinical and laboratory

features of a COVID-19 associated immunopathology with elevated D-dimers and ferritin

levels, persistent and refractory fever ≥ 38.5°C without conclusive pathogenic evidence and

despite anti-infectious treatment. TPE was first initiated without prior screening for IFN-AABs

within a median of six days (IQR 1-10) after hospital admission and the median number of

TPE sessions per patient was 3 (IQR 2-5). TPE was performed using a continuous-flow

centrifugation blood cell separator. Plasma with enclosed cytokines and immunoglobulins

are separated from blood cells by gravity due to different densities of the respective blood

components (Nusshag et al. 2021).

Focusing on patients of the WHO group 6-8, survival of IFN-AAB-negative patients in

the CSC cohorts and TPEC was similar (p = 0.34). The proportion of neutralizing IFN-AAB-

positive patients from the TPEC that survived in hospital was higher than of those patients

from the CSC who did not undergo TPE (60%, 3/5 patients from the TPEC survived versus

7.7%, 1/13 patients from the CSC, p=0.0412) (Fig. 6a).

Longitudinal analysis of sera revealed that three (D011, D017, D018) out of five

patients responded to TPE with decreasing IFN-AAB levels below the cut-off and to a level

that coincided with absence of neutralizing activity. In addition, a sustained reduction of IFN-

AAB quantities was achieved only by repetitive TPE (Fig. 5b, Fig. S8). In contrast to IFN-

AABs, quantities of SARS-CoV-2 Spike IgG and IgA were less, if at all, affected by TPE

(note the logarithmic scale for IFN-AABs versus the linear scale for SARS-CoV-2-IgG/IgA).

Overall, our findings in a limited number of patients suggest that TPE could positively affect

the survival of critically ill IFN-AAB-positive patients. This needs to be corroborated in future,

adequately powered clinical investigations. Potentially, a sustained and significant reduction

of peripheral IFN-AAB levels must be achieved to prevent death.

Diagnostic algorithm for rapid identification of neutralizing IFN-AAB-positive patients

Given that IFN-AABs are risk factors for a worse clinical outcome in hospitalized patients

with COVID-19, future rapid identification of IFN-AAB-positive patients after hospital

admission seems key for the potential implementation and success of specific interventions

such as TPE. In the CSC cohort, the patient number needed to screen (NNS) without

preselection in order to identify one patient with neutralizing IFN-AAB was 31.0 (403/13). We

hypothesized that applying clinical pre-selection criteria which strongly co-present with the

neutralizing IFN-AAB positivity diminishes the NNS. Using univariate analyses, we

established that temperature (>38.5 °C or self-reported) before or upon hospital admission

and the need for supplemental oxygen within the first 72 hours after admission correlated

best with presence of IFN-AAB positivity in the screening assay in all hospitalized COVID-19

patients (fever: p=0.0079; supplemental oxygen p=0.0528). Using fever and need for

supplemental oxygen, the NNS was reduced to 15.6 (172/11). Selection of patients

exceeding the cut-off for ELISA positivity (in our cohort 97.5th percentile of the HC) for

further testing by competition assay would adjust the NNS in the competition assay to 1.4

(15/11). Therefore, in order to increase sensitivity, we propose to consider the need for

supplemental oxygen within 72 hours after admission and fever as pre-selection criteria for

patients that undergo ELISA screening (Fig. 7).

Sera from all patients exceeding the LSC value of 35,639 (13/13) in the screening

ELISA assay demonstrated neutralizing activity against IFN-α (Fig. 2c). We therefore

propose to conduct the IFN neutralization assay only in case of an LSC value lower than

35,639, whereas patients with sera exceeding this value can be considered positive for

neutralizing IFN-AABs without further testing (Fig. 7).

DISCUSSION

IFN-AABs strongly associate with adverse clinical outcome of SARS-CoV-2 infection

(Bastard et al. 2020; Koning et al. 2021; Troya et al. 2021; Goncalves et al. 2021; Bastard et

al. 2021; Lemarquis et al. 2021; de Prost et al. 2021). Unfortunately, the inability to identify

IFN-AAB-positive COVID-19 patients early enough before deterioration delays their access

to specific, yet untested therapies that may require implementation directly upon hospital

admission. In several studies, detection and quantification of IFN-AABs in sera from COVID-

19 patients relies on ELISA and multiplex particle-based assay. While these assays are

amenable to high-throughput and are highly sensitive, they result in a small proportion of

false-positive results (Güven et al. 2014; Sahud et al. 2012), highlighting the ongoing need to

reanalyze positive-tested patient material in functional assays demonstrating the

neutralization activity. However, such assays are sophisticated and time-consuming. They

include luciferase-based interferon-stimulated response element (ISRE) promoter reporter

assays (Troya et al. 2021; Goncalves et al. 2021), flow cytometry-based analyses of STAT

phosphorylation (Bastard et al. 2020; Goncalves et al. 2021; Meisel et al. 2021), and virus

infection-based assays (de Prost et al. 2021; Meisel et al. 2021). The latter allows probing

the activity of the IFN-AABs in the context of infection-inhibitory concentrations of IFN-α and

IFN-ω. Here, we applied and cross-validated previously established assays comprising an

ELISA for sensitive identification, a specificity-validating competition assay, and a functional

neutralization assay (Meisel et al. 2021) using a large collection of serum samples obtained

from three cross-sectional cohorts. Equally important, we identified clinical parameters that

co-present with IFN-AAB positivity at hospital admission. Combination of clinical parameters

and targeted diagnostic testing results in an algorithm for early, sensitive, specific

identification of IFN-AAB-positive patients which can be easily introduced into clinical routine

diagnostics.

Surprisingly, sera from two patients were found to neutralize exogenous IFN-ω

despite negative ELISA results. Presence of neutralization activity in the absence of

detectable IFN-AABs has been reported (Bastard et al. 2021). Explanations for this

phenomenon could include technical aspects of the detection method, including the

possibility that IFN-AABs may be concealed by the binding of the cytokine to the plate or

biotinylation of the cytokine (Meyer et al. 2016).

Here, we calibrated our ELISA cut-off based on the 97.5th percentile in a cohort of

uninfected individuals. Although this strategy may be inexact, the absence of prevalence

IFN-AABs in a cohort of younger and predominantly female healthcare workers supports an

age-dependent increase of IFN-AAB prevalence in uninfected individuals (Bastard et al.

2021). Furthermore, the prevalence of 3.2% (13/403) of patients with neutralizing AABs

against IFN-α and/or IFN-ω in our cross-sectional patient cohort (median max. WHO score

6) and 18.5% (5/27) in critically affected patients (median max. WHO-Score 7) is in line with

reported prevalences of 6-17% in severely (Bastard et al. 2020; Koning et al. 2021; Troya et

al. 2021; van der Wijst et al. 2021) and 11-19% in critically ill (Bastard et al. 2020; van der

Wijst et al. 2021; Goncalves et al. 2021) individuals with COVID-19. Moreover, IFN-AAB

positivity is associated with a worse clinical outcome and a decreased survival probability of

hospitalized patients in our cohorts, confirming previous reports (Bastard et al. 2020; Koning

et al. 2021; Troya et al. 2021; Goncalves et al. 2021; Bastard et al. 2021; Lemarquis et al.

2021; de Prost et al. 2021). Interestingly, a single study to date (Abers et al. 2021)

suggested that survival was not adversely affected by the presence of type-I IFN-AABs,

while confirming the widely accepted association with an increased risk of admission to the

intensive care unit.

We failed to identify a clear association of IFN-AABs with previously described

demographic parameters in our cross-sectional cohort, including male sex or advanced age

(Bastard et al. 2020, 2021), probably due to the relatively limited sample size of our cohorts.

However, the presence of neutralizing IFN-AABs was associated with fever and need for

supplementary oxygen within 72 hours post hospital admission, as well as with elevated

soluble and cellular markers of acute-phase reaction including elevated levels of CRP,

procalcitonin, LDH and ferritin and elevated total neutrophil and leukocyte counts within the

first three days of admission in our CSC. Higher CRP values constitute a biomarker for a

severe disease course, are included in a widely-used clinical risk score for mortality of

COVID-19 (Knight et al. 2020), and associate with neutralizing IFN-AABs along with lower

lymphocyte counts in severely affected patients (Troya et al. 2021). As hospital admission

and thus clinical deterioration occurred at a median of five days PSO in the CSC, we

propose fever and need for supplemental oxygen therapy up to 72 hours post hospital

admission as suitable and simple clinical criteria to identify patients at risk for a severe

disease course.

Our ability to detect IFN-AABs as early as four days PSO in sera from most patients

that present with IFN-AABs at the peak of their disease suggest that they were present prior

to the infection, or alternatively, but less likely, were induced very early post infection. Our

data are in agreement with recently demonstrated presence of IFN-AABs at the day of

hospital admission (de Prost et al. 2021) and in 4% of uninfected individuals >70 years old

(Bastard et al. 2021), underlining the idea that they can serve as biomarkers for

predisposition for a severe course of SARS-CoV-2 infection. Future studies are required to

elucidate the biological mechanisms that lead to elicitation of IFN-AABs in an age-dependent

manner.

Currently discussed therapeutic options tailored to IFN-AAB-positive COVID-19

patients comprise mainly TPE and IFN-β administration. TPE in the context of COVID-19

has been analyzed in individual case reports and case-control studies, including IFN-AAB-

positive and negative patients (Lemarquis et al. 2021; Nusshag et al. 2021; de Prost et al.

2021; Fernandez et al. 2020) and might efficiently remove soluble circulating Fc gamma

receptor-activating immune complexes (Ankerhold et al. 2021). TPE effectively decreased

circulating IFN-AAB, but not SARS-CoV-2 antibody concentrations in four IFN-AAB-positive,

severely ill patients (de Prost et al. 2021) and in a child with APS-1 suffering from severe

COVID-19 (Lemarquis et al. 2021). In our study, TPE reduced circulating IFN-AABs with

patient-specific efficiency and appeared to increase the chances of in-hospital survival,

underlining the rationale to initiate large-scale, adequately powered clinical trials in order to

corroborate the potential benefit of TPE in a general cohort of adult, critically ill COVID-19

patients. Interestingly, SARS-CoV-2-IgG and IgA quantities were less affected by TPE for

unknown reasons, which may include their rapid replenishment by highly abundant

plasmablasts or an extravascular-to-intravascular rebound since immunoglobulins have a

substantial extravascular distribution.

Given the reported absence of detectable IFN-β-AABs, IFN-β administration may

substitute neutralized IFN-α and -ω. While IFN-β therapy failed to result in a detectable

clinical benefit in the SOLIDARITY trial (WHO Solidarity Trial Consortium et al. 2021),

specifically IFN-AAB-positive patients may benefit from IFN-β therapy, a patient group that

might have been under-represented in this study. Furthermore, the benefit of IFN-β

administered by different routes should be systematically explored in this patient group.

CONCLUSIONS

Generation of comprehensive data on the clinical benefit of TPE and IFN-β therapy

are hampered by the difficulty of identifying IFN-AAB-positive patients for enrollment in large

clinical studies. Our diagnostic algorithm for their identification may support clinicians to

promptly identify patients carrying this additional risk factor for progression to severe disease

upon hospital admission. It is based on clinical preselection of patients presenting with fever

and need for supplemental oxygen therapy within 72 hours after admission to be further

screened by an ELISA-based assay for circulating IFN-AABs and an easily implementable

ELISA-based validation assay which may be skipped in case of high positivity in the

screening assay (Fig. 7). Screening criteria should be validated in the future in a large

prospective observational cohort. Ideally, methodological complex gold-standard assays can

be substituted by high-throughput diagnostic tests and stratification by clinical parameters,

facilitating clinical implementation. Our algorithm may allow identifying at-risk patients upon

presentation in the emergency room to be directly allocated to TPE and/or IFN-β therapy

within larger clinical trials, which are urgently needed to determine clinical effectiveness of

these targeted therapies.

Acknowledgements

We thank Mahtab Maleki for support with IFN-AAB ELISAs and competition assays. We

thank Patricia Tscheak, Marie Luisa Schmidt, and Tatjana Schwarz for support with sample

logistics and SARS-CoV-2 antibody testing. V.M.C is a participant of the Charité Clinician

Scientist program funded by Charité – Universitätsmedizin Berlin and the Berlin Institute of

Health. We thank Karin Reiter for the anti-phospholipid antibody measurements. We thank

the patients for their ongoing trust and collaboration. We thank the involved biobanks

(ZeBanC Berlin, FREEZE-Biobank Freiburg) for their support. DD and AL are members of

SFB1425, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) – Project #422681845. NS is partly funded by the Deutsche

Forschungsgemeinschaft (DFG, Research Grant No. SCHA1838/4-2).

Pa-COVID study Group

Set up study platform: Stefan Hippenstiel, Martin Witzenrath, Norbert Suttorp, Elisa T.

Helbig, Lena J. Lippert, Pinkus Tober-Lau, David Hillus, Sarah Steinbrecher, Sascha S.

Haenel, Alexandra Horn, Willi M. Koch, Nadine Olk, Rosa C. Schuhmacher, Katrin K.

Stoyanova, Lisa Ruby, Claudia Zensen, Mirja Mittermaier, Fridolin Steinbeis, Tilman

Lingscheid, Bettina Temmesfeld-Wollbrück, Thomas Zoller, Holger Müller-Redetzky,

Alexander Uhrig, Daniel Grund, Christoph Ruwwe-Glösenkamp, Miriam S. Stegemann,

Katrin M. Heim, Ralf H. Hübner, Bastian Opitz, Kai-Uwe Eckardt, Martin Möckel, Felix

Balzer, Claudia Spies, Steffen Weber-Carstens, Frank Tacke, Chantip Dang-Heine, Michael

Hummel, Georg Schwanitz, Uwe D. Behrens, Maria Rönnefarth, Sein Schmidt, Alexander

Krannich, Saskia Zvorc, Jenny Kollek and Christof von Kalle

Inclusion of patients and clinical data curation: Linda Jürgens, Malte Kleinschmidt, Sophy

Denker, Moritz Pfeiffer, Belén Millet Pascual-Leone, Luisa Mrziglod, Felix Machleidt,

Sebastian Albus, Felix Bremer, Jan-Moritz Doehn, Tim Andermann, Carmen Garcia, Philipp

Knape, Philipp M. Krause, Liron Lechtenberg, Yaosi Li, Panagiotis Pergantis, Till Jacobi,

Teresa Ritter, Berna Yedikat, Lennart Pfannkuch, Christian Zobel, Ute Kellermann, Susanne

Fieberg, Laure Bosquillon de Jarcy, Anne Wetzel, Christoph Tabeling, Markus C. Brack,

Moritz Müller-Plathe, Jan M. Kruse, Daniel Zickler, Andreas Edel, Britta Stier, Roland Körner,

Nils B. Müller, and Philipp Enghard, Lucie Kretzler, Lil A. Meyer-Arndt, Linna Li, and Isabelle

Wirsching

Biobanking and -sampling: Denise Treue*, Dana Briesemeister*, Jenny Schlesinger*,

(*Central Biobank Charité/BIH; ZeBanC) Birgit Sawitzki, Lara Bardtke, Kai Pohl, Philipp

Georg, Daniel Wendisch, Anna L. Hiller, Sophie Brumhard, Marie Luisa Schmidt, Leonie

Meiners, and Patricia Tscheak

Author contributions

Conceptualization: Florian Kurth; Horst von Bernuth; Christian Meisel; Christine Goffinet

Methodology: Bengisu Akbil; Tim Meyer; Olga Staudacher

Formal analysis and investigation: Bengisu Akbil; Tim Meyer; Paula Stubbemann; Charlotte

Thibeault; Olga Staudacher; Jenny Jansen; Barbara Mühlebach; Jan-Moritz Doehn;

Christoph Tabeling; Christian Nusshag, Cédric Hirzel; David Sökler Sanchez; Daniel

Duerschmied, Achim Lother; Nils Schallner; Jan Nikolaus Lieberum; Dietrich August;

Siegbert Rieg; Valeria Falcone; Uwe Kölsch; Nadine Unterwalder; Thibaud Spinetti; Joerg

Schefold; Thomas Dörner; Victor M. Corman; Uta Merle

Writing - original draft preparation: Paula Stubbemann; Charlotte Thibeault; Christine

Gofffinet

Writing - review and editing: Bengisu Akbil; Tim Meyer; Paula Stubbemann; Charlotte

Thibeault; Olga Staudacher; Daniela Niemeyer; Barbara Mühlemann; Christian Nusshag;

Daniel Duerschmied, Achim Lother; Terry C. Jones; Klaus Warnatz; Thibaud Spinetti; Joerg

C. Schefold; Leif Sander; Victor M. Corman; Uta Merle; Florian Kurth; Horst von Bernuth;

Christian Meisel; Christine Goffinet

Funding acquisition: Joerg Schefold; Klaus Warnatz; Uta Merle; Florian Kurth; Horst von

Bernuth; Christian Meisel; Christine Goffinet

Resources: Daniela Niemeyer, Alexandra Nieters; Christian Drosten; Leif Sander; Florian

Kurth; Horst von Bernuth; Christian Meisel; Christine Goffinet

Supervision: Hartmut Hengel; Ralf-Harto Hübner; Norbert Suttorp; Klaus Warnatz; Victor M.

Corman; Florian Kurth; Horst von Bernuth; Christian Meisel; Christine Goffinet

Funding

This work was supported by the Innovationsfond of Labor Berlin (to H.V.B, C.M., and C.G.);

by funding from the Deutsche Forschungsgemeinschaft (DFG) Collaborative Research

Centre CRC900 "Microbial Persistence and its Control," project number 158989968, project

C8 (awarded to C.G.); by funding from the Berlin Institute of Health (BIH) (to C.G.). T.C.J. is

in part funded through NIAID-NIH CEIRS contract HHSN272201400008C. Parts of the work

were funded by the European Union's Horizon 2020 research and innovation program

through project RECOVER (GA101003589) to C.D.; the German Ministry of Research

through the projects VARIPath (01KI2021) to V.M.C., and NaFoUniMedCovid19 - COVIM,

FKZ: 01KX2021 to C.D., V.M.C., L.E.S., F.K. and H.H. The Pa-COVID-19 Study is funded by

BIH and the German Ministry of Research NaFoUniMedCovid19 NUM-NAPKON FKZ:

01KX2021. The COV-IMMUN study is supported by the Berlin University Alliance and the

BIH Clinical Study Center (BIH-CSC). KW has received funding by the Deutsche

Forschungsgemeinschaft (DFG, Research Grant No. WA 1597/6-1 and WA 1597/7-1).

Disclosure of Conflicts of interest

V.M.C is named together with Euroimmun GmbH on a patent application filed recently

regarding the diagnostic of SARS-CoV-2 by antibody testing. Technische Universität Berlin,

Freie Universität Berlin and Charité - Universitätsmedizin have filed a patent application for

siRNAs inhibiting SARS-CoV-2 replication with D.N. as co-author. J.C.S, T.S (full

departmental disclosure): the department of Intensive Care Medicine has/had research

and/or development/consulting contracts with (full disclosure) Orion Corporation, Abbott

Nutrition International, B. Braun Medical AG, CSEM SA, Edwards Lifesciences Services

GmbH/SA, Kenta Biotech Ltd, Maquet Critical Care AB, Omnicare Clinical Research AG,

Phagenesis Ltd, Cytel, and Nestlé. No personal financial gains resulted from respective

development/consulting contracts and/or educational grants.

Research involving human participants

All studies were conducted according to the Declaration of Helsinki and Good Clinical

Practice principles (ICH 1996).

The Pa-COVID-19 study was approved by the ethics committee of Charité –

Universitätsmedizin Berlin (EA2/066/20) and is registered in the German and WHO

international clinical trials registry (DRKS00021688).

The Covimmun-study was approved by the ethics committee of Charité –

Universitätsmedizin Berlin (EA1/068/20).

The study "Evaluation der humoralen und zellulären Immunantwort und der

thromboembolischen Komplikationen bei Infektion mit dem neuartigen Coronavirus

(nCoV19)" was approved by the Kantonale Ethikkommission KEK, Bern, Switzerland, Nr.

2020-00877 and is registered under the name "Characterizing the Immune Response and

Neuronal Damage in COVID-19" at http://clinicaltrials.gov (NCT04510012).

The three studies (COVID-19 Register Freiburg; FREEZE-Covid19 cohort and Biomarker

COVID-19) were approved by the ethics committee of the University of Freiburg (EK-FR

153/20; FREEZE-Biobank EK-FR 383/19; EK-FR 225/20) and are registered in the German

and WHO international clinical trials registry (DRKS00021522 for Biomarker COVID-19).

The Covid-Immun study was approved by the local Ethics Committee of the Medical Faculty

of Heidelberg (S148/2020) and is registered in the German clinical trials registry

(DRKS00021810).

Informed consent

Informed consent was obtained from all individual participants included in the studies.

Data Availability

The datasets generated during and/or analysed during the current study are available from

the corresponding authors on reasonable request.

Legends

Figure 1. Prevalence of AABs against IFN-α2 and IFN-ω in patients with COVID-19.

(**a**) ECLIA-based assay for detection of IgG AABs against IFN-α2 and IFN-ω in sera from

hospitalized patients with COVID-19 from four different university hospital cohorts (Center A,

n=266; Center B, n=50; Center C, n=87; Center D, n=27), in patients with APS-1 (n=6), and

healthy health care workers (HC) without documented SARS-CoV-2 infection (n=667).

Dotted lines indicate the 97.5th percentile of the ECLIA assay LSC in sera from the HC

cohort. Dots indicate samples containing AABs scoring specific (red) or unspecific (blue) for

IFN-α2 and IFN-ω binding in the competition assay (see b), respectively. The prevalence of

sera with specifically binding type I IFN-AABs in each cohort is given in percent.

(**b**) Specificity of the ECLIA assay signal for IFN-α2- and IFN-ω-AABs was tested in an

competition assay by preincubation of sera with increasing concentrations of unlabeled IFN-

α2 and IFN-ω protein (0 - 2.5 μg/ml) before analysis. Samples showing a decrease in assay

signal by at least 75% in the presence of the highest competitor concentration were defined

as specific for type I IFN antibody reactivity and are indicated with red lines (IFN-α2 n=18,

IFN-ω n=11). Samples showing no decrease in the presence of excess unlabeled type I IFN

protein were regarded as unspecific for type I IFN antibody reactivity and are indicated with

blue lines (IFN-α2 n=62, IFN-ω n=39).

Figure 2. IFN-AABs neutralize exogenous IFN in a virus infection-based assay.

(**a-b**) Selected sera were analyzed for IFN neutralization activity in a SARS-CoV-2 infection-

based assay. The ability of individual sera to neutralize exogenous IFN-α2 (a) and IFN-ω (b)

is shown by the rescue of susceptibility to infection as judged by quantification of viral RNA

(x-axis) and infectivity (y-axis) in the supernatant. The infection condition in the absence of

serum and IFN is set to 1.

(**c-d**) The LSC value for individual sera, grouped into non-neutralizing and neutralizing sera.

Dots indicate sera containing AABs scoring specific (red) or unspecific (blue) for IFN-α2 and

IFN-ω binding in the competition assay (see b), respectively. Black dots indicate samples

that scored below the threshold of the ELISA. Black dotted lines indicate the 97.5th

percentile of the ECLIA assay LSC in sera from the healthy health care workers (HC) cohort

(see Fig. 1). Neutralization ability of IFN-α and IFN-ω can be predicted at 100% for sera

displaying LSCs above the respective red dotted lines (IFN-α: 35,639; IFN-ω: 12,603).

Figure 3: Clinical phenotype of COVID-19 patients displaying type-I IFN-AABs.

Values of C-reactive protein (CRP), procalcitonin, ferritin, lactate dehydrogenase (LDH),

absolute leukocyte and neutrophil count and neutrophil-to-lymphocyte ratio (NLR) of patients

with (N=5-6) and without neutralizing IFN-AABs (N=200-265) from the cross-sectional cohort

(CSC, all WHO scores). For each patient, the first available parameter within 72 hours of

hospital admission is shown. Statistical testing was performed with Mann-Whitney *U* test.

Figure 4. In IFN-AAB-positive patients, high quantities of neutralizing IFN-α2-AABs

were present both soon PSO and at the peak of disease.

(**a**) Time course of antibody quantities in patient sera that scored IFN-AAB positive at the

peak of disease (N=8). The dotted line indicates the 97.5th percentile of the ECLIA assay

LSC in sera from the HC cohort (see Fig. 1).

(**b-c**) The ability of the sera to neutralize exogenous IFN-α2 is shown by the rescue of

susceptibility to infection as judged by quantification of viral RNA (b) and infectivity (c) in the

supernatant. The infection condition in the absence of serum and IFN is set to 1.

Figure 5: Clinical outcome of COVID-19 patients with neutralizing IFN-AABs.

(**a**)   Median max. WHO score in hospital. Statistical testing was performed using the Mann-

Whitney *U* test.

(**b**) Proportion of patients requiring invasive mechanical ventilation (IMV) after hospital

admission. Statistical testing was performed using the Chi-Square test.

(**c**) Probability of survival of patients with and without neutralizing IFN-AABs from the cross-

sectional cohort (CSC) from symptom onset until discharge (up to 150 days), death or

transferral (p<0.0001). Statistical testing was performed using a log-rank test.

Neutralizing (N=13), non-neutralizing (N=372).

Figure 6. Inter-individual effect of therapeutic plasma exchange on IFN-AABs and

SARS-CoV-2 antibodies.

(**a**) Probability of survival of neutralizing IFN-AAB-positive and -negative patients with critical

COVID-19 (max. WHO score 6-8) with and without plasma exchange (CSC and TPEC) from

symptom onset until discharge, death or transferral (p=0.04, neutralizing CSC versus

neutralizing TPEC). Statistical testing was performed using a log-rank test. Neutralizing CSC

(N=13), non-neutralizing CSC (N=184), neutralizing TEPC (N=5) and non-neutralizing TPEC

(N=22).

(**b**) Antibody profile in serum from individual COVID-19 patients of the TPEC subjected to

plasma exchange. The quantity of IFN-α2- and IFN-ω-AABs, SARS-CoV-2-IgG and -IgA,

and the IFN-α2 and IFN-ω neutralization status are given for various time points. The patient

identifier is given in red. Viral load profiles were only available for patients D011 and D018

and are shown in Supplementary Figure 7.

Figure 7. Diagnostic algorithm for rapid identification of neutralizing IFN-AAB-positive

patients.

The number needed to screen (NNS) is based on results from the cross-sectional cohort

(CSC). ELISA for IFN-AAB detection was considered to be positive if it exceeded the 97.5th

percentile of the healthy control cohort. (1) NNS of all hospitalized Covid-19 patients without

preselection was 31.0 (403 patients in total, 13 patients with neutralizing IFN-AABs). (2)

Prescreening of patients using the clinical criteria of fever at admission and need for

supplemental oxygen within the first 72 hours after hospitalization diminished the NNS in the

IFN-AAB ELISA (3) by half, to 15.6 (172/11). For patients identified as positive in the

screening ELISA, the NNS in the competition assay to confirm the presence of IFN-specific

AABs is reduced to 1.4 (15/11) (4). For patients with high-titer IFN-AABs (light signal count >

35.639), the competition assay can be omitted. For patients highly positive in the IFN-AAB

ELISA and those with specific results in the competition assay, we propose to consider

therapeutic plasma exchange within clinical studies (5).

	Individual cohorts				CSC, Center A-C			
	Center A, Berlin cohort	Center B, Bern cohort	Center C, Freiburg cohort	Center D, Heidelberg cohort (TPEC)	Center A-C	IFN-AB Neutralizing	IFN-AB Non-neutralizing	p-value
Number of patients	266	50	87	27	403	13	390	/

IFN-AAB
  Neutralizing IFN-alpha
  Neutralizing IFN-omega
	2.6 (7/266)
	2.6 (7/266)
	1.1 (3/266)
		6.0 (3/50)
		6.0 (3/50)
		6.0 (3/50)
			3.5 (3/87)
			3.5 (3/87)
			2.3 (2/87)
				18.5 (5/27)
				18.5 (5/27)
				11.1 (3/27)
					3.2 (13/403)
					3.2 (13/403)
					2.0 (8/403)
						100 (13/13)
						100 (13/13)
						61.5 (8/13)
							/	/

Age (Median, IQR, available n)	61 (50-71), 266	67.3 (56.8-74.5), 50	59 (53-67), 87	65 (56-72), 27	61 (52-71), 403	69.4 (52.5-75.6), 13	61.0 (52-70.2), 390	0.19

Sex
  Female
  Male
	27.8 (74/266)
	72.2 (193/266)
		18.0 (9/50)
		82.0 (41/50)
			33.3 (29/87)
			66.7 (58/87)
				25.9 (7/27)
				74.1 (20/27)
					27.8 (112/403)
					72.2 (291/403)
						15.4 (2/13)
						84.6 (11/13)
							28.2 (110/390)
							71.8 (280/390)
								Male:
								0.31

BMI (kg/m^2, Median, IQR, available n)	28.4 (24,9-32,5), 245	27 (26-31), 47	27.7 (25.2-32.2), 56	31.5 (25.8-40.1), 27	28 (25-32.3), 348	27.4 (25.5-29.5), 10	28 (24.9-32.4), 338	0.73

Comorbidities
  CCI (Median, IQR, available n)
  Chronic heart disease (%)
  Chronic pulmonary disease (%)
  Diabetes (%)
  Obesity (%)
  Autoimmune disease (%)
	2 (1-3.75), 265
	58.9 (155/263)
	18.5 (47/254)
	26.7 (70/262)
	39.2 (96/245)
	2.8 (7/251)
		4 (2-6.5), 33
		30.3 (10/33)
		27.3 (9/33)
		42.4 (14/33)
		31.9 (15/47)
		2.0 (1/50)
			3 (2-5), 83
			23.0 (20/87)
			10.4 (9/87)
			26.4 (23/87)
			28.6 (16/56)
			5.7 (5/87)
				4 (3-5), 27
				77.8 (21/27)
				3.7 (1/27)
				44.4 (12/27)
				55.6 (15/27)
				3.7 (1/27)
					3 (1-4), 380
					48.3 (185/383)
					17.4 (65/374)
					28.0 (107/382)
					36.5 (127/348)
					3.4 (13/388)
						3 (1.5-4), 13
						46.2 (6/13)
						23.1 (3/13)
						23.1 (3/13)
						20.0 (2/10)
						0 (0/13)
							3 (1-4), 367
							48.4 (179/370)
							17.2 (62/361)
							28.2 (104/369)
							37.0 (125/338)
							3.5 (13/375)
								0.54
								0.69
								0.58
								0.54
								0.27
								0.72

Symptoms: Fever	57.0 (151/265)	80.0 (40/50)	77.5 (55/71)	85.2 (23/27)	63.7 (246/386)	100 (12/12)	62.6 (234/374)	0.0079
Days between symptom onset and admission (Median, IQR, available n)	6 (2-9), 233	4 (2-8), 50	6 (3-9), 66	6 (3-8), 27	5 (2-9), 349	4 (3-8), 11	5 (2-9), 338	0.81
Need for supplementary oxygen within first 72 h after admission	78.4 (189/241)	74.0 (37/50)	72 (36/50)	100 (12/12)	76,8 (262/341)	100 (12/12)	76.0 (250/329)	0.0528

IMV	46.2 (117/253)	60,0 (30/50)	39,0 (32/82)	100 (27/27)	46,4 (179/385)	100 (13/13)	44.5 (166/372)	0.0001
Days between symptom onset and intubation (Median, IQR, available n)	8 (4-10), 117	7 (3,75-10), 30	8 (3-11), 31	10 (7-15), 27	7 (4-10), 178	6 (5-11,5), 13	8 (4-10), 165	0.91
Days between hospital admission and intubation (Median, IQR, available n)	2 (0-4), 117	1,5 (0-3,25), 30	2 (1-7), 31	4 (0-6), 27	2 (0-4), 178	1 (0.5-4.5), 13	2 (0-4), 165	0.92
Length of ventilation in days (Median, IQR, available n)	32,5 (18,25-56,5), 116	10,5 (5-18,5), 30	16 (6-21), 23	24 (14-37), 27	23 (11-45), 169	20 (10.75-29.25), 12	24 (11-47,5), 157	0.40
Length of hospital stay in days (Median, IQR, available n)	20 (10-44), 262	13,5 (5-23,25), 50	16 (7-33), 85	41 (24-63), 27	19 (9-38), 397	25 (16-47), 13	17 (9-37,8), 384	0.045
Medication/treatment Dexamethasone Remdesivir	46.9 (123/262) 9,2 (16/174)	48.0 (24/50) 0 (0/50)	21.0 (17/87) 12.6 (11/87)	74,1 (20/27) 29.6 (8/27)	41.7 (164/393) 8.7 (27/311)	46.2 (6/13) 0 (0/13)	41.6 (158/380) 9.0 (27/300)	0.74 0.52
Renal replacement therapy	29.3 (74/253)	26.0 (13/50)	25.3 (20/79)	77.8 (21/27)	28.0 (107/382)	69.2 (9/13)	26.6 (98/369)	0.0008
ECMO	17.4 (44/253)	0 (0/50)	24.4 (20/82)	3.7 (1/27)	16.6 (64/385)	46.2 (6/13)	15.6 (58/372)	0.0036
Plasmapheresis	0 (0/266)	0 (0/50)	0 (0/87)	100 (27/27)	0 (0/403)	0 (0/13)	0 (0/390)	/
max. WHO score (Median, IQR, available n)	5 (4-7), 266	7 (4-8), 50	6 (4-8), 83	7 (7-8), 27	6 (4-7), 399	8 (8-8), 13	6 (4-7), 386	0.0001
Outcome Discharged or transferred Deceased Unknown	81.0 (205/253) 19.0 (48/253) /	72.0 (36/50) 28.0 (14/50) /	72.0 (59/82) 25.6 (21/82) 2.4 (2/82)	51.8 (14/27) 48.2 (13/27) /	77.9 (300/385) 21.6 (83/385) 0.5 (2/385)	7.7 (1/13) 92.3 (12/13) /	80.4 (299/372) 19.1 (71/372) 0.5 (2/372)	0.0001

738 **Table 1: Baseline patient characteristics**

739 Data are shown in % (N/n) unless otherwise indicated.

740 IMV – invasive mechanical ventilation, IQR – interquartile range, CCI – Charlson's comorbidity index

741 Patients with DNI/DNR were excluded for IMV, RRT, ECMO, and Outcome

REFERENCES

Abers, Michael S., Lindsey B. Rosen, Ottavia M. Delmonte, Elana Shaw, Paul Bastard, Luisa Imberti, Virginia Quaresima, et al. 2021. "Neutralizing Type-I Interferon Autoantibodies Are Associated with Delayed Viral Clearance and Intensive Care Unit Admission in Patients with COVID-19." *Immunology and Cell Biology*, July. https://doi.org/10.1111/imcb.12495.

Ankerhold, Jakob, Sebastian Giese, Philipp Kolb, Andrea Maul-Pavicic, Reinhard E. Voll, Nathalie Göppert, Kevin Ciminski, et al. 2021. "Circulating Multimeric Immune Complexes Drive Immunopathology in COVID-19." *bioRxiv*. https://doi.org/10.1101/2021.06.25.449893.

Bastard, Paul, Elizaveta Orlova, Leila Sozaeva, Romain Lévy, Alyssa James, Monica M. Schmitt, Sebastian Ochoa, et al. 2021. "Preexisting Autoantibodies to Type I IFNs Underlie Critical COVID-19 Pneumonia in Patients with APS-1." *The Journal of Experimental Medicine* 218 (7). https://doi.org/10.1084/jem.20210554.

Bastard, Paul, Lindsey B. Rosen, Qian Zhang, Eleftherios Michailidis, Hans-Heinrich Hoffmann, Yu Zhang, Karim Dorgham, et al. 2020. "Autoantibodies against Type I IFNs in Patients with Life-Threatening COVID-19." *Science* 370 (6515). https://doi.org/10.1126/science.abd4585.

Beccuti, G., L. Ghizzoni, V. Cambria, V. Codullo, P. Sacchi, E. Lovati, S. Mongodi, G. A. Iotti, and F. Mojoli. 2020. "A COVID-19 Pneumonia Case Report of Autoimmune Polyendocrine Syndrome Type 1 in Lombardy, Italy: Letter to the Editor." *Journal of Endocrinological Investigation* 43 (8): 1175–77.

Carpino, Andrea, Raffaele Buganza, Patrizia Matarazzo, Gerdi Tuli, Michele Pinon, Pier Luigi Calvo, Davide Montin, Francesco Licciardi, and Luisa De Sanctis. 2021. "Autoimmune Polyendocrinopathy-Candidiasis-Ectodermal Dystrophy in Two Siblings: Same Mutations but Very Different Phenotypes." *Genes* 12 (2). https://doi.org/10.3390/genes12020169.

Corman, Victor M., Olfert Landt, Marco Kaiser, Richard Molenkamp, Adam Meijer, Daniel Kw Chu, Tobias Bleicker, et al. 2020. "Detection of 2019 Novel Coronavirus (2019-nCoV) by Real-Time RT-PCR." *Euro Surveillance: Bulletin Europeen Sur Les Maladies Transmissibles = European Communicable Disease Bulletin* 25 (3). https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045.

COVID-19 Host Genetics Initiative. 2021. "Mapping the Human Genetic Architecture of COVID-19." *Nature*, July. https://doi.org/10.1038/s41586-021-03767-x.

Fernandez, Javier, Jordi Gratacos-Ginès, Pol Olivas, Montserrat Costa, Susana Nieto, Dolors Mateo, María Belén Sánchez, et al. 2020. "Plasma Exchange: An Effective Rescue Therapy in Critically Ill Patients With Coronavirus Disease 2019 Infection." *Critical Care Medicine* 48 (12): e1350–55.

Ferré, Elise M. N., Monica M. Schmitt, Sebastian Ochoa, Lindsey B. Rosen, Elana R. Shaw, Peter D. Burbelo, Jennifer L. Stoddard, et al. 2021. "SARS-CoV-2 Spike Protein-Directed Monoclonal Antibodies May Ameliorate COVID-19 Complications in APECED Patients." *Frontiers in Immunology* 12 (August): 720205.

Goncalves, David, Mehdi Mezidi, Paul Bastard, Magali Perret, Kahina Saker, Nicole Fabien, Rémi Pescarmona, et al. 2021. "Antibodies against Type I Interferon: Detection and Association with Severe Clinical Outcome in COVID-19 Patients." *Clinical & Translational Immunology* 10 (8): e1327.

Güven, Esin, Karen Duus, Magnus Christian Lydolph, Charlotte Sværke Jørgensen, Inga Laursen, and Gunnar Houen. 2014. "Non-Specific Binding in Solid Phase Immunoassays for Autoantibodies Correlates with Inflammation Markers." *Journal of Immunological Methods* 403 (1-2): 26–36.

"Information on COVID-19 Treatment, Prevention and Research." n.d. Accessed September 25, 2021. https://www.covid19treatmentguidelines.nih.gov.

Jones, Terry C., Guido Biele, Barbara Mühlemann, Talitha Veith, Julia Schneider, Jörn Beheim-Schwarzbach, Tobias Bleicker, et al. 2021. "Estimating Infectiousness throughout SARS-CoV-2 Infection Course." *Science* 373 (6551). https://doi.org/10.1126/science.abi5273.

Kim, You-Me, and Eui-Cheol Shin. 2021. "Type I and III Interferon Responses in SARS-CoV-2 Infection." *Experimental & Molecular Medicine* 53 (5): 750–60.

Knight, Stephen R., Antonia Ho, Riinu Pius, Iain Buchan, Gail Carson, Thomas M. Drake, Jake Dunning, et al. 2020. "Risk Stratification of Patients Admitted to Hospital with Covid-19 Using the ISARIC WHO Clinical Characterisation Protocol: Development and Validation of the 4C Mortality Score." *BMJ* 370 (September): m3339.

Koning, Rutger, Paul Bastard, Jean-Laurent Casanova, Matthijs C. Brouwer, Diederik van de Beek, and with the Amsterdam U.M.C. COVID-19 Biobank Investigators. 2021. "Autoantibodies against Type I Interferons Are Associated with Multi-Organ Failure in COVID-19 Patients." *Intensive Care Medicine*.

Kurth, F., M. Roennefarth, C. Thibeault, V. M. Corman, H. Muller-Redetzky, M. Mittermaier, C. Ruwwe-Glosenkamp, et al. 2020. "Studying the Pathophysiology of Coronavirus Disease 2019: A Protocol for the Berlin Prospective COVID-19 Patient Cohort (Pa-COVID-19)." *Infection* 48 (4): 619–26.

Lemarquis, Andri, Tessa Campbell, Maribel Aranda-Guillén, Viktoria Hennings, Petter Brodin, Olle Kämpe, Kaj Blennow, et al. 2021. "Severe COVID-19 in an APS1 Patient with Interferon Autoantibodies Treated with Plasmapheresis." *The Journal of Allergy and Clinical Immunology* 148 (1): 96–98.

Meisel, Christian, Bengisu Akbil, Tim Meyer, Erwin Lankes, Victor M. Corman, Olga Staudacher, Nadine Unterwalder, et al. 2021. "Mild COVID-19 despite Autoantibodies to Type I IFNs in Autoimmune-Polyendocrine-Syndrome Type 1 (APS-1)." *The Journal of Clinical Investigation*. https://www.jci.org/articles/view/150867.

Meyer, Steffen, Martin Woodward, Christina Hertel, Philip Vlaicu, Yasmin Haque, Jaanika Kärner, Annalisa Macagno, et al. 2016. "AIRE-Deficient Patients Harbor Unique High-Affinity Disease-Ameliorating Autoantibodies." *Cell* 166 (3): 582–95.

Nusshag, Christian, Christian Morath, Claudius Speer, Florian Kaelble, Martin Zeier, Monica Boxberger, Elisa Schulze-Schleithoff, Mascha O. Fiedler, Markus A. Weigand, and Uta Merle. 2021. "Plasma Exchange in Patients With Severe Coronavirus Disease 2019: A Single-Center Experience." *Critical Care Explorations* 3 (8): e0517.

Okba, Nisreen M. A., Marcel A. Müller, Wentao Li, Chunyan Wang, Corine H. GeurtsvanKessel, Victor M. Corman, Mart M. Lamers, et al. 2020. "Severe Acute Respiratory Syndrome Coronavirus 2-Specific Antibody Responses in Coronavirus Disease Patients." *Emerging Infectious Diseases* 26 (7): 1478–88.

Pfortmueller, Carmen A., Thibaud Spinetti, Richard D. Urman, Markus M. Luedi, and Joerg C. Schefold. 2021. "COVID-19-Associated Acute Respiratory Distress Syndrome (CARDS): Current Knowledge on Pathophysiology and ICU Treatment - A Narrative Review." *Best Practice & Research. Clinical Anaesthesiology* 35 (3): 351–68.

Prost, Nicolas de, Paul Bastard, Romain Arrestier, Slim Fourati, Mathieu Mahévas, Sonia Burrel, Karim Dorgham, et al. 2021. "Plasma Exchange to Rescue Patients with Autoantibodies Against Type I Interferons and Life-Threatening COVID-19 Pneumonia." *Journal of Clinical Immunology* 41 (3): 536–44.

Sah, Pratha, Meagan C. Fitzpatrick, Charlotte F. Zimmer, Elaheh Abdollahi, Lyndon Juden-Kelly, Seyed M. Moghadas, Burton H. Singer, and Alison P. Galvani. 2021. "Asymptomatic SARS-CoV-2 Infection: A Systematic Review and Meta-Analysis." *Proceedings of the National Academy of Sciences of the United States of America* 118 (34). https://doi.org/10.1073/pnas.2109229118.

Sahud, M., O. Zhukov, K. Mo, J. Popov, and J. Dlott. 2012. "False-Positive Results in ELISA-Based Anti FVIII Antibody Assay May Occur with Lupus Anticoagulant and Phospholipid Antibodies." *Haemophilia: The Official Journal of the World Federation of Hemophilia* 18 (5): 777–81.

Schwarz, Tatjana, Pinkus Tober-Lau, David Hillus, Elisa T. Helbig, Lena J. Lippert, Charlotte

Thibeault, Willi Koch, et al. 2021. "Delayed Antibody and T-Cell Response to BNT162b2 Vaccination in the Elderly, Germany." *Emerging Infectious Diseases* 27 (8): 2174–78.

Spinetti, Thibaud, Cedric Hirzel, Michaela Fux, Laura N. Walti, Patrick Schober, Frank Stueber, Markus M. Luedi, and Joerg C. Schefold. 2020. "Reduced Monocytic Human Leukocyte Antigen-DR Expression Indicates Immunosuppression in Critically Ill COVID-19 Patients." *Anesthesia and Analgesia* 131 (4): 993–99.

Stuckrad, Sae Lim von, Jens Klotsche, Robert Biesen, Mareike Lieber, Julia Thumfart, Christian Meisel, Nadine Unterwalder, and Tilmann Kallinich. 2020. "SIGLEC1 (CD169) Is a Sensitive Biomarker for the Deterioration of the Clinical Course in Childhood Systemic Lupus Erythematosus." *Lupus* 29 (14): 1914–25.

Troya, Jesús, Paul Bastard, Laura Planas-Serra, Pablo Ryan, Montse Ruiz, María de Carranza, Juan Torres, et al. 2021. "Neutralizing Autoantibodies to Type I IFNs in >10% of Patients with Severe COVID-19 Pneumonia Hospitalized in Madrid, Spain." *Journal of Clinical Immunology*. https://doi.org/10.1007/s10875-021-01036-0.

WHO Solidarity Trial Consortium, Hongchao Pan, Richard Peto, Ana-Maria Henao-Restrepo, Marie-Pierre Preziosi, Vasee Sathiyamoorthy, Quarraisha Abdool Karim, et al. 2021. "Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results." *The New England Journal of Medicine* 384 (6): 497–511.

Wijst, Monique G. P. van der, Sara E. Vazquez, George C. Hartoularos, Paul Bastard, Tianna Grant, Raymund Bueno, David S. Lee, et al. 2021. "Type I Interferon Autoantibodies Are Associated with Systemic Immune Alterations in Patients with COVID-19." *Science Translational Medicine* 13 (612): eabh2624.

Williamson, Elizabeth J., Alex J. Walker, Krishnan Bhaskaran, Seb Bacon, Chris Bates, Caroline E. Morton, Helen J. Curtis, et al. 2020. "Factors Associated with COVID-19-Related Death Using OpenSAFELY." *Nature* 584 (7821): 430–36.

Wölfel, Roman, Victor M. Corman, Wolfgang Guggemos, Michael Seilmaier, Sabine Zange, Marcel A. Müller, Daniela Niemeyer, et al. 2020. "Virological Assessment of Hospitalized Patients with COVID-2019." *Nature* 581 (7809): 465–69.

Zhang, Qian, Paul Bastard, Zhiyong Liu, Jérémie Le Pen, Marcela Moncada-Velez, Jie Chen, Masato Ogishi, et al. 2020. "Inborn Errors of Type I IFN Immunity in Patients with Life-Threatening COVID-19." *Science* 370 (6515). https://doi.org/10.1126/science.abd4570.

Abers, Michael S., Lindsey B. Rosen, Ottavia M. Delmonte, Elana Shaw, Paul Bastard, Luisa Imberti, Virginia Quaresima, et al. 2021. "Neutralizing Type-I Interferon Autoantibodies Are Associated with Delayed Viral Clearance and Intensive Care Unit Admission in Patients with COVID-19." Immunology and Cell Biology, July. https://doi.org/10.1111/imcb.12495.

Bastard, Paul, Elizaveta Orlova, Leila Sozaeva, Romain Lévy, Alyssa James, Monica M. Schmitt, Sebastian Ochoa, et al. 2021. "Preexisting Autoantibodies to Type I IFNs Underlie Critical COVID-19 Pneumonia in Patients with APS-1." The Journal of Experimental Medicine 218 (7). https://doi.org/10.1084/jem.20210554.

Bastard, Paul, Lindsey B. Rosen, Qian Zhang, Eleftherios Michailidis, Hans-Heinrich Hoffmann, Yu Zhang, Karim Dorgham, et al. 2020. "Autoantibodies against Type I IFNs in Patients with Life-Threatening COVID-19." Science 370 (6515). https://doi.org/10.1126/science.abd4585.

Beccuti, G., L. Ghizzoni, V. Cambria, V. Codullo, P. Sacchi, E. Lovati, S. Mongodi, G. A. Iotti, and F. Mojoli. 2020. "A COVID-19 Pneumonia Case Report of Autoimmune Polyendocrine Syndrome Type 1 in Lombardy, Italy: Letter to the Editor." Journal of Endocrinological Investigation 43 (8): 1175–77.

Carpino, Andrea, Raffaele Buganza, Patrizia Matarazzo, Gerdi Tuli, Michele Pinon, Pier Luigi Calvo, Davide Montin, Francesco Licciardi, and Luisa De Sanctis. 2021. "Autoimmune Polyendocrinopathy-Candidiasis-Ectodermal Dystrophy in Two Siblings: Same Mutations but Very Different Phenotypes." Genes 12 (2). https://doi.org/10.3390/genes12020169.

Corman, Victor M., Olfert Landt, Marco Kaiser, Richard Molenkamp, Adam Meijer, Daniel Kw Chu, Tobias Bleicker, et al. 2020. "Detection of 2019 Novel Coronavirus (2019-nCoV) by Real-Time RT-PCR." Euro Surveillance: Bulletin Europeen Sur Les Maladies Transmissibles = European Communicable Disease Bulletin 25 (3). https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045.

COVID-19 Host Genetics Initiative. 2021. "Mapping the Human Genetic Architecture of COVID-19." Nature, July. https://doi.org/10.1038/s41586-021-03767-x.

Fernandez, Javier, Jordi Gratacos-Ginès, Pol Olivas, Montserrat Costa, Susana Nieto, Dolors Mateo, María Belén Sánchez, et al. 2020. "Plasma Exchange: An Effective Rescue Therapy in Critically Ill Patients With Coronavirus Disease 2019 Infection." Critical Care Medicine 48 (12): e1350–55.

Ferré, Elise M. N., Monica M. Schmitt, Sebastian Ochoa, Lindsey B. Rosen, Elana R. Shaw, Peter D. Burbelo, Jennifer L. Stoddard, et al. 2021. "SARS-CoV-2 Spike Protein-Directed Monoclonal Antibodies May Ameliorate COVID-19 Complications in APECED Patients." Frontiers in Immunology 12 (August): 720205.

Goncalves, David, Mehdi Mezidi, Paul Bastard, Magali Perret, Kahina Saker, Nicole Fabien, Rémi Pescarmona, et al. 2021. "Antibodies against Type I Interferon: Detection and Association with Severe Clinical Outcome in COVID-19 Patients." Clinical & Translational Immunology 10 (8): e1327.

Güven, Esin, Karen Duus, Magnus Christian Lydolph, Charlotte Sværke Jørgensen, Inga Laursen, and Gunnar Houen. 2014. "Non-Specific Binding in Solid Phase Immunoassays for Autoantibodies Correlates with Inflammation Markers." Journal of Immunological Methods 403 (1-2): 26–36.

"Information on COVID-19 Treatment, Prevention and Research." n.d. Accessed September 25, 2021. https://www.covid19treatmentguidelines.nih.gov.

Jones, Terry C., Guido Biele, Barbara Mühlemann, Talitha Veith, Julia Schneider, Jörn Beheim-Schwarzbach, Tobias Bleicker, et al. 2021. "Estimating Infectiousness throughout SARS-CoV-2 Infection Course." Science 373 (6551). https://doi.org/10.1126/science.abi5273.

Kim, You-Me, and Eui-Cheol Shin. 2021. "Type I and III Interferon Responses in SARS-CoV-2 Infection." Experimental & Molecular Medicine 53 (5): 750–60.

Knight, Stephen R., Antonia Ho, Riinu Pius, Iain Buchan, Gail Carson, Thomas M. Drake, Jake Dunning, et al. 2020. "Risk Stratification of Patients Admitted to Hospital with Covid-19 Using the ISARIC WHO Clinical Characterisation Protocol: Development and Validation of the 4C Mortality Score." BMJ 370 (September): m3339.

Koning, Rutger, Paul Bastard, Jean-Laurent Casanova, Matthijs C. Brouwer, Diederik van de Beek, and with the Amsterdam U.M.C. COVID-19 Biobank Investigators. 2021. "Autoantibodies against Type I Interferons Are Associated with Multi-Organ Failure in COVID-19 Patients." Intensive Care Medicine.

Lemarquis, Andri, Tessa Campbell, Maribel Aranda-Guillén, Viktoria Hennings, Petter Brodin, Olle Kämpe, Kaj Blennow, et al. 2021. "Severe COVID-19 in an APS1 Patient with Interferon Autoantibodies Treated with Plasmapheresis." The Journal of Allergy and Clinical Immunology 148 (1): 96–98.

Meisel, Christian, Bengisu Akbil, Tim Meyer, Erwin Lankes, Victor M. Corman, Olga Staudacher, Nadine Unterwalder, et al. 2021. "Mild COVID-19 despite Autoantibodies to Type I IFNs in Autoimmune-Polyendocrine-Syndrome Type 1 (APS-1)." The Journal of Clinical Investigation. https://www.jci.org/articles/view/150867.

Meyer, Steffen, Martin Woodward, Christina Hertel, Philip Vlaicu, Yasmin Haque, Jaanika Kärner, Annalisa Macagno, et al. 2016. "AIRE-Deficient Patients Harbor Unique High-Affinity Disease-Ameliorating Autoantibodies." Cell 166 (3): 582–95.

Nusshag, Christian, Christian Morath, Claudius Speer, Florian Kaelble, Martin Zeier, Monica Boxberger, Elisa Schulze-Schleithoff, Mascha O. Fiedler, Markus A. Weigand, and Uta Merle. 2021. "Plasma Exchange in Patients With Severe Coronavirus Disease 2019: A Single-Center Experience." Critical Care Explorations 3 (8): e0517.

Okba, Nisreen M. A., Marcel A. Müller, Wentao Li, Chunyan Wang, Corine H.

GeurtsvanKessel, Victor M. Corman, Mart M. Lamers, et al. 2020. "Severe Acute Respiratory Syndrome Coronavirus 2-Specific Antibody Responses in Coronavirus Disease Patients." Emerging Infectious Diseases 26 (7): 1478–88.

Prost, Nicolas de, Paul Bastard, Romain Arrestier, Slim Fourati, Mathieu Mahévas, Sonia Burrel, Karim Dorgham, et al. 2021. "Plasma Exchange to Rescue Patients with Autoantibodies Against Type I Interferons and Life-Threatening COVID-19 Pneumonia." Journal of Clinical Immunology 41 (3): 536–44.

Sahud, M., O. Zhukov, K. Mo, J. Popov, and J. Dlott. 2012. "False-Positive Results in ELISA-Based Anti FVIII Antibody Assay May Occur with Lupus Anticoagulant and Phospholipid Antibodies." Haemophilia: The Official Journal of the World Federation of Hemophilia 18 (5): 777–81.

Schwarz, Tatjana, Pinkus Tober-Lau, David Hillus, Elisa T. Helbig, Lena J. Lippert, Charlotte Thibeault, Willi Koch, et al. 2021. "Delayed Antibody and T-Cell Response to BNT162b2 Vaccination in the Elderly, Germany." Emerging Infectious Diseases 27 (8): 2174–78.

Spinetti, Thibaud, Cedric Hirzel, Michaela Fux, Laura N. Walti, Patrick Schober, Frank Stueber, Markus M. Luedi, and Joerg C. Schefold. 2020. "Reduced Monocytic Human Leukocyte Antigen-DR Expression Indicates Immunosuppression in Critically Ill COVID-19 Patients." Anesthesia and Analgesia 131 (4): 993–99.

Stuckrad, Sae Lim von, Jens Klotsche, Robert Biesen, Mareike Lieber, Julia Thumfart, Christian Meisel, Nadine Unterwalder, and Tilmann Kallinich. 2020. "SIGLEC1 (CD169) Is a Sensitive Biomarker for the Deterioration of the Clinical Course in Childhood Systemic Lupus Erythematosus." Lupus 29 (14): 1914–25.

Troya, Jesús, Paul Bastard, Laura Planas-Serra, Pablo Ryan, Montse Ruiz, María de Carranza, Juan Torres, et al. 2021. "Neutralizing Autoantibodies to Type I IFNs in >10% of Patients with Severe COVID-19 Pneumonia Hospitalized in Madrid, Spain." Journal of Clinical Immunology. https://doi.org/10.1007/s10875-021-01036-0.

WHO Solidarity Trial Consortium, Hongchao Pan, Richard Peto, Ana-Maria Henao-Restrepo, Marie-Pierre Preziosi, Vasee Sathiyamoorthy, Quarraisha Abdool Karim, et al. 2021. "Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results." The New England Journal of Medicine 384 (6): 497–511.

Wijst, Monique G. P. van der, Sara E. Vazquez, George C. Hartoularos, Paul Bastard, Tianna Grant, Raymund Bueno, David S. Lee, et al. 2021. "Type I Interferon Autoantibodies Are Associated with Systemic Immune Alterations in Patients with COVID-19." Science Translational Medicine 13 (612): eabh2624.

Williamson, Elizabeth J., Alex J. Walker, Krishnan Bhaskaran, Seb Bacon, Chris Bates, Caroline E. Morton, Helen J. Curtis, et al. 2020. "Factors Associated with COVID-19-Related Death Using OpenSAFELY." Nature 584 (7821): 430–36.

Wölfel, Roman, Victor M. Corman, Wolfgang Guggemos, Michael Seilmaier, Sabine Zange, Marcel A. Müller, Daniela Niemeyer, et al. 2020. "Virological Assessment of Hospitalized Patients with COVID-2019." Nature 581 (7809): 465–69.

Zhang, Qian, Paul Bastard, Zhiyong Liu, Jérémie Le Pen, Marcela Moncada-Velez, Jie Chen, Masato Ogishi, et al. 2020. "Inborn Errors of Type I IFN Immunity in Patients with Life-Threatening COVID-19." Science 370 (6515). https://doi.org/10.1126/science.abd4570.