Paratype: A genotyping framework and an open-source tool for *Salmonella* Paratyphi A

Arif M. Tanmoy[1,2*], Yogesh Hooda[1,3*], Mohammad S. I. Sajib[1,4], Kesia E. da Silva[5], Junaid Iqbal[6], Farah N. Qamar[6], Stephen P. Luby[5], Gordon Dougan[7], Zoe A. Dyson[7,8,9,12], Stephen Baker[8], Denise O. Garrett[10], Jason R. Andrews[5], Samir K. Saha[1,11, $], Senjuti Saha[1,$,#]

[1] Child Health Research Foundation, Dhaka, Bangladesh. arif.tanmoy@chrfbd.org, yhooda@chrfbd.org, saiful.sajib@chrfbd.org, samir@chrfbd.org, senjutisaha@chrfbd.org

[2] Department of Medical Microbiology and Infectious Diseases, Erasmus University Medical Center, Rotterdam, the Netherlands.

[3] MRC-Laboratory Molecular Biology, Cambridge, UK.

[4] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow, UK.

[5] Division of Infectious Diseases and Geographic Medicine, Stanford University School of Medicine, Stanford, California, USA. kesiaeds@stanford.edu, sluby@stanford.edu, jandr@stanford.edu

[6] Department of Paediatrics and Child Health, Aga Khan University, Karachi, Pakistan. junaid.iqbal@aku.edu, farah.qamar@aku.edu

[7] Wellcome Trust Sanger Institute, Hinxton, Cambridge CB10 1SA, UK. gd312@medschl.cam.ac.uk

[8] Cambridge Institute of Therapeutic Immunology and Infectious Disease, Department of Medicine, University of Cambridge, Cambridge, United Kingdom. sgb47@medschl.cam.ac.uk

[9] Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK. Zoe.Dyson@lshtm.ac.uk

[10] Applied Epidemiology Team, Sabin Vaccine Institute, Washington, DC, USA. Denise.Garrett@Sabin.org

28 [11] Department of Microbiology, Bangladesh Institute of Child Health, Dhaka Shishu Hospital,

29 Dhaka, Bangladesh.

30 [12] Department of Infectious Diseases, Central Clinical School, Monash University,
31 Melbourne, Victoria 3004, Australia.
32

33 *, $ Equal contribution;

34 # Corresponding author

Abstract

Background: *Salmonella enterica* serovar Paratyphi A (*Salmonella* Paratyphi A) is the primary causative agent of paratyphoid fever, which is responsible for an estimated 3.4 million infections annually. However, little genomic information is available on population structure, antimicrobial resistance (AMR), and spatiotemporal distribution of the pathogen. With rising antimicrobial resistance and no licensed vaccines, genomic surveillance is important to track the evolution of this pathogen and monitor transmission.

Results: We performed whole-genome sequencing of 817 *Salmonella* Paratyphi A isolates collected from Bangladesh, Nepal, and Pakistan and added publicly available 562 genomes to build a global database representing 37 countries, covering 1917-2019. To track the evolution of *Salmonella* Paratyphi A, we used the existing lineage scheme, developed earlier based on a small dataset, but certain sub-lineages were not homologous, and many isolates could not be assigned a lineage. Therefore, we developed a single nucleotide polymorphism based genotyping scheme, Paratype, a tool that segregates *Salmonella* Paratyphi A into three primary and nine secondary clades, and 18 genotypes. Each genotype has been assigned a unique allele definition located on a conserved gene. Using Paratype, we identified genomic variation between different sampling locations and specific AMR markers, and mutations in the O2-polysaccharide synthesis locus, a candidate for vaccine development.

Conclusions: This large-scale global analysis proposes the first genotyping tool for *Salmonella* Paratyphi A. Paratype has already been released (https://github.com/CHRF-Genomics/Paratype) as an open-access, command-line tool and is being adopted for large scale genomic analysis. This tool will assist future genomic surveillance and help inform prevention and treatment strategies.

75   **Keywords:** *Salmonella* Paratyphi A; Paratyphoid fever; Paratyphi A genotyping; Genomics;

76   Antimicrobial resistance; Global analysis; Epidemiology; Enteric fever; Neglected tropical

77   disease.

Background

Paratyphoid fever, caused by *Salmonella enterica* subspecies *enterica* serovar Paratyphi A (*Salmonella* Paratyphi A) is a systemic febrile illness that affects an estimated 3.4 million people each year, and causes 19,100 deaths globally [1]. The disease is clinically indistinguishable from typhoid fever, caused by *Salmonella enterica* subspecies *enterica* serovar Typhi (*Salmonella* Typhi). Much like typhoid, paratyphoid fever is also endemic in many low- and middle-income countries of South Asia and Sub-Saharan Africa, due to fecal contamination of water, food and the environment. However, barring a few countries (e.g., China, Myanmar), paratyphoid fever is usually less prevalent than typhoid fever [2,3]. *Salmonella* Paratyphi A continues to be an inadequately studied pathogen [4] hampering the implementation of evidence-based policies for the treatment and prevention of paratyphoid fever.

Relative to *Salmonella* Typhi, little genomic information is available on population structure, antimicrobial resistance (AMR), and spatiotemporal distribution of *Salmonella* Paratyphi A. The first *Salmonella* Paratyphi A genome was published in 2004 and had a size of 4.5 Mb, with ~4,200 genes. To determine the global diversity of *Salmonella* Paratyphi A isolates, Bayesian analysis was conducted on a set of 149 *Salmonella* Paratyphi A genomes, which identified that the last common ancestor of all *Salmonella* Paratyphi A existed for at least 450 years prior to differentiating into at least seven distinct lineages (A to G) which have circulated globally [5]. Whole genome sequencing was also used to characterize clonal paratyphoid outbreaks in Cambodia [6] and China [7] and further extend the lineage scheme to include sub-lineages within Lineage A and C. However, very few studies have characterized isolates from countries in South Asia, which contributes over 80% of all paratyphoid infections [8,9]. Available studies are sporadic, and either focused on genomes

from a specific geographical location or provide no information on antimicrobial resistance

markers, potential vaccine targets, and other virulence factors.

To address this data gap, we performed whole-genome sequencing of 817 *Salmonella*

Paratyphi A isolates collected from Bangladesh, Nepal, and Pakistan and combined them

with whole-genome sequence data of another 562 isolates reported in the literature to build a

global database of 1,379 *Salmonella* Paratyphi A isolates. To track the evolution of

Salmonella Paratyphi A over a century, we used the existing lineage scheme and found that

certain sub-lineages were not homologous, and many isolates could not be assigned a specific

lineage. This motivated us to develop a single nucleotide polymorphism (SNP) based

genotyping scheme, called Paratype. The scheme is phylogenetically informative and

successfully segregates the global population structure into three primary, seven secondary,

and 18 distinct subclades/genotypes. We also identified the specific antimicrobial resistance

genes, mutations, and plasmids present in *Salmonella* Paratyphi A genomes and correlated

these with the different genotypes.

<u>Results</u>

<u>Whole-genome sequencing and compilation of global *Salmonella* Paratyphi A genomes</u>

The Child Health Research Foundation (CHRF) has been conducting typhoid and

paratyphoid fever surveillance in Bangladesh since 1999 and has generated a biobank of

1,123 *Salmonella* Paratyphi A isolates from 1999-2018 [10–12]. We selected 528 of these

isolates, covering all age groups, years of isolation, and hospitalization status

(hospitalized/out-patient), and performed whole-genome sequencing on these isolates

(Additional file 1: Table S1). Of these, 180 *Salmonella* Paratyphi A isolates were collected as

part of the Surveillance of Enteric Fever in Asia Project (SEAP, 2014 - 2019) study, a multi-

country international effort to better understand the epidemiology and impact of enteric fever

in South Asia [13]. In addition to Bangladesh, 133 isolates were sequenced from the SEAP

study conducted in Pakistan, and 156 from Nepal.

To contextualize these genomes, we conducted a literature search to compile all publicly

available *Salmonella* Paratyphi A genomes (for which raw reads were available) to build a

database of 560 additional isolates from 10 studies (Additional file 1: Table S2). Two

reference genomes (ATCC 9150 and AKU_12601) were also included. The largest dataset

consisted of 254 isolates, published by Public Health England as part of their *Salmonella*

surveillance [8,14]; 164 of these isolates were linked to travel, most commonly to South Asia.

In our study, we assigned these isolates to the countries where the patient acquired the

infection. Our final data, including the genomes we sequenced, consisted of a total of 1,379

isolates from 37 different countries, spanning over 103 years - 1917 to 2019. Most of the

isolates (1,112/ 1,379; 81%) were from countries in South Asia (541 from Bangladesh, 268

from Nepal, 187 from Pakistan and 115 from India). South Asian countries also bear a

disproportionately high burden of paratyphoid fever; of the estimated 3.4 million global

paratyphoid infections in 2019, 2.8 (82%) million are estimated to have occurred in South

Asia [1].

Following assembly from raw reads, the pan-genome analysis identified 6,983 genes, of

which 4,114 (59% of all genes) were conserved in more than 95% of isolates (Additional file

1: Figure S1). The average genome size was 4.5 Mb with ~4,300 genes, and the pan genome

does not appear to be closed (decay parameter, alpha = 0.67). Overall, 2,550 genes were

found to be present in less than 15% of isolates, and these included genes often found in

prophages and other mobile regions, and genes encoding adhesins, antimicrobial resistance

markers, and several hypothetical proteins.

Genotyping scheme for *Salmonella* Paratyphi A

To investigate the genomic diversity of *Salmonella* Paratyphi A, we identified 8,346 single

nucleotide polymorphisms (SNPs) in the 1,379 isolates. These were used in RAxML [15] to

generate a Maximum-likelihood phylogenetic tree of the global collection of *Salmonella*

Paratyphi A isolates (Figure 1). A previously reported lineage scheme, proposed for

Salmonella Paratyphi A by Zhou et al. [5] and extended by subsequent studies [6,7,9,16,17]

was overlaid on the RAxML tree. This highlighted the insufficiency of the current lineage

scheme to fully capture the diversity of *Salmonella* Paratyphi A present. First, while the

isolates from lineages B & D - G clustered together, several isolates previously assigned to

lineages A and C in the scheme did not. Second, some sequences belonged to clades that

diverged from isolates before the exitance of the most recent common ancestor for lineages A

and B, indicating that these isolates should be considered to be in a different lineage. This

was not surprising considering that when this scheme was devised, there were limited number

of sequenced *Salmonella* Paratyphi A genomes available, particularly from South Asia.

To build a genotyping scheme based on a larger number of representative samples, first, we

used fastBAPS [18] to generate a potential list of clusters in the RAxML tree (Additional file

1: Figure S2). Next, we selected a set of 315 isolates that included two isolates per year for all

fastBAPS clusters selected randomly and performed phylodynamic analysis using the

Bayesian Evolutionary Analysis by Sampling Trees (BEAST) software [19] (Figure 2). Based

on these analyses, we devised a genotyping scheme with three primary clades, nine secondary

clades, and 18 genotypes that have circulated globally in the last 100 years.

To aid further genomic epidemiological studies, we identified 18 additional alleles

(Additional file 1: Table S3) that are unique to each of the 18 *Salmonella* Paratyphi A

genotypes. These alleles were present in conserved genes involved in essential cellular

functions such as protein synthesis, DNA replication, or metabolism. Identification of these

genotype-specific alleles allowed us to write a Python script – "Paratype" – that assigns

genotypes to *Salmonella* Paratyphi A genomes using fastq, bam, or vcf files obtained during

whole-genome sequencing and variant calling. The Paratype software tool (available at:

https://github.com/CHRF-Genomics/Paratype/) has 100% sensitivity and specificity and was

able to assign the correct genotype to all the 1,379 genomes that were present in our database.

<u>Temporal and geographic distribution of different genotypes</u>

Upon the establishment of the "Paratype" scheme, we considered the geographical

distribution of the different genotypes (Figure 3). Genotype 0.1 under primary clade 0 was

phylogenetically unique (matches with lineage H of Zhou et al [5]); there was only one

isolate belonging to this genotype/primary clade that was isolated in Hong Kong in 1971. The

genome of this isolate was distinct from all other genomes obtained thus far, contained 1288

unique SNPs, and may represent a lineage that is now extinct, or present at very low numbers

in areas that have not been sampled. The other two primary clades, clades 1 and 2, emerged

between 1700-1800 and contain genomes that have been collected in the last two decades.

Clade 1 contains strains largely from lineage F, and fastBAPS predicted two sub-clusters

within this clade. One of these clusters was largely found in Bangladesh and has been

assigned secondary clade 1.2, then sub-divided into genotypes 1.2.1 and 1.2.2 which appear

to have diverged in the 1950s. Both these genotypes are currently present in Bangladesh and

other South Asian countries (Figure 2). The other cluster with13 genomes from Bangladesh

that were first isolated in 1999 have been assigned to genotype 1.1. The remaining 10

genomes were obtained between 1917 to 1963 and have been assigned genotype 1.0.

Most *Salmonella* Paratyphi A genomes (1254/1379; 91%) have been assigned to primary

clade 2, which contains genomes belonging to the lineage A-E of the previous scheme.

Genomes that belonged to lineages B, D, and E have now been assigned to genotypes 2.4,

2.2, and 2.0, respectively. Within genotype 2.0, 13 unique and recent isolates from Pakistan

were identified and have been assigned as genotype 2.0.1. Genotype 2.1 contains isolates

from Nepal that were sampled during the SEAP study, yet the genotype emerged in the 1800s

and is distinct from all other isolates in clade 2. Two clusters in fastBAPS, comprising of

strains largely from what was formerly C lineage are now assigned to genotype 2.3. Genotype

2.3 has been subdivided into genotypes 2.3.1 to 2.3.3, each of which belongs to a distinct

geographical location: 2.3.1 is found predominantly in Cambodia and South-East Asia; 2.3.2

and 2.3.3 are found largely in South Asia. An outbreak of paratyphoid fever in China during

2010 – 2011 [7] was caused by isolates of genotype 2.3.3, and these likely originated in

South Asia. The former lineages A and B have been assigned genotype 2.4, which is further

divided into 2.4.1 to 2.4.4. While genotypes 2.4.1 and 2.4.2 have been observed in different

countries in South Asia, genotype 2.4.4 is predominantly found in Bangladesh, and 2.4.3 is

largely present in Nepal.

Different countries in South Asia had unique genotype distributions. Predominant genotypes

present in Bangladesh were 2.4.4 (56%) followed by 1.2.2 (14%) and 2.3.3 (13%). In Nepal,

2.4.3 (47%), 2.3.3 (16%) and 2.4.1 (14%) were three most common genotypes. Pakistan had

genotypes 2.3.3 (25%), 2.3 (16%) and 2.4 (15%). In India, genotypes 2.4.2 (22%), 2.4 (20%),

2.4.1 (19%), 2.3.3 (17%), and 2.3 (16%) were commonly identified.

<u>Antimicrobial resistance markers in *Salmonella* Paratyphi A</u>

To characterize genomic determinants of antimicrobial resistance in *Salmonella* Paratyphi A,

we screened the 1,379 genomes for the presence of antimicrobial genes and markers using

ResFinder [20] (Figure 4a) and plasmids using PlasmidFinder [21] (Figure 4b). Of the 1,379

isolates, 1,015 (74%) isolates showed no predicted plasmids and 1356/1379 had no predicted

antimicrobial resistance genes. Five genomes with the IncHI1 plasmid were identified, two

genomes (both from India) contained resistance genes for trimethoprim and chloramphenicol,

and the other three genomes contained genes for trimethoprim, chloramphenicol and

ampicillin designated as MDR isolates (one each from India, Pakistan, and Thailand). All five

genomes belonged to genotype 2.3 and the strains were isolated between 1999-2004. We also

identified a genome belonging to genotype 2.4.4 containing *bla*CTX-M-15 and *bla*TEM-1B

on an IncI1-I plasmid; the originating strain was isolated from a patient who contracted the

infection in Bangladesh in 2017 [22]. There were 14 isolates from the genotype 2.3.1 that

contain *bla*TEM-116, which can lead to resistance to ampicillin; all 14 were reported from

Cambodia[6]. Another isolate from genotype 2.3.3 (from Pakistan, 2015) contained a *qnrB19*

gene on a Col(pHAD28) plasmid, which has been shown to lead to quinolone resistance in

other *Salmonella* species [23].

In addition to antimicrobial resistance genes, we also identified chromosomal mutations in

the *acrB* gene and the quinolone resistance determining region (QRDR) to identify isolates

resistant to azithromycin and ciprofloxacin respectively. Six of 1,379 genomes contained an

AcrB R717 mutation, all from Bangladesh and these belonged to genotypes 2.3.3 (1/6) and

2.4.4 (5/6) [24,25]. The first azithromycin resistant *Salmonella* Paratyphi A isolate was

identified in 2014, and this resistance has emerged independently at least twice in two

different genotypes. On the other hand, a majority (1177/1397; 84%) of genomes had

mutations in the QRDR region. The most commonly found single mutation was gyrA-S83F

(941/1379), followed by gyrA-S83Y (205/1379). Two isolates contained double mutations in

the QRDR region; one of them belonged to genotype 2.0.1 (gyrA-S83F & D87N, Pakistan,

2017) and another belonged to genotype 2.3.3 (gyrA-S83F & D87G, UK, 2016). Barring

genotype 0.1, 1.0 and 2.2, all other genotypes had at least one genome with a QRDR

mutation (Figure 4c). The first QRDR mutation was identified in 1997 in India in genotype

2.4 and their prevalence have increased over time. In 2012 and 2013, there was an outbreak

in Cambodia caused by a strain from genotype 2.3.1 that did not have any QRDR mutation

leading to a temporary increase in proportion of *Salmonella* Paratyphi A with no QRDR

mutations during these two years (Additional file 1: Figure S3).

<u>Characterization of mutations in the O2-antigen biosynthetic gene cluster</u>

The majority of the vaccines being developed for *Salmonella* Paratyphi A use the O2-antigen

that is unique to this serovar conjugated to a carrier protein [26]. Recently, through in-silico

metabolic reconstruction, an 18.9 kb region containing genes involved in O-antigen

biosynthesis was identified as important for determining the specific molecular features of the

O2-antigen found in *Salmonella* Paratyphi A [27]. We identified the SNPs in the O2-antigen

biosynthesis genes found in the 1,379 genomes to investigate the conservation of this

genomic loci. In total, 84 SNPs were found, of which 13 were present in more than 10

genomes. The most common SNP was at genomic location 8,68,444 (G> C; synonymous

mutation in *prt* gene encoding paratose synthase), which was found in 17% (239/1,379) of all

isolates. Out of those 13 common SNPs (n≥10), seven led to non-synonymous mutations

(Additional file 1: Figure S4) that could potentially change the O2-antigen structure and

chemistry.

<u>Discussion</u>

Salmonella Paratyphi A is the causative agent of paratyphoid fever, a neglected tropical

disease with a high burden and mortality in low-and-middle-income countries. Limited

information is available regarding its genomic diversity, especially from South Asian

countries that collectively are responsible for over 80% of all paratyphoid cases. As genomic

surveillance becomes more prominent, there is a need for a coherent and easy-to-use scheme

that can be deployed by public health researchers that do not require extensive compute

resources.

We sequenced 817 isolates from Bangladesh, Pakistan and Nepal collected over the last 20

years and compiled a collection of all genomes of *Salmonella* Paratyphi A publicly available

thus far. We describe a genotyping framework for *Salmonella* Paratyphi A using 1,379

isolates obtained from 1917 through 2019. Rather than being guided by a single approach, we

combined ML-based phylogenetics with BAPS and Bayesian analysis via BEAST to design a

genotyping scheme for *Salmonella* Paratyphi A. The scheme divided the *Salmonella*

Paratyphi A population into 18 different genotypes, and each can be identified by the

presence of an allele that is located on the coding sequence of a conserved gene, involved in

housekeeping functions. We only found 8,346 SNPs from all 1,379 isolates, with minimal

recombination, and thus, this genotyping scheme based on SNP alleles can support robust

genotyping and accommodate future evolution of *Salmonella* Paratyphi A. And to assist with

that, we have developed Paratype, an open-source Python script for genotyping of *Salmonella*

Paratyphi A genomes. Paratype can detect the genotype of *Salmonella* Paratyphi A genomes

directly from raw fastq read data. It can also detect mutations in the acrB efflux pump

(determinant of macrolide resistance) and in the QRDR region (determinant of ciprofloxacin

non-susceptibility).

In this genotyping scheme, we propose three primary clades 0, 1, and 2, which diverged

before the 1800s (Figure 2). While only a single isolate of primary clade 0 was obtained in

1971, isolates belonging to clade 1 and 2 have been routinely identified over the past two

decades. Clade 2 is the most abundant and has been subdivided into four secondary clades:

2.1 - 2.4, which probably emerged in the 1800s or the early 1900s. Clade 2.3 could be

subdivided into 2.3.1 - 2.3.3, each with distinct geographic distribution. Clade 2.4 was also

sub-divided into genotypes 2.4.1 - 2.4.4. Genotype 2.4.4 was the most abundant and was

predominantly present in Bangladesh. This genotype emerged in the late 1990s to early 2000s

and possesses high rates of ciprofloxacin resistance (Figure 2). Five of the isolates from this

genotype also contained AcrB-R717Q mutation that leads to azithromycin resistance, while

one was found to harbor a plasmid containing extended-spectrum beta-lactamase gene

(*bla*CTX-M-15) [22].

In line with findings of previous studies, the rates of acquisition of antimicrobial resistance

markers in *Salmonella* Paratyphi A is lower relative to *Salmonella* Typhi (Figure 4) [6,9].

Although a few isolates did acquire the IncHI1 plasmid in the late 1990s to early 2000s

(Figure 4a), no massive spread across the globe was noted; this unlike *Salmonella* Typhi

lineage H58 (genotype 4.3.1) carrying the IncHI1 plasmid spread and became the dominant

lineage in the last 30 years[28]. This is also true for chromosomal mutations such as QRDR

and AcrB mutations, which are overall less prevalent in *Salmonella* Paratyphi A than in

Salmonella Typhi [28,29]. Considering the genetic similarities between *Salmonella* Typhi

and Paratyphi A, and the fact that they occupy the same environmental niche, the differences

in the presence of AMR genes between these typhoidal *Salmonella* serovars warrants further

investigation.

The specific O-antigen in the *Salmonella* Paratyphi A is thought to be conserved (assigned to

serogroup O2) and several vaccine candidates are currently under development, utilizing the

O2 antigen conjugated to a carrier protein as the main vaccine antigen. We compared the 18.9

kbp region responsible for the synthesis of the O2 antigen in this serovar [27] and found 83

SNPs in this region, of which 7 non-synonymous mutations were present in >10 isolates.

While it is not clear if these mutations affect the O2-antigen chemistry, the low mutation rate

and no observed recombination events in the cluster suggests that the O2 antigen vaccine will

have a broadly protective response against all the *Salmonella* Paratyphi A genotypes sampled

thus far. However, any variations in this region should be carefully monitored through

genomic surveillance.

The conclusions that we can draw from this analysis are subject to certain limitations. First,

the available genomes are an incomplete sample; *Salmonella* Paratyphi A is a neglected

pathogen, and hence the available genomes and might not have broad representativeness

across geographies or time. Specifically, a small proportion of genomes were available from

countries in sub-Saharan Africa and India. Second, while the tool has high sensitivity and

specificity on our dataset, as more genomes become available over time and novel

mechanism of AMR emerge, this tool will require updates from the bigger scientific

community. Like all genotyping tools, Paratype is a living tool that will require updates. Our

diverse group of authors plans to continually monitor the library of publicly available

genomes, accept update requests via GitHub, and incorporate any required updates in the

Paratype scheme accordingly.

Conclusions

This study reports the first large-scale global analysis of *Salmonella* Paratyphi A genomes

and proposes the first genotyping tool for this pathogen. Paratype, which has already been

released (https://github.com/CHRF-Genomics/Paratype) as an open-access, easy-to-use,

command-line tool, is being tested and adopted by researchers for large scale genomic

analysis (https://doi.org/10.5281/zenodo.5520408). This tool will assist future genomic

surveillance studies and will help inform prevention and treatment strategies for this

neglected pathogen.

Methods

Study site and isolate selection

Child Health Research Foundation in Bangladesh has been preserving invasive *Salmonella*

isolates since 1999 and maintains a biobank of >9000 typhoidal *Salmonella* isolates, largely

from children (<18 years of age) that were isolated from the blood of the patients in two

different settings: in-patient (hospitalized), and out-patient (community) facility [30]. Clinical

and epidemiological data were collected for all hospitalized patients. From this biobank,

among 640 *Salmonella* Paratyphi A isolates collected till December 2016, 348 were

randomly selected for whole-genome sequencing (WGS) (Additional file 1: Table S1). A set

of 469 *Salmonella* Paratyphi A isolates were also added to this collection, isolated under the

Surveillance for Enteric Fever in Asia (SEAP) project from three different typhoid-endemic

countries, Bangladesh (n= 180), Nepal (n= 156), and Pakistan (n=133). The SEAP-

Bangladesh isolates (n=180) were selected using randomization to represent 483 isolates

collected between 2016 and 2018. In contrast, SEAP-Nepal isolates included all pre-SEAP

isolates (2014 – 2016) and randomly selected SEAP isolates (2017 – 2019). The SEAP-

Pakistan isolates were selected prioritizing the availability of geographic information and

susceptibility profile during 2016 – 2018.

To add to all the isolates sequenced in this study, we also collected raw fastq data of 560

Salmonella Paratyphi A isolates from 37 different countries and 10 published articles

(Additional file 1: Table S2). Complete chromosomal sequences of *Salmonella* Paratyphi A

ATCC 9150 (NC_006511) and AKU_12601 (NC_011147) were also included [31,32]. For

travel-related paratyphoid cases, the country of "traveling from" was considered as the

country of origin. If no travel data is available, the country of "reported from" was considered

as the country. Overall, for globally distributed 562 *Salmonella* Paratyphi A, year and country

data were available for 507 and 536 respectively (Additional file 1: Table S2). In total, we

obtained a global collection of 1,379 *Salmonella* Paratyphi A covering a timeline of 1917 –

2019 and 37 countries [see Additional file 2 for more details].

<u>Whole-genome sequencing</u>

Salmonella Paratyphi A isolates from 1999-2016 (before the start of the SEAP project) from

Bangladesh (n=348) were subcultured on MacConkey agar media and kept overnight at 37°C.

In case of any visible contamination, a single colony was picked and subcultured again.

Later, all colonies were swapped and resuspended into 1 ml of water. From this suspension,

400 μL was used for DNA extraction using the QIAamp DNA Mini Kit (Qiagen, Hilden,

Germany) and sent to Novogene (NovogeneAIT, Singapore) for WGS on Novaseq 6000

platform (PE150). All SEAP isolates were extracted using the same protocol and were

sequenced on Illumina HiSeq 4000 platform (PE150) at the Welcome Sanger Institute,

Cambridge, UK.

Systematic literature review of existing *Salmonella* Paratyphi A genomes

To contextualize the genomes sequenced in this study, we conducted a systematic search to

compile all publicly available *Salmonella* Paratyphi A genomes (for which raw reads and

metadata were available) to build a database of 560 additional isolates from 10 studies

(Additional file 1: Table S2). First, the search terms "(Salmonella Paratyphi A) AND

(Molecular Epidemiology)" "Salmonella Paratyphi A genome" and "(Salmonella Paratyphi

A) AND (Genomic Epidemiology)" were used in PubMed advanced search builder. Next, the

hits were filtered by selecting dates between 1900 and 2019 and the total number of

publications remaining were 231. After screening the abstracts and titles manually and

eliminating duplicated, only 7 studies were found to have any kind of genome/metadata

available for further analysis. In addition, three studies [8,9,22] that meets our criteria

(published and both metadata and raw reads available) but missed/not published during the

initial PubMed search were incorporated from European Nucleotide Archive (ENA) database,

taking the final number of incorporated publications to 10.

<u>Quality check, genome assembly, annotation, and pan-genome analysis</u>

Raw fastq reads of all *Salmonella* Paratyphi A were quality-checked using FastQC and

trimmed using Trimmomatic if necessary[33]. All 1,377 sets of raw fastq reads were

assembled using Unicycler v0.4.8 (*default with --min_fasta_length 200*)[34]. The assembled

contigs (n = 1,377) and downloaded complete chromosomes (n = 2) were annotated using

Prokka (*--gcode 11 --mincontiglen 200*) [35]. The annotated GFF files of all 1,379 isolates

were used to build a pan- and core-genome of *Salmonella* Paratyphi A using Roary v3.3 (

options: -t 11 -e --mafft -n)[36]. The gene_presence_absence matrix output was used to

perform the Heap's law analysis to understand the open/closedness of the pan-genome (*heaps*

function of *micropan* library on R; 1000 permutations).

<u>SNP-based phylogenetic analyses</u>

For the complete "global+SEAP" raw data collection, fastq reads of 1,377 *Salmonella*

Paratyphi A and fasta of two RefSeq chromosomes (NC_006511 and NC_011147) were

mapped against the *Salmonella* Paratyphi A AKU_12601 (FM200053.1) using Bowtie2

v2.3.5.1 [37]. Candidate SNPs were identified using SAMtools (v1.10) and BCFtools

(v1.10.2) [38]. Only the homozygous, unambiguous SNPs with a Phred-quality score of >20

were selected using a customized Python script. SNPs were discarded if they had strand bias

p <0.001, mapping bias p <0.001 or tail bias p <0.001 (using vcfutils.pl script, from

SAMtools). SNPs located in phage or repeat regions (118.9 kb for *Salmonella* Paratyphi A

AKU_12601 as described in Sajib et al. [25]) were also excluded using a customized python

script. Gubbins v2.3.4 was used to detect the recombinant regions [39] and SNPs in those

regions were excluded as well using the same python script, resulting in a set of 8,346

chromosomal SNPs positions for the "global+SEAP" collection (n= 1,379). All SNP alleles

were extracted (fasta) using a customized python script and merged to produce SNP

alignment.

Maximum likelihood trees (MLT) were built from the chromosomal SNP alignments using

RAxML v8.2.12 (with the Generalized Time-Reversible model and a Gamma distribution to

model site-specific rate variation; GTRGAMMA in RAxML) [15]. Support for the MLT was

calculated using 100 bootstrap pseudo-analyses of the alignment. The MLT was outgroup-

rooted by including the pseudo-alleles from *Salmonella* Typhi CT18 (NC_003198.1) in the

alignment. Tree visualization was done using iTol v5.5 [40], including the previous Paratyphi

A lineages proposed by Zhou *et al* [5].

<u>Bayesian analysis and identifying phylogenetically informative clades and subclades</u>

In addition to SNP-based MLT, we investigated the population structure of the global

Salmonella Paratyphi A collection using a Bayesian approach, implemented with the SNP

alignment using fastBaps[40]. To maintain compatibility with the phylogeny, some minor

modifications were made to the clustering pattern proposed by the least conservative

Dirichlet prior hyperparameters on fastbaps, *optimise.baps*. This eventually resulted in a total

of 16 different clusters. A customized python script was used to randomly select two

isolates/year/cluster to represent this global collection of *Salmonella* Paratyphi A, leading to

two independent sample sets of 315 isolates each. The alignment of SNP-alleles for this

representative sample set was used to understand the evolutionary diverging pattern of

different *Salmonella* Paratyphi A clusters over time using BEAST v1.10.4 [19]. The

GTR+Γ(4) substitution model was selected for this analysis with the exponential unrelated

relaxed clock as clock type and Bayesian skyline coalescent model as tree prior. The analysis

considered the year of isolation as tip dates and continued for 500 million steps with

sampling every 50,000 iterations. The BEAST analysis was run twice each on the two

independently generated sets of isolates. The resulting log files and model parameters were

analyzed on Tracer v1.7.1. TreeAnnotator v1.10 was used to generate the maximum–clade-

credibility (MCC) tree [41]. The tree was visualized on FigTree v1.4.4 with a time scale. For

the model with the highest posterior values (joint effective sample size (ESS) of 544) used for

further analysis, time to last common ancestor (MRCA) was calculated to be 1407 AD (95%

highest posterior density (HPD) interval [721.0, 1637.3]). Based on the diverging patterns

suggested by the MCC tree, we assigned the clusters (defined as described above) into

primary clades, secondary clades, and subclades on the MLT. However, a few visible clusters

on the MLT could not be assigned to specific subclades due to a lack of clustering

information from fastBaps, likely due to the low number of SNPs unique to these clusters.

SNP-based genotyping scheme and paratype

We further divided the 16 clusters obtained from fastBAPS into 18 genotypes and identified a

set of 18 SNP alleles, located in a coding sequence for conserved genes to define each

assigned secondary clade and sub-clades. Each SNP allele was unique to only one subclade

or, to one secondary clade and its corresponding subclades (if any). Therefore, we assigned

the term "genotype" to each of the 18 secondary clades or subclades. Sorted read alignment

(BAM) files generated during the SNP analysis were used to assign the genotypes for each

isolate using a customized Python script, named Paratype (available at

https://github.com/CHRF-Genomics/Paratype). Briefly, under BAM mode *(--mode bam)*,

Paratype uses *samtools index* (if bam file is not indexed), *samtools mpileup,* and *bcftools call*

to extract the consensus base calls at those 18 SNP loci from the BAM file. The resulting

variant call format (VCF) file is then processed to identify the presence of the defining SNP

alleles and follow cladistic logic to assign the genotype of the isolate, as well as the primary

clade, secondary clades, and subclade information. Paratype only considers high-quality SNP

alleles (Phred score >20 and 75% read_ratio for the allele) to assign genotypes. Read_ratio is

calculated by the number of high-quality alternative-allele reads on both strands, divided by

the total number of high-quality reads. In addition, Paratype also has fastq mode (*--mode*

fastq) where a user can provide a set of paired-end raw fastq data file (can be gzipped) and

Paratype performs reference mapping (against the *Salmonella* Paratyphi AKU_12601

genome) using Bowtie and SAMtools and follows the same steps described above to detect

the genotype of the isolates. Although the bam mode is the default for the tool, the fastq

mode is more accurate and should be user-friendly to non-coding specializing researchers;

however, it is more time-consuming. Paratype also runs on vcf mode (*--mode vcf*) which is

faster, but also the least accurate if the provided SNPs are not highly trusted.

Plasmid, resistance gene, and mutation analysis

All assembled contigs were screened with PlasmidFinder v2.1 [21] and ResFinder v3.2 [20]

to detect plasmid amplicons and acquired AMR genes respectively. Both results were parsed

using customized python scripts. To detect mutations in *gyrA* and *acrB* genes, we used the

same Paratype script. It uses the same files used for genotyping and produces gene- and

position-specific non-silent and silent mutation results.

We also explored the genomic region where the genes related to O2-antigen biosynthesis are

located (860,008 – 878,865 of AKU_12601 genome). We detected all SNPs in that region

with the number of isolates having those and their corresponding amino-acid changes using

the Paratype. Two additional python scripts were used to count position-specific SNPs and

mutations for the 18.9 kbp region.

<u>Data visualization and statistical analysis</u>

R (v4.0.4) base function and several packages including dplyr, ggplot2, micropan and

scatterpie were used for data visualization and statistical analysis.

<u>Declarations</u>

<u>Ethics approval and consent to participate</u>

Ethical approval for the parent studies were obtained from the Bangladesh Institute of Child

Health Ethical Review Committee, Nepal Health Research Council, Aga Khan University

Hospital Ethics Committee and Pakistan National Ethics Committee, Stanford University

Institutional Review Board, and U.S. Centers for Disease Control and Prevention. Informed

written consent and clinical information were taken from adult participants and legal

guardians of child participants.

<u>Consent for publication</u>

Not applicable (No data from individual person was used for analysis).

<u>Availability of data and materials</u>

The genome dataset supporting the conclusions of this article are available in the European

Nucleotide Archive (ENA) under study accession ERP132884. The genotyping tool for

552 *Salmonella* Paratyphi A, Paratype is available at https://github.com/CHRF-

553 Genomics/Paratype (https://doi.org/10.5281/zenodo.5520408). Customized Python scripts

554 and color scheme used in the manuscript are available at https://github.com/CHRF-

555 Genomics/CHRF_Paratyphi_scripts. The metadata supporting the conclusions of this article

556 is included in Additional file 2.

558 Competing interests

559 The authors declare no competing interests.

561 Funding

562 This study was partially supported by the Bill and Melinda Gates Foundation (grant numbers

563 INV-023821 and INV-008335). None of the funding bodies had any role in the design of the

564 study, analysis, and interpretation of data or, in writing the manuscript.

566 Authors' contributions

567 AMT, YH, MSIS, SKS and SS were involved in conceptualization and design of the study.

568 MSIS performed the DNA extraction for sequencing in Bangladesh and the literature review

569 for the global database construction. AMT, YH and MSIS performed bioinformatic analysis

570 under supervision of SS and SKS. JRA provided continuous guidance during bioinformatic

571 analysis. AMT and YH designed the genotyping scheme and AMT wrote the Paratype script.

572 YH and MSIS conducted the statistical analyses and visualization. KES, JI, ZAD, SB and

573 JRA reviewed the results. ZAD and SB reviewed genotyping scheme and the Paratype tool.

574 AMT, YH, MSIS and SS wrote the first draft of the manuscript. KES, JI, FNQ, SPL, GD,

ZAD, SB, DOG, JRA and SKS reviewed the manuscript. All authors reviewed and approved

the final manuscript.

Acknowledgements

We are thankful to Mr. Hafizur Rahman, Mr. Dipu Chandra Das and Ms. Nusrat Alam of the

Child Health Research Foundation for their help with the wet-lab procedures. We are also

thankful to the entire SEAP team for their unwavering support and coordination between the

teams.

References

1. Stanaway JD, Reiner RC, Blacker BF, Goldberg EM, Khalil IA, Troeger CE, et al. The

global burden of typhoid and paratyphoid fevers: a systematic analysis for the Global Burden

of Disease Study 2017. The Lancet Infectious Diseases [Internet]. [cited 2019 Feb 28];0.

Available from: https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(18)30685-

6/abstract

2. Crump JA, Mintz ED. Global trends in typhoid and paratyphoid fever. Clin Infect Dis.

2010;50:241–6.

3. Lu X, Li Z, Yan M, Pang B, Xu J, Kan B. Regional Transmission of *Salmonella* Paratyphi

A, China, 1998–2012. Emerg Infect Dis. 2017;23:833–6.

4. Furuse Y. Analysis of research intensity on infectious disease by disease burden reveals

which infectious diseases are neglected by researchers. PNAS. National Academy of

Sciences; 2019;116:478–83.

5. Zhou Z, McCann A, Weill F-X, Blin C, Nair S, Wain J, et al. Transient Darwinian

selection in *Salmonella enterica* serovar Paratyphi A during 450 years of global spread of

enteric fever. Proc Natl Acad Sci U S A. 2014;111:12199–204.

6. Kuijpers LMF, Le Hello S, Fawal N, Fabre L, Tourdjman M, Dufour M, et al. Genomic

analysis of *Salmonella enterica* serotype Paratyphi A during an outbreak in Cambodia, 2013–

2015. Microb Genom [Internet]. 2016 [cited 2019 Aug 25];2. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5320704/

7. Yan M, Yang B, Wang Z, Wang S, Zhang X, Zhou Y, et al. A Large-Scale Community-

Based Outbreak of Paratyphoid Fever Caused by Hospital-Derived Transmission in Southern

China. PLOS Neglected Tropical Diseases. Public Library of Science; 2015;9:e0003859.

8. Ashton PM, Nair S, Peters TM, Bale JA, Powell DG, Painset A, et al. Identification of

Salmonella for public health surveillance using whole genome sequencing. PeerJ. PeerJ Inc.;

2016;4:e1752.

9. Britto CD, Dyson ZA, Duchene S, Carter MJ, Gurung M, Kelly DF, et al. Laboratory and

molecular surveillance of paediatric typhoidal Salmonella in Nepal: Antimicrobial resistance

and implications for vaccine policy. PLOS Neglected Tropical Diseases. 2018;12:e0006408.

10. Saha S, Islam M, Uddin MJ, Saha S, Das RC, Baqui AH, et al. Integration of enteric fever

surveillance into the WHO-coordinated Invasive Bacterial-Vaccine Preventable Diseases (IB-

VPD) platform: A low cost approach to track an increasingly important disease. PLOS

Neglected Tropical Diseases. 2017;11:e0005999.

11. Saha S, Islam M, Saha S, Uddin MJ, Rahman H, Das RC, et al. Designing

Comprehensive Public Health Surveillance for Enteric Fever in Endemic Countries:

Importance of Including Different Healthcare Facilities. J Infect Dis. 2018;218:S227–31.

12. Saha S, Islam MS, Sajib MSI, Saha S, Uddin MJ, Hooda Y, et al. Epidemiology of

Typhoid and Paratyphoid: Implications for Vaccine Policy. Clin Infect Dis. 2019;68:S117–

23.

13. Barkume C, Date K, Saha SK, Qamar FN, Sur D, Andrews JR, et al. Phase I of the

Surveillance for Enteric Fever in Asia Project (SEAP): An Overview and Lessons Learned. J

Infect Dis. 2018;218:S188–94.

14. Day MR, Doumith M, Do Nascimento V, Nair S, Ashton PM, Jenkins C, et al.

Comparison of phenotypic and WGS-derived antimicrobial resistance profiles of *Salmonella*

enterica serovars Typhi and Paratyphi. Journal of Antimicrobial Chemotherapy.

2018;73:365–72.

15. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 2014;30:1312–3.

16. Britto CD, Dyson ZA, Mathias S, Bosco A, Dougan G, Jose S, et al. Persistent circulation

of a fluoroquinolone-resistant *Salmonella enterica* Typhi clone in the Indian subcontinent.

Journal of Antimicrobial Chemotherapy. 2020;75:337–41.

17. Sherchan JB, Morita M, Matono T, Izumiya H, Ohnishi M, Sherchand JB, et al.

Molecular and Clinical Epidemiology of *Salmonella* Paratyphi A Isolated from Patients with

Bacteremia in Nepal. The American Journal of Tropical Medicine and Hygiene. The

American Society of Tropical Medicine and Hygiene; 2017;97:1706–9.

18. Tonkin-Hill G, Lees JA, Bentley SD, Frost SDW, Corander J. Fast hierarchical Bayesian

analysis of population structure. Nucleic Acids Res. 2019;47:5539–49.

19. Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees.

BMC Evolutionary Biology. 2007;7:214.

20. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder 4.0

for predictions of phenotypes from genotypes. Journal of Antimicrobial Chemotherapy.

2020;75:3491–500.

21. Carattoli A, Hasman H. PlasmidFinder and In Silico pMLST: Identification and Typing

of Plasmid Replicons in Whole-Genome Sequencing (WGS). Methods Mol Biol.

2020;2075:285–94.

22. Nair S, Day M, Godbole G, Saluja T, Langridge GC, Dallman TJ, et al. Genomic

surveillance detects *Salmonella enterica* serovar Paratyphi A harbouring blaCTX-M-15 from

a traveller returning from Bangladesh. PLOS ONE. Public Library of Science;

2020;15:e0228250.

23. Jibril AH, Okeke IN, Dalsgaard A, Menéndez VG, Olsen JE. Genomic Analysis of

Antimicrobial Resistance and Resistance Plasmids in *Salmonella* Serovars from Poultry in

Nigeria. Antibiotics. Multidisciplinary Digital Publishing Institute; 2021;10:99.

24. Hooda Y, Sajib MSI, Rahman H, Luby SP, Bondy-Denomy J, Santosham M, et al.

Molecular mechanism of azithromycin resistance among typhoidal Salmonella stains in

Bangladesh identified through passive pediatric surveillance. PLOS Neglected Tropical

Diseases. 2019;13:e0007868.

25. Sajib MSI, Tanmoy AM, Hooda Y, Rahman H, Andrews JR, Garrett DO, et al. Tracking

the Emergence of Azithromycin Resistance in Multiple Genotypes of Typhoidal Salmonella.

mBio [Internet]. American Society for Microbiology; 2021 [cited 2021 Feb 16];12. Available

from: https://mbio.asm.org/content/12/1/e03481-20

26. Sahastrabuddhe S, Carbis R, Wierzba TF, Ochiai RL. Increasing rates of *Salmonella*

Paratyphi A and the current status of its vaccine development. Expert Review of Vaccines.

Taylor & Francis; 2013;12:1021–31.

27. Seif Y, Monk JM, Machado H, Kavvas E, Palsson BO. Systems Biology and Pangenome

of *Salmonella* O-Antigens. mBio [Internet]. American Society for Microbiology; 2019 [cited

2020 Aug 11];10. Available from: https://mbio.asm.org/content/10/4/e01247-19

28. Wong VK, Baker S, Pickard DJ, Parkhill J, Page AJ, Feasey NA, et al. Phylogeographical

analysis of the dominant multidrug-resistant H58 clade of *Salmonella* Typhi identifies inter-

and intracontinental transmission events. Nat Genet. 2015;47:632–9.

29. Wong VK, Baker S, Connor TR, Pickard D, Page AJ, Dave J, et al. An extended

genotyping framework for *Salmonella enterica* serovar Typhi, the cause of human typhoid.

Nature Communications. 2016;7:12827.

30. Saha SK, Baqui AH, Hanif M, Darmstadt GL, Ruhulamin M, Nagatake T, et al. Typhoid

fever in Bangladesh: implications for vaccination policy. The Pediatric Infectious Disease

Journal. 2001;20:521–4.

31. McClelland M, Sanderson KE, Clifton SW, Latreille P, Porwollik S, Sabo A, et al.

Comparison of genome degradation in Paratyphi A and Typhi, human-restricted serovars of

Salmonella enterica that cause typhoid. Nature Genetics. Nature Publishing Group;

2004;36:1268–74.

32. Holt KE, Thomson NR, Wain J, Langridge GC, Hasan R, Bhutta ZA, et al. Pseudogene

accumulation in the evolutionary histories of *Salmonella enterica* serovars Paratyphi A and

Typhi. BMC Genomics. 2009;10:36.

33. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 2014;30:2114–20.

34. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. PLOS Computational Biology. Public

Library of Science; 2017;13:e1005595.

35. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.

2014;30:2068–9.

36. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid

large-scale prokaryote pan genome analysis. Bioinformatics. 2015;31:3691–3.

37. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods.

Nature Publishing Group; 2012;9:357–9.

38. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

39. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid

phylogenetic analysis of large samples of recombinant bacterial whole genome sequences

using Gubbins. Nucleic Acids Research. 2015;43:e15–e15.

40. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic Acids Research. 2019;47:W256–9.

41. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian

phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evolution

[Internet]. 2018 [cited 2021 Mar 28];4. Available from: https://doi.org/10.1093/ve/vey016

Figure legends

Figure 1: Genotyping scheme for *Salmonella* Paratyphi A. The scheme is composed of three primary, nine secondary and 18 genotypes on a phylogenetic tree of 1,379 isolates. The 9 secondary clades as highlighted by the coloring of the inner ring. 18 genotypes identified and are shown in the colored middle ring of the figure. The previously proposed lineage system is shown in the outer ring.

Figure 2: Maximum clade credibility tree of 315 representative *Salmonella* Paratyphi A isolates. The tree shows the last common ancestor of all *Salmonella* Paratyphi A existed at least 600 years ago (tMRCA - 1407 AD). The different genotypes are temporally resolved. Countries with greater than or equal to 5 isolates are also included.

Figure 3: Geographical distribution of *Salmonella* Paratyphi A genotypes. The country of isolation for 1378 sequenced *Salmonella* Paratyphi A isolates is shown. The distribution of genotypes per country is shown as scattered pie charts. The size of the each pie chart represents the number of sequences available. A difference in circulating genotypes is observed indicating local populations differ in several endemic countries. Further details are provided in Additional file 2.

Figure 4: Presence of antimicrobial resistance genes, plasmids, and chromosomal mutations linked to quinolone resistance across different *Salmonella* Paratyphi A genotypes. The diversity of **a)** Antimicrobial resistance genes **b)** Plasmids and **c)** quinolone resistance determining region (QRDR) mutations present *Salmonella* Paratyphi A is shown.

Tree scale: 0.01

① **Secondary Clades**

⬛ 0.1	🟦 2.1		
🟪 1.0	🟦 2.2		
🟪 1.1	🟩 2.3		
🟪 1.2	🟧 2.4		
🟦 2.0			

② **Genotypes**

⬛ 0.1	🟦 2.0	🟩 2.3.1	🟧 2.4.2
🟪 1.0	🟦 2.0.1	🟩 2.3.2	🟥 2.4.3
🟪 1.1	🟦 2.1	🟨 2.3.3	🟥 2.4.4
🟪 1.2.1	🟦 2.2	🟧 2.4	
🟪 1.2.2	🟩 2.3	🟧 2.4.1	

③ **Lineages**

🟦 A	🟨 C5	⬜ NA
🟦 A1	🟧 D	
🟥 B	🟧 E	
🟩 C	🟪 F	
🟩 C4	🟪 G	

Genotypes

■ 0.1	■ 1.2.1	■ 2.0.1	■ 2.3	■ 2.3.3	■ 2.4.2
■ 1.0	■ 1.2.2	■ 2.1	■ 2.3.1	■ 2.4	■ 2.4.3
■ 1.1	■ 2.0	■ 2.2	■ 2.3.2	■ 2.4.1	■ 2.4.4