A booster dose of an inactivated SARS-CoV-2 vaccine increases neutralizing antibodies and T cells that recognize Delta and Omicron variants of concern.

Bárbara M Schultz[1,#], Felipe Melo-González[1,#], Luisa F Duarte[1,#], Nicolás MS Gálvez[1,#], Gaspar A Pacheco[1,#], Jorge A Soto[1,#], Roslye V Berríos-Rojas[1,2], Liliana A González[1,2], Daniela Moreno-Tapia[1,2], Daniela Rivera-Pérez[1,2], Mariana Ríos[1,2], Yaneisi Vázquez[1,2], Guillermo Hoppe-Elsholz[1,2], Omar P Vallejos[1,2], Carolina Iturriaga[3], Marcela Urzua[3], María S Navarrete[4], Álvaro Rojas[4], Rodrigo Fasce[5], Jorge Fernández[5], Judith Mora[5], Eugenio Ramírez[5], Aracelly Gaete-Argel[1,6], Mónica Acevedo[1,6], Fernando Valiente-Echeverría[1,6], Ricardo Soto-Rifo[1,6], Daniela Weiskopf[7], Alba Grifoni[7], Alessandro Sette[7,8], Gang Zeng[9], Weining Meng[9], CoronaVac03CL Study Group, José V González-Aramundiz[10], Pablo A González[1,2], Katia Abarca[1,3], Alexis M Kalergis*[1,2,11], Susan M Bueno*[1,2]

Affiliations:

[1]Millennium Institute on Immunology and Immunotherapy, Santiago, Chile.
[2]Departamento de Genética Molecular y Microbiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile.
[3]Departamento de Enfermedades Infecciosas e Inmunología Pediátrica, División de Pediatría, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile. [4]Departamento de Enfermedades Infecciosas del Adulto, División de Medicina, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile. [5]Departamento de Laboratorio Biomédico, Instituto de Salud

Pública de Chile. [6]Laboratorio de Virología Molecular y Celular, Programa de

Virología, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de

Chile, Santiago, Chile. [7]Center for Infectious Disease and Vaccine Research, La

Jolla Institute for Immunology (LJI), La Jolla, CA 92037, USA. [8]Department of

Medicine, Division of Infectious Diseases and Global Public Health, University of

California, San Diego (UCSD), La Jolla, CA 92037, USA. [9]Sinovac Biotech, Beijing,

China. [10]Departamento de Farmacia, Facultad de Química y de Farmacia,

Pontificia Universidad Católica de Chile, Santiago, Chile. [11]Departamento de

Endocrinología, Facultad de Medicina, Escuela de Medicina, Pontificia Universidad

Católica de Chile, Santiago, Chile.

[#] These authors contributed equally to this work.

*Corresponding authors.

Keywords:

CoronaVac; Phase 3 clinical trial; SARS-CoV-2; COVID-19; inactivated vaccine,

Booster dose.

Abstract

Background: CoronaVac® is an inactivated SARS-CoV-2 vaccine approved by the

World Health Organization. Previous studies reported increased levels of

neutralizing antibodies and specific T cells two- and four-weeks after two doses of

CoronaVac®, but the levels of neutralizing antibodies are reduced at six to eight

months after two doses. Here we report the effect of a booster dose of CoronaVac®

on the anti-SARS-CoV-2 immune response generated against variants of concern

(VOC) Delta and Omicron in adults participating in a phase 3 clinical trial in Chile.

Methods: Volunteers immunized with two doses of CoronaVac® in a four-week

interval received a booster dose of the same vaccine between twenty-four and

thirty weeks after the 2nd dose. Four weeks after the booster dose, neutralizing

antibodies and T cell responses were measured. Neutralization capacities and T

cell activation against VOC Delta and Omicron were detected at four weeks after

the booster dose. **Findings**: We observed a significant increase in neutralizing

antibodies at four weeks after the booster dose. We also observed an increase in

CD4$^+$ T cells numbers over time, reaching a peak at four weeks after the booster

dose. Furthermore, neutralizing antibodies and SARS-CoV-2 specific T cells

induced by the booster showed activity against VOC Delta and Omicron.

Interpretation: Our results show that a booster dose of CoronaVac® increases the

anti-SARS-CoV-2 humoral and cellular immune responses in adults. Immunity

induced by a booster dose of CoronaVac® is active against VOC, suggesting an

effective protection.

Background

The ongoing pandemic caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) has promoted the rapid development of safe, immunogenic, and effective vaccines against SARS-CoV-2 to be used by the general population, which have successfully reduced the transmission of the disease burden. CoronaVac® is an inactivated SARS-CoV-2 vaccine developed by Sinovac Life Sciences Co., Ltd. (Beijing, China) and is among the current vaccines approved by the World Health Organization (WHO) to combat coronavirus disease 2019 (COVID-19) and one of the most used vaccines worldwide[1,2]. Phase I/II clinical trials in China demonstrated that this vaccine induces cellular and humoral response upon immunization [3–5]. Furthermore, the ongoing phase 3 clinical trial in Chile has described an increase in the levels of IgG and neutralizing antibodies in adults aged 18-59 years and $\geq$ 60 years two- and four-weeks after the second dose of CoronaVac® [56]. In addition, this vaccination promotes the activation of a T cell immune response against SARS-CoV-2 antigens in a 0-14 immunization schedule [5] (two-weeks interval), being an effective vaccine to prevent COVID-19 [7,8]. In Chile, 93.7% of the target population has received a first vaccine dose, and 91.4% were fully vaccinated with CoronaVac® on December 10th of 2021 in a 0-28 days vaccination schedule [9]. Although this primary immunization schedule induces neutralizing antibody present in the serum of vaccinated people [10], these titers decrease in time [6,11,12] and have lower levels of neutralization against highly transmissible variants of concern (VOC) as compared to the original vaccine strain, potentially decreasing the effectiveness of these vaccines as new variants emerge

[13–17]. For these reasons, the use of booster doses was approved in Chile in August

2021 in Chile for high-risk populations and adults at five months after

administration of the second dose [18]. In this sense, a report published in October

2021 in Chile, showed that the effectiveness of CoronaVac® against COVID-19

increase from 56% to 80% fourteen days after the application of the booster dose

[19]. Notably, a previous study performed in adults aged 18-59 years old

demonstrated that a booster dose of CoronaVac®, applied six months after the first

dose to individuals that previously received two doses of this vaccine, increased

the levels of antibodies 3-5-fold as compared to the levels observed four weeks

after the second dose [12]. Here, we further extend these findings by reporting the

levels of neutralizing antibodies and specific T cells against SARS-CoV-2 and its

activity against VOC Delta and Omicron in adults ≥18 years old that participated in

the phase 3 clinical trial carried out in Chile, who were vaccinated in a 0-28-days

vaccination schedule and received a booster dose five months after the second

dose.

Materials and methods

Volunteers and sample collection

Blood samples were obtained from volunteers recruited in the clinical trial

CoronaVac03CL (clinicaltrials.gov #NCT04651790) carried out in Chile starting in

November 2020. The Institutional Scientific Ethical Committee of Health Sciences

reviewed and approved the study protocol at the Pontificia Universidad Católica de

Chile (#200708006). Trial execution was approved by the Chilean Public Health

Institute (#24204/20) and was conducted according to the current Tripartite

Guidelines for Good Clinical Practices, the Declaration of Helsinki [20], and local

regulations. Informed consent was obtained from all volunteers upon enrollment.

Volunteers receive two doses of CoronaVac® (3 µg or 600SU of inactivated SARS-

CoV-2 inactivated along with alum adjuvant) in a four-week interval (0-28-day

immunization schedule) and then a booster dose five months after the second

dose. A complete inclusion and exclusion criteria list have been reported [5]. On

November 11th, 2021, one hundred and eighty-six volunteers in the immunogenicity

branch received the booster dose. The antibody and cell mediated immune

responses were evaluated volunteers that had completed all their previous visits in

one of the centers of the study (**Figure 1A**). Blood samples were obtained from all

the volunteers before administration of the first dose (pre-immune), two, four, and

twenty weeks (or five months) after the second dose, and four weeks after the

booster dose (**Figure 1B**).

Procedures

The presence of antibodies against RBD with neutralizing capacities were

measured in sera from seventy-seven volunteers that had completed all their study

visits, including one month after the booster dose of CoronaVac®. The neutralizing

capacities of circulating antibodies were evaluated by a surrogate virus

neutralization test (sVNT) (Genscript Cat#L00847-A). Samples were two-fold

serially diluted starting at a 4-fold dilution until reaching a 512-fold dilution. Assays

were performed according to the instructions of the manufacturer and as reported

previously [5]. Neutralizing antibody titers were determined as the last fold dilution in

which the interaction between hACE2 and RBD was inhibited by 30% or more.

Samples with a percentage of inhibition ≤30% at the lowest dilution (1:4) were

assigned as seronegative with a titer of 2. A sample was considered seropositive

when its titer was higher than the pre-immune titer. The percentage of inhibition

was determined as: 100 * [OD450nm value of negative control - OD450nm value of

sample] / [OD450nm of negative control]. A standard curve was used to plot the

neutralization response in the samples as international units (IU) using the WHO

International Standard for SARS-CoV-2 antibody (NIBSC code 20/136), which was

prepared according to the instructions of the manufacturer [21]. Data were analyzed

using a sigmoidal curve model with a logarithm transformation of the concentration,

and the final concentration for each sample was the average of the product of the

interpolated IU from the standard curve and the sample dilution factor required to

reach the OD450 value that falls within the linear range determined for each

sample. Samples with undetermined concentration at the lowest dilution tested

(1:4) were assigned to the lower limit of quantification (16.4 IU). The Geometric

Mean Units (GMU) and titers (GMT) were represented in **Figure 2** and

Supplementary Figure 1, respectively**. Table 2** shows comparisons among the

visits.

Conventional virus neutralization tests (cVNT) were performed in sixty-two

of the previous seventy-seven volunteers and evaluated as previously reported [5].

Briefly, Vero E6 cells were infected with a SARS-CoV-2 strain obtained by viral

isolation in tissue cultures (33782CL-SARS-CoV-2 strain, D614G variant).

Neutralization assays were carried out by the reduction of cytopathic effect (CPE)

in Vero E6 cells (ATCC CRL-1586). The titer of neutralizing antibodies was defined

as the highest serum dilution that neutralized virus infection, at which the CPE was

absent as compared with the virus control wells (cells with CPE). Vero E6 cells

were seeded in 96-well plates (4×10^4 cells/well). For neutralization assays, 100 μL

of 33782CL-SARS-CoV-2 (at a dose of 100 $TCID_{50}$) were incubated with serial

dilutions of heat-inactivated sera samples (dilutions of 1:4, 1:8, 1:16, 1:32, 1:64,

1:128, 1:256, and 1:512) from participants for 1h at 37 °C. Cytopathic effect on

Vero E6 cells was analyzed 7 days after infection.

A pseudotyped virus neutralization test (pVNT) assay was performed to

assess the capacity of the antibodies against SARS-CoV-2 VOC in samples from

thirty volunteers of the seventy-seven previously analyzed by sVNT. As previously

reported [14], a HIV-1 backbone expressing firefly luciferase as a reporter gene and

pseudotyped with the SARS-CoV-2 spike glycoproteins (HIV-1-SΔ19) from from

lineage B.1 (D614G) or variants Delta (T19R, del157/158, L452R, T478K, D614G,

P681R, D950N) and Omicron (A67V, ΔH69-V70, T95I, Y145D, ΔG142 -V143-

Y144, ΔN211, EPE 213-214, G339D, S371L, S373P, S375F, K417N, N440K,

G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, T547K, D614G,

H655Y, N679K, P681H, N764K, N865K, Q954H, N969K, L981F) was prepared as

previously described [22]. Serum samples were two-fold diluted starting at 1:10 or 1:4

and the estimation of the ID80 was obtained using a 4-parameter nonlinear

regression curve fit measured as the percent of neutralization determined by the

difference in average relative light units (RLU) between test samples and

pseudotyped virus controls. Also, cVNT assays were performed to assess the

capacity of the antibodies against SARS-CoV-2 Delta variant in samples from

nineteen volunteers of the seventy-seven previously analyzed by sVNT.

ELISPOT and flow cytometry assays were performed to evaluate the cellular

immune response in forty volunteers of the seventy-seven previously analyzed

(**Figure 1A**), stimulating PBMCs with four Mega Pools (MPs) of peptides derived

from the proteome of SARS-CoV-2 [23]: peptides from the S protein of SARS-CoV-2

(MP-S), the remaining proteins of the viral particle (excluding S protein peptides)

(MP-R), and shorter peptides from the whole proteome of SARS-CoV-2 (MP-CD8-

A and MP-CD8-B) [23]. Thirty of the previously analyzed volunteers were also

stimulated with three Mega Pools (MP) of VOC, provided by La Jolla Institute for

Immunology. MP derived from the S protein of SARS-CoV-2 WT, SARS-CoV-2

B1.617.2 MP 4326 (Delta variant), and SARS-CoV-2 B1.1.529 MP 4359 (Omicron

variant) [14] were used to evaluate T cell activation at four weeks after the booster

dose. Positive and negative controls were held for each assay. The number of Spot

Forming Cells (SFC) for interferon gamma (IFN-γ) were determined by ELISPOT

and the expression of Activation-Induced Markers (AIM) by T cells was evaluated

by flow cytometry. Assays were performed according to the instructions of the

manufacturer and as reported previously [5]. Further details on the ELISPOT assay,

antibodies used for flow cytometry, and the respective protocols can be found in

the Supplementary information and **Supplementary Table 1**.

Interleukin 2 (IL-2) and IFN-γ secretion were evaluated in the supernatants

of twenty-two volunteers previously stimulated for 20h with SARS-CoV-2 MP of

peptides derived from the Spike protein of VOC, using a Luminex 200 xMap

multiplex system (Luminex Corporation, Austin, TX). The limit of detection for the

cytokine measured ranged from 4.2 to 13,390 pg/mL, according to manufacturer's

instructions. Further experimental details can be found in **Supplementary**

Information.

Statistical analyses

Statistical differences for the immunogenicity results considered repeated measures ANOVA with the Geisser-Greenhouse correction and Dunnet's *a posteriori* multiple tests to compare between the booster dose and the other visits. Analyses were performed over the base 10 logarithms of the data for neutralizing antibody by sVNT, cVNT and pVNT. Cellular immune responses were analyzed by a Friedman test for repeated measures for ELISPOT and flow cytometry for the comparisons between booster dose and the other visits. Secretion of cytokines were compared between the secretion induced by the WT strain against the VOC Delta and Omicron by repeated measures ANOVA. The significance level was set at 0.05 for all the analyses. All data were analyzed with GraphPad Prism 9.0.1.

Results

A booster dose of CoronaVac® induces a significant increase in antibody titers with neutralizing capacity in adults.

One hundred and eighty-six volunteers from the immunogenicity branch that received a booster dose of the CoronaVac® were included in this study. The first dose of the vaccine was inoculated from January to March of 2021, and the second dose was inoculated 28 days after the first one. The neutralizing capacity of serum antibodies was evaluated in seventy-seven and sixty-two volunteers by sVNT and cVNT, respectively, at the five different time-points indicated in **Figure 1B**.

As shown in **Figures 2A and D**, the peak level of antibodies with neutralizing capacity in the total population evaluated, tested by sVNT and cVNT,

is reached at two weeks after the second dose (GMU 168.0, 95% CI=19.5-34.2

and GMT, 95% CI=) and four weeks after the second dose (GMU 124.8, 95%

CI=96.3-161.7 and GMT 13.5, 95% CI=9.6-19.2). However, this neutralizing

capacity significantly decreased twenty weeks after the second dose (GMU 39.0,

95% CI=32.4-47.0 and GMT 8.3, 95% CI=9.6-19.2), which is in line with previous

reports where the immunity against SARS-CoV-2 wanes six months after infection

or vaccination [24,25]. After the booster dose, the neutralizing capacity of the

antibodies increased even more than the one reported two weeks after the second

dose (GMU 499.0, 95% CI=370.6-673.0 and GMT 89.5 ± 64.0-125.2). Overall, we

observed that four weeks after the booster dose the neutralizing capacity increased

more than 12-fold (sVNT) and 10-fold (cVNT) as compared to the response at

twenty weeks after the second dose, and almost 3-fold as compared to two weeks

after the second dose (**Figures 2A and D**).

In adults 18-59 years old, the neutralizing capacity of circulating antibodies

tested by sVNT and cVNT (**Figures 2B and E**, respectively) reached its maximum

four weeks after the booster dose (GMU 918.8, 95% CI=623.4-1354 and 176.9,

95% CI=111.7-280.1) increasing more than 18- and 12-fold as compared to five

months after the second dose (GMU 48.9, 95% CI=37.6-63.5 and GMT 14.2, 95%

CI=7.1-28.4) and more than 4-fold as compared to two weeks after the second

dose (GMU 220.2, 95% CI=150.7-321.7 and GMT 17.5, 95% CI=9.8-31.3)

(**Figures 2B and E**). The seropositivity rate in this group reached 100% at four

weeks after the booster dose (**Table 2**). On the other hand, 53.2% of the total

volunteers analyzed here were adults ≥60 years. In this group, the same tendency

was observed, as seen in **Figure 2C and F**, observing an increase in the level of

neutralizing antibodies evaluated by both techniques of more than 9-fold at four

weeks after the booster dose (GMU 300.5, 95% CI=203.5-443.6 and GMT 47.3,

95% CI=32.1-69.5) as compared to the response observed twenty weeks after the

second dose (GMU 32.4, 95% CI=25.1-41.8 and GMT 5.0, 95% CI=3.5-7.0).

Equivalent to the 18-59 years old group, the seropositivity rate in this age group

reached 100% four weeks after the booster dose (**Table 2**). The seropositivity rate

achieved at four weeks after the booster dose was the highest when compared

with the other visits in this study in the total vaccinated group and in both groups

analyzed.

A booster dose of CoronaVac® induces a robust cellular immune

response in adults.

The cellular responses following a booster dose of CoronaVac® were

evaluated in 40 volunteers. We observed that CD4$^+$ T cell activation was increased

twenty weeks after the second dose as compared to the other time points in both

age groups, suggesting that CoronaVac® can stimulate CD4$^+$ T cell responses that

are sustained over time (**Figure 3A-C**). Importantly, we observed a significantly

further increase in CD4$^+$ T cell activation in both age groups following the booster

dose, as compared to the pre-immune sample and samples obtained at two and

four weeks after the second dose (**Figure 3A-C**). However, this difference was not

significant as compared to the sample obtained twenty weeks after the second

dose (**Figure 3A-C)**. Moreover, we did not observe a significant increase in the

expression of AIM by CD8$^+$ T cells following the booster dose as compared to any

other time point, suggesting that specific CD8$^+$ T cell responses induced by

CoronaVac® are not detected with the current methodologies, even after a third

dose (**Supp. Figure 2A and C**). Accordingly, we observed an increase in IFN-γ

production upon stimulation with mega-pools of peptides (MPs) S and R by

ELISPOT at four weeks after the booster dose for both groups, as compared to the

pre-immune sample (**Figure 3D-F**). As in the case of flow cytometry, we did not

observe significant increase of IFN-γ$^+$ SFCs upon stimulation with CD8 MPs at any

time point (**Supp. Figure 2**). These results suggest that although humoral

responses decrease over time following vaccination with CoronaVac®, CD4$^+$ T cell

responses remain significantly increased as compared to pre-immune samples and

the booster dose promotes a small increase both IFN-γ production and CD4$^+$ T cell

activation that is not significantly different as compared to the levels observed 20

weeks after the second dose.

Neutralizing antibodies and specific T cells induced by a booster dose

of CoronaVac® recognize Delta and Omicron variants of SARS-CoV-2.

As we observed that the neutralization capacity and the T cell responses

increased significantly with the booster dose and knowing that vaccinated

volunteers exhibit decreased neutralization against VOC[14], we proceeded to

evaluate the neutralizing capacities of antibodies in the serum from thirty booster-

vaccinated individuals in pseudotyped virus neutralization test (pVNT) assay

against two variants of concern of SARS-CoV-2, comparing with the level obtained

for the D614G SARS-CoV-2 variant (B.1 lineage, **Figure 4A-B**). We observed that

the titers of antibodies with neutralizing capacities against Delta and Omicron

variant show a significant reduction as compared to the levels achieved for the

D61G variant (D614G: GMT 241.8, CI=155.7-375.6, Delta: 159.2, CI=99.1-256.0

and Omicron: GMT 50.7, CI=30.4-84.8), with a reduction of 1.5 for Delta and 4.8

for Omicron (**Figure 4A**). However, when we compared the changes in

seropositivity for Delta and Omicron (**Figure 4B**), we observed a 93% and 76.7%,

respectively, following the booster dose (**Table 3**). Neutralization assays against

Delta variant with a cVNT in a different group of nineteen volunteers also show that

antibodies induced four weeks after the booster dose have reduced capacity to

neutralize this VOC (**Supp Figure 4A**), although the seropositivity rate observed is

84% (**Supp Figure 4B**).

The cellular responses for VOC following a booster dose of CoronaVac®

were also evaluated in thirty volunteers using MPs of peptides derived from the

Spike protein of Delta and Omicron variants. We observed equivalent numbers of

AIM by CD4$^+$ T cells after four weeks of the booster dose upon stimulation with MP-

S of SARS-CoV-2 WT, Delta, or Omicron variant (**Figure 5A**), with no significant

differences between the response against the MP-S of the variants as compared to

the MP-S of the WT strain. IFN-γ secreting T cells were also analyzed in these

samples and no differences were observed **(Figure 5B)**. We also quantified the

production of different cytokines in the supernatant of PBMCs stimulated with the

MP-s of WT, Delta, and Omicron variants, observing that at four weeks after the

booster dose the stimulated cells secrete equivalent levels of IL-2 and IFN-γ

(**Figure Supp 3**). These results suggest that although the humoral response

measured as neutralization capacities and seroconversion against these VOC is

lower as compared to the humoral response against the D614G strain, the cellular

responses against SARS-CoV-2 VOC is equivalent to the responses elicited by the

wild type strain in volunteers vaccinated with booster dose.

Discussion

In this report we evaluated the humoral and cellular immune response

generated four weeks after the application of a booster dose of inactivated

CoronaVac® vaccine in a cohort of volunteers enrolled in the phase 3 clinical trial

held in Chile. The data reported here showed that although there was an adequate

humoral response after two doses of CoronaVac®, with a 65.9% of effectiveness in

preventing COVID-19 [8], both the sVNT and cVNT assays showed a decrease in

the GMT of neutralizing capacities of circulating antibodies against SARS-CoV2

twenty weeks after the second dose (**Figure 2**). Due to this decrease in

neutralizing capacities, a booster dose of CoronaVac® was evaluated in a clinical

study in China, showing promising results in enhanced humoral immune

responses[12,26]. The evaluation of the immune response reported here shows that

after the booster dose, the neutralizing titers and seroconversion rates increase in

the whole group, to a higher extent than two weeks after the second dose, where

the peak in neutralization was previously observed, which is in line with the

observed by *Clemens et a*l.,[15] . Also, we observed a steady activation of the CD4$^+$

T cells and secretion of IFN-γ during the time-points evaluated (**Figure 3**).

Since the neutralizing antibody titers correlated with protection against

SARS-CoV-2 infection [10], these results likely imply a better outcome and protection

against COVID-19, as reported in previous studies performed in Israel that showed

a decrease in the transmission and the disease severity disease by this virus

twelve or more days after booster inoculation. In Chile, the effectiveness and

prevention in hospitalization increased when assessed fourteen days after the

booster dose of CoronaVac® [19,27]. Another study, performed with a booster dose of

CoronaVac®, showed that an additional dose result in good neutralization capacity

against parental SARS-CoV-2 and against Delta variant four weeks after the

booster dose, generating a long-lasting humoral response, which was due to an

enhancement of the memory immune response generated by B cells [26].

Adults ≥60 years old produced lower levels of antibodies with neutralizing

capacities than the whole group during this study, which was also described

previously (**Figure 2C and F**) [5]. This result is in line with previous data reported for

a population vaccinated in Chile [6], among hospital workers who received two

doses of CoronaVac® [28], and in a study with the mRNA-1273 vaccine [29]. In this

sense, our results are equivalent to those described in a phase I/II of the clinical

trial performed with CoronaVac® in China, showing that the neutralizing antibody

titers in this group decreased at five months after the second dose and that a

booster dose is required 6-8 months after the first vaccination to rapidly increase

and maintain the neutralizing antibody titers [30].

In the case of the T cell response (**Figure 3**), other studies have shown that

Pfizer BNT162b2 and mRNA-1273 induce durable CD4$^+$ T cell activation and

cytokine production up to six months following vaccination, but it remains to be

elucidated whether expression of AIM by CD4[+] T cells and cytokine production

further increase following a booster dose with these vaccines [31,32]. Here, we

observed that the activation of CD4[+] T cells and IFN-γ production stays increased

up to twenty weeks after the second dose, and after the booster dose both

parameters increased in the 18-59 years old group and was maintained at the

levels observed twenty weeks after the second dose in adults ≥60 years old. In

contrast to BNT162b2 and mRNA-1273 vaccines, CoronaVac® delivers not only

the Spike protein upon immunization but also other viral antigens, which may

explain why vaccinated individuals still display AIM[+] CD4[+] T cells five months after

the second dose, regardless of a third dose.

When the neutralization capacity analyzed by pVNT of the VOC Delta and

Omicron was evaluated four weeks after the booster dose, we observed

differences in the neutralization capacity as compared to the D614G variant

(lineage B.1), which does not exhibit mutations in the RBD of the S1 protein

(**Figure 4 and table 3**). We previously reported that CoronaVac® is able to induce

neutralization against the Delta variant at 4 weeks after the second dose, although

to a lesser extent compared to the WT strain [14]. Another study recently reported a

significant increase in the neutralizing capacity after a booster dose with

CoronaVac® for the Delta variant, as compared to the levels observed in volunteers

vaccinated with two doses [26,33]. Although we did not observe similar levels of

enhanced neutralization against the variants Delta after the booster dose using

pVNT (**Figure 4A**), the seropositivity against Delta variant is almost 100% (**Figure

4B**)[33]. Here, we also show that a booster dose induces neutralization against the

variant Omicron, which has rapidly spread worldwide and is the predominant

circulating variant to date [34]. The high number of mutations described in the RBD of

this variant has been associated with increased evasion of neutralizing responses

in either unvaccinated or vaccinated subjects [34,35]. Although the neutralization

observed in subjects vaccinated with a booster dose of CoronaVac® is significantly

lower to the observed against the D614G variant, we observed a seropositivity of

76.7% following the booster, suggesting some degree of protection in most of the

vaccinees. In this sense, it has been reported that a heterologous schedule of

vaccination may induce a higher neutralization ability and a better neutralization

against variants of concern as Delta [36] and Omicron[16]. In line with this, a

heterologous vaccination with CoronaVac® and a booster dose of Pfizer BNT162b2

showed a good neutralization titer against VOC Delta and Omicron, with respect to

the ancestral virus [16]. Similarly, a comparison between heterologous and

homologous booster schedules after the vaccination with CoronaVac®, shows an

increase in neutralization against the VOC Delta and Omicron[15]. There are

discrepancies between the results in neutralization titers, which can be attributed to

the neutralization assays performed and/or the study population; however,

important booster responses are observed in these studies, and seropositivity

reached after the booster dose of CoronaVac® against VOC are also similar [16].

In the case of the cellular response, this is the first report to characterize

CD4+ T cell responses following a booster dose of CoronaVac® against the

Omicron variant of SARS-CoV2. Previous studies using the same MP from VOC

evaluated here have shown that CD4+ T cells respond to VOC in a similar extent

as compared to the ancestral strain in individuals vaccinated with CoronaVac® [14,37]

and mRNA vaccines, which has been explained by the high conservation of T cell

epitopes. In this sense, the booster dose of CoronaVac® induces the expression of

CD4$^+$ T cell activation markers and secretion of IFN-γ and IL-2 against the VOC

Delta and Omicron, which is comparable to the response generated against the

WT strain (**Figure 5**). In line with this, a recent study has shown that T cell

responses against the ancestral strain are cross-reactive against the Omicron

variant in convalescent individuals and volunteers vaccinated with Pfizer

BNT162b2 [38], supporting the idea that the induction of T cell responses against the

ancestral strain may be protective against the Omicron variant.

Our report shows that the booster dose of CoronaVac® in a 0-28 days

schedule induces antibodies with neutralizing capacities, which are higher than the

levels observed at 2- and 4-weeks after the second dose, generating an increased

humoral response even in adults ≥60 years old. Besides this, our results suggest

that a third dose of CoronaVac® supports CD4$^+$ T cell activation, which may confer

either protection or enhanced immune responses against the virus and prevent

severe disease following exposure to SARS-CoV-2 exposure. Importantly, the

humoral and cellular immune response promoted by a booster dose of CoronaVac

shows activity against Delta and Omicron variants and probably results in better

effectiveness of this vaccine during predominance of these VOC.

Limitations

This study presents several limitations, such as the reduced sample size for

the assays and the absence of data for neutralization against Omicron variant with

a conventional viral neutralization test. The assessment of total antibody response

against Spike proteins and other SARS-CoV-2 proteins would also add additional

information about the humoral immune response against SARS-CoV-2 after the

booster dose. Due to the limit of quantification of the technique, samples with

undetermined concentration at the lowest dilution tested (1:4) were assigned the

lower limit of quantification (16.4 IU) and other neutralization assays.

Funding

The CoronaVac03CL Study was funded by The Ministry of Health,

Government of Chile, the Confederation of Production and Commerce (CPC),

Chile and SINOVAC Biotech. NIH NIAID, under Contract 75N93021C00016,

supports AS and Contract 75N9301900065 supports AS, AG and DW. The

National Agency for Research and Development (ANID) through the Fondo

Nacional de Desarrollo Científico y Tecnológico (FONDECYT) grants Nº 1190156

and Nº 1211547 supports RSR and FVE, respectively. The Millennium Institute on

Immunology and Immunotherapy, ANID - Millennium Science Initiative Program

ICN09_016 (former P09/016-F) supports RSR, VFE, SMB, KA, PAG and AMK; The

Innovation Fund for Competitiveness FIC-R 2017 (BIP Code: 30488811-0)

supports SMB, PAG and AMK.

Competing interests

ZG and MW are SINOVAC Biotech employees and contributed to the

conceptualization of the study (clinical protocol and eCRF design) and did not

participate in the analysis or interpretation of the data presented in the manuscript.

A.S. is a consultant for Gritstone, Flow Pharma, Arcturus, Immunoscape,

CellCarta, OxfordImmunotech and Avalia. La Jolla Institute for Immunology (LJI)

has filed for patent protection for various aspects of T cell epitope and vaccine

design work. All other authors declare no conflict of interest.

Acknowledgments

We would like to thank the support of the Ministry of Health, Government of Chile;

Ministry of Science, Technology, Knowledge, and Innovation, Government of Chile;

The Ministry of Foreign Affairs, Government of Chile, and the Chilean Public Health

Institute (ISP). We also would like to thank Rami Scharf, Jessica White, Jorge

Flores and Miren Iturriza-Gomara from PATH for their support on experimental

design and discussion. Alex Cabrera and Sergio Bustos from the Flow Cytometry

Facility at Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile

for support with flow cytometry. We also thank the Vice Presidency of Research

(VRI), the Direction of Technology Transfer and Development (DTD), the Legal

Affairs Department (DAJ) of the Pontificia Universidad Católica de Chile. We are

also grateful to the Administrative Directions of the School of Biological Sciences

and the School of Medicine of the Pontificia Universidad Católica de Chile for their

administrative support. We would also like to thank the members of the

independent data safety monitoring committee (members in the SA) for their

oversight, and the subjects enrolled in the study for their participation and

commitment to this trial. This project has been funded in whole or in part with

Federal funds from the National Institute of Allergy and Infectious Diseases,

National Institutes of Health, Department of Health and Human Services, under

Contract No. Contract No. 75N9301900065 to A.S, A.G. and D.W.

Note: Members of the CoronaVac03CL Study Group are listed in the

Supplementary Appendix (SA).

References

1. Mallapaty S. WHO approval of Chinese CoronaVac COVID vaccine will be crucial to curbing pandemic [Internet]. Nature. 2021 [cited 2021 Oct 6];594(7862):161–2. Available from: https://pubmed.ncbi.nlm.nih.gov/34089030/

2. Q G, L B, H M, et al. Development of an inactivated vaccine candidate for SARS-CoV-2. Science [Internet] 2020 [cited 2021 Oct 6];369(6499):77–81. Available from: https://pubmed.ncbi.nlm.nih.gov/32376603/

3. Z W, Y H, M X, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy adults aged 60 years and older: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. Lancet Infect Dis [Internet] 2021 [cited 2021 Oct 6];21(6):803–12. Available from: https://pubmed.ncbi.nlm.nih.gov/33548194/

4. Y Z, G Z, H P, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18-59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. Lancet Infect Dis [Internet] 2021 [cited 2021 Oct 6];21(2):181–92. Available from: https://pubmed.ncbi.nlm.nih.gov/33217362/

5. SM B, K A, PA G, et al. Safety and Immunogenicity of an Inactivated SARS-CoV-2 Vaccine in a Subgroup of Healthy Adults in Chile. Clin Infect Dis [Internet] 2021 [cited 2021 Oct 9];Available from: https://pubmed.ncbi.nlm.nih.gov/34537835/

6. Sauré D, O'Ryan M, Torres JP, Zuniga M, Santelices E, Basso LJ. Dynamic IgG seropositivity after rollout of CoronaVac and BNT162b2 COVID-19

vaccines in Chile: a sentinel surveillance study. Lancet Infect Dis [Internet]

2021 [cited 2021 Oct 11];0(0). Available from:

http://www.thelancet.com/article/S1473309921004795/fulltext

7.   Duarte LF, Gálvez NMS, Iturriaga C, et al. Immune Profile and Clinical

Outcome of Breakthrough Cases After Vaccination With an Inactivated

SARS-CoV-2 Vaccine. Front Immunol [Internet] 2021 [cited 2021 Oct

11];0:3884. Available from:

https://www.frontiersin.org/articles/10.3389/fimmu.2021.742914/full

8.   Jara A, Undurraga EA, González C, et al. Effectiveness of an Inactivated

SARS-CoV-2 Vaccine in Chile. https://doi.org/101056/NEJMoa2107715

[Internet] 2021 [cited 2021 Oct 7];385(10):875–84. Available from:

https://www.nejm.org/doi/full/10.1056/NEJMoa2107715

9.   Chile ha administrado más de 41 millones 980 mil dosis de vacuna contra

SARS-CoV-2 - Ministerio de Salud - Gobierno de Chile [Internet]. [cited 2021

Dec 10];Available from: https://www.minsal.cl/chile-ha-administrado-mas-de-

41-millones-980-mil-dosis-de-vacuna-contra-sars-cov-2/

10.   Khoury DS, Cromer D, Reynaldi A, et al. Neutralizing antibody levels are

highly predictive of immune protection from symptomatic SARS-CoV-2

infection. Nat Med 2021 277 [Internet] 2021 [cited 2021 Oct 6];27(7):1205–

11. Available from: https://www.nature.com/articles/s41591-021-01377-8

11.   Protection duration after vaccination or infection – Swiss National COVID-19

Science Task Force [Internet]. [cited 2021 Oct 6];Available from:

https://sciencetaskforce.ch/en/policy-brief/protection-duration-after-

vaccination-or-infection/

12.   Pan H, Wu Q, Zeng G, et al. Immunogenicity and safety of a third dose, and

immune persistence of CoronaVac vaccine in healthy adults aged 18-59

years: interim results from a double-blind, randomized, placebo-controlled

phase 2 clinical trial. medRxiv [Internet] 2021 [cited 2021 Oct

6];2021.07.23.21261026. Available from:

https://www.medrxiv.org/content/10.1101/2021.07.23.21261026v1

13.   Cao Y, Yisimayi A, Bai Y, et al. Humoral immune response to circulating

SARS-CoV-2 variants elicited by inactivated and RBD-subunit vaccines. Cell

Res 2021 317 [Internet] 2021 [cited 2021 Oct 8];31(7):732–41. Available

from: https://www.nature.com/articles/s41422-021-00514-9

14.   Melo-González F, Soto JA, González LA, et al. Recognition of Variants of

Concern by Antibodies and T Cells Induced by a SARS-CoV-2 Inactivated

Vaccine. Front Immunol [Internet] 2021;12. Available from:

https://www.frontiersin.org/articles/10.3389/fimmu.2021.747830/full

15.   Clemens SAC, Weckx L, Clemens R, et al. Heterologous versus homologous

COVID-19 booster vaccination in previous recipients of two doses of

CoronaVac COVID-19 vaccine in Brazil (RHH-001): a phase 4, non-

inferiority, single blind, randomised study. Lancet [Internet] 2022 [cited 2022

Jan 24];0(0). Available from:

http://www.thelancet.com/article/S0140673622000940/fulltext

16.   Pérez-Then E, Lucas C, Monteiro VS, et al. Neutralizing antibodies against

the SARS-CoV-2 Delta and Omicron variants following heterologous

CoronaVac plus BNT162b2 booster vaccination. Nat Med 2022 [Internet]

2022 [cited 2022 Jan 24];1–1. Available from:

https://www.nature.com/articles/s41591-022-01705-6

17.   Acevedo M, Alonso-Palomares L, Pantano S, Pasteur I, Navarrete M, Soto-

Rifo R. Neutralization of the emerging SARS-CoV-2 variant Lambda by

antibodies elicited by inactivated virus and mRNA vaccines. 2021 [cited 2022

Jan 31];Available from: https://www.researchsquare.com

18.   Depto Inmunizaciones D, De Prevención D, Control, Y Enfermedades S,

Pública M de salud. DOSIS DE REFUERZO EN LA CAMPAÑA DE

VACUNACIÓN CONTRA SARS-COV-2 EN CHILE. 2021 [cited 2021 Oct

6];Available from: https://www.minsal.cl/wp-

19.   Immunization Campaign against SARS-CoV-2 Early estimates of the

effectiveness of booster shots in Chile [Internet]. 2021 [cited 2021 Dec

10];Available from: https://www.minsal.cl/wp-content/uploads/2021/10/2021-

10-07-EFECTIVIDAD-DOSIS-DE-REFUERZO_ENG.pdf

20.   World medical association. World Medical Association declaration of

Helsinki: Ethical principles for medical research involving human subjects

[Internet]. JAMA - J. Am. Med. Assoc. 2013 [cited 2021 Oct 6];310(20):2191–

4. Available from: https://pubmed.ncbi.nlm.nih.gov/24141714/

21.   WHO/BS.2020.2403 Establishment of the WHO International Standard and

Reference Panel for anti-SARS-CoV-2 antibody [Internet]. [cited 2021 Oct

8];Available from: https://www.who.int/publications/m/item/WHO-BS-

2020.2403

22.   Acevedo ML, Alonso-Palomares L, Bustamante A, et al. Infectivity and

immune escape of the new SARS-CoV-2 variant of interest Lambda.

medRxiv [Internet] 2021 [cited 2022 Jan 24];2021.06.28.21259673. Available

from: https://www.medrxiv.org/content/10.1101/2021.06.28.21259673v1

23. Grifoni A, Weiskopf D, Ramirez SI, et al. Targets of T Cell Responses to

SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and

Unexposed Individuals. Cell 2020;181(7):1489-1501.e15.

24. Lumley SF, O'Donnell D, Stoesser NE, et al. Antibody Status and Incidence

of SARS-CoV-2 Infection in Health Care Workers.

https://doi.org/101056/NEJMoa2034545 [Internet] 2020 [cited 2021 Oct

8];384(6):533–40. Available from:

https://www.nejm.org/doi/10.1056/NEJMoa2034545

25. Sabino EC, Buss LF, Carvalho MPS, et al. Resurgence of COVID-19 in

Manaus, Brazil, despite high seroprevalence. Lancet 2021;397(10273):452–

5.

26. Wang K, Cao Y, Zhou Y, et al. A third dose of inactivated vaccine augments

the potency, breadth, and duration of anamnestic responses against SARS-

CoV-2. medRxiv [Internet] 2021 [cited 2021 Oct 8];2021.09.02.21261735.

Available from:

https://www.medrxiv.org/content/10.1101/2021.09.02.21261735v1

27. Bar-On YM, Goldberg Y, Mandel M, et al. BNT162b2 vaccine booster dose

protection: A nationwide study from Israel. medRxiv [Internet] 2021 [cited

2021 Oct 8];2021.08.27.21262679. Available from:

https://www.medrxiv.org/content/10.1101/2021.08.27.21262679v1

28. Seyahi E, Bakhdiyarli G, Oztas M, et al. Antibody response to inactivated

COVID-19 vaccine (CoronaVac) in immune-mediated diseases: a controlled

study among hospital workers and elderly. Rheumatol Int 2021 418 [Internet]

2021 [cited 2021 Oct 8];41(8):1429–40. Available from:

https://link.springer.com/article/10.1007/s00296-021-04910-7

29. Widge AT, Rouphael NG, Jackson LA, et al. Durability of Responses after

SARS-CoV-2 mRNA-1273 Vaccination.

https://doi.org/101056/NEJMc2032195 [Internet] 2020 [cited 2021 Oct

8];384(1):80–2. Available from:

https://www.nejm.org/doi/full/10.1056/NEJMc2032195

30. Li M, Yang J, Wang L, et al. A booster dose is immunogenic and will be

needed for older adults who have completed two doses vaccination with

CoronaVac: a randomised, double-blind, placebo-controlled, phase 1/2

clinical trial. medRxiv [Internet] 2021 [cited 2021 Oct

8];2021.08.03.21261544. Available from:

https://www.medrxiv.org/content/10.1101/2021.08.03.21261544v1

31. Mateus J, Dan JM, Zhang Z, et al. Low-dose mRNA-1273 COVID-19 vaccine

generates durable memory enhanced by cross-reactive T cells. Science (80-

) [Internet] 2021 [cited 2021 Nov 11];374(6566). Available from:

https://www.science.org/doi/abs/10.1126/science.abj9853

32. Guerrera G, Picozza M, D'Orso S, et al. BNT162b2 vaccination induces

durable SARS-CoV-2 specific T cells with a stem cell memory phenotype.

Sci Immunol [Internet] 2021 [cited 2021 Nov 11];Available from:

https://www.science.org/doi/abs/10.1126/sciimmunol.abl5344

33. Cao Y, Hao X, Wang X, et al. Humoral immunogenicity and reactogenicity of

CoronaVac or ZF2001 booster after two doses of inactivated vaccine. Cell

Res 2021 [Internet] 2021 [cited 2021 Dec 10];1–3. Available from:

https://www.nature.com/articles/s41422-021-00596-5

34.   Viana R, Moyo S, Amoako DG, et al. Rapid epidemic expansion of the

SARS-CoV-2 Omicron variant in southern Africa. Nat 2022 [Internet] 2022

[cited 2022 Jan 31];1–10. Available from:

https://www.nature.com/articles/s41586-022-04411-y

35.   Hoffmann M, Krüger N, Schulz S, et al. The Omicron variant is highly

resistant against antibody-mediated neutralization: Implications for control of

the COVID-19 pandemic. Cell [Internet] 2021 [cited 2022 Jan 25];Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0092867421014951

36.   Ai J, Zhang H, Zhang Q, et al. Recombinant protein subunit vaccine booster

following two-dose inactivated vaccines dramatically enhanced anti-RBD

responses and neutralizing titers against SARS-CoV-2 and Variants of

Concern. Cell Res 2021 [Internet] 2021 [cited 2021 Dec 10];1–4. Available

from: https://www.nature.com/articles/s41422-021-00590-x

37.   A T, J S, N M, et al. Impact of SARS-CoV-2 variants on the total CD4 + and

CD8 + T cell reactivity in infected or vaccinated individuals. Cell reports Med

[Internet] 2021 [cited 2021 Oct 25];2(7). Available from:

https://pubmed.ncbi.nlm.nih.gov/34230917/

38.   Gao Y, Cai C, Grifoni A, et al. Ancestral SARS-CoV-2-specific T cells cross-

recognize the Omicron variant. Nat Med 2022 [Internet] 2022 [cited 2022 Jan

25];1–9. Available from: https://www.nature.com/articles/s41591-022-01700-

x

Figures legends

Figure 1: Study profile, enrolled volunteers, and cohort included in the study by November 11[th], 2021. A. From the one hundred and eighty-six vaccinated individuals that received the booster dose, seventy-seven volunteers that received two doses of CoronaVac® in a 28 days interval (0-28 days schedule of vaccination), were selected from the center assigned for the immunogenicity study. Seventy-seven volunteers were tested for neutralizing antibodies by sVNT, sixty-two were selected to analyze neutralizing antibodies by cVNT and forty were selected to analyze cellular immunity. Analyses for immunity against SARS-CoV-2 variants were performed in 30 volunteers for sVNT, pVNT and T cells assays. **B.** Timeline of 0–28 days schedule of vaccination and booster dose immunization. Text in red denotes timepoints at which blood draws occurred.

Figure 2: Quantification of circulating antibodies inhibiting the interaction between the S1-RBD and hACE2 and in live SARS-CoV-2 in volunteers that received the booster dose of CoronaVac®. A-C. Inhibiting antibodies were detected in serum of volunteers immunized with CoronaVac® using a surrogate Viral Neutralization Test (sVNT), which quantifies the interaction between S1-RBD and hACE2 on ELISA plates. Results were obtained from a total of seventy-seven volunteers (**A**), thirty-six of them were adults between 18-59 years old (**B**), and forty-one of them were ≥ 60 years old (**C**). Data is represented as WHO arbitrary units/mL, the numbers above each set of individual data points show the Geometric Mean Units (GMU), the error bars indicate the 95% CI, and the number

at the right represents the fold increase of the GMU four weeks after the third dose

as compared with the respective times after administration of the second dose. **D-**

F. Neutralizing antibodies were detected in serum of volunteers that received a

booster dose of CoronaVac® twenty weeks after the second dose, using a

conventional Viral Neutralization Test (cVNT), which quantifies the reduction of

cytopathic effect (CPE) in Vero E6 cells infected with SARS-CoV-2. Results were

obtained from 62 volunteers (**D**), 30 of them were adults between 18-59 years old

(**E**), and 32 of them were ≥ 60 years old (**F**). Data are expressed as the reciprocal

of the highest serum dilution preventing 100% cytopathic effect, the numbers

above each set of individual data points show the Geometric Mean Titer (GMT),

the error bars indicate the 95% CI, and the number at the right represents the fold

increase of the GMU the third dose + 4 weeks as compared with the respective

times after administration of the second dose. CI were not adjusted for multiplicity

and should not be used for inference. A repeated measures One-Way ANOVA test

assessed statistical differences to compare all times against 3^{rd} dose + four weeks.

****p<0.0001.

Figure 3: Changes in activation-induced markers (AIMs) expression in CD4$^+$ T

cells and in the number of IFN-γ-secreting cells specific for SARS-CoV-2 after

a booster dose of CoronaVac®. **A-C.** AIM$^+$ CD4$^+$ T cells were quantified in

peripheral blood mononuclear cells of volunteers that received a booster dose of

CoronaVac® twenty weeks after the second dose by flow cytometry, upon

stimulation with mega-pools of peptides derived from SARS-CoV-2 proteins. The

percentage of activated AIM$^+$ CD4$^+$ T cells (OX40$^+$, CD137$^+$) were determined upon

stimulation for 24h with MP-S+R in samples obtained at pre-immune, two weeks

after the second dose, four weeks after the second dose, twenty weeks the second

dose, and four weeks after the booster dose. Data from flow cytometry was

normalized against DMSO and analyzed separately by a Friedman test against the

booster dose. Results were obtained from a total of forty volunteers (**A**), twenty-

one were of them were adults between 18-59 years old (**B**), and nineteen of them

were ≥ 60 years old (**C**). Changes in the secretion of IFN-γ were quantified as the

number of Spot Forming Cells (SFCs) in peripheral blood mononuclear cells of

volunteers that received a booster dose of CoronaVac® 20 weeks after the second

dose. **D-F**. Data was obtained upon stimulation with MP-S+R for 48h in samples

obtained at pre-immune, two weeks after the second dose, four weeks after the

second dose, twenty weeks the second dose, and four weeks after the booster

dose. Results were obtained from a total of 40 volunteers (**D**), 21 were of them

were adults between 18-59 years old (**E**), and 19 of them were ≥ 60 years old (**F**).

Data from ELISPOT were analyzed separately by Friedman test against the

booster dose *p<0.05; **p<0.005; ***p< < 0.001; ****p<0.0001.

Figure 4: Quantification of circulating neutralizing antibodies against SARS-

CoV-2 variants in volunteers that received the booster dose of CoronaVac®.

A. Neutralizing antibodies were detected in the serum of thirty volunteers, four

weeks after the booster dose of CoronaVac®, using a pseudotyped virus

neutralization test (pVNT). Data are expressed as the reciprocal of the highest

dilution preventing 80% of the infection (ID80). Numbers above the bars show the

Mean, and the error bars indicate the 95% CI. The number at the right represents

the fold decrease of the GMT four weeks after the booster dose as compared with

the response of D614G **B**. Seropositivity rate of neutralizing antibodies is shown for

each time point analyzed. Numbers above the bars show the percentage of

seropositivity rate in the respective graphs. Numbers above the bars show the

percentage of seropositivity rate in the respective graphs. A repeated measures

One-Way ANOVA test assessed statistical differences of the GMT to compare

each variant against D614G. $*p<0.05$; $***p < 0.001$; $****p<0.0001$

Figure 5: A booster dose of CoronaVac® induce changes in the number of

IFN-γ-secreting cells and in activation-induced markers (AIMs) expression in

CD4⁺ T cells specific for the Spike protein of SARS-CoV-2 variants. A.

Changes in the secretion of IFN-γ, determined as the number of Spot Forming

Cells (SFCs) were determined. Data was obtained upon stimulation of PBMC with

MP-S of variant of concern of SARS-CoV-2 for 48h in samples obtained four weeks

after the booster dose. Data shown represent mean + 95%CI. Data from thirty

volunteers were analyzed at four weeks after the booster dose to compare among

the MP-S of the variant of concern. Data from ELISPOT were analyzed separately

by Friedman test against the WT MP-S. No significant differences were obtained.

B. AIM⁺ CD4⁺ T cells were quantified in peripheral blood mononuclear cells of thirty

volunteers four weeks after that received a booster dose of CoronaVac® by flow

cytometry, upon stimulation with mega-pools of peptides derived from proteins of

variant of concern of SARS-CoV-2. The percentage of activated AIM$^+$ CD4$^+$ T cells

(OX40$^+$, CD137$^+$) were determined upon stimulation for 24h with MP-S+R in

samples obtained four weeks after the booster dose. Data shown represent mean

+ 95%CI. Data from flow cytometry was normalized against DMSO. No significant

differences were obtained between WT and the variant MP stimulation.

Table

Table 1: Demographic and clinical data of seventy-seven volunteers
analyzed.

763
764

	Age group	AHT N (%)	AR N (%)	MD N (%)	Obesity N (%)	Insulin resistance N (%)	COPD N (%)	HT N (%)
Female sex N (%)	41 (53.2)	11 (14.3)	8 (10.4)	1 (1.3)	6 (7.8)	6 (7.8)	3 (3.9)	7 (9.1)
18-59 years old	18 (23.4)	3 (3.9)	6 (7.8)	0	2 (2.6)	3 (3.9)	0	2 (2.6)
≥60 years old	23 (30.0)	8 (10.4)	2 (2.6)	1 (1.3)	4 (5.2)	3 (3.9)	3 (3.9)	5 (6.5)
Male sex N (%)	36 (46.8)	11 (14.3)	8 (10.4)	3 (3.9)	11 (14.3)	1 (1.3)	0	1 (1.3)
18-59 years old	17 (22.0)	4 (5.2)	4 (5.2)	2 (2.6)	4 (5.2)	0	0	0
≥60 years old	19 (24.6)	7 (9.1)	4 (5.2)	1 (1.3)	7 (9.1)	1 (1.3)	0	1 (1.3)

765 *Arterial hypertension: AHT; Chronic obstructive pulmonary disease: COPD; Mellitus diabetes: MD;*
766 *Hypothyroidism: HT; Allergic rhinitis: AR*
767

Table 2: Seropositivity rates, Geometric Mean Titer (GMT), and Geometric Mean Units (GMU) of circulating neutralizing antibodies against SARS-CoV-2 RBD.

Methodology	Age group	Indicators	2nd dose + 2 weeks	2nd dose + 4 weeks	2nd dose + 20 weeks	3rd dose + 4 weeks
sVNT	Total Vaccine	Seropositivity n/N	72/77	73/77	38/77	75/77
		(%)	93.5	94.8	49.4	97.4
		GMU	168.0	124.8	39.0	499.4
		95% CI	126.8-222.5	96.3-161.7	32.4-47.0	370.6-673.0
		GMT	25.8	16.6	3.5	53.0
		95% CI	19.5-34.2	13.1-21.0	3.0-4.1	40.8-68.8
	18-59	Seropositivity n/N	35/36	36/36	24/36	36/36
		(%)	97.2	97.2	66.7	100
		GMU	220.2	155.0	48.9	918.8
		95% CI	150.7-321.7	108.0-222.6	37.6-63.5	623.4-1354
		GMT	33.3	19.1	4.3	82.8
		95% CI	23.4-47.3	14.0-26.1	3.4-5.4	59.7-114.8
	≥60	Seropositivity n/N	38/41	39/42	15/42	40/42
		(%)	90.5	92.9	35.7	95.2
		GMU	134.1	104.1	32.4	300.5
		95% CI	89.2-201.6	71.8-151.0	25.1-41.8	203.5-443.6
		GMT	20.8	14.7	2.4	36.5
		95% CI	13.6-31.9	10.3-21.0	2.4-3.5	25.3-52.7
cVNT	Total Vaccine	Seropositivity n/N	49/62	51/62	44/62	62/72
		(%)	79.0	82.3	71.0	100
		GMT	12.8	13.5	8.3	89.5
		95% CI	8.8-18.5	9.6-19.2	5.6-12.2	64.0-125.2
	18-59	Seropositivity n/N	25/30	27/30	23/30	30/30
		(%)	83.3	90.0	76.7	100

		GMT	17.5	18.8	14.2	176.9
		95% CI	9.8-31.3	11.2-31.7	7.1-28.4	111.7-280.1
	≥60	Seropositivity n/N	24/32	24/32	21/32	32/32
		(%)	75.0	75.0	65.6	100
		GMT	9.5	9.9	5.0	47.3
		95% CI	5.8-15.4	6.2-15.8	3.5-7.0	32.1-69.5

771   *sVNT: Surrogate Virus Neutralization; cVNT: Conventional Virus Neutralization; GMT: Geometric*
772   *mean titer; GMU: Geometric mean units.*

773 **Table 3: Seropositivity rates, Geometric Mean Titer (GMT) of circulating**
774 **neutralizing antibodies against SARS-CoV-2 RBD of D614G and variants of**
775 **concern (Delta and Omicron).**

	Variant	D614G	Delta (B.1.617.2)	Omicron (B.1.1.529)
pVNT	Indicators	3rd dose + 4 weeks	3rd dose + 4 weeks	3rd dose + 4 weeks
	Seropositivity n/N	30/30	28/30	23/30
	(%)	100	93.3	76.6
	GMT	241.8	159.2	50.7
	95% CI	155.7-375.6	99.1-256.0	30.4-84.8

778 *GMT: Geometric mean titer.*

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5