

Studies of parenchymal texture added to mammographic breast density and risk of breast cancer: a systematic review of the methods used in the literature

Akila Anandarajah¹
Yongzhen Chen²
Graham A. Colditz¹
Angela Hardi¹
Carolyn Stoll¹
Shu Jiang¹

1. Washington University School of Medicine
2. Saint Louis University School of Medicine

Correspondence to Shu Jiang: Jiang.shu@wustl.edu

Abstract

This systematic review aimed to assess the methods used to classify mammographic breast parenchymal features in relation to prediction of future breast cancer including the time from mammogram to diagnosis of breast cancer, and methods for the identification of texture features and selection of features for inclusion in analysis. The databases including Medline (Ovid) 1946-, Embase.com 1947-, CINAHL Plus 1937-, Scopus 1823-, Cochrane Library (including CENTRAL), and Clinicaltrials.gov. were searched through October 2021 to extract published articles in English describing the relationship of parenchymal texture features with risk of breast cancer. Twenty-eight articles published since 2016 were included in the final review. Of these, 7 assessed texture features from film mammograms images, 3 did not report details of the image used, and the others used full field mammograms from Hologic, GE and other manufacturers. The identification of parenchymal texture features varied from using a predefined list to machine-driven identification. Reduction in number of features chosen for analysis in relation to cancer incidence then varied across statistical approaches and machine learning methods. The variation in approach and number of features identified for inclusion in analysis precluded generating a quantitative summary or meta-analysis of the value of these to improve predicting risk of future breast cancers. This updated overview of the state of the art revealed research gaps; based on these, we provide recommendations for future studies using parenchymal features for mammogram images to make use of accumulating image data, and external validation of prediction models that extend to 5 and 10 years to guide clinical risk management. By following these recommendations, we expect to improve risk classification and risk prediction for women to tailor screening and prevention strategies to level of risk.

Funding statement

This work was supported by Breast Cancer Research Foundation grant number (BCRF 21-028), and in part by NCI (R37 CA256810).

Competing interests The authors declare they have no competing interests.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Data availability statement

Data are available upon reasonable request by contacting the corresponding author. All data relevant to the study are included in the article or uploaded as supplementary information.

Introduction

Evolving technology from film mammograms to digital images has changed the sources of data and ease of access to study a range of summary measures from breast mammograms and risk of breast cancer.^{1,2} These include more extreme measures of density and also measures of breast texture features. In particular, as women have repeated mammograms as part of a regular screening program,³⁻⁵ access to repeated images including changing texture features has become more feasible in real time for risk classification and counseling women for their risk management.⁶⁻⁸ Using these features may facilitate improvement in risk classification⁹ and hence more fully support precision prevention for breast cancer.^{8,10}

The leading measure for risk categorization extracted from mammograms is breast density.^{11,12} This is now widely used and reported with many states requiring return of mammographic breast density measures to women as part of routine screening. Mammographic breast density is a strong reproducible risk factor for breast cancer across different approaches used to measure it (clinical judgement or semi/automated estimation), and across regions of the world.¹¹ Other texture features within mammograms have been much less frequently studied for risk stratification and risk prediction. However, growing access to the large data from digital mammograms encourages exploration of additional texture features, complimentary to mammogram density, to assess risk of subsequent breast cancer.^{13,14}

While patterns of breast parenchymal complexity, formed by the x-ray attenuation of fatty, fibroglandular, and stromal tissues, are known to be associated with breast cancer,^{15,16} mammogram density only aims to measure the relative amount of fibroglandular tissue in the breast,¹⁷ which limits the ability to fully capture heterogeneity between patients in the breast tissue. In 2016 Gastounioti and colleagues summarized the literature at that time to classify approaches to parenchymal texture classification: 1) grey-level features – skewness; kurtosis; entropy; and sum intensity, 2) co-occurrence features – entropy; inertia; difference moment; and coarseness, 3) run-length measures grey-level non-uniformity and run-length non-uniformity, 4) structural patterns measures lacunarity, fractal dimension, and 5) multi-resolution spectral features.¹⁸ They conclude from this review that multiparametric texture features may be more effective in predicting breast cancer than single group features. To address use of more comprehensive summaries of these features since their review in 2016, we conducted a systematic review of published studies.

We aim to summarize the methods used to classify mammographic breast parenchymal features in relation to prediction of future breast cancer, the time from mammogram to diagnosis of breast cancer, and the analysis of data from either one or both breasts (averaged or assessed individually). We then identify gaps in evidence to prioritize future studies and speed us to better support precision prevention of breast cancer.

Methods

Eligibility Criteria

Population: We considered all studies of adult women (at least eighteen years old) involving original data. Abstract-only papers, review articles, and conference papers were excluded.

Intervention: We included studies measuring at least one non-density mammographic feature. A study had to explicitly define the mammographic features used to be included. Studies that did not do this, such as those which used a deep model, were excluded.

Outcomes: Our primary outcome of interest was risk of breast cancer, including both invasive and in situ cancers. Risk of breast cancer was required to be dichotomized (yes/no), and analysis of other risks (e.g., risk of interval vs. screen-detected cancer) were not included.

Studies examining the association between mammographic features and other risk factors were excluded to narrow the scope of our paper.

Only studies available in English were included. Additionally, only studies published from 2016 onwards were included to avoid overlap with previous reviews.

Information Sources

The published literature was searched using strategies designed by a medical librarian for the concepts of breast density, mammography, and related synonyms. These strategies were created using a combination of controlled vocabulary terms and keywords, and were executed in Medline (Ovid) 1946-, Embase.com 1947-, CINAHL Plus 1937-, Scopus 1823-, Cochrane Library (including CENTRAL), and Clinicaltrials.gov. Results were limited to English using database-supplied filters. Letters, comments, notes, and editorials were also excluded from the results using publication type filters and limits.

Search Strategy

An example search is provided below (for Embase).

```
('breast density')/exp OR ((breast NEAR/3 densit*):ti,ab,kw OR (mammary NEAR/3  
densit*):ti,ab,kw OR (mammographic NEAR/3 densit*):ti,ab,kw)) AND ('mammography'/deOR  
mammograph*:ti,ab,kwOR mammogram*:ti,ab,kwOR mastrography:ti,ab,kwOR 'digital breast  
tomosynthesis':ti,ab,kwOR 'x-ray breast tomosynthesis':ti,ab,kw)NOT('editorial'/it OR 'letter'/it  
OR 'note'/it) AND [english]/lim
```

The search was completed for the first time on September 9, 2020, and was run again on October 14, 2021 to retrieve citations that were published since the original search. The second search was dated limited to 2020-present (October 2021). Full search strategies are provided in the appendix.

Selection Process

Two reviewers (AA, CS) worked independently to review the titles and abstracts of the records. Next, the two reviewers independently screened the full text of the articles that they did not reject to determine which were eligible for inclusion. Any disagreements of which articles to include were resolved by consensus. Two reviewers (AA, YC) then went through this subset independently and excluded the ones without explicitly defined features.

Data Collection Process

We created a data extraction sheet which two reviewers (AA, YC) used to independently extract data from the included studies. Disagreements were resolved by a third reviewer. If included studies were missing any desired information, any additional papers from the works cited, such as previous reports, methods papers, or protocols, were reviewed for this information.

Data Items

Any estimate of risk of breast cancer was eligible to be included. Risk models could combine multiple texture features or examine texture features individually. Predictive ability could be evaluated using an area under the curve, odds ratio, matched concordance index, hazard ratio, or p value. No restrictions on follow-up time were placed. For studies that reported multiple risk estimates, we prioritized the area under the curve with the most non-mammogram covariates included from the validation study if applicable. If the study did not report an area under the curve, we listed the primary models which were discussed in the results section of the paper.

We collected data on:

the report: author, publication year

the study: location/institution, number of cases, number of controls

the research design and features: lapsed time from mammogram to diagnosis

the mammogram: machine type, mammogram view(s), breast(s) used for analysis

the model: how density was measured, number of texture features extracted, types of texture features extracted, whether feature extraction was machine or human, whether all features were used in analysis, how features for analysis were chosen, non-mammogram covariates included, prediction horizon

Risk of Bias

The objective of this review is to summarize the methods and analysis techniques used to assess parenchymal texture features and risk of breast cancer rather than to quantitatively synthesize the results of the studies. Therefore, a risk of bias assessment, while typically performed in a systematic review, would not serve the objective and was not performed.

Human subjects

This study did not involve human subjects and therefore oversight from an Institutional Review Board was not required.

Registration and protocol

This review was not registered and a protocol was not prepared.

Results

The search and study selection process is shown in Figure 1. A total of 11,111 results were retrieved from the initial database literature search and imported into Endnote. 11 citations from ClinicalTrials.gov were retrieved and added to an Excel file library. After removing duplicates, 4,863 unique citations remained for analysis. The search was run again in October 2021 to retrieve citations that were published since the original search. A total of 1,633 results were retrieved and imported to Endnote. After removing duplicates, including duplicates from the original search, 466 unique citations were added to the pool of results for screening. Between the two searches, a total of 11,577 results were retrieved, and there were 5,329 unique citations.

Of the 5,329 unique citations, 5,124 were excluded based on review of title and abstract. 205 full-text reports were retrieved and assessed for eligibility by two readers. Of these, 177 were excluded for reasons such as not measuring a non-density feature, not having risk of breast cancer as an outcome, being an abstract or a duplicate paper, or not being published in English.

We identified 28 studies published since 2016 that met eligibility criteria as set out in the selection flow chart.¹⁹⁻⁴⁶.

Of the 28 studies, only 7 were based on digitized analogue film images, 3 did not provide details, and the others used full field digital mammograms from Hologic, GE, or other manufacturers, or did not report details (see Table 1). The number of cases included in studies ranged from 20 up to 1900. Of the 28 studies, 8 included fewer than 100 cases.

The approach to assessment of breast parenchymal texture and side of body (ipsilateral or contralateral breast) varied across studies. Almost half of the studies (13 out of 28) used BIRADS as the baseline measure of density. Others used Volpara and machine-derived density

or Cumulus-like approaches. Time from mammogram to diagnosis of breast cancer ranged from under 24 months on average up to 82 months.

Table 2 summarizes how the texture features were identified in the mammographic images. These methods included defined masses or calcifications²³, or a predefined list of texture features such as 34²⁵ or 44²⁶ or even 112⁴⁰ or 944³⁹ initial features. Machine-driven identification of features was also reported. After machine identification, the features were reduced for analysis often based on statistical rules.

We next assessed the value added from addition of texture features to prediction models for breast cancer. Many papers only reported on the association of texture with risk of breast cancer using an odds ratio or relative risk. These were often contemporaneous with diagnosis (measured on contralateral breast).^{33,40,47-49} Model building details and results were not routinely reported. Table 3 gives the AUC for a baseline model without texture and then the value for the model with texture when these were reported separately. Studies were not comparable across design, time horizon, and baseline models. Hence, we did not proceed to a numerical summary such as meta-analysis of AUC values. However, within studies we saw that those that reported concordance for models primarily reported performance of models with MD and then MD plus texture features. In these studies, we saw an increase in reported concordance when texture features were added. Addition of parenchymal features often increased the AUC by 0.05 though Pertuz noted an increase in AUC from 0.609 to 0.786 when using texture features in the contralateral breast at the time of cancer diagnosis.⁴⁸ Furthermore as noted in Table 2, there was substantial variation in the number of texture features included and the method of their identification for inclusion (human defined or machine identified).

The prediction horizon was only defined for 3 of the studies (see Table 3) and was usually the time to the next routine screening mammogram, but less than 3 years on average. Examples of several studies are summarized to give more context to the details in the tables. Eriksson⁵⁰ evaluated data from the KARMA cohort that followed 70,877 women for up to 3 years after baseline mammograms. Median time from the screening mammogram to breast cancer diagnosis was 1.74 years and 433 breast cancers were diagnosed. In their analysis, they used both 2- and 3-year horizons. The AUC improved from 0.64 for the model that included age, MD, and BMI to 0.71 after adding calcifications. Heidari 2018²⁶ chose 4 features associated with asymmetry of mammogram images from a pool of 44 machine-identified features. The time horizon was 12 to 18 months, defined as time to the next screening mammogram. A machine learning classifier was built to predict breast cancer on the subsequent mammogram, reducing the features to a vector with 4 features. The AUC improved from 0.62 to 0.68.

Not all studies reported concordance statistics for their analysis. We summarized 6 studies in Table 4 that reported association measures for texture features with breast cancer.

Discussion

We identified 28 studies that evaluated the value of adding texture features to predict future breast cancer incidence or reported the association between these features and risk. Of these, 7 were based on film images, 3 did not report details, and the others used digital mammograms. Among these, we observed 5 studies using measures of texture that were measured or defined at time of cancer diagnosis and from the contralateral breast. However, 5 studies did not report a time horizon for the interval from mammogram to cancer diagnosis. Those studies evaluating future risk are limited to less than 3 years, on average. When evaluated as a contributor to risk of future breast cancer diagnosis, the common measure of discrimination, the AUC, increases when breast texture measures are added to mammographic breast density. We did not identify

any study using mammography texture features to predict 5- or 10-year risk of breast cancer that would be sufficient to guide prevention.^{6,51-54}

There is substantial variation in the methods used for defining and summarizing the measures of texture. No consistent approach is used to reduce the large number of predefined features, or machine-identified features, to a subset or summary for analysis. Given this variation, we did not combine data across studies but note that comparison within studies shows that texture features are related to breast cancer incidence and improved concordance or AUC. Texture features are important contributors to breast cancer risk beyond mammographic breast density.

In the 2016 review, Gastounioti et al. reviewed features of automated parenchymal texture analysis in relation to breast cancer risk.¹⁸ This included details of methods to classify texture in mammographic images and its contribution to discrimination in case-control studies. Based on the review, they concluded that further research including large prospective studies is needed to establish the predictive value of parenchymal texture for ultimate inclusion in breast cancer risk prediction models. Such prediction models that extend to 5 and 10 years then require external validation to support clinical risk management.

Regarding our methods for this updated systematic review, we do not use risk of bias given the inconsistent approaches and the lack of quantitative data to estimate a summary measure of change in AUC. There are several limitations with the current review. Heterogeneity of the data did not allow for a meta-analysis. Additionally, systematic reviews are always subject to possible publication bias if all relevant studies have not been published. We used several strategies to reduce the risk of this including using a thorough search strategy designed by a medical librarian with expertise in searching for systematic reviews, and searching clinicaltrials.gov for any ongoing studies.

For clinical use to guide precision prevention we must identify both high-risk women for a range of risk reduction strategies⁶ and low-risk women to consider frequency of screening.⁵⁵ From this systematic literature review we identify gaps in evidence to prioritize future studies. They include: i) details on prediction horizon for risk of breast cancer; ii) other statistical approaches might also be used to assess risk of breast cancer from time of image acquisition that include the texture features to the diagnosis of breast cancer such as survival analysis.

Conclusion

Despite current limitations in the literature, the more widespread use of digital mammography and availability of digital images including parenchymal features offer growing opportunity to more uniformly assess image texture features and incorporate these into risk prediction models that can improve risk classification and risk prediction for women.




Figure 1: PRISMA 2020 Flow Diagram

Table 1. Studies of breast texture features classified from mammograms included in systematic review (sorted by year published)

Author	Year	City/ institution	Machine type	View used (CC/MLO/both)	# cases	# controls	Time mmg to dx ca
Choi ²²	2016	University of Ulsan	General Electric Senographe DS and film	both	240	0	mean = 9.7 months (range 6–15 months)
Malkov ³²	2016	USA	film	CC	1171	1659	mean = 5.1 years
Tan ³⁸	2016	University of Pittsburgh Medical Center	digital (not specified further)	both	159	176	The average elapsed time between the “current” and each of “prior” #1, #2 and #3 studies was 1.16 ± 0.41 , 2.30 ± 0.55 and 3.44 ± 0.72 years, respectively.
Winkel ⁴⁴	2016	Bispebjerg Hospital	film	both	121	259	average = 26 months (range 4–45 months)
Ali ²⁰	2017	Sweden	film for main analysis, GE Senographe Essential for validation study	MLO	1170 for main analysis, 69 for validation study	1283 for main analysis, 231 for validation study	less than 3 years before diagnosis (and at latest, at date of diagnosis)
Eriksson ²³	2017	Sweden	digital (not specified further)	both	433 for model development. An additional 137 women lacking information were included in calculating	1732 when comparing study participant characteristics and mammographic features,	median = 1.74 years

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

					the absolute risk estimates, whereby missing data were replaced with the average risk of that risk factor.	60237 for calculating the absolute risk estimate	
Wang ⁴⁰	2017	National Health Service	GE Senographe system	CC	264 for training case-control study, 317 for validation case-control study	787 for training, 931 for validation	Training study: diagnosed at the same time as mammogram. Validation study: average = 3.0 years.
Winkel ⁴³	2017	Copenhagen, Denmark	film	both	288	288	average = 82.0 months, median = 75.5 months, range = 5 to 192 months
Yan ⁴⁵	2017 (August)	NR	Hologic Selenia	CC	83	85	The interval between the prior (negative) and current (cancer detected) examinations are 410.0 +/- 51.7 days for cases.
Yan ⁴⁶	2017 (October)	NR	Hologic Selenia	CC	83	85	12-36 months
Gastounioti ²⁵	2018	University of Pennsylvania	Hologic Selenia Dimensions	MLO	115	460	Average = 1.9 years ± 0.7. Cases had negative screening mammograms at least one year prior

It is made available under a CC-BY-NC-ND 4.0 International license.

							to their diagnosis.
Heidari ²⁶	2018	NR	digital (not specified further)	CC	250	250	the time interval between the “prior” and “current” mammography screenings ranged from 12 to 18 months
Li ³¹	2018	Fudan University Shanghai Cancer Center	NR	CC	84	987	NR
Schmidt ³⁵	2018	Australia and Hawaii, USA	film	CC	1236	2931	Melbourne: Cases were diagnosed, on average, 8 years after MCCS baseline interview (range, 3 months–16 years), and mammography was performed, on average, 2.8 years (standard deviation, 2.6 years; range, 0–14 years) after MCCS baseline. Australia: average 4 years for 32% of cases, and for the other affected women we used the mammogram from the opposite side to that in which the cancer was diagnosed. Hawaii:

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

							The mean time between the earliest mammogram and the breast cancer diagnosis was 6.3 years, while the earliest and the latest mammogram were, on average, 5.1 years apart for cases.
Tagliafico ^{37*}	2018	Italy	Hologic Selenia Dimensions	NR	20	20	cancer was detected at tomosynthesis
Ward ⁴¹	2018	National Health Service	Hologic Selenia	both	34	746	NR
Evans ²⁴	2019	Bradford (UK) Teaching Hospitals NHS Foundation Trust	digital (not specified further)	mix of CC and MLO	58 images from 35 patients with cancer. Experiment 1D included an additional 50 abnormal mammograms with visible cancerous lesions taken from 50 patients.	58 images from 35 patients without cancer. Experiment 1D included an additional 50 normal mammograms taken from 50 patients.	Mammograms used were acquired 3 years prior to the mammograms that had revealed visible and actionable cancer.
Hsu ²⁸	2019	University of California, Los Angeles	NR	NR	463 biopsy results	1675 biopsy results	A false-positive biopsy recommendation was defined by the lack of cancer

It is made available under a CC-BY-NC-ND 4.0 International license.

							within 1 year of the screening examination.
Kontos ²⁹	2019	University of Pennsylvania for case-control sample for evaluating associations to breast cancer, NR for screening sample for phenotype identification.	Hologic Selenia Dimensions for screening sample, GE Senograph 2000D and DS for case-control sample	both for screening sample, NR for case-control sample	Screening sample included 18 detected cases with 12 in the training sample and 6 in the testing sample. 76 cases were in case-control sample.	Screening sample included 2011 controls with 1327 in the training sample and 684 in the testing sample. 158 controls were in case-control sample.	For screening sample, within 1 year. Not specified for case-control sample.
Pérez-Benito ³³	2019	Valencian Community, Spain	FUJIFILM, IMS s.r.l., Giotto IRE, HOLOGIC, SIEMENS, or unknown	both	808 cases with 606 in training set and 202 in test set	755 with 566 in training set and 189 in test set	NR
Pertuz ³⁴	2019	Tampere University Hospital	Philips MicroDose SI or General Electric Senograph Essential	CC	114	114	NR
Tan ³⁹	2019	Subang Jaya Medical Center	Hologic Selenia	CC	250	250	within a year
Abdolell ¹⁹	2020	NR	Siemens MAMMOMAT Inspiration or MAMMOMAT Novation DRimaging	both	1882	5888	NR

It is made available under a CC-BY-NC-ND 4.0 International license.

			system				
Ma ³⁰	2020	Nanfang Hospital	Hologic Selenia Dimensions	both	608 for risk model development, 203 for validation	1261 for risk model development, 421 for validation	at least 1 year later for validation
Sorin ³⁶	2020	NR	GE Senograph Essential	both	53	463	Cancer cases were defined as all cancers detected at the time of CESM imaging as well as cancers diagnosed during the follow-up period. Controls had at least 1 year follow up.
Azam ²¹	2021	Sweden	General Electric, Philips, Sectram Hologic, and Siemens	both	676	52597	The median number of years between the last negative mammogram and the date of diagnosis was 2.8 years.
Heine ²⁷	2021	Mayo Clinic for first case-control study, the San Francisco Mammography Registry for generalization study	Hologic Selenia	CC	514 for first study, 1474 for generalization study	1377 for first study, 2942 for generalization study	at least 6 months
Warner ⁴²	2021	USA	film	CC	1900	3921	median = 4.1 years

*Tagliafico 2018 used digital breast tomosynthesis

It is made available under a CC-BY-NC-ND 4.0 International license.

Table 2. Features of mammographic breast images used to assess breast texture features in addition to mammographic breast density and breast cancer risk (sorted by year).

Author	Year	Side used (L/R/bot h/av)	Density (BIRAD categories/continuous)	Number of texture features extracted (other than density)	Types of Texture features extracted (list all)	Machine or human extraction?	All texture feature used in the model (yes/no)	How features for analysis are chosen
Choi	2016	NR	BIRADS	8	normal appearing tissue, benign-appearing calcification, mass, calcification, architecture distortion, focal asymmetry	human	yes	N/A
Malkov	2016	avg	Cumulus and custom software comparable to Cumulus	46	first- and second-order features, Fourier transform, and fractal dimension analysis	machine	no	AUCs for each feature individually given
Tan	2016	both	computer-aided detection scheme	158	mammographic density, structural similarity, and texture based image features	machine	no	stepwise regression analysis
Winkel	2016	both	BIRADS	1	Tabár classification of parenchymal	human	yes	N/A

It is made available under a CC-BY-NC-ND 4.0 International license.

					patterns			
Ali	2017	contralateral for cases, random side chosen for controls	Cumulus and automated measure of area PD	13	spatial organisation of dense vs. fatty regions of the breast	machine	yes for AUC given, no for further analysis	stepwise selection procedure
Eriksson	2017	both	BIRADS and STRATUS	2	calcifications and masses	machine	yes	N/A
Wang	2017	Training: contralateral for cases and the left for controls. Validation: contralateral for cases and the same side for controls.	Volpara	112	features based on a grey-level co-occurrence matrix, neighborhood grey-tone difference matrix, form and shape of breast boundary, run-length, and grey-level size zone matrix, and statistical moments of pixel values	machine	no	selected from training set using least absolute shrinkage and selection operator
Winkel	2017	both	BIRADS and Cumulus-like approach	1	Tabár classification of parenchymal patterns	human	yes	N/A
Yan	2017 (August)	both	BIRADS	148	bilateral mammographic tissue asymmetry	machine	no	WEKA data mining and machine learning

It is made available under a CC-BY-NC-ND 4.0 International license.

					maximum features			software package
Yan	2017 (October)	both	mutual threshold	220	asymmetry, mean and maximum features	machine	no	WEKA data mining and machine learning software package
Gastounioti	2018	avg	BIRADS, LIBRA, and Quantus	34	anatomically-oriented texture features	machine	no	Identified pairs of features with absolute Pearson correlation greater than 0.90 and for each pair removed the feature with the lowest variability in terms of its interquartile range. Starting from the remaining features, elastic net regression with nested cross-validation was used to build a parsimonious logistic regression model with the most discriminatory

It is made available under a CC-BY-NC-ND 4.0 International license.

								subset of covariates.
Heidari	2018	both	BIRADS	44	bilateral asymmetry of mammographic tissue density distribution	machine	no	locally preserving projection based feature combination algorithm
Li	2018	both	AutoDensity	1	breast area	machine	yes	N/A
Schmidt	2018	avg	Cumulus	20	gray-level co-occurrence matrix textural features	machine	yes	N/A
Tagliafico	2018	avg	BIRADS	104	radiomics features including skewness, energy, entropy, kurtosis, 90percentile and dissimilarity	machine	no	selected to reduce the risk of over-fitting and according to features previously used to associate breast parenchymal patterns with cancer risk
Ward	2018	both	Volpara	1	patterns of parenchymal tissue	human	yes	N/A
Evans	2019	both	scale similar to BIRADS used	1	non-localizable global gist signal	human	yes	N/A
Hsu	2019	NR	BIRADS	5	presence of lump, mass, calcification, architecture distortion, asymmetry	human	no	presence of lump included in final model. mass, calcifications, architecture

It is made available under a CC-BY-NC-ND 4.0 International license.

								distortion, asymmetry examined individually with PPV values given.
Kontos	2019	avg	BIRADS and LIBRA	29	phenotypes of mammographic parenchymal complexity based on four main types of features: histogram, co-occurrence, run-length, and structural	machine	no	excluded features with extremely low variation and those with extreme skewness
Pérez-Benito	2019	contralateral	DMScan	23	geometrical features and a global feature based on local histograms of oriented gradients	machine	yes	N/A
Pertuz	2019	contralateral for cases, right for controls	Cumulus-like approach	37	parenchymal features including computational features and imaging parameters related to the mammographic system (compressed breast)	machine	yes	N/A

It is made available under a CC-BY-NC-ND 4.0 International license.

					thickness, compression force, X-ray tube voltage peak and target-filter combination)			
Tan	2019	contralat- eral	Volpara	944	grey-level co- occurrence matrix features, structural/patter- n measures, grey-level intensity/histogr- am features, run-length features, and multiresolution/ spectral features	machine	no	stepwise regression analysis
Abdolell	2020	NR	Densitas	1	breast volume	machine	yes	N/A
Ma	2020	both	BIRADS	1	normalized average glandular dose	machine	yes	N/A
Sorin	2020	both	BIRADS	1	background parenchymal enhancement	human	yes	N/A
Azam	2021	both	STRATUS	1	microcalcificatio- n clusters	machine	yes	N/A
Heine	2021	avg	Volpara	4	variation measures	machine	no	the two variants of V produced similar findings so only one was discussed in the results
Warner	2021	both	Cumulus and	1	gray-scale	machine	yes	N/A

			automated computer algorithm		variation			
--	--	--	------------------------------------	--	-----------	--	--	--

It is made available under a CC-BY-NC-ND 4.0 International license.

Table 3. Analytical models used for breast cancer risk with addition of breast texture features in addition to mammographic breast density (sorted by year).

Author	Year	non mammogram covariates included (e.g., age, parity, etc)	Prediction horizon (<5?/5/10yr)	AUC (baseline model)	Overall AUC (with texture features added)
Malkov	2016	adjusted for age, body mass index, first-degree family history, percent density, study	NR	0.617	0.621
Tan	2016	none	NR	NR	0.730
Winkel	2016	adjusted for age	NR	BI-RADS density = 0.63	0.65
Ali	2017	age, BMI, density, hormone replacement therapy status, parity, age at first birth	NR	0.687 for apparent, 0.634 for honest	0.703 for apparent, 0.643 for honest
Eriksson	2017	percentage mammographic density, age at mammography, BMI, family history of breast cancer, HRT use	2 years (for main model) and 3 years (relative risks given)	0.64	0.71 with density and interaction between percentage density and masses also included
Wang	2017	adjusted for age, BMI	NR	mC = 0.57	mC = 0.58
Winkel	2017	adjusted for birth year, age at false-positive-screen, invitation round at false-positive-screen	NR	BI-RADS density = 0.65, percentage mammographic density = 0.62	0.63
Yan	2017 (August)	none	next sequential mammography screening	NR	0.816
Yan	2017 (October)	none	next sequential sequencing	NR	0.830
Gastounioti	2018	density, BMI, age	NR	0.62	0.67
Heidari	2018	none	12 to 18 months	NR	0.68
Schmidt	2018	adjusted for age, BMI	NR	percent mammographic density = 0.620	0.662
Tagliafico	2018	none	NR	NR	0.567

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Evans	2019	none	3 years	NR	0.54
Hsu	2019	BI-RADS, density, age, race, BMI, age at first live birth, noticeable changes in breast, number of risk factors, 5-year Gail risk $\geq 1.67\%$	NR	cv-AUC = 0.83 with BI-RADS and density only	cv-AUC = 0.84
Kontos	2019	density, BMI	NR for AUC model. 5-year risk from Gail and Breast Cancer Surveillance Consortium models compared by phenotype.	0.80	0.84
Pérez-Benito	2019	percent density	NR	0.560	0.614
Pertuz	2019	none and with age, percent density	NR	density only = 0.609	0.786
Tan	2019	none, with age only, and with age and BMI	NR	density = 0.52	0.68
Abdolell	2020	age, percent mammographic density	tailored estimates of current breast cancer risk	0.584	0.597
Ma	2020	age, age at menarche, menopausal status, age at first birth, parity, family history of breast cancer, breast density	NR	0.61 for training set and 0.56 for test set	0.77 for training set and 0.75 for test set
Heine	2021	adjusted for study, age, body mass index and also with dense volume included	NR	volumetric breast density = 0.61 for first study and 0.59 for generalization study	$V = 0.61, P_1 = 0.61, p_1 = 0.60$ for first study. $V = 0.59, P_1 = 0.57, p_1 = 0.58$ for generalization study.

It is made available under a CC-BY-NC-ND 4.0 International license.

Table 4. Studies without AUC (sorted by year).

Author	Year	non mammogram covariates included (e.g., age, parity, etc)	Prediction horizon (<5?/5/10yr)	Risk other than AUC
Choi	2016	N/A	N/A	In the minimal sign group, the most common finding was normal appearing tissue (61/78), followed by benign-appearing calcification (17/78).
Li	2018	none	NR	OR = 1.018
Ward	2018	none	NR	There was a significant correlation between a diagnosis of cancer and nodular parenchymal pattern compared to not nodular parenchymal pattern ($p = 0.043$).
Sorin	2020	adjusted for age, family history, breast density	NR	The odds for breast cancer significantly increased with increased BPE (OR = 2.24).
Azam	2021	adjusted for BMI, baseline mammographic density, smoking status, alcohol consumption, age at menarche, age at first birth, number of children, breastfeeding duration, oral contraceptive use, menopausal hormone therapy use, family history of breast cancer	NR	Each additional microcalcification cluster was associated with 20% increased risk of breast cancer (HR = 1.20). Women with ≥ 3 microcalcification clusters had an overall 2-fold increased risk of breast cancer compared to women with no clusters (HR = 2.17).
Warner	2021	adjusted for age, fasting status, time of blood draw, body mass index, menopausal status, hormone therapy use, mammography read batch and also with either percent mammographic density or automated percent density	NR	V was positively associated with breast cancer risk (OR = 1.15 with percent mammographic density, 1.18 with automated percent density).

Literature cited

- 1 van Ravesteyn, N. T. *et al.* Transition from film to digital mammography: impact for breast cancer screening through the national breast and cervical cancer early detection program. *Am J Prev Med* **48**, 535-542, doi:10.1016/j.amepre.2014.11.010 (2015).
- 2 Hopper, J. L. *et al.* Going Beyond Conventional Mammographic Density to Discover Novel Mammogram-Based Predictors of Breast Cancer Risk. *J Clin Med* **9**, doi:10.3390/jcm9030627 (2020).
- 3 Oeffinger, K. C. *et al.* Breast Cancer Screening for Women at Average Risk: 2015 Guideline Update From the American Cancer Society. *JAMA* **314**, 1599-1614, doi:10.1001/jama.2015.12783 (2015).
- 4 Siu A. L. & U. S. Preventive Services Task Force. Screening for Breast Cancer: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med* **164**, 279-296, doi:10.7326/M15-2886 (2016).
- 5 Qaseem, A. *et al.* Screening for Breast Cancer in Average-Risk Women: A Guidance Statement From the American College of Physicians. *Ann Intern Med* **170**, 547-560, doi:10.7326/M18-2147 (2019).
- 6 Bevers, T. B. *et al.* Breast Cancer Risk Reduction, Version 2.2015. *J Natl Compr Canc Netw* **13**, 880-915, doi:10.6004/jnccn.2015.0105 (2015).
- 7 Louro, J. *et al.* A systematic review and quality assessment of individualised breast cancer risk prediction models. *Br J Cancer* **121**, 76-85, doi:10.1038/s41416-019-0476-8 (2019).
- 8 Brooks, J. D. *et al.* Personalized Risk Assessment for Prevention and Early Detection of Breast Cancer: Integration and Implementation (PERSPECTIVE I&I). *J Pers Med* **11**, doi:10.3390/jpm11060511 (2021).
- 9 Jiang, S., Cao, J., Rosner, B. & Colditz, G. Supervised two-dimensional functional principal component analysis with time-to-event outcomes and mammogram imaging data. *Biometrics* In press (2021).
- 10 Pashayan, N. *et al.* Personalized early detection and prevention of breast cancer: ENVISION consensus statement. *Nat Rev Clin Oncol* **17**, 687-705, doi:10.1038/s41571-020-0388-9 (2020).
- 11 Boyd, N. F., Guo, H., Martin, L.J., Sun, L., Stone, J., Fishell, E., Jong, R.A, Hislop, G., Chiarelli, A., Minkin, S., Yaffe, M.J. Mammographic density and the risk and detection of breast cancer. *N Engl J Med* **356**, 227-236 (2007).
- 12 Boyd, N. F. *et al.* Mammographic breast density as an intermediate phenotype for breast cancer. *Lancet Oncol* **6**, 798-808, doi:10.1016/S1470-2045(05)70390-9 (2005).
- 13 Salazar, A. S., Rakhmankulova, M., Simon, L. E. & Toriola, A. T. Chemoprevention Agents to Reduce Mammographic Breast Density in Premenopausal Women: A Systematic Review of Clinical Trials. *JNCI Cancer Spectr* **5**, pkaa125, doi:10.1093/jncics/pkaa125 (2021).
- 14 Brentnall, A. R. *et al.* Mammographic density change in a cohort of premenopausal women receiving tamoxifen for breast cancer prevention over 5 years. *Breast Cancer Res* **22**, 101, doi:10.1186/s13058-020-01340-4 (2020).

- 15 Kaufhold, J., Thomas, J., Eberhard, J., Galbo, C. & Trotter, D. G. A calibration approach to glandular tissue composition estimation in digital mammography. *Medical physics* **29**, 1867-1880 (2002).
- 16 Wolfe, J. N. Breast patterns as an index of risk for developing breast cancer. *American Journal of Roentgenology* **126**, 1130-1137 (1976).
- 17 Yaffe, M. J. Mammographic density. Measurement of mammographic density. *Breast Cancer Research* **10**, 209 (2008).
- 18 Gastounioti, A., Conant, E. F. & Kontos, D. Beyond breast density: a review on the advancing role of parenchymal texture analysis in breast cancer risk assessment. *Breast Cancer Res* **18**, 91, doi:10.1186/s13058-016-0755-8 (2016).
- 19 Abdolell, M. P., J. I. : Caines, J. : Tsuruda, K. : Barnes, P. J. : Talbot, P. J. : Tong, O. : Brown, P. : Rivers-Bowerman, M. : Iles, S. Assessing breast cancer risk within the general screening population: developing a breast cancer risk model to identify higher risk women at mammographic screening. *European Radiology*, doi:10.1007/s00330-020-06901-x (2020).
- 20 Ali, M. A. C., K. : Eriksson, L. : Hall, P. : Humphreys, K. Breast tissue organisation and its association with breast cancer risk. *Breast Cancer Research* **19**, doi:10.1186/s13058-017-0894-6 (2017).
- 21 Azam, S. et al. Mammographic microcalcifications and risk of breast cancer. *Br J Cancer*, doi:10.1038/s41416-021-01459-x (2021).
- 22 Choi, W. J. C., J. H. : Kim, H. H. : Shin, H. J. : Chae, E. Y. Analysis of prior mammography with negative result in women with interval breast cancer. *Breast Cancer* **23**, 583-589, doi:10.1007/s12282-015-0606-y (2016).
- 23 Eriksson, M. C., K. : Pawitan, Y. : Leifland, K. : Darabi, H. : Hall, P. A clinical model for identifying the short-term risk of breast cancer. *Breast Cancer Research* **19**, doi:10.1186/s13058-017-0820-y (2017).
- 24 Evans, K. K. C., A. M. : Wolfe, J. M. Detecting the "GIST" of breast cancer in mammograms three years before localized signs of cancer are visible. *British Journal of Radiology* **92**, doi:10.1259/bjr.20190136 (2019).
- 25 Gastounioti, A. H., M. K. : Cohen, E. : Pantalone, L. : Conant, E. F. : Kontos, D. Incorporating Breast Anatomy in Computational Phenotyping of Mammographic Parenchymal Patterns for Breast Cancer Risk Estimation. *Scientific reports* **8**, 17489, doi:10.1038/s41598-018-35929-9 (2018).
- 26 Heidari, M. et al. Prediction of breast cancer risk using a machine learning approach embedded with a locality preserving projection algorithm. *Phys Med Biol* **63**, 035020, doi:10.1088/1361-6560/aaa1ca (2018).
- 27 Heine, J. et al. Mammographic Variation Measures, Breast Density, and Breast Cancer Risk. *AJR Am J Roentgenol* **217**, 326-335, doi:10.2214/AJR.20.22794 (2021).
- 28 Hsu, W. Z., X. : Petrusse, A. : Chau, N. : Lee-Felker, S. : Hoyt, A. : Wenger, N. : Elashoff, D. : Naeim, A. Role of Clinical and Imaging Risk Factors in Predicting Breast Cancer Diagnosis Among BI-RADS 4 Cases. *Clinical Breast Cancer* **19**, e142-e151, doi:10.1016/j.clbc.2018.08.008 (2019).

- 29 Kontos, D. et al. Radiomic Phenotypes of Mammographic Parenchymal Complexity: Toward Augmenting Breast Density in Breast Cancer Risk Assessment. *Radiology* **290**, 41-49, doi:10.1148/radiol.2018180179 (2019).
- 30 Le Ma, Y. C., Xiaojia Lin, Zilong He, Hui Zeng, Weigu Chen, Genggeng Qin. Association of the Differences in Average Glandular Dose with Breast Cancer Risk. *BioMed Research International* **2020**, Article ID 8943659, doi:10.1155/2020/8943659 (2020).
- 31 Li, T. T., L. : Gandomkar, Z. : Heard, R. : Mello-Thoms, C. : Shao, Z. : Brennan, P. Mammographic density and other risk factors for breast cancer among women in China. *Breast J* **24**, 426-428, doi:<https://dx.doi.org/10.1111/tbj.12967> (2018).
- 32 Malkov, S. S., J. A. : Scott, C. G. : Tamimi, R. M. : Ma, L. : Bertrand, K. A. : Couch, F. : Jensen, M. R. : Mahmoudzadeh, A. P. : Fan, B. : Norman, A. : Brandt, K. R. : Pankratz, V. S. : Vachon, C. M. : Kerlikowske, K. Mammographic texture and risk of breast cancer by tumor type and estrogen receptor status. *Breast Cancer Research* **18**, doi:10.1186/s13058-016-0778-1 (2016).
- 33 Pérez-Benito, F. J. S., F. : Pérez-Cortés, J. C. : Pollán, M. : Pérez-Gómez, B. : Salas-Trejo, D. : Casals, M. : Martínez, I. : Llobet, R. Global parenchymal texture features based on histograms of oriented gradients improve cancer development risk estimation from healthy breasts. *Computer Methods and Programs in Biomedicine* **177**, 123-132, doi:10.1016/j.cmpb.2019.05.022 (2019).
- 34 Pertuz, S. S., A. : Karivaara-Mäkelä, M. : Holli-Helenius, K. : Lääperi, A. L. : Rinta-Kiikka, I. : Arponen, O. : Kämäräinen, J. K. Micro-parenchymal patterns for breast cancer risk assessment. *Biomedical Physics and Engineering Express* **5**, doi:10.1088/2057-1976/ab42f4 (2019).
- 35 Schmidt, D. F. et al. Cirrus: An Automated Mammography-Based Measure of Breast Cancer Risk Based on Textural Features. *JNCI Cancer Spectr* **2**, pky057, doi:10.1093/jncics/pky057 (2018).
- 36 Sorin, V. Y., Y. : Shalmon, A. : Gotlieb, M. : Faermann, R. : Halshtok-Neiman, O. : Sklair-Levy, M. Background Parenchymal Enhancement at Contrast-Enhanced Spectral Mammography (CESM) as a Breast Cancer Risk Factor. *Acad. Radiol.* **27**, 1234-1240, doi:10.1016/j.acra.2019.10.034 (2020).
- 37 Tagliafico, A. S. V., F. : Mariscotti, G. : Durando, M. : Nori, J. : La Forgia, D. : Rosenberg, I. : Caumo, F. : Gandolfo, N. : Houssami, N. : Calabrese, M. An exploratory radiomics analysis on digital breast tomosynthesis in women with mammographically negative dense breasts. *Breast* **40**, 92-96, doi:10.1016/j.breast.2018.04.016 (2018).
- 38 Tan, M., Zheng, B., Leader, J. K. & Gur, D. Association Between Changes in Mammographic Image Features and Risk for Near-Term Breast Cancer Development. *IEEE Trans Med Imaging* **35**, 1719-1728, doi:10.1109/TMI.2016.2527619 (2016).
- 39 Tan, M. M., S. : Yip, C. H. : Ng, K. H. : Teo, S. H. A novel method of determining breast cancer risk using parenchymal textural analysis of mammography images on an Asian cohort. *Physics in medicine and biology* **64**, 035016, doi:10.1088/1361-6560/aafabd (2019).
- 40 Wang, C. B., A. R. : Cuzick, J. : Harkness, E. F. : Evans, D. G. : Astley, S. A novel and fully automated mammographic texture analysis for risk prediction: Results from two case-control studies. *Breast Cancer Research* **19**, doi:10.1186/s13058-017-0906-6 (2017).

- 41 Ward, L. H., S. : Hudson, S. : Wilkinson, L. Parenchymal pattern in women with dense breasts. Variation with age and impact on screening outcomes: observations from a UK screening programme. *European Radiology* **28**, 4717-4724, doi:10.1007/s00330-018-5420-4 (2018).
- 42 Warner, E. T. *et al.* Automated percent mammographic density, mammographic texture variation, and risk of breast cancer: a nested case-control study. *NPJ Breast Cancer* **7**, 68, doi:10.1038/s41523-021-00272-2 (2021).
- 43 Winkel, R. R. E.-C., M. V. : Lynge, E. : Diao, P. : Lillholm, M. : Kallenberg, M. : Forman, J. L. : Nielsen, M. B. : Uldall, W. Y. : Nielsen, M. : Vejborg, I. Risk stratification of women with false-positive test results in mammography screening based on mammographic morphology and density: A case control study. *Cancer Epidemiology* **49**, 53-60, doi:10.1016/j.canep.2017.05.006 (2017).
- 44 Winkel, R. R. v. E.-C., M. : Nielsen, M. : Petersen, K. : Lillholm, M. : Nielsen, M. B. : Lynge, E. : Uldall, W. Y. : Vejborg, I. Mammographic density and structural features can individually and jointly contribute to breast cancer risk assessment in mammography screening: A case-control study. *BMC Cancer* **16**, doi:10.1186/s12885-016-2450-7 (2016).
- 45 Yan, S. W., Y. : Aghaei, F. : Qiu, Y. : Zheng, B. Improving Performance of Breast Cancer Risk Prediction by Incorporating Optical Density Image Feature Analysis. An Assessment. *Acad. Radiol.*, doi:10.1016/j.acra.2017.08.007 (2017).
- 46 Yan, S. W., Y. : Aghaei, F. : Qiu, Y. : Zheng, B. Applying a new bilateral mammographic density segmentation method to improve accuracy of breast cancer risk prediction. *International Journal of Computer Assisted Radiology and Surgery* **12**, 1819-1828, doi:10.1007/s11548-017-1648-8 (2017).
- 47 Ali, M. A., Czene, K., Eriksson, L., Hall, P. & Humphreys, K. Breast Tissue Organisation and its Association with Breast Cancer Risk. *Breast Cancer Res* **19**, 103, doi:10.1186/s13058-017-0894-6 (2017).
- 48 Pertuz, S. S., A. : Holli-Helenius, K. : Kämäräinen, J. : Rinta-Kiikka, I. : Lääperi, A. L. : Arponen, O. Clinical evaluation of a fully-automated parenchymal analysis software for breast cancer risk assessment: A pilot study in a Finnish sample. *Eur J Radiol* **121**, doi:10.1016/j.ejrad.2019.108710 (2019).
- 49 Tan, M., Mariapun, S., Yip, C. H., Ng, K. H. & Teo, S. H. A novel method of determining breast cancer risk using parenchymal textural analysis of mammography images on an Asian cohort. *Phys Med Biol* **64**, 035016, doi:10.1088/1361-6560/aafabd (2019).
- 50 Eriksson, M. *et al.* A clinical model for identifying the short-term risk of breast cancer. *Breast Cancer Res* **19**, 29, doi:10.1186/s13058-017-0820-y (2017).
- 51 Brentnall, A. R., Cuzick, J., Buist, D. S. & Bowles, E. J. A. Long-term accuracy of breast cancer risk assessment combining classic risk factors and breast density. *JAMA oncology* **4**, e180174-e180174 (2018).
- 52 Evans, D. G. R. *et al.* Breast cancer risk feedback to women in the UK NHS breast screening population. *British journal of cancer* **114**, 1045-1052 (2016).
- 53 Brentnall, A. R. *et al.* Mammographic density adds accuracy to both the Tyrer-Cuzick and Gail breast cancer risk models in a prospective UK screening cohort. *Breast Cancer Research* **17**, 147 (2015).

- 54 Visvanathan, K. *et al.* Use of Endocrine Therapy for Breast Cancer Risk Reduction: ASCO Clinical Practice Guideline Update. *J Clin Oncol* **37**, 3152-3165, doi:10.1200/JCO.19.01472 (2019).
- 55 Pace, L. E. & Keating, N. L. Should Women at Lower-Than-Average Risk of Breast Cancer Undergo Less Frequent Screening? *J Natl Cancer Inst* **113**, 953-954, doi:10.1093/jnci/djaa219 (2021).

Appendix

Complete Search Strategies:

Search strategies designed and executed by Angela Hardi, MLIS

Embase.com

=3,919 results on 9/9/2020 (Limited to English; editorials, letters, and notes excluded from results)

Updated search (date limited to 2020-present): 602 on 10/14/2021

('breast density'/exp OR ((breast NEAR/3 densit*):ti,ab,kw OR (mammary NEAR/3 densit*):ti,ab,kw OR (mammographic NEAR/3 densit*):ti,ab,kw)) AND ('mammography'/de OR mammograph*:ti,ab,kw OR mammogram*:ti,ab,kw OR mastrography:ti,ab,kw OR 'digital breast tomosynthesis':ti,ab,kw OR 'x-ray breast tomosynthesis':ti,ab,kw) NOT ('editorial'/it OR 'letter'/it OR 'note'/it) AND [english]/lim

Ovid Medline All

= 2694 results on 9/9/2020 (Limited to English; editorials, comments, and letters excluded)

Updated search (date limited to 2020-present): 440 results on 10/14/2021

(Breast Density/ OR (breast adj3 densit*).ti,ab. OR (mammary adj3 densit*).ti,ab. OR (mammographic adj3 densit*).ti,ab.) AND (Mammography/ OR mammograph*.ti,ab. OR mammogram*.ti,ab. OR mastrography.ti,ab. OR "digital breast tomosynthesis".ti,ab. OR "x-ray breast tomosynthesis".ti,ab.) NOT (comment.pt. OR editorial.pt. OR letter.pt.)

CINAHL Plus

=978 results on 9/9/2020; (Limited to English and these publication types: Clinical Trial, Corrected Article, Journal Article, Meta Analysis, Meta Synthesis, Practice Guidelines, Proceedings, Protocol, Randomized Controlled Trial, Research, Review, Systematic Review)

Updated search (dated limited to 2020-present): 135 results on 10/14/2021

((MH "Breast Tissue Density") OR AB(breast N3 densit*) OR TI(breast N3 densit*) OR AB(mammary N3 densit*) TI(mammary N3 densit*) OR AB(mammographic N3 densit*) OR TI(mammographic N3 densit*)) AND ((MH "Mammography") OR AB(mammograph*) OR TI(mammograph*) OR AB(mammogram*) OR TI(mammogram*) OR AB(mastrography) OR TI(mastrography) OR AB("digital breast tomosynthesis") OR TI("digital breast tomosynthesis") OR AB("x-ray breast tomosynthesis") OR TI("x-ray breast tomosynthesis"))

Scopus

=3,162 results on 9/9/2020 (Limited to English; editorials, notes, letters, and book chapters excluded from results)

Updated search (date limited to 2020-present): 423 results on 10/14/2021

TITLE-ABS ((breast W/3 densit*) OR (mammary W/3 densit*) OR (mammographic W/3 densit*)) AND TITLE-ABS (mammograph* OR mammogram* OR mastrography OR "digital breast tomosynthesis" OR "x-ray breast tomosynthesis") AND (EXCLUDE (DOCTYPE , "no") OR EXCLUDE (DOCTYPE , "ch") OR EXCLUDE (DOCTYPE , "le") OR EXCLUDE (DOCTYPE , "ed")) AND (LIMIT-TO (LANGUAGE , "English"))

Cochrane Library

=358 results on 9/9/2020 (1 Cochrane Protocol and 357 results from CENTRAL Trials)

Updated Search (date limited 2020-present in CENTRAL Trials) = 32 results on 10/14/2021

ID Search

#1 MeSH descriptor: [Breast Density] explode all trees

#2 ((breast NEAR/3 densit*) OR (mammary NEAR/3 densit*) OR (mammary NEAR/3 densit*) OR (mammographic NEAR/3 densit*)):ti,ab,kw

```
#3    #1 OR #2
#4    MeSH descriptor: [Mammography] explode all trees
#5    (mammograph* OR mammogram* OR mastrography OR "digital breast tomosynthesis"
OR "xray breast tomosynthesis"):ti,ab,kw
#6    #4 OR #5
#7    #3 AND #6
```

ClinicalTrials.gov

= 11 results (searched the "Other terms" field) on 9/9/2020

Updated Search = 12 results on 10/14/2021 (1 new result, added to the Excel library)
("breast density" OR "mammary density") AND (mammograph* OR mammogram* OR
mastrography OR "digital breast tomosynthesis" OR "x-ray breast tomosynthesis")



PRISMA 2020 Checklist

Section and Topic	Item #	Checklist item	Location where item is reported
TITLE			
Title	1	Identify the report as a systematic review.	Page 1
ABSTRACT			
Abstract	2	See the PRISMA 2020 for Abstracts checklist.	Page 1
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	Page 3
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	Page 3
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	Pages 3-4
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	Page 4
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	Appendix
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	Page 4
Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	Page 4
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	Page 4
	10b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	Page 5
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	Page 5
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results.	Page 4
Synthesis methods	13a	Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)).	N/A
	13b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	N/A
	13c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	N/A
	13d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	N/A
	13e	Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression).	N/A
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	N/A
Reporting bias	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	N/A



PRISMA 2020 Checklist

Section and Topic	Item #	Checklist item	Location where item is reported
assessment			
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	N/A
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram.	Page 5 and Figure 1
	16b	Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded.	Page 5
Study characteristics	17	Cite each included study and present its characteristics.	Pages 5-6
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	N/A
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	Tables 1-4
Results of syntheses	20a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	N/A
	20b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	N/A
	20c	Present results of all investigations of possible causes of heterogeneity among study results.	N/A
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	N/A
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	N/A
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	N/A
DISCUSSION			
Discussion	23a	Provide a general interpretation of the results in the context of other evidence.	Pages 6-7
	23b	Discuss any limitations of the evidence included in the review.	Pages 6-7
	23c	Discuss any limitations of the review processes used.	Page 7
	23d	Discuss implications of the results for practice, policy, and future research.	Page 7
OTHER INFORMATION			
Registration and protocol	24a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	Page 5
	24b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	Page 5
	24c	Describe and explain any amendments to information provided at registration or in the protocol.	N/A
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	Page 1
Competing interests	26	Declare any competing interests of review authors.	Page 1



PRISMA 2020 Checklist

Section and Topic	Item #	Checklist item	Location where item is reported
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	Page 2

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ 2021;372:n71. doi: 10.1136/bmj.n71

For more information, visit: <http://www.prisma-statement.org/>