COVID-19 Health Care Behaviour in The Gambia: a cross-sectional survey of

205 adults who went through mandatory institutional quarantine

Authors:

Penda Johm[1], Oluwatosin Nkereuwem[1], Aji-Matty Manjang[1], Omar Ceesay[1], Lamin Leigh[1], Amie

Ceesay[1], Mustapha Bittaye[2], Adeyemi Roberts[2], Buba Manjang[2], Sana Sambou[2], Sainey Sanneh[2],

Lamin Saidy[1], Binta Saidy[1], Beate Kampmann[1,3]

Affiliations:

1.   Vaccines and Immunity Theme, Medical Research Council Unit The Gambia at the London School

of Hygiene and Tropical Medicine, Fajara, The Gambia

2.   Ministry of Health, Quadrangle, Banjul, The Gambia

3.   The Vaccine Centre, Faculty of Infectious and Tropical Diseases, London School of Hygiene and

Tropical Medicine, London, UK

*Corresponding author:

Professor Beate Kampmann; bkampmann@mrc.gm

Vaccines and Immunity Theme, Medical Research Council Unit The Gambia at the London School of

Hygiene and Tropical Medicine, Fajara, The Gambia

Abstract: (290 words)

Background: To control the spread of the novel Coronavirus disease 2019 (COVID-19) caused by

severe acute respiratory syndrome novel Coronavirus-2 (SARS-CoV-2), countries around the world

subsequently implemented several public health measures, including the adoption of mandatory

institutional quarantine for close contacts. This study explored the experiences of individuals who

underwent institutional quarantine in The Gambia to inform government measures to increase its

effectiveness and reduce its associated negative impacts.

Methods: Questionnaires were administered via mobile phone call with data collectors calling and

directly recording participant responses on a tablet in an electronic online form developed in

REDCap (Research Electronic Data Capture). The questionnaire contained questions on COVID-19

related knowledge, health care behaviour, attitudes, perceptions and stigma. Data were analysed

using STATA v.13 (Stata Corp, College Station, TX, USA).

Results: In total, 205 adults who observed the mandatory institutional quarantine were interviewed.

There was varied knowledge of COVID-19 causes, spread, symptoms, diagnosis, treatment, and

severity. Participants believed the purpose of quarantine was monitoring for signs and symptoms of

coronavirus disease, testing for SARS-CoV-2, separation from the community, and protection from

coronavirus disease. While a majority reported positive experiences while in quarantine, some

expressed prominent dissatisfaction related to the essential services and quality of care provided.

Different forms of stigma were also experienced before, during and after the quarantine experience.

Conclusion: This study provides important information on quarantine experiences in The Gambia

during the global COVID-19 pandemic. The Ministry of Health in The Gambia and other countries

could improve the experience of quarantined individuals by consistently providing psychosocial

support, compensation for loss of earnings, and timely provision of SARS-CoV-2 test results.

Furthermore, stigma experiences and practices should be addressed during and after individuals stay

in quarantine via the provision of psychosocial support.

INTRODUCTION

On the 11[th] of March 2020, the World Health Organization (WHO) declared Coronavirus disease 2019

(COVID-19) a pandemic. COVID-19 is a novel disease caused by severe acute respiratory syndrome

novel Coronavirus-2 (SARS-CoV-2) (1). As of the 20[th] of August 2021, there have been more than 212

million cases and over 4.4 million deaths recorded worldwide (2). Over 147,000 new cases were

reported in the African region from the 23[rd] to 29[th] of August 2021 (3). In The Gambia, according to

the Ministry of Health (MoH), 359th Situation Report, there were 9,470 total confirmed cases, 9049

recoveries and 301 deaths as of that same date. The Gambia received its first shipment of 36,000

doses of COVID-19 vaccines through the COVAX facility on the 3[rd] of March 2021 with rollout of

vaccinations commencing in the days thereafter. Regarding COVID-19 vaccination status, of the total

national target population (≥ 18 years), 132,756 have received a single dose of Johnson & Johnson's

Janssen vaccine, 35597 received a first dose of Oxford, AstraZeneca vaccine and 4,388 a first dose of

Sinopharm vaccine (4).

Several public health measures have been put in place globally to reduce the transmission of COVID-

19 and to minimize the impact of the disease. This includes risk communication, surveillance, contact

tracing, social/physical distancing and mandatory institutional quarantine. Contact tracing has been

used to quickly determine whether any secondary cases of COVID-19 had occurred due to an

infected person encountering other people. Those contacts are then given information on how to

maintain preventive measures during quarantine and what to do if they develop symptoms of

COVID-19, including where to be tested (5). Institutional quarantine has been used to restrict the

activities of individuals who are not ill with COVID-19 but may have been in close contact with a

person with signs and symptoms of COVID-19 or travelled from an area with high transmission of

COVID-19, with the aim of protecting unexposed members of the community from contracting the

disease (6). In The Gambia, quarantined individuals stayed isolated from others except key hotel and

MoH staff at a facility identified by the government for a mandatory period of 14 days. Those held at

the facilities were provided with food, shelter and medical assistance at the expense of the

government. All quarantined individuals were required to be tested twice during quarantine, and if

found positive, they were transferred to an isolation centre identified by the MoH. If found negative

after 14 days, they were discharged and provided with a medical certificate. A recent study in China

highlights the success of institutional quarantine in minimizing the risk of community spread of

COVID-19 (7).

As the pandemic continues to evolve, it is important to understand health care behaviour in

different geographical and economic settings, evaluate adherence to epidemic prevention and

treatment responses and disease related stigma across the world at different stages of the global

response. Adhering to national COVID-19 guidance is not necessarily feasible for everyone. Those

who live in overcrowded conditions may not be able to practice social distancing and those without

access to clean water and soap will not be able to wash their hands frequently (8). Other individuals

may not know or understand the specific steps to follow or may not be convinced of the need to

practice these behaviours. Another barrier to engagement in health care and adherence to

treatment for health conditions is stigma (9). Stigma is a powerful element in determining health

behaviour and is one reason for social isolation and exclusion. Stigma stems from individual

characteristics as well as context-specific cultural values, norms, and attitudes. Once a stigma is

applied, it manifests in a range of stigma experiences and practices such as verbal abuse or gossip

(10).

Taking these factors into consideration, we documented the experiences of individuals in The

Gambia who completed the government mandated institutional quarantine to develop effective

recommendations for the MoH in The Gambia and other similar cultural settings to better support

people before, during and after quarantine. We explored their experiences and perceptions of

COVID-19 behaviour change measures, including adherence to mitigation gestures such as self-

isolation and social/physical distancing and factors influencing COVID-19 related stigma.

METHODS

Ethical considerations

This study received approval from the Gambia Government/MRC Joint Ethics Committee (Ref.

22271) on the 11[th] of August 2020 and the London School of Hygiene & Tropical Medicine (LSHTM)

Observational/Interventions Research Ethics Committee (Ref. 22271) on the 17[th] of August 2020.

Joining the study was voluntary and participants personal information remained confidential. Unique

identifiers were allocated to each person prior to being sensitized and names were not recorded on

study documents but only on encrypted databases. After sensitization and before consenting, data

collectors asked participants if they understood everything read out to them and gave them the

opportunity to ask questions and discuss answers. The survey had an introductory text stating the

survey was completely anonymous, and consent was implied by respondents agreeing to participate

and completing the survey. Participants also carried out an Assessment of Understanding (AoU),

wherein they needed to answer at least three out of four questions correctly with 89.76% success

rate at first attempt and 10.24% success rate at second attempt.

Study setting

The Gambia is the smallest country on mainland Africa, specifically located in West Africa, and

bordered by Senegal on three sides. The nation has two main religions and eight main ethnic groups

(11). According to The Gambia Multiple Indicator Cluster Survey (MICS) 2018 (12), the percentage of

women and men aged 15-24 years who were able to read a short simple statement about everyday

life or who attended secondary or higher education was 64.3 and 68.0 respectively. 74% of women

and 85% of men aged 15-49 years owned a mobile phone. In the latest Gambia Demographic and

Health Survey (GDHS) 2019-20 (13), a total of 11,865 women and 4,636 men were interviewed,

representing a response rate of 95% female and 87% male respectively. 47% of women and 67% of

men aged 15-49 were literate. Only 2% of women and 7% of men accessed the newspaper,

television, and radio on a weekly basis. These indicators affect how Gambian people respond in a

pandemic such as COVID-19 that requires knowledge and effective communication.

Study design and population

We conducted a cross-sectional mobile phone survey from April to September 2020 to capture

quantitative data from a list of individuals who had been institutionally quarantined. This password-

protected person list and contact details were obtained through the MoH with approval from the

Director of Health Services. The list included the name, arrival date, gender, age, address, phone

number, room number, nationality, country and date of departure of every individual known to had

come into The Gambia by air or land travel and through contact tracing. A contact was defined as

any individual who encountered a person with known COVID-19 in the 48 hours preceding onset of

their own symptoms or sample tested positive to 14 days after. During this time, a total of 863

individuals were put in compulsory institutional quarantine in accordance with the government

regulation in The Gambia, accommodated in four hotels usually housing tourists and based at the

coast. We filtered the list based on study inclusion criteria and attempted to reach every single

individual through phone calls.

Criteria for sample selection

We enrolled all individuals who completed compulsory quarantine for COVID-19 in The Gambia,

aged 18 or above, residing in The Gambia, with a valid phone number, and documented consent to

take part in the study.

Patient and Public Involvement

Our study participants were not involved in setting up the research question, outcome measures,

conduct and/or implementation of the survey. Study results will be designed into an infographic for

ease of sharing and disseminated to all participants to read and retain via WhatsApp and/or email.

Data collection

Questionnaires were administered via mobile phone calls with participant responses recorded on a

tablet in an electronic online form developed in REDCap (Research Electronic Data Capture) and

hosted at the MRCG at LSHTM in the Gambia. Data collectors (5 males, 4 females) with experience

conducting surveys in The Gambia and fluent in English and at least one local language called every

person fitting inclusion criteria to sensitize and recruit them, noting those who refused to participate

or were unreachable. During this initial call, a summary of the study information sheet in either

English, Mandinka, Wolof, or Fula were read out and the full study information sheet was sent to

retain via WhatsApp. The study interviewer would re-attempt the phone call two more times on two

separate days if initially unreachable. A third unsuccessful attempt was noted down as a non-

response. At the time of data collection, all persons had exited the quarantine centres.

Questionnaire answers were collected remotely with a pretested questionnaire to avoid any face-to-

face contact due to the rising cases of COVID-19 at the time. The survey was pre-tested and included

questions covering individual knowledge, attitudes, and perceptions of their quarantine experience.

Demographic characteristics were also captured including sex, age, and highest level of education.

Variables and data analysis

The REDCap data dictionary codebook contained a total of 76 survey questions/fields with items of

different formats; closed-ended (multiple choice, dichotomous, likert scale, self-assessment with an

alternative titled "other") and open-ended to obtain in-depth information on opinions and attitudes

(Supplementary file 1). "Other" responses were evaluated and reclassified into one of the existing or

created new categories. Data were analysed using STATA v.13 (Stata Corp, College Station, Texas,

USA) for descriptive summary statistics (frequencies and percentages). Responses to the open-

ended questions were analysed thematically using an inductive framework approach. This involved

reading and re-reading the dataset to develop familiarity of the data. We then iteratively developed

coding and themes that identified important elements related to the survey questions (14, 15).

Initial themes were determined based on the prevalence of recurring responses across the dataset

and from there the overarching themes and corresponding sub-themes for each question were

identified.

Theoretical frameworks

The frameworks of health seeing behaviour and stigma informed the development of our

questionnaire. Behaviour is a complex phenomenon, influenced by factors within the individual as

well as without (family and peer networks, community, and society). The Partners for Applied Social

Sciences (PASS) model describes the path people follow when seeking care and focuses on the

factors involved in each step that hinder or facilitate access to care (16). The Health Stigma and

Discrimination Framework shows the stigmatization process as it unfolds across the social ecological

spectrum, including drivers and facilitators such as race, class, gender, sexual orientation, or

occupation. Once a stigma is applied, it manifests in a range of stigma experiences and practices.

Stigma experiences can include experienced discrimination, which refers to stigmatizing behaviours

that fall within the purview of the law, or stigmatizing behaviours that fall outside the purview of the

law such as verbal abuse or gossip. Another experience of stigma is self-stigma, wherein a

stigmatized person adopts society's negative beliefs and feelings regarding their status (10). These

frameworks were selected as they are not only theoretical but facilitate an understanding of the

factors that enable health care behaviour, adherence, and stigma.

RESULTS

Participant characteristics

Out of the 429 individuals eligible, 201 were categorized as non-response (123 switched off, 51 no

answer, 13 wrong numbers, 12 unavailable, 1 died, 1 one mentally unstable), 21 refused and 2

withdrew. 228 were contacted and sensitized, 205 accepted and were interviewed (Figure 1). A total

of 205 individuals were interviewed between September to December 2020 (Figure 1). Participants

were mostly males (81.1%), Gambian nationals (84.4%) and Muslims (92.7%). The socio-

demographic characteristics of participants are shown in Table 1.

Table 1: Socio-Demographic Characteristics of Participants (N=205)

Characteristic	N	%
Gender		
Male	167	81.5
Female	38	18.5
Nationality		
Gambian	173	84.4
Non-Gambian	32	15.6
Age group		
18 – 29	63	30.7
30 – 39	55	26.8
40 – 49	49	23.9
>49	38	18.5
Religion		
Islam	190	92.7
Christianity	15	7.3
Marital status		
Married	156	76.1
Not married	49	23.9
Highest level of education		
None	21	10.2
Primary (Grades 1-6)	24	11.7
Junior Secondary (Grades 7-9)	17	8.3
Higher secondary (Grades 10-12)	43	21.0
Tertiary (University education)	44	21.5
Other	56	27.3
Ethnicity		
Mandinka	52	25.4
Fula	36	17.6
Wolof	48	23.4

	Other	69	33.7

Knowledge and perceptions of coronavirus disease 2019, COVID-19

The initial questions probed the individuals' knowledge base (Table 2). Social and mass media were

identified as the most reported first source of information regarding COVID-19 and almost all

participants trusted this first source of information. When asked about the mode of acquiring

COVID-19 was, nearly half of participants mentioned droplets from infected people when they cough

or sneeze. About a quarter said direct contact with an infected person. Almost half of our

participants identified fever and coughing as symptoms of COVID-19. Most of our participants said

they believed that COVID-19 was deadly and about half identified elderly people as the most at-risk

population of being infected with the disease. Almost all said they worried that they were at risk of

contracting coronavirus disease.

Table 2: Knowledge and Perceptions of Coronavirus Disease 2019, COVID-19 (N=205)

Variable	Category	N	%
From whom or from where did you FIRST hear about coronavirus disease, also known as COVID-19? (choose one)	Family member	11	5.4
	Health staff/Health centre	14	6.8
	Mass media/Social media	158	77.1
	Religious centre	2	1.0
	Workplace	4	2.0
	Other	16	7.8
Did you trust the source from which you FIRST heard about coronavirus disease?	Yes	168	82.0
	No	37	18.0
How does coronavirus disease spread/how can someone get infected with coronavirus disease? (multiple choice)	Don't know	11	2.0
	Droplets from infected people when they cough or sneeze	265	49.1
	Direct contact with an infected person	143	26.5
	Touching objects/surfaces contaminated by an infected person	102	18.9
	Other (please specify)	19	3.5
What are the symptoms of coronavirus disease, COVID-19? (multiple choice)	Don't know	14	1.9
	Fever	160	21.6
	Sore throat	90	12.1

	Coughing	156	21.1
	Breathing difficulties	95	12.8
	Body pains	66	8.9
	Headache	78	10.5
	Loss of smell	26	3.5
	Loss of taste	30	4.0
	Other	26	3.5
Do you believe that coronavirus disease 2019, COVID-19, is deadly?	Yes	181	88.3
	No	24	11.7
Who do you think is most at risk of being infected with coronavirus disease? (choose one)	Don't know	8	3.9
	Elderly	103	50.2
	Persons with underlying conditions	27	13.2
	Those that did not follow guidelines	5	2.4
	Health workers	6	2.9
	Everyone	44	21.5
	Other	12	5.9
Do you worry that you are at risk of contracting coronavirus disease?	Yes	162	79.0
	No	43	21.0
What do you consider important actions to prevent the spread of coronavirus disease in your community? (multiple choice)	Staying at home	48	7.95
	Washing hands regularly using hand soap and water or hand sanitizer	136	22.52
	Testing people for coronavirus disease	52	8.61
	Community sensitization	107	17.72
	Everyone wearing a mask when out in public	111	18.38
	Social/physical distancing	112	18.54
	People get vaccinated against coronavirus disease	15	2.48
	Other	23	3.81

COVID-19 Health Care Behaviour

We assessed the way participants had sought health care for themselves before and during

quarantine (Table 3). Most of our participants came to be quarantined following land travel into the

country or the MoH contact tracing. When asked what they understood to be the purpose of the

quarantine process, a majority mentioned monitoring for signs and symptoms of COVID-19, with

some saying they were there to be tested for COVID-19. As we further explored the health care

behaviour of our consenting participants, almost half reported that they would visit a hospital or

health facility as their first point of care if they experienced symptoms of COVID-19. Exploring

adherence, (Appendix 1), participants were asked which quarantine measures they did or did not

comply with during their stay at the quarantine center. Almost all (99%) reported that they stayed at

their place of quarantine every day. 66% did not allow any visitors compared to 34% who did. The

majority separated themselves from others at the place where they were being quarantined (93%),

did not share food and utensils with others (81%), and wore a face mask if they had to leave their

room (76%).

Table 3: COVID-19 Health Care Behaviour (N=205)

Variable	Category	N	%
How did you come to be quarantined? (choose one)	Came in via air travel	21	10.2
	Came in via land travel	97	47.3
	Contact tracing	72	23.4
	Directed by the national coronavirus helpline, 1025	6	2.9
	Other	9	16.1
What do you understand to be the purpose of you having been quarantined? (choose one)	Monitoring me for signs and symptoms of coronavirus disease	170	41.7
	Testing me for coronavirus disease	105	25.7
	Separating me from the community	65	15.9
	Protecting me from coronavirus disease	45	11.0
	Political reasons	3	0.7
	Other	20	4.9
What would you do FIRST if you experience symptoms of coronavirus disease today? (choose one)	I will go to the hospital / health facility	100	48.8
	I will call the national coronavirus helpline on 1025	58	28.3
	I will quarantine myself at home	16	7.8
	I will drink local herbal remedies	8	3.9
	I will go to buy medicines at the pharmacy	2	1.0
	Other	21	10.2

COVID-19 Related Stigma

Participants were asked questions about stigma regarding their experience before, during and after

going into quarantine (Table 4). COVID-19 related stigma was hardly experienced during quarantine.

However, a significant number (40%) had people suggesting they were to go into quarantine prior to

being admitted. More experienced stigma after quarantine, with people gossiping about them

(38%), family and friends refusing to visit them at home (19.5), anyone who called thinking they had

been treated for coronavirus disease (27.8%), and even being called names to make them feel

ashamed (25.4). Participants also experienced enacted stigma (overt discrimination against the

stigmatized) after quarantine leading to feelings of shame, self-blame, loneliness, stress, anger and

withdrawal from daily activities.

259
260   *Table 4: COVID-19 Related Stigma (N=205)*

Variable	Category		N	%
Did you experience any of these before going into quarantine?	People kept suggesting you go into quarantine	YES	82	40
		NO	123	60
	Family and friends refusing to visit you at home	YES	38	18.5
		NO	167	81.5
	Being called names to make you feel ashamed	YES	35	17.1
		NO	170	82.9
	Feeling ashamed because of how you were taken into quarantine	YES	41	20
		NO	164	80
	Other	YES	13	6.3
		NO	192	93.7
Did you experience any of these while you were in quarantine?	People kept gossiping about you	YES	36	17.6
		NO	169	82.4
	People keeping their distance (out of fear)	YES	53	26
		NO	152	74
	People had a judgmental attitude	YES	31	15.1
		NO	174	84.9
	Anyone who called thought you were being treated for coronavirus disease	YES	40	19.5
		NO	165	80.5
	Other	YES	5	2.4
		NO	200	97.6

Did you experience any of these after completing quarantine?	People kept gossiping about you	YES	78	38
		NO	127	62
	Family and friends refused to visit you at home	YES	40	19.5
		NO	165	80.5
	Anyone who called thought you had been treated for coronavirus disease	YES	57	27.8
		NO	148	72.2
	Being called names to make you feel ashamed	YES	52	25.4
		NO	153	74.6
	Other	YES	12	5.9
		NO	193	94.1
Did you experience any of these effects of stigma after completing quarantine?	Shame	YES	30	14.6
		NO	175	85.4
	Self-blame	YES	15	7.3
		NO	190	92.7
	Loneliness	YES	46	22.4
		NO	159	77.6
	Stress	YES	38	18.5
		NO	167	81.5
	Anger	YES	34	16.6
		NO	171	83.4
	Withdrawal from daily activities	YES	30	14.6
		NO	175	85.4
	Strained relationship with family	YES	19	9.3
		NO	186	90.7
	Other	YES	101	49.3
		NO	104	50.7

Attitudes towards COVID-19 related stigma were measured using a Likert scale, wherein

respondents rated their levels of agreement with different statements (Figure 2). Over 80% strongly

agreed or agreed that judging and shaming is hurtful and can negatively affect mental health. Almost

90% strongly agreed or agreed that recovering from coronavirus disease is nothing to be ashamed

about or blamed for. Over 80% strongly agreed or agreed that people may refuse to go into

quarantine for fear of being accused of having coronavirus disease. There was discrepancy regarding

whether public judgement or shaming of others is caused by the publics fear of being infected with

coronavirus disease. 70% strongly agreed or agreed whereas 28% disagreed. There was also

discrepancy regarding whether people who recover from coronavirus disease will be shamed and

blamed for having it. More than half strongly disagreed or disagreed compared to over 40% who

strongly agreed or agreed.

Perceptions of quarantine experience

Participants were also asked to evaluate their quarantine experience (Table 5). Most participants

(68.29%) shared positive quarantine experiences in the facilities where they were accommodated.

While some considered their stay a good experience, some said they had time to relax after many

years of working without rest, others had time for spiritual activities, and most commended the

efforts of the facility staff who they considered friendly. Almost a third (29.27%) of the participants

expressed negative experiences, some participants were upset while sharing their experiences with

our team. Listed benefits of quarantine were testing for and knowing their COVID-19 status;

separating potentially positive cases from others in the community; reading the Quran and having

more time to worship God; care provided by health workers and hotel staff; relaxation; adequate

food and housing; and receiving a certificate after completion of quarantine. Disadvantages were

cited as being away from and missing family; not being able to provide financially for the family as

the household head; poor taste and limited quantity of food served at the hotel; loneliness, lack of

socialisation and social activities; no receipt of COVID-19 test results; lots of mosquitoes and no bed

nets; and time wasted with no remuneration.

Table 5: Perceptions of Quarantine Experience

Positive experiences	Negative experiences
"It was good experience for me due to the quarantine services received from health care workers and supporting staff in Badala Park Hotel." **(50-59 Gambian man, identified via contact tracing)**	*"It was painful leaving my family behind; horrible; traumatised; like a prison."* **(18-29 Gambian man, identified via contact tracing)**

"I relaxed a lot. In my 20 years of work experience, I have not just had a break. I slept a lot and did some paperwork as well." **(40-49 Senegalese man, came in via land travel)**	*"The contact tracing team promised to pick us up, but they never did. We were later informed that there was no fuel available to pick us up."* **(40-49 Gambian man, identified via contact tracing)**
"It was ok with me, and entire health staff and the hotel staff were friendly on their services." **(60-69 Gambian man, came in via air travel)**	*"Very frustrating. Needed medications for body pains and could not get them in time even though I requested from the Ministry of Health staff, they kept delaying the arrival. No psychosocial support."* **(30-39 Gambian man, identified via contact tracing)**

DISCUSSION

As a response to the COVID pandemic, many nations implemented mandatory institutional

quarantine with the aim of protecting unexposed individuals from contracting the disease, but there

is little insight into the experiences of individuals who had been through these procedures. Our

interview-based study explored the topics of COVID-19 knowledge, risk perceptions, attitudes,

adherence, and stigma amongst individuals who had been quarantined in The Gambia to inform

governmental practices that could increase effectiveness of such pandemic control measures and

reduce associated negative impacts.

While exploring knowledge, we found considerable similarities to data from a recent online survey

on the knowledge, attitude, and practice of Sudanese people towards COVID-19 (17), with top listed

sources of knowledge acquisition of COVID-19 being mass media and social media. Predominantly

listed COVID-19 related symptoms identified by our participants such as fever, coughing, breathing

difficulties and sore throat were consistent with widespread information circulating at the time.

Over 60% of participants identified droplets from coughs and sneezes as sources of infection. This

finding is in line with a recently published study conducted in Ethiopia on community COVID-19 risk

perception and health-seeking behaviour (18), wherein 60.3% respondents said COVID-19 could be

transmitted via inhalation of a droplet from infected people and 58.0% reported airborne as a

transmission route for COVID-19. Regarding perceived severity of COVID-19, 88.3% of our

participants believed that coronavirus disease was deadly, like a recent study in Ethiopia (19) and in

Sudan where 85.6 % and 89.5% respectively understood COVID-19 to be a dangerous disease (17).

To nudge the public towards positive epidemic prevention behaviours, individuals must perceive

their susceptibility to and the severity of COVID-19 (20). Risk perceptions give us important insight

into how individuals view personal risks, which in turn is an important determinant of their

behaviour and adherence to protective measures (21, 22). When gauging personal risk perceptions,

it was evident that 79% of our respondents worried that they were at risk of contracting coronavirus

disease, similar to 81% in a recent study in Ethiopia (18). Exactly half of our participants said the

elderly were most at risk of being infected with coronavirus disease, followed by everyone else,

again like a study in Sudan wherein 81.7% said those at risk of developing a severe form of COVID-19

were the elderly, followed by everyone else (36%) (17). In contrast, in another study in Ethiopia

examining myths, beliefs, and perceptions about COVID-19, 45.1% and 62.2% of respondents

perceived that children and youth, respectively, are at a moderate risk of COVID-19 (23). This

variation may be due to the difference sources and timely access of risk information of respondents.

Several studies have explored the knowledge and adoption of actions considered important to

prevent the spread of coronavirus disease in the general population, (17, 23-25), however, none of

these explored the perceptions of quarantined individuals. Quarantine was not listed as a preventive

measure by our study participants, however, most of our participants identified monitoring for signs

and symptoms of coronavirus disease as the purpose of the mandatory institutional quarantine.

Regarding pathway to seeking care for COVID-19 related symptoms in the future, 48.8% and 28.3%

of our study participants claimed they would either first visit a hospital or health facility or call the

national coronavirus helpline on 1025, respectively, if they experienced symptoms of coronavirus

disease.

Looking at facility related factors during quarantine, there was prominent dissatisfaction reported by

most of our participants regarding the facility environment, specifically cleanliness and the services

provided, specifically meals. Our findings are similar to the experiences of persons in COVID-19

institutional quarantine in Uganda (26). In The Gambia, all quarantine related costs were covered by

the government and therefore our study participants were not financially burdened with the cost of

accommodation, food, testing and certification. Nevertheless, a predominant dissatisfaction related

to cost noted by our study participants was specifically the desire to have been provided with

compensation in the form of cash, food and/or transport for their time spent in quarantine, which

would have benefited their families. In other countries, support measures such as financial support,

employment benefits and/or practical support were adopted to enable people to follow self-

isolation or quarantine guidance. In Germany, all employed people in mandatory self-isolation who

tested positive were provided with 100% remuneration of their salary for up to six weeks and in

South Korea, any person required to quarantine for 14 days was provided with daily necessities,

sanitary kits and financial support (27). Although these findings are from high income countries,

similar measures could be adapted in low- and middle- income countries such as The Gambia.

Adherence to COVID-19 prevention and control measures for quarantined individuals are key to the

success of institutional quarantine. Most participants claimed to have adhered to the measures,

however, few deviated and disregarded guidelines. From the limited literature published globally on

adherence to self-isolation measures, it has been found that public trust, an altruistic attitude and

access to information increase adherence to social/physical distancing protocols (27, 28). We utilized

a conceptual framework for quarantine acceptance and adherence to investigate variables affecting

adherence to mandatory COVID-19 quarantine (29). In relation to the framework, our results

showed that individuals life circumstances and quarantine measures being enforced were the main

predictors of their adherence.

Stigma experiences such as discrimination and behaviours such gossip can negatively affect the

mental health of the stigmatized individual and lead to social exclusion. Stigma practices can include

stereotypes, prejudice, stigmatizing behaviours, and discriminatory attitudes (10). In this study,

COVID-19 related stigma was hardly experienced before and during quarantine but rather following

on from quarantine. Some participants reported people gossiping about them, family and friends

refusing to visit them at home and even being called names to make them feel ashamed. Others had

people call them thinking they were being treated for coronavirus disease at the quarantine facility,

and misunderstandings such as these stem from public fears of being infected by people in

quarantine (7). Our participants also experienced overt discrimination (enacted stigma) after

quarantine leading to feelings of shame, self-blame, loneliness, stress, anger and withdrawal from

daily activities (felt stigma). Unfortunately, this is a reality many other individuals in self-isolation

and quarantine have faced worldwide as found in recent studies (30, 31). It is therefore important to

continually uncover and mitigate the many drivers of stigma to improve the lives of quarantined

individuals.

Study strengths and limitations

One of the key strengths of this study was carrying out interviews via mobile phone call guided by

our REDCap questionnaire displayed on tablets. This method seemed most appropriate over

traditional face-to-face interviewing due to the rise of new cases at the time and government

COVID-19 preventive mandates such as social/physical distancing. Our chosen method ensured we

generated rapid first-hand evidence and helped us save on travel costs. Another key strength of our

study was conducting interviews soon after participants left quarantine to minimise their recall bias

after changing environment.

It is important to note that our study population does not represent the diverse demographics of

The Gambia and not every person who went through institutional quarantine was included. Most of

our participants were Gambian males who are more likely to travel than women during the
pandemic and therefore were more frequently identified for institutional quarantine. Our chosen
methods of phone call surveys reduced our opportunity to take note of participants' nonverbal
communication. We did not explore the stigma experience of associative stigma, meaning any
stigma experienced by family or friends of those who went through institutional quarantine as they
were not the cohort of focus in our study.

CONCLUSION

This study sheds light on the lives of those who had been in institutional quarantine in The Gambia
and provides evidence that can be used to inform the Gambia MoH, policy makers and other key
stakeholders working on epidemic preparedness and response. Such information was missing in the
literature in general at the time this study was conducted. To improve the experience of individuals
in designated institutional quarantine facilities, there should be daily provision of quality food and
healthcare, timely provision of SARS-CoV-2 test results and negligible or complimentary associated
costs of quarantine. Furthermore, stigma experiences and practices should be addressed during and
after individuals stay in quarantine via the provision of psychosocial support.

ACKNOWLEDGMENTS

We express heartfelt gratitude to our study participants for their valuable time and genuine
responses to our mobile phone surveys. We would like to sincerely thank the Ministry of Health staff
who supported our data collection: Baboucarr Jallow, Modou Jallow, and Ebrima Keita.

AUTHORS' CONTRIBUTIONS

PJ conceived and designed the study with feedback from ON, AMM, OC, LL, AC, BK, MB, BM, SS and
SS. AMM, OC, LL, AC, ON, BJ, MJ, EK, PJ collected data. AMM, PJ along with LS resolved data queries
and cleaned the dataset. PJ and ON led data analysis and the literature search, wrote the first draft

of the manuscript, and updated subsequent versions of the paper with co-author comments and

inputs. PJ, ON, BK, AMM, OC, LL, AC contributed to the interpretation of results, revised the

manuscript, suggested policy implications from the findings and edited the final draft. All authors

read and approved the final manuscript.

FUNDING

The authors received no specific funding for this study. However, data collection activities were

funded from discretional MRCG@LSHTM funds awarded to BK. The funders had no role in the design

of the study and collection, analysis, and interpretation of data and in writing the manuscript.

AVAILABILITY OF DATA AND MATERIALS

The datasets used and/or analysed during the current study are available from the corresponding

author on reasonable request.

DECLARATIONS

Consent for publication

Not applicable.

Competing interests

None declared.

ORCID IDs

Penda Johm https://orcid.org/0000-0001-8932-7082

Oluwatosin Nkereuwem https://orcid.org/0000-0001-6194-6228

Beate Kampmann https://orcid.org/0000-0002-6546-4709

LIST OF ABBREVIATIONS

AoU   Assessment of Understanding

ANC   Antenatal care

CDC   Centres for Disease Control and Prevention

COVID-19   Coronavirus disease 2019

GDHS   Gambia Demographic and Health Survey

HSB   Health seeking behaviour

MICS   Multiple Indicator Cluster Survey

MoH   Ministry of Health

MRCG at LSHTM   Medical Research Council Unit The Gambia at the London School of Hygiene

451   and Tropical Medicine

PASS   Partners for Applied Social Sciences

REDCap   Research Electronic Data Capture

SARS-Cov-2   Severe Acute Respiratory Syndrome Novel Coronavirus-2

WHO   World Health Organization

REFERENCES

(1) World Health Organization. WHO Director-General's opening remarks at the media briefing on

COVID-19-11 March 2020.

(2) Worldometer of COVID-19 Coronavirus pandemic, Available at:

https://www.worldometers.info/coronavirus/ (Accessed on 20th August 2021).

(3) World Health Organization. COVID-19 Weekly Epidemiological Update Edition 43, Published 8

June 2021.

(4) The Gambia COVID-19 Outbreak Situational Report #359. (2021, August 20). MINISTRY OF

HEALTH. https://www.moh.gov.gm/wp-content/uploads/2021/08/GMB-COVID-19-Situational-

Report-2021-20th-August-No-359.pdf

(5) Lowe M. Using rapid online survey to assess public perceptions of Covid-19 in Gambia. The Pan

African Medical Journal. 2020;35(Suppl 2).

(6) COVID-19 and Your Health. (2020, February 11). Centers for Disease Control and Prevention.

https://www.cdc.gov/coronavirus/2019-ncov/your-health/quarantine-isolation.html

(7) Lio CF, Cheong HH, Lei CI, Lo IL, Lam C, Leong IH. Minimizing the risk of community spread of

COVID-19 via institutional quarantine of high-risk travelers with serial viral RNA testing: A successful

experience from Macao SAR, China. World journal of clinical cases. 2020 Jul 6;8(13):2674.

(8) The Abdul Latif Jameel Poverty Action Lab (J-PAL). (2020). Increasing adherence to COVID-19

guidelines: Lessons   from   existing   evidence.   (online)   Available   at:

https://www.povertyactionlab.org/blog/4-10-20/increasing-adherence-covid-19-guidelines-lessons-

existing-evidence

(9) Link BG, Phelan JC. Conceptualizing stigma. Annual review of Sociology. 2001 Aug;27(1):363-85.

(10) Stangl AL, Earnshaw VA, Logie CH, van Brakel W, Simbayi LC, Barré I, Dovidio JF. The Health

Stigma and Discrimination Framework: a global, crosscutting framework to inform research,

intervention development, and policy on health-related stigmas. BMC medicine. 2019 Dec;17(1):1-3.

(11) Mwakikagile G. The Gambia and its people: ethnic identities and cultural integration in Africa.

New Africa Press; 2010.

(12) UNICEF. The Gambia multiple indicator cluster survey 2018: survey findings report. InThe

Gambia multiple indicator cluster survey 2018: survey findings report 2019 (pp. 694-694).

(13) Gambia Bureau of Statistics (GBoS) and ICF. 2021. The Gambia Demographic and Health Survey

2019-20. Banjul, The Gambia and Rockville, Maryland, USA: GBoS and ICF.

(14) Gibbs GR. Thematic coding and categorizing. Analyzing qualitative data. 2007;703:38-56.

(15) Braun V, Clarke V. Using thematic analysis in psychology. Qualitative research in psychology.

2006 Jan 1;3(2):77-101.

(16) Muela SH, Ribera JM, Toomer E, Grietens KP. The PASS-model: a model for guiding health-

seeking behavior and access to care research. Malaria Reports. 2012 Dec 13;2(1):e3-.

493 (17) Mohamed AA, Elhassan EA, Mohamed AO, Mohammed AA, Mahgoop MA, Sharif ME, Bashir MI,

494 Abdelrahim RB, Idriss WI, Malik EM. Knowledge, attitude and practice of the Sudanese people

495 towards COVID-19: An online survey. BMC public health. 2021 Dec;21(1):1-7.

496 (18) Eyeberu A, Mengistu DA, Negash B, Alemu A, Abate D, Raru TB, Wayessa AD, Debela A, Bahiru

497 N, Heluf H, Kure MA. Community risk perception and health-seeking behavior in the era of COVID-19

498 among adult residents of Harari regional state, eastern Ethiopia. SAGE Open Medicine. 2021

499 Jul;9:20503121211036132.

500 (19) Asnakew Z, Kerebih Asrese MA. Community risk perception and compliance with preventive

501 measures for COVID-19 pandemic in Ethiopia. Risk Management and Healthcare Policy.

502 2020;13:2887.

503 (20) Birhanu Z, Ambelu A, Fufa D, Mecha M, Zeynudin A, Abafita J, Belay A, Doyore F, Oljira L, Bacha

504 E, Feyisa J. Risk perceptions and attitudinal responses to COVID-19 pandemic: an online survey in

505 Ethiopia. BMC public health. 2021 Dec;21(1):1-7.

506 (21) Rohrmann B. Risk perception, risk attitude, risk communication, risk management: A conceptual

507 appraisal. In15th Internaional Emergency Management Society (TIEMS) Annual Conference 2008 Jun

508 (Vol. 2008).

509 (22) Person B, Sy F, Holton K, Govert B, Liang A. Fear and stigma: the epidemic within the SARS

510 outbreak. Emerging infectious diseases. 2004 Feb;10(2):358.

511 (23) Kebede Y, Yitayih Y, Birhanu Z, Mekonen S, Ambelu A. Knowledge, perceptions and preventive

512 practices towards COVID-19 early in the outbreak among Jimma university medical center visitors,

513 Southwest Ethiopia. PloS one. 2020 May 21;15(5):e0233744.

514 (24) Alkhaldi G, Aljuraiban GS, Alhurishi S, De Souza R, Lamahewa K, Lau R, Alshaikh F. Perceptions

515 towards COVID-19 and adoption of preventive measures among the public in Saudi Arabia: a cross

516 sectional study. BMC public health. 2021 Dec;21(1):1-21.

(25) Mendy PA, Jawo E, Jallow BJ, Ceesay B, Mendy SA, Camara B, Ndaw M. Knowledge, Attitude and

Practice of Gambians towards the Novel Coronavirus (COVID-19) Pandemic in The Gambia: A cross-

sectional study. African Journal of Health Sciences. 2021 Jun 10;34(1):134-47.

(26) Ndejjo R, Naggayi G, Tibiita R, Mugahi R, Kibira SP. Experiences of persons in COVID-19

institutional quarantine in Uganda: a qualitative study. BMC public health. 2021 Dec;21(1):1-1.

(27) Patel J, Fernandes G, Sridhar D. How can we improve self-isolation and quarantine for covid-19?.

bmj. 2021 Mar 10;372.

(28) Webster RK, Brooks SK, Smith LE, Woodland L, Wessely S, Rubin GJ. How to improve adherence

with quarantine: rapid review of the evidence. Public Health. 2020 May 1;182:163-9.

(29) Sopory P, Novak JM, Noyes JP. Quarantine acceptance and adherence: qualitative evidence

synthesis and conceptual framework. Journal of Public Health. 2021 Apr 16:1-1.

(30) Lohiniva AL, Dub T, Hagberg L, Nohynek H. Learning about COVID-19-related stigma, quarantine

and isolation experiences in Finland. PloS one. 2021 Apr 14;16(4):e0247962.

(31) Turner-Musa J, Ajayi O, Kemp L. Examining social determinants of health, stigma, and COVID-19

disparities. InHealthcare 2020 Jun (Vol. 8, No. 2, p. 168). Multidisciplinary Digital Publishing Institute.

*Switched off 123; No answer 51; Wrong number 5; Busy/unavailable 12; Died 1; Mentally unstable

1; Wrong person 8

Figure 1. Participant recruitment process based on study inclusion criteria.

Figure 2. Likert scale gauging attitudes towards COVID-19 related stigma.