

1        **COVID-19 Health Care Behaviour in The Gambia: a cross-sectional survey of**  
2                    **205 adults who went through mandatory institutional quarantine**

3

4        **Authors:**

5        Penda Johm<sup>1</sup>, Oluwatosin Nkereuwem<sup>1</sup>, Aji-Matty Manjang<sup>1</sup>, Omar Ceesay<sup>1</sup>, Lamin Leigh<sup>1</sup>, Amie  
6        Ceesay<sup>1</sup>, Mustapha Bittaye<sup>2</sup>, Adeyemi Roberts<sup>2</sup>, Buba Manjang<sup>2</sup>, Sana Sambou<sup>2</sup>, Sainey Sanneh<sup>2</sup>,  
7        Lamin Saidy<sup>1</sup>, Binta Saidy<sup>1</sup>, Beate Kampmann<sup>1,3</sup>

8

9        **Affiliations:**

- 10        1. Vaccines and Immunity Theme, Medical Research Council Unit The Gambia at the London School  
11            of Hygiene and Tropical Medicine, Fajara, The Gambia  
12        2. Ministry of Health, Quadrangle, Banjul, The Gambia  
13        3. The Vaccine Centre, Faculty of Infectious and Tropical Diseases, London School of Hygiene and  
14            Tropical Medicine, London, UK

15

16        \*Corresponding author:

17        Professor Beate Kampmann; [bkampmann@mrc.gm](mailto:bkampmann@mrc.gm)

18        Vaccines and Immunity Theme, Medical Research Council Unit The Gambia at the London School of  
19        Hygiene and Tropical Medicine, Fajara, The Gambia

20

21

22

23

24

25

26 **Abstract: (290 words)**

27 **Background:** To control the spread of the novel Coronavirus disease 2019 (COVID-19) caused by  
28 severe acute respiratory syndrome novel Coronavirus-2 (SARS-CoV-2), countries around the world  
29 subsequently implemented several public health measures, including the adoption of mandatory  
30 institutional quarantine for close contacts. This study explored the experiences of individuals who  
31 underwent institutional quarantine in The Gambia to inform government measures to increase its  
32 effectiveness and reduce its associated negative impacts.

33

34 **Methods:** Questionnaires were administered via mobile phone call with data collectors calling and  
35 directly recording participant responses on a tablet in an electronic online form developed in  
36 REDCap (Research Electronic Data Capture). The questionnaire contained questions on COVID-19  
37 related knowledge, health care behaviour, attitudes, perceptions and stigma. Data were analysed  
38 using STATA v.13 (Stata Corp, College Station, TX, USA).

39

40 **Results:** In total, 205 adults who observed the mandatory institutional quarantine were interviewed.  
41 There was varied knowledge of COVID-19 causes, spread, symptoms, diagnosis, treatment, and  
42 severity. Participants believed the purpose of quarantine was monitoring for signs and symptoms of  
43 coronavirus disease, testing for SARS-CoV-2, separation from the community, and protection from  
44 coronavirus disease. While a majority reported positive experiences while in quarantine, some  
45 expressed prominent dissatisfaction related to the essential services and quality of care provided.  
46 Different forms of stigma were also experienced before, during and after the quarantine experience.

47

48 **Conclusion:** This study provides important information on quarantine experiences in The Gambia  
49 during the global COVID-19 pandemic. The Ministry of Health in The Gambia and other countries  
50 could improve the experience of quarantined individuals by consistently providing psychosocial  
51 support, compensation for loss of earnings, and timely provision of SARS-CoV-2 test results.

52 Furthermore, stigma experiences and practices should be addressed during and after individuals stay  
53 in quarantine via the provision of psychosocial support.

54

## 55 **INTRODUCTION**

56 On the 11<sup>th</sup> of March 2020, the World Health Organization (WHO) declared Coronavirus disease 2019  
57 (COVID-19) a pandemic. COVID-19 is a novel disease caused by severe acute respiratory syndrome  
58 novel Coronavirus-2 (SARS-CoV-2) (1). As of the 20<sup>th</sup> of August 2021, there have been more than 212  
59 million cases and over 4.4 million deaths recorded worldwide (2). Over 147,000 new cases were  
60 reported in the African region from the 23<sup>rd</sup> to 29<sup>th</sup> of August 2021 (3). In The Gambia, according to  
61 the Ministry of Health (MoH), 359<sup>th</sup> Situation Report, there were 9,470 total confirmed cases, 9049  
62 recoveries and 301 deaths as of that same date. The Gambia received its first shipment of 36,000  
63 doses of COVID-19 vaccines through the COVAX facility on the 3<sup>rd</sup> of March 2021 with rollout of  
64 vaccinations commencing in the days thereafter. Regarding COVID-19 vaccination status, of the total  
65 national target population ( $\geq 18$  years), 132,756 have received a single dose of Johnson & Johnson's  
66 Janssen vaccine, 35597 received a first dose of Oxford, AstraZeneca vaccine and 4,388 a first dose of  
67 Sinopharm vaccine (4).

68

69 Several public health measures have been put in place globally to reduce the transmission of COVID-  
70 19 and to minimize the impact of the disease. This includes risk communication, surveillance, contact  
71 tracing, social/physical distancing and mandatory institutional quarantine. Contact tracing has been  
72 used to quickly determine whether any secondary cases of COVID-19 had occurred due to an  
73 infected person encountering other people. Those contacts are then given information on how to  
74 maintain preventive measures during quarantine and what to do if they develop symptoms of  
75 COVID-19, including where to be tested (5). Institutional quarantine has been used to restrict the  
76 activities of individuals who are not ill with COVID-19 but may have been in close contact with a  
77 person with signs and symptoms of COVID-19 or travelled from an area with high transmission of

78 COVID-19, with the aim of protecting unexposed members of the community from contracting the  
79 disease (6). In The Gambia, quarantined individuals stayed isolated from others except key hotel and  
80 MoH staff at a facility identified by the government for a mandatory period of 14 days. Those held at  
81 the facilities were provided with food, shelter and medical assistance at the expense of the  
82 government. All quarantined individuals were required to be tested twice during quarantine, and if  
83 found positive, they were transferred to an isolation centre identified by the MoH. If found negative  
84 after 14 days, they were discharged and provided with a medical certificate. A recent study in China  
85 highlights the success of institutional quarantine in minimizing the risk of community spread of  
86 COVID-19 (7).

87

88 As the pandemic continues to evolve, it is important to understand health care behaviour in  
89 different geographical and economic settings, evaluate adherence to epidemic prevention and  
90 treatment responses and disease related stigma across the world at different stages of the global  
91 response. Adhering to national COVID-19 guidance is not necessarily feasible for everyone. Those  
92 who live in overcrowded conditions may not be able to practice social distancing and those without  
93 access to clean water and soap will not be able to wash their hands frequently (8). Other individuals  
94 may not know or understand the specific steps to follow or may not be convinced of the need to  
95 practice these behaviours. Another barrier to engagement in health care and adherence to  
96 treatment for health conditions is stigma (9). Stigma is a powerful element in determining health  
97 behaviour and is one reason for social isolation and exclusion. Stigma stems from individual  
98 characteristics as well as context-specific cultural values, norms, and attitudes. Once a stigma is  
99 applied, it manifests in a range of stigma experiences and practices such as verbal abuse or gossip  
100 (10).

101

102 Taking these factors into consideration, we documented the experiences of individuals in The  
103 Gambia who completed the government mandated institutional quarantine to develop effective

104 recommendations for the MoH in The Gambia and other similar cultural settings to better support  
105 people before, during and after quarantine. We explored their experiences and perceptions of  
106 COVID-19 behaviour change measures, including adherence to mitigation gestures such as self-  
107 isolation and social/physical distancing and factors influencing COVID-19 related stigma.

108

## 109 **METHODS**

### 110 **Ethical considerations**

111 This study received approval from the Gambia Government/MRC Joint Ethics Committee (Ref.  
112 22271) on the 11<sup>th</sup> of August 2020 and the London School of Hygiene & Tropical Medicine (LSHTM)  
113 Observational/Interventions Research Ethics Committee (Ref. 22271) on the 17<sup>th</sup> of August 2020.  
114 Joining the study was voluntary and participants personal information remained confidential. Unique  
115 identifiers were allocated to each person prior to being sensitized and names were not recorded on  
116 study documents but only on encrypted databases. After sensitization and before consenting, data  
117 collectors asked participants if they understood everything read out to them and gave them the  
118 opportunity to ask questions and discuss answers. The survey had an introductory text stating the  
119 survey was completely anonymous, and consent was implied by respondents agreeing to participate  
120 and completing the survey. Participants also carried out an Assessment of Understanding (AoU),  
121 wherein they needed to answer at least three out of four questions correctly with 89.76% success  
122 rate at first attempt and 10.24% success rate at second attempt.

123

### 124 **Study setting**

125 The Gambia is the smallest country on mainland Africa, specifically located in West Africa, and  
126 bordered by Senegal on three sides. The nation has two main religions and eight main ethnic groups  
127 (11). According to The Gambia Multiple Indicator Cluster Survey (MICS) 2018 (12), the percentage of  
128 women and men aged 15-24 years who were able to read a short simple statement about everyday  
129 life or who attended secondary or higher education was 64.3 and 68.0 respectively. 74% of women

130 and 85% of men aged 15-49 years owned a mobile phone. In the latest Gambia Demographic and  
131 Health Survey (GDHS) 2019-20 (13), a total of 11,865 women and 4,636 men were interviewed,  
132 representing a response rate of 95% female and 87% male respectively. 47% of women and 67% of  
133 men aged 15-49 were literate. Only 2% of women and 7% of men accessed the newspaper,  
134 television, and radio on a weekly basis. These indicators affect how Gambian people respond in a  
135 pandemic such as COVID-19 that requires knowledge and effective communication.

136

### 137 **Study design and population**

138 We conducted a cross-sectional mobile phone survey from April to September 2020 to capture  
139 quantitative data from a list of individuals who had been institutionally quarantined. This password-  
140 protected person list and contact details were obtained through the MoH with approval from the  
141 Director of Health Services. The list included the name, arrival date, gender, age, address, phone  
142 number, room number, nationality, country and date of departure of every individual known to had  
143 come into The Gambia by air or land travel and through contact tracing. A contact was defined as  
144 any individual who encountered a person with known COVID-19 in the 48 hours preceding onset of  
145 their own symptoms or sample tested positive to 14 days after. During this time, a total of 863  
146 individuals were put in compulsory institutional quarantine in accordance with the government  
147 regulation in The Gambia, accommodated in four hotels usually housing tourists and based at the  
148 coast. We filtered the list based on study inclusion criteria and attempted to reach every single  
149 individual through phone calls.

150

### 151 **Criteria for sample selection**

152 We enrolled all individuals who completed compulsory quarantine for COVID-19 in The Gambia,  
153 aged 18 or above, residing in The Gambia, with a valid phone number, and documented consent to  
154 take part in the study.

155

156 **Patient and Public Involvement**

157 Our study participants were not involved in setting up the research question, outcome measures,  
158 conduct and/or implementation of the survey. Study results will be designed into an infographic for  
159 ease of sharing and disseminated to all participants to read and retain via WhatsApp and/or email.

160

161 **Data collection**

162 Questionnaires were administered via mobile phone calls with participant responses recorded on a  
163 tablet in an electronic online form developed in REDCap (Research Electronic Data Capture) and  
164 hosted at the MRCG at LSHTM in the Gambia. Data collectors (5 males, 4 females) with experience  
165 conducting surveys in The Gambia and fluent in English and at least one local language called every  
166 person fitting inclusion criteria to sensitize and recruit them, noting those who refused to participate  
167 or were unreachable. During this initial call, a summary of the study information sheet in either  
168 English, Mandinka, Wolof, or Fula were read out and the full study information sheet was sent to  
169 retain via WhatsApp. The study interviewer would re-attempt the phone call two more times on two  
170 separate days if initially unreachable. A third unsuccessful attempt was noted down as a non-  
171 response. At the time of data collection, all persons had exited the quarantine centres.  
172 Questionnaire answers were collected remotely with a pretested questionnaire to avoid any face-to-  
173 face contact due to the rising cases of COVID-19 at the time. The survey was pre-tested and included  
174 questions covering individual knowledge, attitudes, and perceptions of their quarantine experience.  
175 Demographic characteristics were also captured including sex, age, and highest level of education.

176

177 **Variables and data analysis**

178 The REDCap data dictionary codebook contained a total of 76 survey questions/fields with items of  
179 different formats; closed-ended (multiple choice, dichotomous, likert scale, self-assessment with an  
180 alternative titled “other”) and open-ended to obtain in-depth information on opinions and attitudes  
181 (Supplementary file 1). “Other” responses were evaluated and reclassified into one of the existing or

182 created new categories. Data were analysed using STATA v.13 (Stata Corp, College Station, Texas,  
183 USA) for descriptive summary statistics (frequencies and percentages). Responses to the open-  
184 ended questions were analysed thematically using an inductive framework approach. This involved  
185 reading and re-reading the dataset to develop familiarity of the data. We then iteratively developed  
186 coding and themes that identified important elements related to the survey questions (14, 15).  
187 Initial themes were determined based on the prevalence of recurring responses across the dataset  
188 and from there the overarching themes and corresponding sub-themes for each question were  
189 identified.

190

### 191 **Theoretical frameworks**

192 The frameworks of health seeking behaviour and stigma informed the development of our  
193 questionnaire. Behaviour is a complex phenomenon, influenced by factors within the individual as  
194 well as without (family and peer networks, community, and society). The Partners for Applied Social  
195 Sciences (PASS) model describes the path people follow when seeking care and focuses on the  
196 factors involved in each step that hinder or facilitate access to care (16). The Health Stigma and  
197 Discrimination Framework shows the stigmatization process as it unfolds across the social ecological  
198 spectrum, including drivers and facilitators such as race, class, gender, sexual orientation, or  
199 occupation. Once a stigma is applied, it manifests in a range of stigma experiences and practices.  
200 Stigma experiences can include experienced discrimination, which refers to stigmatizing behaviours  
201 that fall within the purview of the law, or stigmatizing behaviours that fall outside the purview of the  
202 law such as verbal abuse or gossip. Another experience of stigma is self-stigma, wherein a  
203 stigmatized person adopts society's negative beliefs and feelings regarding their status (10). These  
204 frameworks were selected as they are not only theoretical but facilitate an understanding of the  
205 factors that enable health care behaviour, adherence, and stigma.

206

### 207 **RESULTS**

208 **Participant characteristics**

209 Out of the 429 individuals eligible, 201 were categorized as non-response (123 switched off, 51 no  
210 answer, 13 wrong numbers, 12 unavailable, 1 died, 1 one mentally unstable), 21 refused and 2  
211 withdrew. 228 were contacted and sensitized, 205 accepted and were interviewed (Figure 1). A total  
212 of 205 individuals were interviewed between September to December 2020 (Figure 1). Participants  
213 were mostly males (81.1%), Gambian nationals (84.4%) and Muslims (92.7%). The socio-  
214 demographic characteristics of participants are shown in Table 1.

215

216 *Table 1: Socio-Demographic Characteristics of Participants (N=205)*

Characteristic	N	%
Gender		
Male	167	81.5
Female	38	18.5
Nationality		
Gambian	173	84.4
Non-Gambian	32	15.6
Age group		
18 – 29	63	30.7
30 – 39	55	26.8
40 – 49	49	23.9
>49	38	18.5
Religion		
Islam	190	92.7
Christianity	15	7.3
Marital status		
Married	156	76.1
Not married	49	23.9
Highest level of education		
None	21	10.2
Primary (Grades 1-6)	24	11.7
Junior Secondary (Grades 7-9)	17	8.3
Higher secondary (Grades 10-12)	43	21.0
Tertiary (University education)	44	21.5
Other	56	27.3
Ethnicity		
Mandinka	52	25.4
Fula	36	17.6
Wolof	48	23.4

217 Other 69 33.7

218 **Knowledge and perceptions of coronavirus disease 2019, COVID-19**

219 The initial questions probed the individuals’ knowledge base (Table 2). Social and mass media were  
 220 identified as the most reported first source of information regarding COVID-19 and almost all  
 221 participants trusted this first source of information. When asked about the mode of acquiring  
 222 COVID-19 was, nearly half of participants mentioned droplets from infected people when they cough  
 223 or sneeze. About a quarter said direct contact with an infected person. Almost half of our  
 224 participants identified fever and coughing as symptoms of COVID-19. Most of our participants said  
 225 they believed that COVID-19 was deadly and about half identified elderly people as the most at-risk  
 226 population of being infected with the disease. Almost all said they worried that they were at risk of  
 227 contracting coronavirus disease.

228

229 *Table 2: Knowledge and Perceptions of Coronavirus Disease 2019, COVID-19 (N=205)*

Variable	Category	N	%
From whom or from where did you FIRST hear about coronavirus disease, also known as COVID-19? (choose one)	Family member	11	5.4
	Health staff/Health centre	14	6.8
	Mass media/Social media	158	77.1
	Religious centre	2	1.0
	Workplace	4	2.0
	Other	16	7.8
Did you trust the source from which you FIRST heard about coronavirus disease?	Yes	168	82.0
	No	37	18.0
How does coronavirus disease spread/how can someone get infected with coronavirus disease? (multiple choice)	Don't know	11	2.0
	Droplets from infected people when they cough or sneeze	265	49.1
	Direct contact with an infected person	143	26.5
	Touching objects/surfaces contaminated by an infected person	102	18.9
	Other (please specify)	19	3.5
What are the symptoms of coronavirus disease, COVID-19? (multiple choice)	Don't know	14	1.9
	Fever	160	21.6
	Sore throat	90	12.1

	Coughing	156	21.1
	Breathing difficulties	95	12.8
	Body pains	66	8.9
	Headache	78	10.5
	Loss of smell	26	3.5
	Loss of taste	30	4.0
	Other	26	3.5
Do you believe that coronavirus disease 2019, COVID-19, is deadly?	Yes	181	88.3
	No	24	11.7
Who do you think is most at risk of being infected with coronavirus disease? (choose one)	Don't know	8	3.9
	Elderly	103	50.2
	Persons with underlying conditions	27	13.2
	Those that did not follow guidelines	5	2.4
	Health workers	6	2.9
	Everyone	44	21.5
	Other	12	5.9
Do you worry that you are at risk of contracting coronavirus disease?	Yes	162	79.0
	No	43	21.0
What do you consider important actions to prevent the spread of coronavirus disease in your community? (multiple choice)	Staying at home	48	7.95
	Washing hands regularly using hand soap and water or hand sanitizer	136	22.52
	Testing people for coronavirus disease	52	8.61
	Community sensitization	107	17.72
	Everyone wearing a mask when out in public	111	18.38
	Social/physical distancing	112	18.54
	People get vaccinated against coronavirus disease	15	2.48
	Other	23	3.81

230

### 231 COVID-19 Health Care Behaviour

232 We assessed the way participants had sought health care for themselves before and during  
 233 quarantine (Table 3). Most of our participants came to be quarantined following land travel into the  
 234 country or the MoH contact tracing. When asked what they understood to be the purpose of the  
 235 quarantine process, a majority mentioned monitoring for signs and symptoms of COVID-19, with  
 236 some saying they were there to be tested for COVID-19. As we further explored the health care  
 237 behaviour of our consenting participants, almost half reported that they would visit a hospital or  
 238 health facility as their first point of care if they experienced symptoms of COVID-19. Exploring

239 adherence, (Appendix 1), participants were asked which quarantine measures they did or did not  
 240 comply with during their stay at the quarantine center. Almost all (99%) reported that they stayed at  
 241 their place of quarantine every day. 66% did not allow any visitors compared to 34% who did. The  
 242 majority separated themselves from others at the place where they were being quarantined (93%),  
 243 did not share food and utensils with others (81%), and wore a face mask if they had to leave their  
 244 room (76%).

245

246 *Table 3: COVID-19 Health Care Behaviour (N=205)*

Variable	Category	N	%
How did you come to be quarantined? (choose one)	Came in via air travel	21	10.2
	Came in via land travel	97	47.3
	Contact tracing	72	23.4
	Directed by the national coronavirus helpline, 1025	6	2.9
	Other	9	16.1
What do you understand to be the purpose of you having been quarantined? (choose one)	Monitoring me for signs and symptoms of coronavirus disease	170	41.7
	Testing me for coronavirus disease	105	25.7
	Separating me from the community	65	15.9
	Protecting me from coronavirus disease	45	11.0
	Political reasons	3	0.7
What would you do FIRST if you experience symptoms of coronavirus disease today? (choose one)	Other	20	4.9
	I will go to the hospital / health facility	100	48.8
	I will call the national coronavirus helpline on 1025	58	28.3
	I will quarantine myself at home	16	7.8
	I will drink local herbal remedies	8	3.9
	I will go to buy medicines at the pharmacy	2	1.0
	Other	21	10.2

247

248

249 **COVID-19 Related Stigma**

250 Participants were asked questions about stigma regarding their experience before, during and after  
 251 going into quarantine (Table 4). COVID-19 related stigma was hardly experienced during quarantine.  
 252 However, a significant number (40%) had people suggesting they were to go into quarantine prior to  
 253 being admitted. More experienced stigma after quarantine, with people gossiping about them  
 254 (38%), family and friends refusing to visit them at home (19.5), anyone who called thinking they had  
 255 been treated for coronavirus disease (27.8%), and even being called names to make them feel  
 256 ashamed (25.4). Participants also experienced enacted stigma (overt discrimination against the  
 257 stigmatized) after quarantine leading to feelings of shame, self-blame, loneliness, stress, anger and  
 258 withdrawal from daily activities.

259  
 260 *Table 4: COVID-19 Related Stigma (N=205)*

Variable	Category		N	%
Did you experience any of these before going into quarantine?	People kept suggesting you go into quarantine	YES	82	40
		NO	123	60
	Family and friends refusing to visit you at home	YES	38	18.5
		NO	167	81.5
	Being called names to make you feel ashamed	YES	35	17.1
		NO	170	82.9
	Feeling ashamed because of how you were taken into quarantine	YES	41	20
		NO	164	80
	Other	YES	13	6.3
		NO	192	93.7
Did you experience any of these while you were in quarantine?	People kept gossiping about you	YES	36	17.6
		NO	169	82.4
	People keeping their distance (out of fear)	YES	53	26
		NO	152	74
	People had a judgmental attitude	YES	31	15.1
		NO	174	84.9
	Anyone who called thought you were being treated for coronavirus disease	YES	40	19.5
		NO	165	80.5
	Other	YES	5	2.4
		NO	200	97.6

Did you experience any of these after completing quarantine?	People kept gossiping about you	YES	78	38
		NO	127	62
	Family and friends refused to visit you at home	YES	40	19.5
		NO	165	80.5
	Anyone who called thought you had been treated for coronavirus disease	YES	57	27.8
		NO	148	72.2
	Being called names to make you feel ashamed	YES	52	25.4
		NO	153	74.6
	Other	YES	12	5.9
		NO	193	94.1
Did you experience any of these effects of stigma after completing quarantine?	Shame	YES	30	14.6
		NO	175	85.4
	Self-blame	YES	15	7.3
		NO	190	92.7
	Loneliness	YES	46	22.4
		NO	159	77.6
	Stress	YES	38	18.5
		NO	167	81.5
	Anger	YES	34	16.6
		NO	171	83.4
	Withdrawal from daily activities	YES	30	14.6
		NO	175	85.4
	Strained relationship with family	YES	19	9.3
		NO	186	90.7
	Other	YES	101	49.3
		NO	104	50.7

261

262 Attitudes towards COVID-19 related stigma were measured using a Likert scale, wherein  
 263 respondents rated their levels of agreement with different statements (Figure 2). Over 80% strongly  
 264 agreed or agreed that judging and shaming is hurtful and can negatively affect mental health. Almost  
 265 90% strongly agreed or agreed that recovering from coronavirus disease is nothing to be ashamed  
 266 about or blamed for. Over 80% strongly agreed or agreed that people may refuse to go into  
 267 quarantine for fear of being accused of having coronavirus disease. There was discrepancy regarding  
 268 whether public judgement or shaming of others is caused by the publics fear of being infected with

269 coronavirus disease. 70% strongly agreed or agreed whereas 28% disagreed. There was also  
270 discrepancy regarding whether people who recover from coronavirus disease will be shamed and  
271 blamed for having it. More than half strongly disagreed or disagreed compared to over 40% who  
272 strongly agreed or agreed.

273

#### 274 **Perceptions of quarantine experience**

275 Participants were also asked to evaluate their quarantine experience (Table 5). Most participants  
276 (68.29%) shared positive quarantine experiences in the facilities where they were accommodated.  
277 While some considered their stay a good experience, some said they had time to relax after many  
278 years of working without rest, others had time for spiritual activities, and most commended the  
279 efforts of the facility staff who they considered friendly. Almost a third (29.27%) of the participants  
280 expressed negative experiences, some participants were upset while sharing their experiences with  
281 our team. Listed benefits of quarantine were testing for and knowing their COVID-19 status;  
282 separating potentially positive cases from others in the community; reading the Quran and having  
283 more time to worship God; care provided by health workers and hotel staff; relaxation; adequate  
284 food and housing; and receiving a certificate after completion of quarantine. Disadvantages were  
285 cited as being away from and missing family; not being able to provide financially for the family as  
286 the household head; poor taste and limited quantity of food served at the hotel; loneliness, lack of  
287 socialisation and social activities; no receipt of COVID-19 test results; lots of mosquitoes and no bed  
288 nets; and time wasted with no remuneration.

289

290 *Table 5: Perceptions of Quarantine Experience*

Positive experiences	Negative experiences
<i>“It was good experience for me due to the quarantine services received from health care workers and supporting staff in Badala Park Hotel.”</i>	<i>“It was painful leaving my family behind; horrible; traumatised; like a prison.”</i>
<b>(50-59 Gambian man, identified via contact tracing)</b>	<b>(18-29 Gambian man, identified via contact tracing)</b>

---

<p><i>"I relaxed a lot. In my 20 years of work experience, I have not just had a break. I slept a lot and did some paperwork as well."</i> <b>(40-49 Senegalese man, came in via land travel)</b></p>	<p><i>"The contact tracing team promised to pick us up, but they never did. We were later informed that there was no fuel available to pick us up."</i> <b>(40-49 Gambian man, identified via contact tracing)</b></p>
<p><i>"It was ok with me, and entire health staff and the hotel staff were friendly on their services."</i> <b>(60-69 Gambian man, came in via air travel)</b></p>	<p><i>"Very frustrating. Needed medications for body pains and could not get them in time even though I requested from the Ministry of Health staff, they kept delaying the arrival. No psychosocial support."</i> <b>(30-39 Gambian man, identified via contact tracing)</b></p>

---

291

292 **DISCUSSION**

293 As a response to the COVID pandemic, many nations implemented mandatory institutional  
294 quarantine with the aim of protecting unexposed individuals from contracting the disease, but there  
295 is little insight into the experiences of individuals who had been through these procedures. Our  
296 interview-based study explored the topics of COVID-19 knowledge, risk perceptions, attitudes,  
297 adherence, and stigma amongst individuals who had been quarantined in The Gambia to inform  
298 governmental practices that could increase effectiveness of such pandemic control measures and  
299 reduce associated negative impacts.

300

301 While exploring knowledge, we found considerable similarities to data from a recent online survey  
302 on the knowledge, attitude, and practice of Sudanese people towards COVID-19 (17), with top listed  
303 sources of knowledge acquisition of COVID-19 being mass media and social media. Predominantly  
304 listed COVID-19 related symptoms identified by our participants such as fever, coughing, breathing  
305 difficulties and sore throat were consistent with widespread information circulating at the time.  
306 Over 60% of participants identified droplets from coughs and sneezes as sources of infection. This  
307 finding is in line with a recently published study conducted in Ethiopia on community COVID-19 risk  
308 perception and health-seeking behaviour (18), wherein 60.3% respondents said COVID-19 could be  
309 transmitted via inhalation of a droplet from infected people and 58.0% reported airborne as a  
310 transmission route for COVID-19. Regarding perceived severity of COVID-19, 88.3% of our

311 participants believed that coronavirus disease was deadly, like a recent study in Ethiopia (19) and in  
312 Sudan where 85.6 % and 89.5% respectively understood COVID-19 to be a dangerous disease (17).

313

314 To nudge the public towards positive epidemic prevention behaviours, individuals must perceive  
315 their susceptibility to and the severity of COVID-19 (20). Risk perceptions give us important insight  
316 into how individuals view personal risks, which in turn is an important determinant of their  
317 behaviour and adherence to protective measures (21, 22). When gauging personal risk perceptions,  
318 it was evident that 79% of our respondents worried that they were at risk of contracting coronavirus  
319 disease, similar to 81% in a recent study in Ethiopia (18). Exactly half of our participants said the  
320 elderly were most at risk of being infected with coronavirus disease, followed by everyone else,  
321 again like a study in Sudan wherein 81.7% said those at risk of developing a severe form of COVID-19  
322 were the elderly, followed by everyone else (36%) (17). In contrast, in another study in Ethiopia  
323 examining myths, beliefs, and perceptions about COVID-19, 45.1% and 62.2% of respondents  
324 perceived that children and youth, respectively, are at a moderate risk of COVID-19 (23). This  
325 variation may be due to the difference sources and timely access of risk information of respondents.

326

327 Several studies have explored the knowledge and adoption of actions considered important to  
328 prevent the spread of coronavirus disease in the general population, (17, 23-25), however, none of  
329 these explored the perceptions of quarantined individuals. Quarantine was not listed as a preventive  
330 measure by our study participants, however, most of our participants identified monitoring for signs  
331 and symptoms of coronavirus disease as the purpose of the mandatory institutional quarantine.  
332 Regarding pathway to seeking care for COVID-19 related symptoms in the future, 48.8% and 28.3%  
333 of our study participants claimed they would either first visit a hospital or health facility or call the  
334 national coronavirus helpline on 1025, respectively, if they experienced symptoms of coronavirus  
335 disease.

336

337 Looking at facility related factors during quarantine, there was prominent dissatisfaction reported by  
338 most of our participants regarding the facility environment, specifically cleanliness and the services  
339 provided, specifically meals. Our findings are similar to the experiences of persons in COVID-19  
340 institutional quarantine in Uganda (26). In The Gambia, all quarantine related costs were covered by  
341 the government and therefore our study participants were not financially burdened with the cost of  
342 accommodation, food, testing and certification. Nevertheless, a predominant dissatisfaction related  
343 to cost noted by our study participants was specifically the desire to have been provided with  
344 compensation in the form of cash, food and/or transport for their time spent in quarantine, which  
345 would have benefited their families. In other countries, support measures such as financial support,  
346 employment benefits and/or practical support were adopted to enable people to follow self-  
347 isolation or quarantine guidance. In Germany, all employed people in mandatory self-isolation who  
348 tested positive were provided with 100% remuneration of their salary for up to six weeks and in  
349 South Korea, any person required to quarantine for 14 days was provided with daily necessities,  
350 sanitary kits and financial support (27). Although these findings are from high income countries,  
351 similar measures could be adapted in low- and middle- income countries such as The Gambia.

352

353 Adherence to COVID-19 prevention and control measures for quarantined individuals are key to the  
354 success of institutional quarantine. Most participants claimed to have adhered to the measures,  
355 however, few deviated and disregarded guidelines. From the limited literature published globally on  
356 adherence to self-isolation measures, it has been found that public trust, an altruistic attitude and  
357 access to information increase adherence to social/physical distancing protocols (27, 28). We utilized  
358 a conceptual framework for quarantine acceptance and adherence to investigate variables affecting  
359 adherence to mandatory COVID-19 quarantine (29). In relation to the framework, our results  
360 showed that individuals life circumstances and quarantine measures being enforced were the main  
361 predictors of their adherence.

362

363 Stigma experiences such as discrimination and behaviours such gossip can negatively affect the  
364 mental health of the stigmatized individual and lead to social exclusion. Stigma practices can include  
365 stereotypes, prejudice, stigmatizing behaviours, and discriminatory attitudes (10). In this study,  
366 COVID-19 related stigma was hardly experienced before and during quarantine but rather following  
367 on from quarantine. Some participants reported people gossiping about them, family and friends  
368 refusing to visit them at home and even being called names to make them feel ashamed. Others had  
369 people call them thinking they were being treated for coronavirus disease at the quarantine facility,  
370 and misunderstandings such as these stem from public fears of being infected by people in  
371 quarantine (7). Our participants also experienced overt discrimination (enacted stigma) after  
372 quarantine leading to feelings of shame, self-blame, loneliness, stress, anger and withdrawal from  
373 daily activities (felt stigma). Unfortunately, this is a reality many other individuals in self-isolation  
374 and quarantine have faced worldwide as found in recent studies (30, 31). It is therefore important to  
375 continually uncover and mitigate the many drivers of stigma to improve the lives of quarantined  
376 individuals.

377

### 378 **Study strengths and limitations**

379 One of the key strengths of this study was carrying out interviews via mobile phone call guided by  
380 our REDCap questionnaire displayed on tablets. This method seemed most appropriate over  
381 traditional face-to-face interviewing due to the rise of new cases at the time and government  
382 COVID-19 preventive mandates such as social/physical distancing. Our chosen method ensured we  
383 generated rapid first-hand evidence and helped us save on travel costs. Another key strength of our  
384 study was conducting interviews soon after participants left quarantine to minimise their recall bias  
385 after changing environment.

386

387 It is important to note that our study population does not represent the diverse demographics of  
388 The Gambia and not every person who went through institutional quarantine was included. Most of

389 our participants were Gambian males who are more likely to travel than women during the  
390 pandemic and therefore were more frequently identified for institutional quarantine. Our chosen  
391 methods of phone call surveys reduced our opportunity to take note of participants' nonverbal  
392 communication. We did not explore the stigma experience of associative stigma, meaning any  
393 stigma experienced by family or friends of those who went through institutional quarantine as they  
394 were not the cohort of focus in our study.

395

## 396 **CONCLUSION**

397 This study sheds light on the lives of those who had been in institutional quarantine in The Gambia  
398 and provides evidence that can be used to inform the Gambia MoH, policy makers and other key  
399 stakeholders working on epidemic preparedness and response. Such information was missing in the  
400 literature in general at the time this study was conducted. To improve the experience of individuals  
401 in designated institutional quarantine facilities, there should be daily provision of quality food and  
402 healthcare, timely provision of SARS-CoV-2 test results and negligible or complimentary associated  
403 costs of quarantine. Furthermore, stigma experiences and practices should be addressed during and  
404 after individuals stay in quarantine via the provision of psychosocial support.

405

## 406 **ACKNOWLEDGMENTS**

407 We express heartfelt gratitude to our study participants for their valuable time and genuine  
408 responses to our mobile phone surveys. We would like to sincerely thank the Ministry of Health staff  
409 who supported our data collection: Baboucarr Jallow, Modou Jallow, and Ebrima Keita.

410

## 411 **AUTHORS' CONTRIBUTIONS**

412 PJ conceived and designed the study with feedback from ON, AMM, OC, LL, AC, BK, MB, BM, SS and  
413 SS. AMM, OC, LL, AC, ON, BJ, MJ, EK, PJ collected data. AMM, PJ along with LS resolved data queries  
414 and cleaned the dataset. PJ and ON led data analysis and the literature search, wrote the first draft

415 of the manuscript, and updated subsequent versions of the paper with co-author comments and  
416 inputs. PJ, ON, BK, AMM, OC, LL, AC contributed to the interpretation of results, revised the  
417 manuscript, suggested policy implications from the findings and edited the final draft. All authors  
418 read and approved the final manuscript.

419

#### 420 **FUNDING**

421 The authors received no specific funding for this study. However, data collection activities were  
422 funded from discretionary MRCG@LSHTM funds awarded to BK. The funders had no role in the design  
423 of the study and collection, analysis, and interpretation of data and in writing the manuscript.

424

#### 425 **AVAILABILITY OF DATA AND MATERIALS**

426 The datasets used and/or analysed during the current study are available from the corresponding  
427 author on reasonable request.

428

#### 429 **DECLARATIONS**

##### 430 **Consent for publication**

431 Not applicable.

432

##### 433 **Competing interests**

434 None declared.

435

#### 436 **ORCID IDs**

437 Penda Johm  <https://orcid.org/0000-0001-8932-7082>

438 Oluwatosin Nkereuwem  <https://orcid.org/0000-0001-6194-6228>

439 Beate Kampmann  <https://orcid.org/0000-0002-6546-4709>

440

441 **LIST OF ABBREVIATIONS**

442	AoU	Assessment of Understanding
443	ANC	Antenatal care
444	CDC	Centres for Disease Control and Prevention
445	COVID-19	Coronavirus disease 2019
446	GDHS	Gambia Demographic and Health Survey
447	HSB	Health seeking behaviour
448	MICS	Multiple Indicator Cluster Survey
449	MoH	Ministry of Health
450	MRCG at LSHTM	Medical Research Council Unit The Gambia at the London School of Hygiene
451		and Tropical Medicine
452	PASS	Partners for Applied Social Sciences
453	REDCap	Research Electronic Data Capture
454	SARS-Cov-2	Severe Acute Respiratory Syndrome Novel Coronavirus-2
455	WHO	World Health Organization

456


457 **REFERENCES**

- 458 (1) World Health Organization. WHO Director-General's opening remarks at the media briefing on  
459 COVID-19-11 March 2020.
- 460 (2) Worldometer of COVID-19 Coronavirus pandemic, Available at:  
461 <https://www.worldometers.info/coronavirus/> (Accessed on 20th August 2021).
- 462 (3) World Health Organization. COVID-19 Weekly Epidemiological Update Edition 43, Published 8  
463 June 2021.
- 464 (4) The Gambia COVID-19 Outbreak Situational Report #359. (2021, August 20). MINISTRY OF  
465 HEALTH. [https://www.moh.gov.gm/wp-content/uploads/2021/08/GMB-COVID-19-Situational-  
466 Report-2021-20th-August-No-359.pdf](https://www.moh.gov.gm/wp-content/uploads/2021/08/GMB-COVID-19-Situational-Report-2021-20th-August-No-359.pdf)

- 467 (5) Lowe M. Using rapid online survey to assess public perceptions of Covid-19 in Gambia. The Pan  
468 African Medical Journal. 2020;35(Suppl 2).
- 469 (6) COVID-19 and Your Health. (2020, February 11). Centers for Disease Control and Prevention.  
470 <https://www.cdc.gov/coronavirus/2019-ncov/your-health/quarantine-isolation.html>
- 471 (7) Lio CF, Cheong HH, Lei CI, Lo IL, Lam C, Leong IH. Minimizing the risk of community spread of  
472 COVID-19 via institutional quarantine of high-risk travelers with serial viral RNA testing: A successful  
473 experience from Macao SAR, China. World journal of clinical cases. 2020 Jul 6;8(13):2674.
- 474 (8) The Abdul Latif Jameel Poverty Action Lab (J-PAL). (2020). Increasing adherence to COVID-19  
475 guidelines: Lessons from existing evidence. (online) Available at:  
476 [https://www.povertyactionlab.org/blog/4-10-20/increasing-adherence-covid-19-guidelines-lessons-](https://www.povertyactionlab.org/blog/4-10-20/increasing-adherence-covid-19-guidelines-lessons-existing-evidence)  
477 [existing-evidence](https://www.povertyactionlab.org/blog/4-10-20/increasing-adherence-covid-19-guidelines-lessons-existing-evidence)
- 478 (9) Link BG, Phelan JC. Conceptualizing stigma. Annual review of Sociology. 2001 Aug;27(1):363-85.
- 479 (10) Stangl AL, Earnshaw VA, Logie CH, van Brakel W, Simbayi LC, Barré I, Dovidio JF. The Health  
480 Stigma and Discrimination Framework: a global, crosscutting framework to inform research,  
481 intervention development, and policy on health-related stigmas. BMC medicine. 2019 Dec;17(1):1-3.
- 482 (11) Mwakikagile G. The Gambia and its people: ethnic identities and cultural integration in Africa.  
483 New Africa Press; 2010.
- 484 (12) UNICEF. The Gambia multiple indicator cluster survey 2018: survey findings report. InThe  
485 Gambia multiple indicator cluster survey 2018: survey findings report 2019 (pp. 694-694).
- 486 (13) Gambia Bureau of Statistics (GBoS) and ICF. 2021. The Gambia Demographic and Health Survey  
487 2019-20. Banjul, The Gambia and Rockville, Maryland, USA: GBoS and ICF.
- 488 (14) Gibbs GR. Thematic coding and categorizing. Analyzing qualitative data. 2007;703:38-56.
- 489 (15) Braun V, Clarke V. Using thematic analysis in psychology. Qualitative research in psychology.  
490 2006 Jan 1;3(2):77-101.
- 491 (16) Muela SH, Ribera JM, Toomer E, Grietens KP. The PASS-model: a model for guiding health-  
492 seeking behavior and access to care research. Malaria Reports. 2012 Dec 13;2(1):e3-

- 493 (17) Mohamed AA, Elhassan EA, Mohamed AO, Mohammed AA, Mahgoop MA, Sharif ME, Bashir MI,  
494 Abdelrahim RB, Idriss WI, Malik EM. Knowledge, attitude and practice of the Sudanese people  
495 towards COVID-19: An online survey. BMC public health. 2021 Dec;21(1):1-7.
- 496 (18) Eyeberu A, Mengistu DA, Negash B, Alemu A, Abate D, Raru TB, Wayessa AD, Debela A, Bahiru  
497 N, Heluf H, Kure MA. Community risk perception and health-seeking behavior in the era of COVID-19  
498 among adult residents of Harari regional state, eastern Ethiopia. SAGE Open Medicine. 2021  
499 Jul;9:20503121211036132.
- 500 (19) Asnakew Z, Kerebih Asrese MA. Community risk perception and compliance with preventive  
501 measures for COVID-19 pandemic in Ethiopia. Risk Management and Healthcare Policy.  
502 2020;13:2887.
- 503 (20) Birhanu Z, Ambelu A, Fufa D, Mecha M, Zeynudin A, Abafita J, Belay A, Doyore F, Oljira L, Bacha  
504 E, Feyisa J. Risk perceptions and attitudinal responses to COVID-19 pandemic: an online survey in  
505 Ethiopia. BMC public health. 2021 Dec;21(1):1-7.
- 506 (21) Rohrmann B. Risk perception, risk attitude, risk communication, risk management: A conceptual  
507 appraisal. In 15th International Emergency Management Society (TIEMS) Annual Conference 2008 Jun  
508 (Vol. 2008).
- 509 (22) Person B, Sy F, Holton K, Govert B, Liang A. Fear and stigma: the epidemic within the SARS  
510 outbreak. Emerging infectious diseases. 2004 Feb;10(2):358.
- 511 (23) Kebede Y, Yitayih Y, Birhanu Z, Mekonen S, Ambelu A. Knowledge, perceptions and preventive  
512 practices towards COVID-19 early in the outbreak among Jimma university medical center visitors,  
513 Southwest Ethiopia. PloS one. 2020 May 21;15(5):e0233744.
- 514 (24) Alkhalidi G, Aljuraiban GS, Alhurishi S, De Souza R, Lamahewa K, Lau R, Alshaikh F. Perceptions  
515 towards COVID-19 and adoption of preventive measures among the public in Saudi Arabia: a cross  
516 sectional study. BMC public health. 2021 Dec;21(1):1-21.

- 517 (25) Mendy PA, Jawo E, Jallow BJ, Ceesay B, Mendy SA, Camara B, Ndaw M. Knowledge, Attitude and  
518 Practice of Gambians towards the Novel Coronavirus (COVID-19) Pandemic in The Gambia: A cross-  
519 sectional study. African Journal of Health Sciences. 2021 Jun 10;34(1):134-47.
- 520 (26) Ndejjo R, Naggayi G, Tibiita R, Mugahi R, Kibira SP. Experiences of persons in COVID-19  
521 institutional quarantine in Uganda: a qualitative study. BMC public health. 2021 Dec;21(1):1-1.
- 522 (27) Patel J, Fernandes G, Sridhar D. How can we improve self-isolation and quarantine for covid-19?.  
523 bmj. 2021 Mar 10;372.
- 524 (28) Webster RK, Brooks SK, Smith LE, Woodland L, Wessely S, Rubin GJ. How to improve adherence  
525 with quarantine: rapid review of the evidence. Public Health. 2020 May 1;182:163-9.
- 526 (29) Sopory P, Novak JM, Noyes JP. Quarantine acceptance and adherence: qualitative evidence  
527 synthesis and conceptual framework. Journal of Public Health. 2021 Apr 16:1-1.
- 528 (30) Lohiniva AL, Dub T, Hagberg L, Nohynek H. Learning about COVID-19-related stigma, quarantine  
529 and isolation experiences in Finland. PloS one. 2021 Apr 14;16(4):e0247962.
- 530 (31) Turner-Musa J, Ajayi O, Kemp L. Examining social determinants of health, stigma, and COVID-19  
531 disparities. InHealthcare 2020 Jun (Vol. 8, No. 2, p. 168). Multidisciplinary Digital Publishing Institute.



\*Switched off 123; No answer 51; Wrong number 5; Busy/unavailable 12; Died 1; Mentally unstable 1; Wrong person 8

**Figure 1. Participant recruitment process based on study inclusion criteria.**

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .




Figure 2. Likert scale gauging attitudes towards COVID-19 related stigma.