

1 **Child marriage and adverse reproductive outcomes among young Afghan women:**
2 **implication for policy and practice in Afghanistan**

3
4 **Authors:**

5 Omid Dadras^{1,2}, Takeo Nakayama³

6
7
8 **Affiliations**

9 ¹ Iranian Research Center for HIV/AIDS (IRCHA). Tehran University of Medical Sciences, Iran.

10 ² Excellent Center for Dengue and Community Public Health (EC for DACH), Walailak
11 University, Thailand

12 ³ Department of Health Informatics, School of Public Health, Kyoto University, Japan.

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

Corresponding authors: Omid Dadras, Iranian Research Center for HIV/AIDS (IRCHA),
Tehran University of Medical Sciences, Iran.

Tel: +989114058914

Email: omiddadras@yahoo.com

35 **Abstract**

36 **Background:** Child marriage is defined as marriage before the age of 18 years and it has been
37 linked to several adverse health and social outcomes. This study aimed to explore the
38 prevalence and determinants of child marriage and its association with adverse reproductive
39 outcomes among a nationally representative sample of young Afghan women.

40 **Methods:** This was a secondary analysis of the 2015 Afghanistan Demographic and Health
41 Survey (DHS) data. Based on the UN recommendation on child marriage study, only women
42 aged 20-24 years old were included. Descriptive statistics and binary logistic regression were
43 employed to determine the distribution of respondents' characteristics and prevalence and
44 determinants of child marriage. Multivariate logistic models examined the association between
45 child marriage and adverse reproductive outcomes accounting for the sociodemographic
46 factors.

47 **Results:** An estimated 52% of the Afghan women aged 20-24 married at ages less than 18
48 years. Poor illiterate women were more likely to marry at early ages (<18). There was a
49 significant negative relationship between child marriage and history of rapid repeat childbirth,
50 delivery by skilled personnel, and institutional delivery. In both adjusted and unadjusted models,
51 women married at age ≤ 14 were more likely to experience terminated or unintended pregnancy
52 (AOR = 1.89, 95% CI: 1.31-2.75 and AOR = 2.20, 95% CI: 1.24-3.91, respectively), inadequate
53 ANC (AOR = 1.71, 95% CI: 1.01-2.90), unmet need for family planning (AOR = 1.44, 95% CI:
54 1.05-1.98), fistula (AOR = 2.36, 95% CI: 1.22-4.57); While, for those married at age 15-17
55 years, only terminated or unintended pregnancy remained significant.

56 **Conclusion:** The younger age of marriage was associated with a higher prevalence of adverse
57 reproductive outcomes among Afghan women. Moreover, poverty and illiteracy proved to be
58 important predictors of child marriage in Afghan women. Strict international law enforcement
59 and advocacy are a need in the current situation of Afghanistan to increase young women's
60 education, promote their civil rights, and improve their autonomy and role in decision-making
61 concerning their health.

62 **Keywords:** Afghan women, child marriage, reproductive outcome, Afghanistan

63
64
65
66
67
68

69

70

71 **Introduction**

72 Child marriage is defined as marriage before the age of 18 years and is increasingly recognized
73 as a violation of human rights and a serious public health issue (1). Although one-fifth of women
74 worldwide were subjected to child marriage in 2020, the geographic distribution of child
75 marriage could vary significantly across the regions and countries and it is mostly concentrated
76 in poor- and middle-income countries (2). For instance, approximately a third of women in Sub-
77 Saharan Africa were subjected to child marriage in 2017 (3); while this figure for South Asia was
78 about 45% the same year with different rates across countries in this region (4). Child marriage
79 could not only harm the gender equality efforts but also the achievement of millennium
80 development goals, particularly goals 1 to 5: eradication of poverty, achievement of universal
81 education, women empowerment, reducing child mortality, and improving maternal health (5).

82 It has been shown that it is associated with a number of adverse health and social outcomes
83 such as unintended and high-risk pregnancy, maternal and infant mortality, human
84 immunodeficiency virus/acquired immunodeficiency syndrome, cross-generational sex, obstetric
85 fistula, high maternal mortality and morbidity, intimate partner violence, and depression (6-8).
86 These adverse health outcomes have been attributed to several factors, including limited
87 access to media and inadequate health education, less autonomy in decision making and
88 inability to actively engage in health promotion programs, limited social engagement, and
89 inability to adequately access health care (9, 10). Empowering young adolescent girls is
90 essential to improve the overall reproductive and child health outcomes and accelerate social
91 and economic development in low- and middle-income countries (11). However, it appeared that
92 the content of reproductive health programs is often not serving this key vulnerable population
93 who may be at a higher risk for poor reproductive health outcomes (1, 12).

94 Despite the current evidence of possible adverse reproductive outcomes among girls who marry
95 at early ages, there is still a gap in knowledge of true association and determinants of child
96 marriage and reproductive outcomes in some of the most afflicted countries such as
97 Afghanistan. The long-standing ignorance of young women's reproductive needs in patriarchal
98 societies such as Afghanistan not only harmed the young women's health but also the health of
99 offspring who may suffer from perinatal complications (13). In addition, it limited the efforts of
100 national and international parties in improving adolescent reproductive health through well-
101 tailored programs that specifically target this vulnerable subgroup in Afghan society. To our
102 knowledge, this is the first study in Afghanistan that explored the prevalence and determinants

103 of child marriage and its association with adverse reproductive outcomes among a nationally
104 representative sample of Afghan women. The results are to inform the policies and interventions
105 of the most recent situation concerning child marriage and adverse reproductive outcomes in
106 Afghanistan.

107 **Methods**

108 **Study design**

109 This was a cross-sectional study using the data from the most recent Demographic and Health
110 Survey (DHS) in Afghanistan conducted in 2015. It is a nationally representative survey
111 implemented by the Central Statistics Organization (CSO) in collaboration with the Afghanistan
112 Ministry of Public Health (MoPH) and funded by the United States Agency for International
113 Development (USAID).

114 **Study population and sampling**

115 Afghanistan DHS 2015 (ADHS 205) is nationally representative that collected the data for
116 women aged 15-49 years. The ADHS 2015 employed a stratified two-stage sample design to
117 estimate the key indicators at the national level, in urban and rural areas, and for each of the 34
118 provinces of Afghanistan. At the first stage, 950 clusters consisting of EAs (enumeration areas)
119 were selected; 260 in urban areas and 690 in rural areas. At the second stage, 25,650
120 households were selected through an equal probability systematic selection process. Weighting
121 components were calculated and applied to obtain a representative estimate at the national
122 level. All the women aged 15-49 years who were either permanent residents of the selected
123 households or visitors who stayed in the households the night before the survey were recruited
124 after informed consent. Our analyses have been restricted to women 20-24 years of age, the
125 United Nations (UN) recommended age group for child marriage assessment (14). Selecting
126 this age group not only enhances the comparability of the results but also minimizes the
127 possibility of errors due to recall which may occur as the participants move to the older age
128 groups (15).

129 **Scales and measurement**

130 A standard DHS questionnaire including the questions related to the demographic and health
131 issues was administered to all eligible women. The questionnaire collected data on women's
132 demographic characteristics, household information, family planning, fertility preferences, child
133 health, sexually transmitted diseases, marriage and sexual behavior, and domestic violence
134 (16). However, for the current study, our primary interest was questions related to child marriage
135 and reproductive outcomes.

136 **Explanatory variables**

137 The main explanatory variable was the age at first marriage which was translated into child
138 marriage if the respondent's age was less than 18 years at the time of marriage. The
139 respondents were further classified into three groups based on the age at the first marriage: ≥ 18
140 years (marriage as an adult), 15-17 years (marriage in middle adolescence), and ≤ 14 years of
141 age (marriage in early adolescence or childhood) to capture the potential vulnerability that may
142 exist for the marriage in very early ages.

143 **Outcomes of interest**

- 144 • **Early fertility:** It was a binary variable indicating whether the woman had a birth within
145 the first 12 months of marriage.
- 146 • **Unmet need for family planning:** It was a binary variable that includes the unmet need
147 for limiting (i.e. women whose most recent pregnancy was not wanted at all, fecund
148 women who did not use contraception despite their desire to have no more children,
149 women who were postpartum amenorrhoeic for 2 years following an unwanted birth and
150 were not using contraception) and spacing (i.e. women whose most recent pregnancy
151 was not wanted initially but wanted later, fecund women not using contraception who
152 were undecided when/if they wanted a to have a child or who wanted a child 2+ years
153 later, and women who were postpartum amenorrhoeic for 2 years following a mistimed
154 birth and were not using contraception) (17).
- 155 • **History of rapid repeat childbirth:** It was defined as having had at least one birth
156 within 24 months of previous childbirth.
- 157 • **History of pregnancy termination:** It was a binary variable indicating whether a
158 woman has ever had a pregnancy ended by miscarriage, abortion, or stillbirth.
- 159 • **Unintended pregnancy:** It was a binary indicator of whether the respondent has had at
160 least one child in the past 5 years that was wanted later or not wanted at all.
- 161 • **Inadequate antenatal care (ANC):** It indicates the number of antenatal visits during the
162 last pregnancy. Based on the World Health Organization guidelines, less than 4 ANC
163 visits were considered inadequate.
- 164 • **Delivery by skilled personnel:** It indicates whether the woman had received
165 assistance from a skilled health care worker in her last birth.
- 166 • **Institutional delivery:** It was a binary variable indicating whether the woman delivered
167 her last child in a health care facility (public or private).
- 168 • **Fistula:** It was a binary variable defined as the involuntary loss of urine and/or feces
169 through the vagina after labour.

170 **Covariates**

171 Based on the literature review, the most relevant
172 sociodemographic factors such as age (continuous in
173 years), education (literate, illiterate), employment
174 (yes/no) area of residence (rural, urban), wealth quintile
175 (poor, middle, rich), husband age difference (≤ 5 , 6-9,
176 ≥ 10), husband education (literate, illiterate), polygamy,
177 listening to radio, watching television, and ethnicity were
178 also collected and accounted for in the multivariate
179 analysis.

180 **Statistical Analysis**

181 All the analyses were restricted to the age group 20-24 years. Descriptive statistics were
182 employed to describe the respondent's characteristics in this age group. Binary logistic
183 regression estimated the odds of child marriage across the different sociodemographic factors.
184 Multivariate Logistic regression models were constructed to examine the association between
185 various outcome variables and age at marriage before and after adjustment for
186 sociodemographic factors. Results are presented as odds ratio (OR) and adjusted odds ratio
187 (AOR) with 95% confidence intervals (95% CI). To avoid the bias in estimates for the history of
188 rapid repeat births, we restricted the sample to women who had a minimum of two children. All
189 the analyses were performed using STATA version 14 and sampling weight and design were
190 defined and accounted for using the STATA command "svy" in all analyses. All tests of
191 hypothesis were 2-tailed, with a type 1 error rate fixed at 5%.

192 **Ethical consideration**

193 The data used in this study could be accessed upon a reasonable request from the DHS
194 website and are publicly available. The ethical necessities for data collection were sought and
195 assured by the institutions that commissioned, funded, or managed the surveys. All DHS are
196 reviewed and approved by ICF International and Institutional Review Board of Ministry of Health
197 in the target country to ensure that the protocols comply with the U.S. Department of Health and
198 Human Services regulations for the protection of human subjects. Therefore, this study did not
199 require further ethical approval.

200

Table 1. The characteristics of Afghan women aged 20-24 years, ADHS 2015 (N=6083)

Variables	n (weighted%)
Age at marriage	
≥ 18 (adult)	1850 (47.97)
15-17 (middle)	1559 (38.92)

201 **Results** Table 2. The sociodemographic
 202 **Table 1** determinants of child marriage among
 203 Afghan aged 20-24 years, ADHS 2015
 204 presents the distribution of sociodemographic factors of
 205 the respondents, A total of 6083 young women aged
 206 20-24 were included in this study to estimate the child
 207 marriage prevalence and explore its association with
 208 adverse reproductive outcomes in Afghan women living
 209 in Afghanistan. The majority of the respondents were
 210 illiterate (72.56%), unemployed (87.03%), and were
 211 living in rural areas (76.73%). Approximately 70% of the
 212 women were married to a man who was no more than 5
 213 years older than them. Half of the husbands were
 214 literate and almost all of them were employed. Almost
 215 an equal proportion of women a fifth were in each
 216 wealth index group. Less than half of the respondents
 217 reported watching television (48.23%) or listening to the
 218 radio (42.31%) at least once in the last week. The
 219 highest proportion of respondents were from Pashtun
 220 (42.75%) and Tajik (32.70%) ethnic groups; while, the
 221 lowest proportions belonged to Baloch (0.52%),
 Nuristani (0.82%), and Pashai (0.96%).

222 **Child marriage prevalence and its**
 223 **sociodemographic determinants**

224 As table 2 illustrates, almost half of the respondents
 225 married at ages less than 18 years, including 38.92%
 226 married as a middle adolescent (15-17 years of age)
 227 and 13.11% married as a young adolescent or a child
 228 (≤ 14 years of age). The odds of child marriage were

229 20% lower in literate women compared to illiterate ones. Women with ≥ 10 spousal age gap
 230 were more likely to endure child marriage compared to the reference group (spousal age gap
 231 ≤ 5). The women from the poorest households were more likely to marry at ages less than 18
 232 years (OR = 1.43, 95% CI:1.09-1.89). The likelihood of early marriage (age < 18) was
 233 significantly higher in Tajik (OR = 1.39, 95% CI:1.07-1.79) and Baloch (OR = 2.92, 95% CI:1.43-
 234 5.95) ethnic groups compared to reference group (Pashtun).

adolescents)	
≤ 14 (young adolescents/children)	514 (13.11)
Age groups (years)	
20	1721 (26.74)
21	736 (12.28)
22	1278 (24.03)
23	1144 (18.01)
24	1104 (18.94)
Education	
Illiterate	4615 (72.56)
Literate	1468 (27.44)
Employment	
Employed	769 (12.97)
Unemployed	5314 (87.03)
Husband age gap (year)	
≤ 5	4427 (70.17)
6-9	985 (17.50)
≥ 10	671 (12.33)
Husband Education	
Illiterate	2926 (50.97)
Literate	3987 (49.03)
Wealth index	
Poorest	1103 (19.29)
Poorer	1388 (21.02)
Middle	1307 (19.44)
Richer	1323 (21.08)
Richest	962 (19.17)
Place of Living	
Urban	1423 (23.27)
Rural	4660 (76.73)
Watch television	
Yes	2685 (48.23)
No	3384 (51.77)
Listen to radio	
Yes	2568 (42.31)
No	3507 (57.69)
Ethnicity	
Pashtun	2715 (42.75)
Tajik	1811 (32.70)
Hazara	496 (8.64)
Uzbek	384 (9.68)
Turkmen	72 (1.91)
Nuristani	323 (0.82)
Baloch	68 (0.52)
Pashai	113 (0.96)
Other	91 (2.01)

*The total number for some variable does not add up to 6083 due to the missing data.

235 **The**
 236 **marriage**
 237 Table 3
 238 and
 239 intervals for
 240 and adverse
 241 marriage
 242 ≤14 (early
 243 adolescent),
 244 variability of
 245 across the
 246 outcomes
 247 1) Early
 248 association
 249 fertility even
 250 factors.
 251 2) Unmet
 252 married at
 253 from the
 254 only before
 255 also after
 256 adjustment for sociodemographic factors. No significant association was observed for those

Variables	Age at marriage <18 OR (95% CI)
Education	
Illiterate	Ref
Literate	0.80 (0.65-0.99)
Employment	
Employed	1.10 (0.84-1.43)
Unemployed	Ref
Husband age gap (year)	
≤5	Ref
6-9	1.19 (0.90-1.59)
≥10	1.82 (1.28-2.61)
Wealth index	
Poorest	1.43 (1.09-1.89)
Poorer	1.06 (0.79-1.42)
Middle	0.95 (0.69-1.44)
Richer	1.11 (0.85-1.44)
Richest	Ref
Place of Living	
Urban	Ref
Rural	0.98 (0.78-1.25)
Polygamy	
Yes	1.35 (0.77-2.36)
No	Ref
Ethnicity	
Pashtun	Ref
Tajik	1.39 (1.07-1.79)
Hazara	1.05 (0.78-1.42)
Uzbek	1.13 (0.74-1.70)
Turkmen	0.72 (0.43-1.21)
Nuristani	0.91 (0.64-1.29)
Baloch	2.92 (1.43-5.95)
Pashai	0.71 (0.45-1.13)
Other	1.08 (0.62-1.88)

OR: Odds ratio, CI: Confidence interval

association between the age of and adverse reproductive outcomes
 represents the odds ratios (adjusted unadjusted) and 95% confidence intervals for the association between child marriage and adverse reproductive outcomes. The age at marriage was classified into three categories: adolescent/child, 15-17 (middle adolescent) and ≥18 (adult) to capture the variability of reproductive outcomes prevalence across the younger age groups. Nine reproductive outcomes were included in the analysis as below:
 1) Early fertility: It appeared that there is no association between age at marriage and early fertility even after adjustment for sociodemographic factors.
 2) Unmet needs for family planning: Women aged ≤14 were more likely to suffer unmet needs for family planning not only before adjustment for sociodemographic factors (OR = 1.37, 95% CI: 0.99-1.90) and but also after adjustment for sociodemographic factors (AOR = 1.44, 95% CI: 1.05-1.98). No significant association was observed for those

Table 3. The association between the age at marriage and adverse reproductive outcomes among Afghan women aged 20-24 years, ADHS 2015

Variables	Age at marriage ≤14 ^a		Age at marriage 15-17 ^a	
	OR (95% CI)	AOR ^b (95% CI)	OR (95% CI)	AOR ^b (95% CI)
Early fertility ^c	0.90 (0.63-1.28)	0.86 (0.60-1.24)	0.94 (0.76-1.15)	0.93 (0.75-1.16)
Unmet needs for family planning	1.37 (0.99-1.90)	1.44 (1.05-1.98)	1.19 (0.93-1.54)	1.23 (0.96-1.57)
History of rapid repeat childbirth ^d	0.34 (0.24-0.48)	0.27 (0.18-0.39)	0.56 (0.42-0.74)	0.45 (0.33-0.61)
History of pregnancy termination	1.81 (1.26-2.59)	1.89 (1.31-2.75)	1.45 (0.98-2.13)	1.57 (1.09-2.66)
Unintended pregnancy	1.94 (1.15-3.26)	2.20 (1.24-3.91)	1.32 (0.85-2.05)	1.61 (1.03-2.56)
Inadequate ANC (<4 ANC visits)	1.92 (1.23-3.00)	1.71 (1.01-2.90)	1.55 (1.07-2.24)	1.42 (0.91-2.23)
Delivery by skilled personnel	0.51 (0.38-0.69)	0.48 (0.34-0.69)	0.67 (0.53-0.86)	0.64 (0.48-0.85)
Institutional delivery	0.49 (0.36-0.66)	0.45 (0.32-0.66)	0.68 (0.54-0.86)	0.65 (0.49-0.85)
Fistula	2.24 (1.10-4.57)	2.36 (1.22-4.57)	1.39 (0.70-2.78)	1.38 (0.69-2.75)

OR: Odds ratio, CI: Confidence interval.

^a The reference group: Age at marriage ≥18.

^b AOR: Adjusted odds ratio (controlled for current age, education, wealth index, ethnicity, husband education, husband age gap, watching tv, listening radio).

^c Within one year after marriage.

^d Defined as birth interval < 24 months in last pregnancy.

* Due to the small sample size of some ethnic groups, this variable was not included in multivariate analysis.

257 married at ages 15-17 years compared to the reference group (married at ≥ 18)

258 3) History of rapid repeat childbirth: It was defined as birth interval < 24 months in the last
259 pregnancy and only women with at least two births were included. It appeared that the likelihood
260 of having rapid repeat pregnancy is significantly lower among women married at both ≤ 14 and
261 15-17 years of age compared to those married at age > 18 years, even after adjustment for
262 sociodemographic factors (AOR = 0.27, 95% CI: 0.18-0.39 and AOR = 0.45, 95% CI: 0.33-0.61,
263 respectively).

264 4) History of pregnancy termination (stillbirth, abortion, miscarriage): There was a significant
265 positive association between child marriage and history of pregnancy termination after
266 adjustment for sociodemographic factors. In the adjusted models, the odds of experiencing
267 terminated pregnancy (stillbirth, abortion, or miscarriage) were almost 90% and 60% higher
268 among those married at ≤ 14 and 15-17 years of age compared to those married at older ages.

269 5) Unintended pregnancy: The odds of experiencing unintended pregnancy among girls married
270 at age ≤ 14 was almost twice the odds of women married at ≥ 18 in both unadjusted and adjusted
271 models (OR = 1.94, 95% CI: 1.15-3.26 & AOR = 2.20, 95% CI: 1.24-3.01). Likewise, in the
272 adjusted model, women married at age 15-17 years were more likely to have an unintended
273 pregnancy (AOR = 1.61, 95% CI: 1.03-2.56).

274 6) Inadequate ANC (< 4 ANC visits): Women married at ages less than 18 were more likely to
275 endure an inadequate ANC during their last pregnancy (OR = 1.92, 95% CI: 1.23-3.00 for
276 women married at ≤ 14 and OR = 1.55, 95% CI: 1.07-2.24 for those married at 15-18); however,
277 this association was weakened for those married at age 15-17 years after adjustment for
278 sociodemographic factors (AOR = 1.42, 95% CI: 0.91-2.23).

279 7) Delivery by skilled personnel: it appeared that likelihood of having a delivery by a skilled
280 health worker at last birth was significantly lower among women married at age < 18 . As Table 3
281 indicated, in both adjusted and unadjusted models, the odds of delivery by a skilled worker were
282 almost 50% and 35% lower among women married at age ≤ 14 and 15-17 compared to those
283 married at age > 18 years, respectively.

284 8) Institutional delivery (public, private): Similar to the delivery by skilled personnel, the odds of
285 having institutional delivery, in both adjusted and unadjusted models, was approximately 50%
286 and 35% lower among women married at age ≤ 14 and 15-17 compared to those married at age
287 > 18 years, respectively.

288 9) Fistula: Women married at age ≤ 14 had significantly higher odds of experiencing involuntary
289 loss of urine and/or feces through the vagina after labour compared to those married at age > 18
290 even after adjustment for sociodemographic factors (OR: 2.24, 95% CI: 1.10-4.57 & AOR =

291 2.36, 95% CI: 1.22-4.57). However, for those married at age 15-17 years, no significant relation
292 was observed

293 **Discussion**

294 Although several studies examined the impact of child marriage on adverse reproductive
295 outcomes in other countries and regions such as South Asian countries (4, 18, 19) or Sub-
296 Saharan African countries (20, 21), to the best of our knowledge, this is the first study that
297 explored such relationship among Afghan women living in Afghanistan. The results showed that
298 an estimated 52% of Afghan women aged 20-24 married at ages less than 18 years.
299 Furthermore, there was a significant association between marriage at early ages and adverse
300 reproductive outcomes among young Afghan women living in Afghanistan. It appeared that
301 women married at age ≤ 14 are more vulnerable and except for early fertility, they were more
302 likely to experience any other type of adverse reproductive outcomes including the history of an
303 unintended or terminated pregnancy, unmet need for family planning, inadequate ANC, delivery
304 by an unskilled worker in a non-institutional place, and fistula; whereas, marrying at age 15-17
305 years was only associated with a history of unintended, or terminated pregnancy and delivery by
306 an unskilled worker in a not-institutional place such as home after adjustment for
307 sociodemographic factors. However, they may also have inadequate ANC (<4) during their
308 pregnancy, even though, this association disappeared after accounting for the effect of
309 sociodemographic factors. We also found that there is a higher prevalence of child marriage
310 among poor, illiterate women who belong to the Tajik and Baloch ethnic groups.

311 Although more than half the women in this study endured child marriage, it appeared that the
312 prevalence of child marriage significantly decreased over the period 2010 to 2015 (76% in DHS
313 2010 vs 52% in DHS 2015 in women aged 20-24 years). This may be due to the national and
314 international campaigns and enormous efforts toward empowering the Afghan women that have
315 been shaped during the last decade. However, women from lower socioeconomic classes with
316 low literacy are still at a greater risk for child marriage. Similar to our study, several studies have
317 reported poverty and illiteracy as risk factors for child marriage (22) and adverse fertility
318 outcomes (23, 24). Poverty and illiteracy appear to limit the women's autonomy in critical
319 decisions and often exclude them from the decisions related to their marriage and reproductive
320 needs (24). In fact, it has been suggested that retaining girls at the school for a longer period
321 not only raise the age at the first marriage but also enhances the autonomy of the young girls
322 through individual development (18). In Afghanistan, child marriage is often a measure of
323 strengthening ties between rival families and tribes and is considered a peaceful solution to
324 settle debts and disputes. Many poor families often sell their adolescent girls in exchange for

325 large dowries from rich people and the husbands are usually much older (25). Similarly, in our
326 study, a higher chance of child marriage was observed in those with ≥ 10 years spousal age
327 gap. Almost all the countries with a high prevalence of child marriage are located in the
328 developing world (3). Child marriage is a common phenomenon in rural areas where poverty
329 and illiteracy are more dominant. Several studies indicated a higher prevalence of child
330 marriage among women living in rural areas (26). However, our study found no significant
331 difference in child marriage between rural and urban respondents, this could be due to the
332 widespread poverty and illiteracy of Afghan women regardless of their place of living. More
333 studies are suggested to scrutinize this issue among rural and urban habitants in Afghanistan.
334 Child marriage has been linked to several maternal and child adverse health outcomes such as
335 low birth weight, sexually transmitted diseases (STDs), unintended pregnancy, abortion,
336 stillbirth, maternal and child mortality (6, 8). There are also negative social impacts such as
337 social isolation and domestic violence (27); however, the findings are inconsistent. The
338 discrepancies in findings across studies have been attributed to the differences in the socio-
339 cultural and economic status of the target society (28). In our study, the odds of all adverse
340 reproductive outcomes, except early fertility and history of rapid repeat pregnancy, were higher
341 in women married at age ≤ 14 compared to those married as an adult (age > 18). We found no
342 significant association between early fertility and age at the marriage, which was in line with
343 previous studies from India, Nepal, and Pakistan (4); however, there was a negative association
344 between the history of rapid repeat pregnancy and age at marriage across all child marriage
345 groups. This has been attributed to the low fecundity at very young ages due to the physical and
346 biological immaturity and probably low coital frequency (29). Several factors could influence the
347 coital frequency at very young ages such as arranged marriage, delayed consummation, and
348 living in an extended family which reduces the chance of intercourse and early fertility (29).

349 An increase in odds of unmet need for family planning was documented among those married at
350 very young ages in the present study. This could partly justify by an increased desire to use
351 modern contraceptives among women married at young ages to prevent further unintended
352 pregnancies (20). The women married at very young ages often attain the desired number of
353 children in their twenties and they may seek contraceptives to avoid further pregnancy. This
354 could be a reason for higher needs for family planning at younger ages compared to those
355 married at older ages who may still want more children (30, 31). The higher unmet need for
356 family planning obviously leads to a higher prevalence of unintended pregnancy, which was
357 observed among women married at age < 18 in our study. Besides, unintended pregnancy often
358 leads to negative pregnancy outcomes such as abortion, miscarriage, and stillbirth, especially at

359 very young ages due to the anatomical and physiological immaturity of the body (32, 33).
360 Moreover, consistent with our finding, it has been shown that young girls aged 10-15 years are
361 more likely to suffer from obstetric fistula due to their small pelvises (34). Therefore, the reasons
362 and determinants of unmet need for family planning services among women married at very
363 young ages in Afghanistan should be a priority for future research.

364 We observed a negative association not only between delivery by a skilled worker at institutional
365 facilities but also between adequate ANC with child marriage. Similar findings have been
366 documented in other countries such as Pakistan, Nepal, and India (35-37). According to
367 previous studies, in Islamic patriarchal societies such as Afghanistan women married at very
368 young ages are more likely to suffer from spousal violence and often controlled by their
369 husbands or in-laws (38, 39). This is mainly due to the low empowerment of women marrying as
370 children and this could limit the autonomy in decision-making and access to reproductive and
371 maternal health services (40-42). Due to the dependence and low status of child brides, they
372 need to seek permission from the head of a family to visit health services (43), otherwise, they
373 may encounter spousal violence and this limits their access to maternal healthcare services
374 (44). This may also explain the high prevalence of other adverse reproductive outcomes in our
375 study.

376 **Limitations**

377 Although our study is the first study that measures the prevalence of child marriage and its
378 association with adverse reproductive outcomes in a nationally representative sample of Afghan
379 women aged 20-24 years, some cautions need to be considered in interpreting the results. The
380 study used self-reported data that are subjected to recall and socially desirable biases. For
381 example, they may be an error in reporting the age of respondent and age of marriage as there
382 was no documented registration system for age. Furthermore, some adverse reproductive
383 outcomes such as unintended or terminated pregnancy may be underreported. Also, pregnancy
384 termination did not differentiate between miscarriage, abortion, or stillbirth and hence
385 determining the relationship of each complication with child marriage was not possible.
386 Moreover, we were not able to distinguish between induced and natural pregnancy termination.
387 Due to the cross-sectional design of DHS surveys, the causal relationship between child
388 marriage and adverse reproductive outcomes may not be assumed. However, since the
389 reproductive outcomes occur following marriage, this assumption may not stand for this
390 particular study. Last, the findings were limited to the age group 20-24 years in Afghanistan and
391 cannot be generalized to other ages. However, the same age group of women has been

392 recommended by the UN and used in other international publications concerning child marriage
393 (19, 20, 45)

394 **Conclusion**

395 This study not only contributes to the current knowledge related to the impact of child marriage
396 on reproductive outcomes but also described the most recent situation of child marriage and its
397 sociodemographic determinants among Afghan women. The findings underscore the critical
398 need for policies and interventions to strive toward increasing the age of marriage among
399 Afghan girls. Although in recent years, Afghanistan has made substantial efforts to reduce child
400 marriage in face of incredible internal conflict, continued support from the international
401 community and human rights advocates are essential as the recent worsening and political
402 conflict could be a backslide to the progress that has been already made over the years. Strict
403 international law enforcement and advocacy are a need in the current situation of Afghanistan to
404 increase young women's education, promote their civil rights, and improve their autonomy and
405 role in decision-making concerning their health.

406 **Abbreviations**

407 UN: United Nations

408 DHS: Demographic and Health Survey

409 CSO: Central Statistics Organization

410 ANC: Antenatal care

411 AOR: Adjusted odds ratio

412 MoPH: Ministry of Public Health

413 STD: Sexually transmitted disease

414 **Ethics approval and consent to participate**

415 The data used in this study could be accessed upon a reasonable request from the DHS
416 website and are publicly available. The ethical necessities for data collection were sought and
417 assured by the institutions that commissioned, funded, or managed the surveys. All DHS are
418 reviewed and approved by ICF International and Institutional Review Board of Ministry of Health
419 in the target country to ensure that the protocols comply with the U.S. Department of Health and
420 Human Services regulations for the protection of human subjects. Therefore, this study did not
421 require further ethical approval.

422 **Consent for publication**

423 Not applicable.

424 **Availability of data and material**

425 The DHS questionnaire that collected the data in Afghanistan's demographic and health survey
426 in 2015 could be reached at DHS official website (<https://www.dhsprogram.com>). The dataset
427 that was used in this study could be available upon a reasonable request from the DHS website.

428 **Conflict of interest**

429 The authors declared no conflict of interest.

430 **Funding**

431 None

432 **Authors' contribution**

433 OD wrote the research protocol and performed the data analysis and prepared the final draft.
434 TN provided critical feedback and comments on the data analysis and results and help in
435 preparing the final draft.

436 **Acknowledgment**

437 We would like to thank the staff at Walailak University Institute of Research for their kind support
438 and technical assistance in this research.

439 **References**

- 440 1. Nour NM. Child marriage: a silent health and human rights issue. *Rev Obstet Gynecol.*
441 2009;2(1):51-6.
- 442 2. WHO. Child marriages-39 000 every day: More than 140 million girls will marry between 2011
443 and 2020. Geneva: WHO; 2013.
- 444 3. UNICEF. Child marriage around the world: UNICEF; 2020 [Available from:
445 <https://www.unicef.org/stories/child-marriage-around-world>].
- 446 4. Godha D, Hotchkiss DR, Gage AJ. Association between child marriage and reproductive health
447 outcomes and service utilization: a multi-country study from South Asia. *J Adolesc Health.*
448 2013;52(5):552-8.
- 449 5. Amin S, Asadullah MN, Hossain S, Wahhaj Z. Eradicating Child Marriage in the Commonwealth: Is
450 Investment in Girls' Education Sufficient? *The Round Table.* 2017;106(2):221-3.
- 451 6. Raj A, Saggurti N, Winter M, Labonte A, Decker MR, Balaiah D, et al. The effect of maternal child
452 marriage on morbidity and mortality of children under 5 in India: cross sectional study of a nationally
453 representative sample. *BMJ.* 2010;340:b4258.
- 454 7. Santhya KG. Early marriage and sexual and reproductive health vulnerabilities of young women:
455 a synthesis of recent evidence from developing countries. *Current Opinion in Obstetrics and Gynecology.*
456 2011;23(5):334-9.
- 457 8. Raj A, Boehmer U. Girl Child Marriage and Its Association With National Rates of HIV, Maternal
458 Health, and Infant Mortality Across 97 Countries. *Violence Against Women.* 2013;19(4):536-51.
- 459 9. Aychiluhm SB, Tesema AK, Tadesse AW. Early Marriage and Its Determinants among Married
460 Reproductive Age Group Women in Amhara Regional State, Ethiopia: A Multilevel Analysis. *Biomed Res*
461 *Int.* 2021;2021:1969721.
- 462 10. Clark S, Bruce J, Dude A. Protecting young women from HIV/AIDS: the case against child and
463 adolescent marriage. *Int Fam Plan Perspect.* 2006;32(2):79-88.
- 464 11. (ICRW) ICFRoW. *How to End Child Marriage: Action Strategies for Prevention and Protection.*
465 Washington, DC: ICRW; 2007.

- 466 12. Mathur S, Greene M, Malhotra A. Too young to wed: The lives, rights and health of young
467 married girls. Washington, DC: International Center for
468 Research on Women; 2003.
- 469 13. Dadras O, Nakayama T, Kihara M, Ono-Kihara M, Seyedalinaghi S, Dadras F. The prevalence and
470 associated factors of adverse pregnancy outcomes among Afghan women in Iran; Findings from
471 community-based survey. PLOS ONE. 2021;16(1):e0245007.
- 472 14. Fund UNP. The State of World Population 2014: The Power of 1.8 Billion. New York, NY: UNFPA;
473 2014.
- 474 15. Kenny L, Koshin H, Sulaiman M, Cislighi B. Adolescent-led marriage in Somaliland and Puntland:
475 A surprising interaction of agency and social norms. J Adolesc. 2019;72:101-11.
- 476 16. Dadras. O, S Chapman. R. Biomass Fuel Smoke and Stunting in Early Childhood: Finding from a
477 National Survey Nepal. Journal of Health Research. 2017;31:s7-s15.
- 478 17. Bradle S, Croft T, Fishel J, Westoff C. Revising Unmet Need for Family Planning. Calverton, MD:
479 ICF International; 2012.
- 480 18. Kamal SMM, Hassan CH. Child Marriage and Its Association With Adverse Reproductive
481 Outcomes for Women in Bangladesh. Asia Pacific Journal of Public Health. 2015;27(2):NP1492-NP506.
- 482 19. Raj A, Saggurti N, Balaiah D, Silverman JG. Prevalence of child marriage and its effect on fertility
483 and fertility-control outcomes of young women in India: a cross-sectional, observational study. Lancet.
484 2009;373(9678):1883-9.
- 485 20. Yaya S, Odusina EK, Bishwajit G. Prevalence of child marriage and its impact on fertility
486 outcomes in 34 sub-Saharan African countries. BMC International Health and Human Rights.
487 2019;19(1):33.
- 488 21. Budu E, Ahinkorah BO, Seidu A-A, Hagan JE, Agbemavi W, Frimpong JB, et al. Child Marriage and
489 Sexual Autonomy among Women in Sub-Saharan Africa: Evidence from 31 Demographic and Health
490 Surveys. International Journal of Environmental Research and Public Health. 2021;18(7):3754.
- 491 22. Parsons J, Edmeades J, Kes A, Petroni S, Sexton M, Wodon Q. Economic Impacts of Child
492 Marriage: A Review of the Literature. The Review of Faith & International Affairs. 2015;13(3):12-22.
- 493 23. Izugbara CO, Ngilangwa DP. Women, poverty and adverse maternal outcomes in Nairobi, Kenya.
494 BMC Women's Health. 2010;10(1):33.
- 495 24. Afifi M. Wealth Index association with gender issues and the reproductive health of Egyptian
496 women. Nursing & Health Sciences. 2009;11(1):29-36.
- 497 25. UNICEF. Child marriage in Afghanistan: changing the narrative. UNICEF; 2018.
- 498 26. Saleheen AAS, Afrin S, Kabir S, Habib MJ, Zinnia MA, Hossain MI, et al. Sociodemographic factors
499 and early marriage among women in Bangladesh, Ghana and Iraq: An illustration from Multiple Indicator
500 Cluster Survey. Heliyon. 2021;7(5):e07111.
- 501 27. Kidman R. Child marriage and intimate partner violence: a comparative study of 34 countries.
502 International Journal of Epidemiology. 2016;46(2):662-75.
- 503 28. The Partnership for Maternal NaCH. PMNCH Knowledge Summary #22 Reaching Child Brides.
504 WHO; 2012.
- 505 29. P. D. Marriage age and fertility dynamics in India. Calverton, MD: Macro International Inc; 2008.
- 506 30. Hampton T. Child marriage threatens girls' health. Jama. 2010;304(5):509-10.
- 507 31. Kamal SM. Decline in child marriage and changes in its effect on reproductive outcomes in
508 Bangladesh. J Health Popul Nutr. 2012;30(3):317-30.
- 509 32. Bearak J, Popinchalk A, Ganatra B, Moller A-B, Tunçalp Ö, Beavin C, et al. Unintended pregnancy
510 and abortion by income, region, and the legal status of abortion: estimates from a comprehensive
511 model for 1990–2019. The Lancet Global Health. 2020;8(9):e1152-e61.
- 512 33. Finer LB, Zolna MR. Unintended pregnancy in the United States: incidence and disparities, 2006.
513 Contraception. 2011;84(5):478-85.

- 514 34. Fund UNCs. *Fistula in Niamey, Niger*. New York: United Nations Children’s Fund; 1998.
- 515 35. Paul P, Chouhan P. Association between child marriage and utilization of maternal health care
516 services in India: Evidence from a nationally representative cross-sectional survey. *Midwifery*.
517 2019;75:66-71.
- 518 36. Sekine K, Carter DJ. The effect of child marriage on the utilization of maternal health care in
519 Nepal: A cross-sectional analysis of Demographic and Health Survey 2016. *PLOS ONE*.
520 2019;14(9):e0222643.
- 521 37. Nasrullah M, Zakar R, Krämer A. Effect of Child Marriage on Use of Maternal Health Care
522 Services in Pakistan. *Obstetrics & Gynecology*. 2013;122(3):517-24.
- 523 38. Raj A. When the mother is a child: the impact of child marriage on the health and human rights
524 of girls. *Arch Dis Child*. 2010;95(11):931-5.
- 525 39. Dadras O, Taghizade Z, Dadras F, Alizade L, Seyedalinaghi S, Ono-Kihara M, et al. “It is good, but I
526 can’t afford it ...” potential barriers to adequate prenatal care among Afghan women in Iran: a
527 qualitative study in South Tehran. *BMC Pregnancy and Childbirth*. 2020;20(1):274.
- 528 40. Hou X, Ma N. Empowering women: The effect of women’s decision-making power on
529 reproductive health services uptakedEvidence from Pakistan. Washington, DC: World Bank South Asia
530 Region, Human Development Unit; 2011.
- 531 41. Moore A, Singh S, Ram U, al. e. Adolescent marriage and childbearing in India: Current situation
532 and recent trends. New York: Guttmacher Institute; 2009.
- 533 42. Dadras O, Dadras F, Taghizade Z, Seyedalinaghi S, Ono-Kihara M, Kihara M, et al. Barriers and
534 associated factors for adequate antenatal care among Afghan women in Iran; findings from a
535 community-based survey. *BMC Pregnancy and Childbirth*. 2020;20(1):427.
- 536 43. Khan YP, Bhutta SZ, Munim S, Bhutta ZA. Maternal health and survival in Pakistan: issues and
537 options. *J Obstet Gynaecol Can*. 2009;31(10):920-9.
- 538 44. Ali TS, Krantz G, Gul R, Asad N, Johansson E, Mogren I. Gender roles and their influence on life
539 prospects for women in urban Karachi, Pakistan: a qualitative study. *Glob Health Action*. 2011;4:7448.
- 540 45. Women ICfRo. New insights on preventing child marriage: A global analysis of factors and
541 programs: ICRW; 2007 [Available from: [http://www.icrw.org/files/publications/New-Insights-](http://www.icrw.org/files/publications/New-Insights-onPreventing-Child-Marriage.pdf)
542 [onPreventing-Child-Marriage.pdf](http://www.icrw.org/files/publications/New-Insights-onPreventing-Child-Marriage.pdf).
543