Best practices for multi-ancestry, meta-analytic transcriptome-wide association studies: lessons

from the Global Biobank Meta-analysis Initiative

Arjun Bhattacharya*[1,2], Jibril B. Hirbo*[3,4], Dan Zhou[3,4], Wei Zhou[5,6,7], Jie Zheng[8], Masahiro Kanai[5,6,7,9,10],

the Global Biobank Meta-analysis Initiative, Mark J. Daly[5,6,7,11], Bogdan Pasaniuc[1,8,9†], Eric R.

Gamazon[3,4,14†], Nancy J. Cox[3,4†]

1. Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, University of
California, Los Angeles, CA, United States
2. Institute of Quantitative and Computational Biosciences, David Geffen School of Medicine, University
of California, Los Angeles, CA, United States
3. Department of Medicine, Division of Genetic Medicine, Vanderbilt University School of Medicine,
Nashville, TN, United States
4. Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Nashville, TN, United States
5. Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA
6. Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA
7. Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, Cambridge, MA
8. MRC Integrative Epidemiology Unit (IEU), Bristol Medical School, University of Bristol, Oakfield
House, Oakfield Grove, Bristol, BS8 2BN, UK
9. Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
10. Department of Statistical Genetics, Osaka University Graduate School of Medicine, Suita 565-0871,
Japan
11. Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
12. Department of Human Genetics, David Geffen School of Medicine, University of California, Los
Angeles, CA, United States
13. Department of Computational Medicine, David Geffen School of Medicine, University of California,
Los Angeles, CA, United States
14. MRC Epidemiology Unit, University of Cambridge, Cambridge, UK
*Corresponding authors: Arjun Bhattacharya (abtbhatt@ucla.edu), Jibril Hirbo (jibril.hirbo@vumc.org)
†These authors contributed equally to this manuscript.

SUMMARY

The Global Biobank Meta-analysis Initiative (GBMI), through its genetic and demographic diversity, provides a valuable opportunity to study population-wide and ancestry-specific genetic associations. However, with multiple ascertainment strategies and multi-ethnic study populations across biobanks, the GBMI provides a distinct set of challenges in implementing statistical genetics methods. Transcriptome-wide association studies (TWAS) are a popular tool to boost detection power for and provide biological context to genetic associations by integrating single nucleotide polymorphism to trait (SNP-trait) associations from genome-wide association studies (GWAS) with SNP-based predictive models of gene expression. TWAS presents unique challenges beyond GWAS, especially in a multi-biobank and meta-analytic setting like the GBMI. In this work, we present the GBMI TWAS pipeline, outlining practical considerations for ancestry and tissue specificity and meta-analytic strategies, as well as open challenges at every step of the framework. Our work provides a strong foundation for adding tissue-specific gene expression context to biobank-linked genetic association studies, allowing for ancestry-aware discovery to accelerate genomic medicine.

KEYWORDS

transcriptome-wide association study; meta-analysis; multi-ancestry genetic analysis; Global Biobank Meta-analysis Initiative

INTRODUCTION

Large population-based or clinical-case based biobanks are a key component of precision medicine efforts and provide opportunities for genetic and genomic research (Abul-Husn and Kenny, 2019). Biobanks offer context to deploy genome-wide associations (GWAS) at scale. Multi-biobank collaborations facilitate well-powered, multi-ethnic genetic research (Swede et al., 2007). In addition, such collaborations can accelerate the elucidation of the biological mechanisms that underlie diseases by in-silico longitudinal genetic studies and examination of pleiotropy.

A key challenge in GWAS is interpreting significant trait-associated loci and these loci to genes or epigenomic features (Gallagher and Chen-Plotkin, 2018; Wijmenga and Zhernakova, 2018). Viable options to add biological interpretation to our understanding of GWAS loci include colocalization (Giambartolomei et al., 2014, 2018; Gleason et al., 2020; He et al., 2013) or Mendelian randomization methods (Hauberg et al., 2017; Pavlides et al., 2016; Smith and Ebrahim, 2003). Another suite of tools include transcriptome-wide association studies (TWAS), which integrate GWAS with expression quantitative trait loci (eQTL) analyses to prioritize gene-trait associations (GTAs) with applications of mediation analysis (Gamazon et al., 2015; Gusev et al., 2016) or Mendelian randomization (Zhang et al., 2020). TWAS involves three general steps. First, per-gene predictive models of gene expression are trained in the eQTL dataset using genetic variants. Then, genetically-regulated expression (GReX) is imputed in the GWAS cohort with individual-level genotypes. Lastly, statistical associations between GReX and trait are estimated (Barbeira et al., 2018; Gamazon et al., 2015; Gusev et al., 2016). TWAS is also viable with GWAS summary statistics by estimating the test statistic of the TWAS association using a proper LD reference panel (Gusev et al., 2016). Generally, most TWAS methods predict expression using SNPs local to the gene within 1 Megabase of the gene body (Barbeira et al., 2018; Gamazon et al., 2015; Gusev et al., 2016; Hu et al., 2019; Nagpal et al., 2019; Parrish et al., 2022; Zhou et al., 2020). Recently, methods that include strong distal-eQTL signals have shown improved prediction and power to detect GTAs (Bhattacharya et al., 2021a; Luningham et al., 2020). Nonetheless, practical and statistical considerations to accurately prioritize GTAs through TWAS still require methodological improvement.

Along with those from traditional GWAS, TWAS introduces new challenges by incorporating gene expression (Wainberg et al., 2019) (**Figure 1A**). On the genetic level, as in GWAS, disentangling signals from complex LD structure, relatedness, and ancestry requires careful modeling considerations (Mbatchou et al., 2020; Zhou et al., 2018). Selection of LD reference is specifically important in multi-ancestry settings, like the GBMI, as LD structure across ancestry groups differs greatly(Shifman et al., 2003). Mismatched LD may lead to gene expression models with reduced predictive power, reduced power to detect GTAs, and increased false positives (Bhattacharya et al., 2020; Geoffroy et al., 2020; Keys et al., 2020). In addition, phenotype acquisition and aggregation are challenging, especially across

multiple biobanks with different healthcare, electronic health record, and case-control definitions.

However, compared to GWAS, a challenge specific to TWAS is the integration of gene expression with

GWAS signal. Not only is it an active topic of methodological research to choose an optimal set of genes

and tissues that best explains the phenotype association at a given genetic locus, the role of context-

specific expression is still being evaluated in trait associations. Dynamic differences in bulk tissue

expression from cell-type- or cell-state-specificity can give additional granularity to gene-trait associations.

The impact of these challenges in a meta-analytic framework has not been previously explored.

Here, we outline a framework for analyzing trans-ancestry, meta-analytic GWAS across multiple biobanks

with TWAS. We review and explore practical considerations for all three steps (**Figure 1B**): ancestry

specificity of expression models and LD reference panels, meta-analytic techniques for detection of

GTAs, and follow-up tests and analyses for biological context. Our framework can be applied to various

phenotypes to study population-wide and ancestry-specific genetic associations mediated by tissue-

specific expression.

RESULTS

Expression models are not portable across ancestry groups

The diversity represented in the GBMI enables uniquely well-powered studies to detect genetic

associations in non-European populations. However, optimal TWAS requires ancestry-matched training

datasets of genetic and tissue-specific gene expression data, which are still lacking in non-European

samples. As Cao *et al* points out (Cao et al., 2021), statistical power to detect GTAs in TWAS is

dependent on expression heritability and the ability of the predictive expression model to recapitulate that

heritable expression in the external GWAS panel. Accordingly, training expression models that perform

well in all ancestry populations is necessary to ensure that discoveries made through TWAS are not

restricted to European populations. For the first GBMI TWAS, we restrict analysis to populations of

European ancestry, due to small sample sizes of non-European ancestry samples (Aguet et al., 2020).

However, as sample sizes for eQTL datasets in non-European populations increase, the TWAS pipeline

will include expression models for these understudied and underserved populations (**STAR Methods**).

Here, we illustrate some challenges in building these expression models across ancestry groups.

We considered the 5 tissues in GTEx with at least 70 samples from both European (EUR) and African

(AFR) ancestry: subcutaneous adipose (abbreviated ADIP, $N = 71$ samples of AFR ancestry and 492

samples of EUR ancestry), tibial artery (ARTERY, $N = 76$ and 489), skeletal muscle (MUSC, $N = 86$ and

602), sun exposed lower leg skin (SKIN, $N = 73$ and 518), and whole blood (BLOOD, $N = 80$ and 574).

For genes with significantly heritable expression in both EUR and AFR GTEx samples (restricted

maximum likelihood-based estimate of heritability > 0 with nominal $P < 0.01$), we trained EUR- and AFR-

specific models using elastic net regularized regression (Friedman et al., 2010) and imputed expression

into the aligned (i.e., training and imputation samples have similar ancestries) and misaligned (i.e.,

training and imputation sample have different ancestries). For context, we also built ancestry-unaware

models, where EUR and AFR samples were pooled together. We calculated predictive performance with

adjusted R^2 to account for sample size, using leave-one-out CV when measuring predicting performance

in an aligned imputation sample (**STAR Methods**).

Across these tissues, models trained in EUR samples performed, on average, 4 times worse (differences

of 0.03-0.04 in median adjusted R^2) in AFR samples compared to models trained in AFR samples (**Figure**

2A, Tables S1-S3), with more than 80% of gene models with stronger performance if trained in AFR

samples. Similar trends hold for ancestry-specific models imputed into down-sampled EUR imputation

samples (**Figure S1-S2, Table S1-S3**), consistent with previous simulation and real-world studies

(Bhattacharya et al., 2020; Keys et al., 2020); here, we considered a randomly selected EUR imputation

sample with equal sample size to that of the AFR sample in the same tissue. In fact, we observed that

ancestry-specific models imputed into a sample with aligned ancestry showed larger predictive R^2 than

ancestry-unaware (individuals of EUR and AFR ancestry in the training sample) models imputed into the

same sample (**Figure 2B**, **Table S4**), despite generally increased sample sizes. This observation also

holds if we further increase the sample size of the training sample by including individuals of other

ancestries (Asian, American Indian, and Unknown ancestries) into the training sample (**Figure S3**). This

observation emphasizes the need for ancestry matching in gene prediction from genetic data and greater recruitment of non-European ancestry patients in eQTL studies.

In these analyses, one reason ancestry-unaware models may perform poorly in AFR samples is due to differences in minor allele frequency (MAF) of highly predictive SNPs between EUR and AFR ancestry populations. It is important to note that this discrepancy is not generally specific to any one ancestry; rather, ancestry imbalance in the training or reference datasets may lead to poor portability of genetic models due to differences in allele frequency. To account for common SNPs in both AFR or EUR ancestry populations, we additionally trained ancestry-unaware and ancestry-specific models using SNPs with minor allele frequency (MAF) exceeding various thresholds in both AFR and EUR samples. Excluding SNPs with MAF < 0.01 improved predictive performance of ancestry-unaware models across all tissues (**Figure S4, Table S5**). However, the gap in predictive performance between ancestry-specific and ancestry-unaware models did not decrease when the MAF cutoff was increased (**Figure 2B, Table S4**). This observation may reflect that dropping ancestry-specific rarer SNPs ignores variants with large ancestry-specific effects on gene expression. Additionally, excluding rare ancestry-specific SNPs does not address the differences in LD across the EUR and AFR samples which leads to different regularization paths and, hence, SNP-gene weights. Addressing the trans-ancestry portability of expression models remains an open study direction; methodology that borrows information from functional annotations or across different cell-type- or cell-state-specific contexts may bridge this gap in predictive performance, similar to recent developments in polygenic risk score prediction for complex traits (Amariuta et al., 2020; Márquez-Luna et al., 2020).

Meta-analytic strategies must be ancestry-aware

Another critical consideration for the GBMI involves meta-analysis while using GWAS summary statistics. TWAS estimates the association between GReX and the phenotype by weighting the standardized SNP-trait effect sizes from GWAS summary statistics by SNP-gene weights from the expression models. To account for the correlation between SNPs, an external LD reference panel, like the 1000Genomes Project (Auton et al., 2015), is used to estimate the standard error of the TWAS association. Accordingly, the

accuracy of this reference panel to the LD structure in the GWAS cohort dictates how aligned the

summary-statistics based TWAS association is to the TWAS association from direct imputation into

individual-level genotypes in the GWAS cohort. Ideally, in-sample LD will give the best estimate of the

TWAS standard error, but several biobanks do not provide this information under their specific genetic

data sharing and privacy policies. Even departures in LD across subgroups of European ancestry

populations may influence the standard error estimate. In addition, as the estimates of SNP-gene weights

are influenced by the LD in the eQTL panel, differences in LD between the eQTL and GWAS panel will

also affect the TWAS effect size.

As LD structure greatly differs across ancestry groups (Shifman et al., 2003), pooling ancestry groups in

TWAS may lead to reduced power. We conducted TWAS for asthma risk using ancestry-unaware and

EUR- and AFR-specific models of whole blood expression (4,782 genes with heritable expression at

nominal P < 0.01 and models with cross-validation R^2 > 0.01 with nominal P < 0.05, trained via elastic net

regression). Ancestry-specific TWAS Z-scores across EUR and AFR ancestry groups were not strongly

correlated ($r = 0.11$), potentially due to differences in sample size and eQTL and GWAS architecture

(**Figure 3A, Figure S5-S6**) (Shang et al., 2020; Wyss et al., 2018). In fact, we detected only two genes

across both EUR and AFR with P < 2.5×10^{-6}. One of these genes, *DFFA*, has been implicated with

asthma risk through GWAS and colocalization in EUR (Vicente et al., 2017). However, the TWAS

associations across EUR and AFR were in opposite directions using blood tissue. In the other 4 tissues

explored, *DFFA* TWAS associations did not reach transcriptome-wide significance but effect directions

were generally concordant (**Figure S7**). In blood, lead local-eQTLs (within 1 Megabase) of *DFFA* show

are in opposite directions, though only nominally significant at P < 0.05 (**Figure S8**). Although they are

within 60 kilobases, the lead eQTLs for DFFA across AFR (rs263526) and EUR (rs903916) are not in LD

(R^2 = 3 x 10^{-4} in AFR, 0.0072 in EUR). The GWAS effect sizes of SNPs local to *DFFA* do not show large

deviations in effect direction and are only nominally significant, as well (**Figure S8**). These differences in

TWAS associations across ancestry motivate careful consideration of meta-analytic strategy to avoid

biasing cross-ancestry associations towards cohorts with larger sample sizes, which still tend to be

predominantly of EUR ancestry.

We investigated 5 different meta-analytic strategies empirically: meta-analyzing across ancestry-specific,

per-biobank GWAS summary statistics using (1) inverse-variance weighting (IVW) and (2) sample-size

weighting (SSW), meta-analyzing across ancestry-specific meta-analyzed GWAS summary statistics

using (3) IVW and (4) SSW, and (5) TWAS using ancestry-unaware models and meta-analyzed GWAS

summary statistics across EUR and AFR ancestry groups (**STAR Methods**). QQ-plots in **Figure 3B** show

earlier departure of Z-scores from the QQ-line for SSW meta-analyzed Z-scores and the ancestry-

unaware strategy, suggesting inflation. This observation is supported with estimates of test statistic bias

and inflation using an empirical Bayes method, bacon (van Iterson et al., 2017), which show the largest

estimated bias and inflation for these SSW and ancestry-unaware methods. IVW strategies show similar

levels of inflation, with IVW meta-analysis across ancestry-specific meta-analyzed GWAS summary

statistics showing minimal bias (**Figure S9**). These results align with intuition – that the more naïve SSW

meta-analysis and ancestry-unaware methods bias towards EUR cohorts, which have the larger sample

sizes, whereas Z-scores from the IVW methods showed positive correlations with Z-scores from AFR

cohorts (**Figure S6**).

However, it is unclear whether ancestry-specific IVW meta-analysis to the per-biobank level is necessary.

As shown in **Figure S10**, Z-scores from these two IVW methods are moderately positively correlated ($r =$

0.51 across 4,152 Z-scores), with this correlation increasing when we consider genes with nominally

significant Z-scores for both strategies ($r = 0.70$ across 564 tests). We observed that top associations

across these IVW meta-analyses often had high degrees of heterogeneity in effect size across biobanks,

as measured by the Higgins-Thompson I^2 statistic (**Figure 3C, Figure S11**) (Higgins and Thompson,

2002). One gene, *A1BG*, that showed directionally concordant transcriptome-wide significant associations

across both IVW strategies had a large degree of heterogeneity in the underlying participating cohorts (I^2

$= 0.77$). In fact, the cross-biobank heterogeneity is often larger than the cross-ancestry heterogeneity for

TWAS associations of *A1BG*. Interestingly, *ZNF665*, another gene with directionally concordant

associations across both IVW strategies showed a low degree of heterogeneity in the per-biobank effect

sizes ($I^2 = 0.26$). However, genes with discordant associations across IVW strategies showed varied

patterns. Two illustrative examples are *MLNR* and *MYOZ3*, both with large degrees of test statistic

heterogeneity (I^2 = 0.91 and 0.82, respectively). Across the two IVW strategies, effect sizes are in

opposite directions, possibly due to large standard error differences across the ancestry-specific per-

biobank associations. A thorough investigation of the power and false discovery rates of these meta-

analysis strategies through simulations is necessary. More sophisticated methods (Hedges and Vevea,

1998; Lee et al., 2017; Shi and Lee, 2016) that can properly incorporate the per-biobank uncertainty into

meta-analyzed TWAS associations must be explored to increase power and properly leverage the large

sample sizes of the GBMI.

In addition to considerations for TWAS in trans-ethnic populations, analyzing genetic data from individuals

of admixed ancestry is also an open area of study. For example, in this analysis, we have used the

1000Genomes AFR LD reference panel as an estimate of the LD for the AFR-ancestry samples from

each biobank. However, most of these AFR-ancestry populations are of admixed ancestry (e.g, African

Americans or African British). A single LD reference panel of AFR-ancestry may not reflect the genetic

diversity in these admixed populations of AFR and EUR ancestries from around the world (Baharian et

al., 2016). As Zhong et al highlights, in multiethnic and admixed populations, using local ancestry

estimates aids in better characterization of heritability of complex traits and more accurate mapping of

genetic associations, especially eQTLs (Zhong et al., 2019). Accordingly, incorporating local ancestry

estimates into both expression model step and the association testing step of TWAS may lead to

increased power and should be explored.

Follow-up tests provide biological and clinical context to TWAS GTAs

TWAS GTAs identified using GWAS summary statistics are subject to several factors that may lead to

false positives. We implement several follow-up tests to provide context to TWAS-identified GTAs. First, a

TWAS GTA could attain transcriptome-wide significance due to only the strong SNP-trait associations

from the underlying GWAS. To quantify the significance of the GTA conditional on the SNP-trait effects at

the locus, we perform a permutation test by permuting the SNP-gene weights from the expression model

to generate a null distribution (**STAR Methods**). Comparing the original TWAS Z-score to this null

distribution assesses how much signal is added by the expression given the specific GWAS architecture

of the locus. As Gusev *et al* point out, this permutation test is highly conservative and intended to

prioritize only associations already significant in the standard TWAS GTA detection (Gusev et al., 2016).

Next, gene expression models for genes in adjacent genomic windows may be built from overlapping

SNPs or SNPs in strong LD. When TWAS detects GTAs in overlapping genomic regions, we apply

Bayesian probabilistic fine-mapping using FOCUS (Mancuso et al., 2019)to estimate a 90% credible set

of genes to explain the observed association signal in a given tissue (**STAR Methods**). However, the

current iteration of FOCUS has limitations. Priors for the correlation matrix between GReX of overlapping

genes are dependent on SNP LD reference panels. Thus, fine-mapping in trans-ancestry settings is

difficult, though recent machinery has been added to FOCUS to account for differences in genetic

architecture across the study sample (Gopalan and Lu et al, in preparation). Another challenge for gene-

level fine-mapping in multi-tissue TWAS is distinguishing between overlapping signals across tissues.

Primarily due to cross-cell-type variation in expression levels and eQTL architecture, TWAS may prioritize

genes in multiple tissues that are overrepresented by the same underlying causal cell-types (Wainberg et

al., 2019). This multi-tissue gene prioritization extends to fine-mapping overlapping TWAS signals across

tissue, as priors for FOCUS are not tissue-dependent. An interesting future direction involves careful

tissue-specific prior elicitation in TWAS fine-mapping – extracting posterior signal that is biologically

consistent and meaningful without allowing the prior to dominate.

The GBMI TWAS model incorporates gene expression models using MOSTWAS, a TWAS extension that

prioritizes distal-eQTLs by testing their mediation effect through local molecular features (**STAR**

Methods). For genes with models trained with MOSTWAS and associated with the trait at transcriptome-

wide significance, we test the additional association signal from the distal-SNPs using an added-last test,

analogous to a group-added-last test in linear regression (Bhattacharya et al., 2021a). This test also

prioritizes sets of genomic or epigenomic features that mediate the predicted distal-eQTLs for subsequent

study of upstream, tissue-specific regulation of GTAs. In one application of MOSTWAS, one prioritized

functional hypothesis was experimentally validated *in vitro* (Bhattacharya et al., 2021b). As distal-eQTLs

are more likely to be tissue- or cell-type-specific (Yang et al., 2017), the association signal from these

distal-eQTLs could also be leveraged in cross-tissue fine-mapping strategies.

Lastly, TWAS suffers from severely reduced power and inflated false positives in the presence of SNP

pleiotropy, where the genetic variants in the gene expression model affect the trait, independent of gene

expression (Veturi and Ritchie, 2018). We encourage estimating the degree of and accounting for SNP

pleiotropy using LDA-MR-Egger (Barfield et al., 2018) or PMR-Egger (Yuan et al., 2020), especially in

settings with individual-level GWAS genotypes. Applications for these methods using GWAS summary

statistics reveals some inflation of standard errors (Zhu et al., 2021), suggesting the need for further

evaluation and development of summary statistics-based methods to account for SNP pleiotropy in

TWAS GTA detection.

Biobanks enable GReX-PheWAS for biological context

Biobanks aggregated in the GBMI provide a rich catalog of phenotypes for analysis, with phenotype

codes (phecodes) aggregated from ICD codes classified into clinically relevant categories (Wei et al.,

2017). This phenotype catalog enables Phenome-Wide Association Studies (PheWAS) as a complement

to GWAS by both replicating GWAS associations and providing a larger set of traits associations with

GWAS variants. To follow-up on novel TWAS-prioritized genes, we may expand the PheWAS framework

to the tissue-specific GReX level in a similarly complementary analysis: GReX-level Phenome-Wide

Association Study (GReX-PheWAS), as previously deployed in biobank settings and similar to the

PredixVU database (Pathak et al., 2020; Unlu et al., 2019, 2020). Not only do these analyses replicate

and detect new TWAS associations, but they can also point to groups of phenotypes that show

enrichments for trait-associations for the gene of interest.

We briefly illustrate an example of GReX-PheWAS using 3 genes (**Figure 4, Figure S12-S13, Table S6**):

TAF7, a novel gene in our TWAS, and *ILRAP18* and *TMEM258*, two genes previously implicated through

GWAS (Johansson et al., 2019; Portelli et al., 2020; Reijmerink et al., 2008, 2010; Zhu et al., 2020).

These genes were prioritized from European-specific TWAS for asthma risk from Zhou *et al* (Zhou et al.,

2021) using lung tissue expression (101,311 cases and 1,118,682 controls): *TAF7* (MOSTWAS model),

IL18RAP (JTI model), and *TMEM258* (JTI model). European-specific TWAS meta-analysis for asthma

detected a negative association with *TAF7* cis-GReX, a gene that did not intersect a GWAS-significant

locus. In TWAS follow-up tests, *TAF7* passed permutation testing and was estimated in the 90% credible

set at the genomic locus via FOCUS with posterior inclusion probability 1. *TAF7* encodes a component of

the TFIID protein complex, which binds to the TATA box in class II promoters and recruits RNA

polymerase II and other factors (Bhattacharya et al., 2014). As the clinically-relevant associations for

TAF7 lung GReX are not characterized, we employed GReX-PheWAS in UKBB European-ancestry

GWAS summary statistics across 731 traits and diseases with sample sizes greater than 100,000,

grouped into 9 categories (**Figure 4, STAR Methods**). We see enrichments for phenotypes of the

hematopoietic and musculoskeletal groups (**Figure 4A**) with hypothyroidism and chronic laryngitis as the

top phenotype associations (**Figure 4B, Table S6**). These phenotypes include multiple inflammations of

organs (e.g., laryngitis, osteitis, meningitis, inflammations of the digestive and respiratory system, etc).

We also detected several associations with related respiratory diseases and traits. Similarly, for the two

previously implicated genes, we find enrichments for respiratory and hematopoietic GTAs for *ILRAP18*

and across multiple categories for *TMEM258*, consistent with the categorized functions and associations

of these genes (**Figure S12-S13, Table S6**). The utility of GBMI's robust roster of phenotypes enables

GReX-PheWAS to add biological and clinical context to novel TWAS associations.

GReX-PheWAS, despite its utility, shares the challenges of PheWAS. Phenotypes within and across

groups may be correlated, leading to a series of dependent tests. Even divergent phenotypes may be

correlated, either clinically or biologically. Thus, simple adjustments of multiple testing burden may not be

appropriate, and methods that account for correlation between phenotypes, like permutation tests, may

be more applicable (Hebbring, 2014; Korthauer et al., 2019; Stevens et al., 2017; Wei et al., 2017). In

addition, covariate adjustments in expression models built for disease-specific analyses may not be

generalizable for multiple phenotypes. Most population-based clinical biobanks lack comprehensive

clinical and lifestyle information of the individuals. Phenotyping information is typically incomplete, mostly

due to gaps in electronic health records. Phenotype groupings may also be deceptive: as most biobanks

follow ICD coding that groups traits and diseases by body systems, GReX-PheWAS enrichments for a

given group may not reflect shared genetic pathways across body systems. In addition, case-control

selection may not be optimal due to differences in exclusion criteria (McGuirl et al., 2020). Lastly,

phenotype acquisition and aggregation across multiple biobanks is challenging, with different healthcare,

electronic health record, and case-control definition assignments. Despite these limitations in phenotype

acquisition, recent methods focusing on identifying shared genetic architecture among multiple

phenotypes (McGuirl et al., 2020) in a phenome-wide approach highlight the advantages of GReX-

PheWAS.

DISCUSSION

Here, we provide a framework for TWAS in a multi-biobank setting across many ancestry groups. Our

work outlines several methodological gaps that should be addressed in the future: (1) training expression

models that are portable across ancestry groups, (2) limiting false discovery in TWAS by properly

modeling differences in LD across ancestry groups, (3) incorporating uncertainty within and heterogeneity

across biobanks to boost TWAS meta-analytic power, and (4) contextualizing TWAS GTAs through

follow-up testing, probabilistic fine-mapping across ancestry groups and expression contexts, and GReX-

PheWAS. Along with the discussed issues with current TWAS methodology, tissue-specific expression

may not provide sufficient granularity needed to discover trait-relevant biological mechanisms. Recent

methods that study the mediation of the SNP-trait relationship by cell-type heterogeneity show that cell-

types are influenced by genetics and predict complex traits (Liu et al., 2021). In turn, single-cell eQTL

datasets can be integrated with GWAS to identify cell-type- or cell-state-specific expression pathways that

are health- or disease-related. Incorporating single cell expression data into a predictive model will

require more sophisticated statistical methodology that relies on modeling cell identity as a spectrum,

rather than a categorical definition (Burkhardt et al., 2021; Verma and Engelhardt, 2020). Furthermore, a

multi-omic approach that incorporates functional data with TWAS may better model the flow of biological

information in a biologically interpretable fashion (Baca et al., 2021; Bhattacharya et al., 2021a), with

Zhao et al, in preparation.

Despite the limitations of this suite of methods, TWAS continues to be a useful tool for interpreting GWAS

associations and independently discovering genetic associations mediated by gene expression. There is

a severe need for an increase in reference eQTL data from individuals of non-European ancestry at parity

with those of European-ancestry individuals. Moreover, more sophisticated integrative computational and

experimental tools to complement improved TWAS and GWAS to understand the biology underlying

health and disease need to be developed.

FUNDING

BP was partially supported by NIH awards R01 HG009120, R01 MH115676, R01 CA251555, R01

AI153827, R01 HG006399, R01 CA244670, U01 HG011715. ERG is supported by the National Institutes

of Health (NIH) grants: NHGRI R35HG010718, NHGRI R01HG011138, NIA AG068026, and NIGMS

R01GM140287. NJC is supported by U01HG009086.

DATA AND CODE AVAILABILITY

The all-biobank and ancestry-specific GWAS summary statistics are publicly available for downloading at

https://www.globalbiobankmeta.org/resources and browsed at the PheWeb Browser

http://results.globalbiobankmeta.org/. 1000 Genome Phase 3 data can be accessed at

ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/1000_genomes_project/data. MOSTWAS can

be accessed from https://github.com/bhattacharya-a-bt/MOSTWAS, and JTI can be accessed from

https://github.com/gamazonlab/MR-JTI. Sample scripts for this manuscript are available at

https://github.com/bhattacharya-a-bt/gbmi_twas.

ACKNOWLEDGEMENTS

We thank Nicholas Mancuso, Michael Love, and Achal Patel for their thoughtful discussion during the

research process. We would like to thank the organizing committee of the International Common Disease

Alliance for intellectual contributions on the set up of the GBMI as a nascent activity to the larger

effort. We would like to thank Daniel King from the Hail team and Sam Bryant from the Stanley Center

Data Management team at the Broad Institute for helping with the Google bucket set up and data

sharing, and Bethany Klunder from the University of Michigan Medical school for helping with the

paper submission.

AUTHOR CONTRIBUTIONS

397 Conceptualization: AB, JH; Methodology: AB, JH, DZ, ERG, BP, NJC; Software: AB, JH, DZ, ERG, BP,

398 NJC; Validation: AB, JH, DZ; Formal analysis: AB, JH; Investigation: all authors; Resources: all authors;

399 Data curation: WZ, MK; Writing - original draft: AB, JH; Writing - Review & Editing: all authors;

400 Visualization: AB, JH; Supervision: EG, BP, NJC; Project administration: AB, JH, ERG, BP, NJC; Funding

401 acquisition: all authors

DECLARATION OF INTERESTS

FIGURE LEGENDS

Figure 1: *Overview of GBMI transcriptome-wide association study (TWAS) pipeline with challenges at*

every data level. **(A)** Each level of data in TWAS introduces a unique set of challenges: (1) genetics data

include confounding from genetic ancestry, population structure and relatedness, and complex linkage

disequilibrium patterns, (2) gene expression data introduces context-specific factors, such as tissue-, cell-

type-, or cell-state-specific expression, and (3) phenotypic data, especially in the meta-analyses of

multiple biobanks, involve challenges in acquiring and aggregating phenotypes, properly defining controls

for phenotypes, and ascertainment and selection bias from non-random sampling. **(B)** An overview of the

GBMI TWAS pipeline: (1) JTI and MOSTWAS for model training, (2) inverse-variance weighted meta-

analysis using per-biobank, per-ancestry group TWAS summary statistics, and (3) various follow-up tests,

including conditional or permutation tests, distal-SNPs added last test, probabilistic fine-mapping using

FOCUS, and tests for SNP horizontal pleiotropy. Dotted lines represent associations that are tested in the

TWAS pipeline, while the solid lines represent a link built through predictive modeling.

Figure 2: *Comparison of predictive performance of genetic models of expression across ancestry.* **(A)** Distribution of difference in adjusted R^2 (Y-axis) when predicting expression in the AFR imputation sample between models trained in EUR and in AFR training samples across tissue (X-axis). **(B)** Distribution of difference in adjusted R^2 between ancestry-specific and ancestry-unaware models imputing into EUR (left) and AFR (right) samples.

Figure 3: *Comparison of meta-analytic strategies for multi-biobank, trans-ancestry TWAS.* **(A)** Scatterplot of per-ancestry meta-analyzed TWAS scores across EUR (X-axis) and AFR ancestry (Y-axis). The dotted horizontal and vertical lines indicate $P < 2.5 \times 10^{-6}$ with a diagonal line for reference. Points are colored based on which ancestry population the TWAS association meets $P < 2.5 \times 10^{-6}$. **(B)** QQ-plot of TWAS Z-scores, colored by meta-analytic strategies. Per ancestry refers to TWAS meta-analysis across meta-analyzed ancestry-specific GWAS summary statistics. Per bank/per ancestry refers to TWAS meta-analysis using all biobank- and ancestry-specific GWAS summary statistics. **(C)** Effect sizes and Bonferroni-corrected confidence intervals (CIs) for TWAS associations across 17 individual biobanks (stratified by ancestry group with EUR in green and AFR in red) and 2 IVW meta-analysis strategies (in yellow) for 5 representative genes. The Higgins-Thompson I^2 statistic for heterogeneity is provided, with the dotted line showing the null.

Figure 4: *GReX-PheWAS for categorizing phenome-wide associations for TAF7 genetically-regulated expression in UKBB.* **(A)** Boxplots of -$\log_{10}$ Benjamini-Hochberg FDR-adjusted P-values of GTAs across 9 phenotype groups. The dotted grey line showed FDR-adjusted P = 0.05. **(B)** Miami plot of TWAS Z-scores (Y-axis) across phenotypes, colored by phecode group. The dotted grey line shows the significance threshold for Benjamini-Hochberg FDR correction and phenotypes are labelled if the association passes Bonferroni correction.

STAR METHODS

We first outline the steps of the TWAS pipeline employed for phenotype available for analysis in the GBMI. Then, we provide details for the analyses presented in **Results**.

The GBMI TWAS Pipeline

Training expression models from genetics

Tissue-specific expression models trained with reference data from the Genotype-Tissue Expression

Project (GTEx) v8 (Aguet et al., 2020) are built using two methods: (1) Joint-Tissue Imputation (JTI),

which leverages shared genetic *cis*-regulation across tissues (Zhou et al., 2020), and (2) MOSTWAS,

which prioritizes tissue-specific distal-SNPs through rigorous mediation analysis to account for additional

expression heritability (Bhattacharya et al., 2021a). Genes with significantly positive expression

heritability (nominal $P < 0.05$) and five-fold cross-validation (CV) adjusted $R^2 \geq 0.01$ with $P < 0.05$ are

considered for TWAS. Ancestry-specific models are trained, excluding SNPs with MAF < 0.01 and

deviated from Hardy-Weinberg at P < 10^{-5} across all 838 GTEx samples. The first iteration of the GBMI

TWAS pipeline focuses on EUR-ancestry models due to larger sample sizes. However, as sample sizes

for other ancestry groups increase, this pipeline can be adapted for these currently underrepresented

ancestries. In addition, models from other data sources using other methods can be incorporated in

subsequent steps.

Hypothesis tests for TWAS

To test for an association between tissue-specific GReX of a gene and a trait of interest, GWAS summary

statistics are integrated with these expression models. For the EUR-specific TWAS, we use EUR-specific

meta-analyzed GWAS summary statistics across all biobanks. JTI and MOSTWAS use two different

approaches to test for a GTA. For MR-JTI, the posterior predictive distribution of GReX is estimated, and

multiple-instrumental-variable causal inference is used to estimate the GTA, controlling for overall

heterogeneity (Zhou et al., 2020). For MOSTWAS, a weighted burden test is constructed, as in FUSION

(Bhattacharya et al., 2021a; Gusev et al., 2016; Pasaniuc et al., 2014). Both of these methods require a

LD reference panel; the GTEx LD matrix is used as a reference. Taken together, these methods provide

effect sizes, standard errors, Z-scores (effect sizes standardized by standard error), and P-values for

GTAs. A GTA is transcriptome-wide significant using a Bonferroni correction across all tests run. The

number of tests run is equal to the sum of the number of significant gene models across all tissues.

Follow-up tests and analyses are then run to provide context to the TWAS GTAs. A permutation test is

run by shuffling the SNP-gene weights 1,000 times and determining the TWAS Z-score at each

permutation, generating a null distribution. The original TWAS Z-score is compared to this null distribution

to generate a permutation P-value; Benjamini-Hochberg FDR correction is used to account for multiple

testing burden here. This test examines whether the SNP-gene relationship provides more information

than just the SNP-trait association. Next, for MOSTWAS, the distal-SNPs added-last test is run to

measure the association from distal-SNPs in the expression models, conditional on the association from

local-SNPs (Bhattacharya et al., 2021a). This test prioritizes sets of mediating molecular features for the

SNP-gene relationship with significant effects on the trait. Lastly, for genes whose models are built using

SNPs from overlapping genomic regions, probabilistic fine-mapping via FOCUS (default parameters and

priors) is employed to determine a 90% credible set of genes that explain the gene-level association

signal at the locus (Mancuso et al., 2019). FOCUS also outputs posterior inclusion probabilities for each

gene in the 90% credible set.

Analysis of ancestry-specific and -unaware models

To show the utility of ancestry-specific models, we train EUR- and AFR-specific models using elastic net

regression for 5 tissues with more than 70 samples from AFR ancestry patients: subcutaneous adipose

($N = 492$ EUR, $N = 71$ AFR), tibial artery ($N = 489$ EUR, $N = 76$ AFR), skeletal muscle ($N = 602$ EUR,

$N = 86$ AFR), sun exposed lower leg skin ($N = 518$ EUR, $N = 73$ AFR), and whole blood ($N = 574$ EUR,

$N = 80$ AFR). To balance sample sizes in the imputation sample, we down-sampled the EUR ancestry

imputation sample to match the AFR imputation sample. We consider only genes with positive expression

heritability in both EUR and AFR training samples (Yang et al., 2011). We also build ancestry-unaware

models, where genotypes for EUR and AFR samples are pooled together in the training sample. We

calculate predictive performance in aligned and misaligned imputation samples based on ancestry; the

aligned imputation sample is one with ancestry that predominantly matches the ancestry of the training

sample. Predictive performance is measured with adjusted R^2 to account for sample size, using an

appropriate linear model between predicted and observed expression. For imputation samples that are

used in training (aligned imputation panel), we use leave-one-out CV when measuring predictive

performance. Lastly, when imputing into AFR and EUR samples using the ancestry-unaware models, we

use leave-one-out CV, as well, but only cross-validating over the AFR or EUR samples, respectively.

Comparison of meta-analytic strategies

We compared 5 different meta-analytic strategies empirically: meta-analyzing across ancestry-specific,

per-biobank GWAS summary statistics using (1) inverse-variance weighting (IVW) and (2) sample-size

weighting (SSW), meta-analyzing across ancestry-specific meta-analyzed GWAS summary statistics

using (3) IVW and (4) SSW, and (5) TWAS using ancestry-unaware models into meta-analyzed GWAS

summary statistics across EUR and AFR ancestry groups. First, we consider three different sets of

GWAS summary statistics: biobank- and ancestry-specific summary statistics, ancestry-specific summary

statistics meta-analyzed across all biobanks, and summary statistics meta-analyzed across biobanks and

ancestry groups. In two former settings, for biobank i and a given gene, we generate $\beta_{TWAS,i}$, the TWAS

effect size, and $SE_{TWAS,i}$, the corresponding standard error. Given B different biobanks, the IVW TWAS Z-

score, $Z_{TWAS,IVW}$, is calculated as:

$$Z_{TWAS,IVW} = \frac{\left(\frac{\sum_{i=1}^{B} \beta_{TWAS,i}/SE_{TWAS,i}}{\sum_{i=1}^{B} SE_{TWAS,i}^{-1}}\right)}{\left(\sum_{i=1}^{B} SE_{TWAS,i}\right)^{1/2}}.$$

523 With $Z_{TWAS,i} = \beta_{TWAS,i}/SE_{TWAS,i}$ and N_i as the sample size of the ith biobank (or pooled sample size

524 across all ancestry-specific biobank summary statistics), the SSW TWAS Z-score, $Z_{TWAS,SSW}$, is calculated

525 as:

$$Z_{TWAS,SSW} = \frac{\sum_{i=1}^{B} N_i Z_{TWAS,i}}{\left(\sum_{i=1}^{B} N_i^2\right)^{1/2}}.$$

529 For the ancestry-unaware TWAS, we use ancestry-unaware elastic net regression models and integrate

530 with GWAS summary statistics meta-analyzed across all ancestry groups and biobanks.

GReX-level phenome-wide association studies (GReX-PheWAS)

Transcriptome-wide significant genes are further prioritized by performing GReX-PheWAS to categorize

associations across a broad spectrum of phenotypes. Using UKBB summary statistics from European

ancestry patients (Bycroft et al., 2018), we tested for GTAs for 731 phenotypes grouped into 9 categories:

dermatologic, digestive, endocrine/metabolic, genitourinary, hematopoietic, musculoskeletal, neoplasms,

neurological, and respiratory. Here, we illustrate GReX-PheWAS using three genes from the European-

specific TWAS for asthma risk using lung tissue expression: *TAF7* (MOSTWAS model), *IL18RAP* (JTI

model), and *TMEM258* (JTI model). A phenome-wide significant association was defined via Bonferroni

correction (P $< \frac{0.05}{3 \times 731} = 2.28 \times 10^{-5}$).

SUPPLEMENTAL INFORMATION

Table S1: *Difference in adjusted R^2 between models trained in aligned and misaligned ancestry samples*

as the ancestry of the imputation sample.

Table S2: *R^2 of ancestry-specific models imputed into EUR imputation sample (training:imputation)*

Table S3: *R^2 of ancestry-specific models imputed into AFR imputation sample (training:imputation)*

Table S4: *Difference in R^2 between ancestry-specific and ancestry-unaware models across MAF*

Table S5: *Cross-validation R^2 of ancestry-unaware models across MAF threshold*

Table S6: *GReX-PheWAS results for 3 representative genes from asthma meta-analytic TWAS in*

Europeans in GBMI that meet Bonferroni correction (P < 0.05/731)

Figure S1: *Ratio of predictive performance of expression models in aligned versus misaligned imputation*

samples across AFR (left) and EUR (right) ancestry in the imputation sample. Here, we down-sample the

EUR imputation sample to match the sample size of the AFR imputation sample.

Figure S2: *Predictive performance of expression models in aligned and misaligned imputation samples.*

Figure S3: *Predictive performance of ancestry-unaware expression models compared to ancestry-*

specific models across 5 tissues. Boxplot of difference in predictive performance in EUR **(A)** and AFR **(B)**

samples between ancestry-aligned models and ancestry-unaware models. We consider (1) individuals of

all ancestry in the training sample of the ancestry-unaware model (gold) or only EUR and AFR individuals in the training sample (grey). The red line indicates a difference of 0.

Figure S4: *Predictive performance of ancestry-unaware expression models across minor allele frequency thresholds.*

Figure S5: *TWAS Miami plots across AFR and EUR ancestry groups for asthma using whole blood gene expression models.*

Figure S6: *Correlation of TWAS Z-scores across ancestry-specific, individual biobank GWAS cohorts and 5 meta-analytic strategies.*

Figure S7: *TWAS associations across EUR and AFR ancestry groups for DFFA across 5 tissues.* The effect size is given with the point (triangle if association is transcriptome-wide significant) with a 95% confidence interval provided.

Figure S8: *Miami plots of DFFA local-eQTLs and GWAS signal for SNPs around DFFA.* In (A), color shows linkage disequilibrium R^2 to lead eQTL SNP. Grey line shows a nominal P-value cutoff of 0.05 (|Z| = 1.96).

Figure S9: *Empirical Bayes estimates of bias and inflation in TWAS Z-scores across meta-analysis strategies.* Estimates of bias (top) and bottom (inflation) with one standard error width around the estimate are given across meta-analysis strategies. The dotted lines provide a reference for the null (0 for bias and 1 for inflation).

Figure S10: *Comparison of two IVW meta-analyzed Z-scores.* Vertical and horizontal dotted lines give a reference for the Bonferroni-corrected threshold for transcriptome-significance. A diagonal line is provided from reference.

Figure S11: *Comparison of meta-analyzed Z-scores with individual biobank TWAS Z-scores.* Ancestry-specific TWAS Z-scores for individual biobanks are shown in the top panel, colored by ancestry. Meta-analyzed Z-scores are shown in the bottom panel with shapes reflecting the different strategies. Dotted lines provide a reference for transcriptome-wide significance.

Figure S12: *UKBB T-PheWAS associations across 5 representative asthma-associated genes through European-only meta-analytic TWAS, grouped by phecode group.* The horizontal dotted line shows FDR-adjusted P = 0.05.

Figure S13: *Miami plots of UKBB T-PheWAS associations across 2 genes previously implicated through*

GWAS and detected in European-only meta-analytic TWAS in GBMI.

REFERENCES

Abul-Husn, N.S., and Kenny, E.E. (2019). Personalized Medicine and the Power of Electronic Health
Records. Cell *177*, 58–69.

Aguet, F., Barbeira, A.N., Bonazzola, R., Brown, A., Castel, S.E., Jo, B., Kasela, S., Kim-Hellmuth, S., Liang,
Y., Oliva, M., et al. (2020). The GTEx Consortium atlas of genetic regulatory effects across human tissues.
Science *369*, 1318–1330.

Amariuta, T., Ishigaki, K., Sugishita, H., Ohta, T., Koido, M., Dey, K.K., Matsuda, K., Murakami, Y., Price,
A.L., Kawakami, E., et al. (2020). Improving the trans-ancestry portability of polygenic risk scores by
prioritizing variants in predicted cell-type-specific regulatory elements. Nature Genetics *52*, 1346–1354.

Auton, A., Abecasis, G.R., Altshuler, D.M., Durbin, R.M., Bentley, D.R., Chakravarti, A., Clark, A.G.,
Donnelly, P., Eichler, E.E., Flicek, P., et al. (2015). A global reference for human genetic variation. Nature
526, 68–74.

Baca, S., Singler, C., Zacharia, S., Seo, J.-H., Morova, T., Hach, F., Ding, Y., Schwarz, T., Flora Huang, C.-C.,
Kalita, C., et al. (2021). Genetic determinants of chromatin reveal prostate cancer risk mediated by
context-dependent gene regulation. BioRxiv 2021.05.10.443466.

Baharian, S., Barakatt, M., Gignoux, C.R., Shringarpure, S., Errington, J., Blot, W.J., Bustamante, C.D.,
Kenny, E.E., Williams, S.M., Aldrich, M.C., et al. (2016). The Great Migration and African-American
Genomic Diversity. PLoS Genetics *12*, e1006059.

Barbeira, A.N., Dickinson, S.P., Bonazzola, R., Zheng, J., Wheeler, H.E., Torres, J.M., Torstenson, E.S.,
Shah, K.P., Garcia, T., Edwards, T.L., et al. (2018). Exploring the phenotypic consequences of tissue
specific gene expression variation inferred from GWAS summary statistics. Nature Communications *9*, 1–
20.

Barfield, R., Feng, H., Gusev, A., Wu, L., Zheng, W., Pasaniuc, B., and Kraft, P. (2018). Transcriptome-wide
association studies accounting for colocalization using Egger regression. Genetic Epidemiology *42*, 418–
433.

Bhattacharya, A., García-Closas, M., Olshan, A.F., Perou, C.M., Troester, M.A., and Love, M.I. (2020). A
framework for transcriptome-wide association studies in breast cancer in diverse study populations.
Genome Biology *21*, 42.

Bhattacharya, A., Li, Y., and Love, M.I. (2021a). MOSTWAS: Multi-Omic Strategies for Transcriptome-
Wide Association Studies. *17*, e1009398.

Bhattacharya, A., Freedman, A.N., Avula, V., Harris, R., Liu, W., Pan, C., Lusis, A.J., Joseph, R.M.,
Smeester, L., Hartwell, H.J., et al. (2021b). Genetic control of fetal placental genomics contributes to
development of health and disease. MedRxiv 2021.04.12.21255170.

Bhattacharya, S., Lou, X., Hwang, P., Rajashankar, K.R., Wang, X., Gustafsson, J.Å., Fletterick, R.J.,
Jacobson, R.H., and Webb, P. (2014). Structural and functional insight into TAF1-TAF7, a subcomplex of
transcription factor II D. Proceedings of the National Academy of Sciences of the United States of
America *111*, 9103–9108.

Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A., Vukcevic, D., Delaneau,
O., O'Connell, J., et al. (2018). The UK Biobank resource with deep phenotyping and genomic data.
Nature *562*, 203–209.

Cao, C., Ding, B., Li, Q., Kwok, D., Wu, J., and Long, Q. (2021). Power analysis of transcriptome-wide
association study: Implications for practical protocol choice. PLoS Genetics *17*, e1009405.

Friedman, J., Hastie, T., and Tibshirani, R. (2010). Regularization Paths for Generalized Linear Models via
Coordinate Descent. Journal of Statistical Software *33*, 1–22.

Gallagher, M.D., and Chen-Plotkin, A.S. (2018). The Post-GWAS Era: From Association to Function.
American Journal of Human Genetics *102*, 717–730.

Gamazon, E.R., Wheeler, H.E., Shah, K.P., Mozaffari, S. v, Aquino-Michaels, K., Carroll, R.J., Eyler, A.E.,
Denny, J.C., Nicolae, D.L., Cox, N.J., et al. (2015). A gene-based association method for mapping traits
using reference transcriptome data. Nature Genetics *47*, 1091–1098.

Geoffroy, E., Gregga, I., and Wheeler, H.E. (2020). Population-Matched Transcriptome Prediction
Increases TWAS Discovery and Replication Rate. IScience *23*, 101850.

Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D., Wallace, C., and Plagnol, V.
(2014). Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary
Statistics. PLoS Genetics *10*, e1004383.

Giambartolomei, C., Liu, J.Z., Zhang, W., Hauberg, M., Shi, H., Boocock, J., Pickrell, J., Jaffe, A.E.,
Pasaniuc, B., and Roussos, P. (2018). A Bayesian framework for multiple trait colocalization from
summary association statistics. Bioinformatics *34*, 2538–2545.

Gleason, K.J., Yang, F., Pierce, B.L., He, X., and Chen, L.S. (2020). Primo: Integration of multiple GWAS
and omics QTL summary statistics for elucidation of molecular mechanisms of trait-associated SNPs and
detection of pleiotropy in complex traits. Genome Biology *21*, 236.

Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W.J.H., Jansen, R., de Geus, E.J.C., Boomsma,
D.I., Wright, F.A., et al. (2016). Integrative approaches for large-scale transcriptome-wide association
studies. Nature Genetics *48*, 245–252.

Hauberg, M.E., Zhang, W., Giambartolomei, C., Franzén, O., Morris, D.L., Vyse, T.J., Ruusalepp, A.,
Fromer, M., Sieberts, S.K., Johnson, J.S., et al. (2017). Large-Scale Identification of Common Trait and
Disease Variants Affecting Gene Expression. American Journal of Human Genetics *100*, 885–894.

He, X., Fuller, C.K., Song, Y., Meng, Q., Zhang, B., Yang, X., and Li, H. (2013). Sherlock: Detecting gene-
disease associations by matching patterns of expression QTL and GWAS. American Journal of Human
Genetics *92*, 667–680.

Hebbring, S.J. (2014). The challenges, advantages and future of phenome-wide association studies.
Immunology *141*, 157–165.

Hedges, L., and Vevea, J. (1998). Fixed- and random-effects models in meta-analysis. Psychological
Methods *3*, 486–504.

Higgins, J.P.T., and Thompson, S.G. (2002). Quantifying heterogeneity in a meta-analysis. Statistics in
Medicine *21*, 1539–1558.

Hu, Y., Li, M., Lu, Q., Weng, H., Wang, J., Zekavat, S.M., Yu, Z., Li, B., Gu, J., Muchnik, S., et al. (2019). A
statistical framework for cross-tissue transcriptome-wide association analysis. Nature Genetics *51*, 568–
576.

van Iterson, M., van Zwet, E.W., Heijmans, B.T., and Heijmans, B.T. (2017). Controlling bias and inflation
in epigenome- and transcriptome-wide association studies using the empirical null distribution. Genome
Biology *18*, 19.

Johansson, Å., Rask-Andersen, M., Karlsson, T., and Ek, W.E. (2019). Genome-wide association analysis
of 350 000 Caucasians from the UK Biobank identifies novel loci for asthma, hay fever and eczema.
Human Molecular Genetics *28*, 4022–4041.

Keys, K.L., Mak, A.C.Y., White, M.J., Eckalbar, W.L., Dahl, A.W., Mefford, J., Mikhaylova, A. v., Contreras,
M.G., Elhawary, J.R., Eng, C., et al. (2020). On the cross-population generalizability of gene expression
prediction models. PLoS Genetics *16*, e1008927.

Korthauer, K., Kimes, P.K., Duvallet, C., Reyes, A., Subramanian, A., Teng, M., Shukla, C., Alm, E.J., and
Hicks, S.C. (2019). A practical guide to methods controlling false discoveries in computational biology.
Genome Biology *20*, 1–21.

Lee, C.H., Eskin, E., and Han, B. (2017). Increasing the power of meta-analysis of genome-wide
association studies to detect heterogeneous effects. In Bioinformatics, (Oxford University Press), pp.
i379–i388.

Luningham, J.M., Chen, J., Tang, S., de Jager, P.L., Bennett, D.A., Buchman, A.S., and Yang, J. (2020).
Bayesian Genome-wide TWAS Method to Leverage both cis- and trans-eQTL Information through
Summary Statistics. American Journal of Human Genetics *107*, 714–726.

Mancuso, N., Freund, M.K., Johnson, R., Shi, H., Kichaev, G., Gusev, A., and Pasaniuc, B. (2019).
Probabilistic fine-mapping of transcriptome-wide association studies. Nature Genetics *51*, 675–682.

Márquez-Luna, C., Gazal, S., Loh, P.R., Kim, S.S., Furlotte, N., Auton, A., and Price, A.L. (2020). LDpred-
funct: incorporating functional priors improves polygenic prediction accuracy in UK Biobank and
23andMe data sets. BioRxiv.

Mbatchou, J., Barnard, L., Backman, J., Marcketta, A., Kosmicki, J.A., Ziyatdinov, A., Benner, C.,
O'Dushlaine, C., Barber, M., Boutkov, B., et al. (2020). Computationally efficient whole genome
regression for quantitative and binary traits. BioRxiv 2020.06.19.162354.

McGuirl, M.R., Smith, S.P., Sandstede, B., and Ramachandran, S. (2020). Detecting shared genetic
architecture among multiple phenotypes by hierarchical clustering of gene-level association statistics.
Genetics *215*, 511–529.

Nagpal, S., Meng, X., Epstein, M.P., Tsoi, L.C., Patrick, M., Gibson, G., de Jager, P.L., Bennett, D.A., Wingo,
A.P., Wingo, T.S., et al. (2019). TIGAR: An Improved Bayesian Tool for Transcriptomic Data Imputation
Enhances Gene Mapping of Complex Traits. American Journal of Human Genetics *105*, 258–266.

Parrish, R.L., Gibson, G.C., Epstein, M.P., and Yang, J. (2022). TIGAR-V2: Efficient TWAS tool with
nonparametric Bayesian eQTL weights of 49 tissue types from GTEx V8. Human Genetics and Genomics
Advances *3*, 100068.

Pasaniuc, B., Zaitlen, N., Shi, H., Bhatia, G., Gusev, A., Pickrell, J., Hirschhorn, J., Strachan, D.P., Patterson,
N., Price, A.L., et al. (2014). Fast and accurate imputation of summary statistics enhances evidence of
functional enrichment. Bioinformatics *30*, 2906–2914.

Pathak, G.A., Singh, K., Miller-Fleming, T.W., Wendt, F., Ehsan, N., Hou, K., Johnson, R., Lu, Z., Gopalan,
S., Dimbou, L.Y., et al. (2020). Integrative analyses identify susceptibility genes underlying COVID-19
hospitalization. MedRxiv.

Pavlides, J.M.W., Zhu, Z., Gratten, J., McRae, A.F., Wray, N.R., and Yang, J. (2016). Predicting gene
targets from integrative analyses of summary data from GWAS and eQTL studies for 28 human complex
traits. Genome Medicine *8*, 84.

Portelli, M.A., Nicole Dijk, F., Ketelaar, M.E., Shrine, N., Hankinson, J., Bhaker, S., Grotenboer, N.S.,
Obeidat, M., Henry, A.P., Billington, C.K., et al. (2020). Phenotypic and functional translation of IL1RL1
locus polymorphisms in lung tissue and asthmatic airway epithelium. JCI Insight *5*.

Reijmerink, N.E., Postma, D.S., Bruinenberg, M., Nolte, I.M., Meyers, D.A., Bleecker, E.R., and
Koppelman, G.H. (2008). Association of IL1RL1, IL18R1, and IL18RAP gene cluster polymorphisms with
asthma and atopy. Journal of Allergy and Clinical Immunology *122*, 651-654.e8.

Reijmerink, N.E., Postma, D.S., and Koppelman, G.H. (2010). The candidate gene approach in asthma:
what happens with the neighbours? European Journal of Human Genetics 17.

Shang, L., Smith, J.A., Zhao, W., Kho, M., Turner, S.T., Mosley, T.H., Kardia, S.L.R., and Zhou, X. (2020).
Genetic Architecture of Gene Expression in European and African Americans: An eQTL Mapping Study in
GENOA. American Journal of Human Genetics *106*, 496–512.

Shi, J., and Lee, S. (2016). A novel random effect model for GWAS meta-analysis and its application to
trans-ethnic meta-analysis. Biometrics *72*, 945–954.

Shifman, S., Kuypers, J., Kokoris, M., Yakir, B., and Darvasi, A. (2003). Linkage disequilibrium patterns of
the human genome across populations. Human Molecular Genetics *12*, 771–776.

Smith, G.D., and Ebrahim, S. (2003). 'Mendelian randomization': can genetic epidemiology contribute to
understanding environmental determinants of disease? International Journal of Epidemiology *32*, 1–22.

Stevens, J.R., Masud, A. al, and Suyundikov, A. (2017). A comparison of multiple testing adjustment
methods with block-correlation positivelydependent tests. PLoS ONE *12*.

Swede, H., Stone, C.L., and Norwood, A.R. (2007). National population-based biobanks for genetic
research. Genetics in Medicine *9*, 141–149.

Unlu, G., Gamazon, E.R., Qi, X., Levic, D.S., Bastarache, L., Denny, J.C., Roden, D.M., Mayzus, I., Breyer,
M., Zhong, X., et al. (2019). GRIK5 Genetically Regulated Expression Associated with Eye and Vascular
Phenomes: Discovery through Iteration among Biobanks, Electronic Health Records, and Zebrafish. The
American Journal of Human Genetics *104*, 503–519.

Unlu, G., Qi, X., Gamazon, E.R., Melville, D.B., Patel, N., Rushing, A.R., Hashem, M., Al-Faifi, A., Chen, R.,
Li, B., et al. (2020). Phenome-based approach identifies RIC1 -linked Mendelian syndrome through
zebrafish models, biobank associations and clinical studies. Nature Medicine 2020 26:1 *26*, 98–109.

Veturi, Y., and Ritchie, M.D. (2018). How powerful are summary-based methods for identifying
expression-trait associations under different genetic architectures? In Pacific Symposium on
Biocomputing, (World Scientific Publishing Co. Pte Ltd), pp. 228–239.

Vicente, C.T., Revez, J.A., and Ferreira, M.A.R. (2017). Lessons from ten years of genome-wide
association studies of asthma. Clinical and Translational Immunology *6*, e165.

Wainberg, M., Sinnott-Armstrong, N., Mancuso, N., Barbeira, A.N., Knowles, D.A., Golan, D., Ermel, R.,
Ruusalepp, A., Quertermous, T., Hao, K., et al. (2019). Opportunities and challenges for transcriptome-
wide association studies. Nature Genetics *51*, 592–599.

Wei, W.Q., Bastarache, L.A., Carroll, R.J., Marlo, J.E., Osterman, T.J., Gamazon, E.R., Cox, N.J., Roden,
D.M., and Denny, J.C. (2017). Evaluating phecodes, clinical classification software, and ICD-9-CM codes
for phenome-wide association studies in the electronic health record. PLoS ONE *12*.

Wijmenga, C., and Zhernakova, A. (2018). The importance of cohort studies in the post-GWAS era.
Nature Genetics *50*, 322–328.

Wyss, A.B., Sofer, T., Lee, M.K., Terzikhan, N., Nguyen, J.N., Lahousse, L., Latourelle, J.C., Smith, A.V.,
Bartz, T.M., Feitosa, M.F., et al. (2018). Multiethnic meta-analysis identifies ancestry-specific and cross-
ancestry loci for pulmonary function. Nature Communications *9*, 1–15.

Yang, F., Wang, J., Pierce, B.L., and Chen, L.S. (2017). Identifying cis-mediators for trans-eQTLs across
many human tissues using genomic mediation analysis. Genome Research *27*, 1859–1871.

Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a tool for genome-wide complex trait
analysis. American Journal of Human Genetics *88*, 76–82.

Yuan, Z., Zhu, H., Zeng, P., Yang, S., Sun, S., Yang, C., Liu, J., and Zhou, X. (2020). Testing and controlling
for horizontal pleiotropy with probabilistic Mendelian randomization in transcriptome-wide association
studies. Nature Communications *11*, 1–14.

Zhang, Y., Quick, C., Yu, K., Barbeira, A., Luca, F., Pique-Regi, R., Kyung Im, H., and Wen, X. (2020).
PTWAS: Investigating tissue-relevant causal molecular mechanisms of complex traits using probabilistic
TWAS analysis. Genome Biology *21*, 1–26.

Zhong, Y., Perera, M.A., and Gamazon, E.R. (2019). On Using Local Ancestry to Characterize the Genetic
Architecture of Human Traits: Genetic Regulation of Gene Expression in Multiethnic or Admixed
Populations. American Journal of Human Genetics *104*, 1097–1115.

Zhou, D., Jiang, Y., Zhong, X., Cox, N.J., Liu, C., and Gamazon, E.R. (2020). A unified framework for joint-
tissue transcriptome-wide association and Mendelian randomization analysis. Nature Genetics *52*,
1239–1246.

Zhou, W., Nielsen, J.B., Fritsche, L.G., Dey, R., Gabrielsen, M.E., Wolford, B.N., LeFaive, J., VandeHaar, P.,
Gagliano, S.A., Gifford, A., et al. (2018). Efficiently controlling for case-control imbalance and sample
relatedness in large-scale genetic association studies. Nature Genetics *50*, 1335–1341.

Zhou, W., Kanai, M., Wu, K.-H.H., Humaira, R., Tsuo, K., Hirbo, J.B., Wang, Y., Bhattacharya, A., Zhao, H.,
Namba, S., et al. (2021). Global Biobank Meta-analysis Initiative: powering genetic discovery across
human diseases. MedRxiv *27*, 2021.11.19.21266436.

Zhu, A., Matoba, N., Wilson, E.P., Tapia, A.L., Li, Y., Ibrahim, J.G., Stein, J.L., and Love, M.I. (2021).
MRLocus: Identifying causal genes mediating a trait through Bayesian estimation of allelic
heterogeneity. PLOS Genetics *17*, e1009455.

Zhu, Z., Guo, Y., Shi, H., Liu, C.L., Panganiban, R.A., Chung, W., O'Connor, L.J., Himes, B.E., Gazal, S.,
Hasegawa, K., et al. (2020). Shared genetic and experimental links between obesity-related traits and
asthma subtypes in UK Biobank. Journal of Allergy and Clinical Immunology *145*, 537–549.

A. Transcriptome-Wide Association Study

SNPs

TGCTAGA
GAGACAT
CTAGAGT
TCATTGT

CHALLENGES:
• Genetic ancestry
• Population structure
• Linkage disequilibrium

Tissue-specific GReX

CHALLENGES:
• Tissue specificity
• Cell-type- and state specificity
• Dynamic trajectories

Trait

Case
Control
Control
Case

CHALLENGES:
• Phenotype acquisition
• Case-control definition
• Ascertainment bias

B. GBMI TWAS Pipeline

SNPs

Step I

TGCTAGA
GAGACAT
CTAGAGT
TCATTGT

Tissue-specific GReX

Step II, III

Step III

Trait (Biobank 1)

Case
Control
Control
Case

...

Trait (Biobank 19)

Case
Control
Control
Control

I. Train expression models from germline genetics
• Cis-SNP, multi-tissue method: **JTI** (Zhou *et al* 2020, *Nat Gen*)
• Distal-SNP-enriched method: **MOSTWAS** (Bhattacharya *et al* 2021, *PLOS Genet*)
• Align genetic ancestry of eQTL and GWAS panels

II. Test for gene-trait associations
• Meta-analyze across biobanks and ancestry groups using properly aligned LD references with inverse-variance weights

III. Conduct follow-up tests for prioritized genes
• Conditional/permutation tests
• **FOCUS** to estimate credible set of causal genes at a locus (Mancuso *et al* 2019, *Nat Gen*)
• Distal SNP test for potential mediators in **MOSTWAS**
• Estimate the degree of SNP horizontal pleiotropy

Figure 1: *Overview of GBMI transcriptome-wide association study (TWAS) pipeline with challenges at every data level.* **(A)** Each level of data in TWAS introduces a unique set of challenges: (1) genetics data include confounding from genetic ancestry, population structure and relatedness, and complex linkage disequilibrium patterns, (2) gene expression data introduces context-specific factors, such as tissue-, cell-type-, or cell-state-specific expression, and (3) phenotypic data, especially in the meta-analyses of multiple biobanks, involve challenges in acquiring and aggregating phenotypes, properly defining controls for phenotypes, and ascertainment and selection bias from non-random sampling. **(B)** An overview of the GBMI TWAS pipeline: (1) JTI and MOSTWAS for model training, (2) inverse-variance weighted meta-analysis using per-biobank, per-ancestry group TWAS summary statistics, and (3) various follow-up tests, including conditional or permutation tests, distal-SNPs added last test, probabilistic fine-mapping using FOCUS, and tests for SNP horizontal pleiotropy. Dotted lines represent associations that are tested in the TWAS pipeline, while the solid lines represent a link built through predictive modeling.

Figure 2: *Comparison of predictive performance of genetic models of expression across ancestry*. **(A)** Distribution of difference in adjusted R^2 (Y-axis) when predicting expression in the AFR imputation sample between models trained in EUR and in AFR training samples across tissue (X-axis). **(B)** Distribution of difference in adjusted R^2 between ancestry-specific and ancestry-unaware models imputing into EUR (left) and AFR (right) samples.

Figure 3: *Comparison of meta-analytic strategies for multi-biobank, trans-ancestry TWAS.* **(A)** Scatterplot of per-ancestry meta-analyzed TWAS scores across EUR (X-axis) and AFR ancestry (Y-axis). The dotted horizontal and vertical lines indicate P < *2.5 × 10⁻⁶* with a diagonal line for reference. Points are colored based on which ancestry population the TWAS association meets P < *2.5 × 10⁻⁶*. **(B)** QQ-plot of TWAS Z-scores, colored by meta-analytic strategies. Per ancestry refers to TWAS meta-analysis across meta-analyzed ancestry-specific GWAS summary statistics. Per bank/per ancestry refers to TWAS meta-analysis using all biobank- and ancestry-specific GWAS summary statistics. **(C)** Effect sizes and Bonferroni-corrected confidence intervals (CIs) for TWAS associations across 17 individual biobanks (stratified by ancestry group with EUR in green and AFR in red) and 2 IVW meta-analysis strategies (in yellow) for 5 representative genes. The Higgins-Thompson I² statistic for heterogeneity is provided, with the dotted line showing the null.

Figure 4: *GReX-PheWAS for categorizing phenome-wide associations for TAF7 genetically-regulated expression in UKBB.* **(A)** Boxplots of -log$_{10}$ Benjamini-Hochberg FDR-adjusted P-values of GTAs across 9 phenotype groups. The dotted grey line showed FDR-adjusted P = 0.05. **(B)** Miami plot of TWAS Z-scores (Y-axis) across phenotypes, colored by phecode group. The dotted grey line shows the significance threshold for Benjamini-Hochberg FDR correction and phenotypes are labelled if the association passes Bonferroni correction.