Genomic epidemiology and longitudinal sampling of ward wastewater

environments and patients reveals complexity of the transmission dynamics of

bla_{KPC}-carbapenemase-producing Enterobacterales in a hospital setting

Stoesser N[a, b,]*, George R[c,]*, Aiken Z[c], Phan HTT[d], Lipworth S[a], Wyllie DH[e], Quan

TP[a], Mathers AJ[f], De Maio N[g], Seale AC[h], Eyre DW[a, b], Vaughan A[a], Swann J[a], Peto

TEA[a, i], Crook DW[a, b, i], Cawthorne J[c], Dodgson A[c], Walker AS[a, b, i], and the TRACE

Investigators' Group

[a] Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

[b] NIHR Health Protection Research Unit in Healthcare Associated Infections and

Antimicrobial Resistance at University of Oxford in partnership with Public Health

England, Oxford, United Kingdom

[c] Manchester University NHS Foundation Trust, Manchester, United Kingdom

[d] University of Southampton, Southampton, United Kingdom

[e] United Kingdom Health Security Agency, United Kingdom

[f] University of Virginia, Charlottesville, United States of America

[g] EMBL-European Bioinformatics Institute, Cambridge, United Kingdom

[h] University of Warwick, Coventry, United Kingdom

[i] NIHR Oxford Biomedical Research Centre, Oxford University Hospitals NHS

Foundation Trust, John Radcliffe Hospital, Oxford, United Kingdom

*Contributed equally

Keywords: Carbapenemase-producing Enterobacterales, genomic epidemiology, *bla*~KPC~, wastewater sites, healthcare-associated

Corresponding author: Nicole Stoesser

Corresponding address: Department of Microbiology (Research), Level 7, John Radcliffe Hospital, Headley Way, Headington, OX3 9DU

Affiliation: Nuffield Department of Medicine, University of Oxford, Oxford UK and Oxford University Hospitals NHS Foundation Trust, Oxford, UK

e-mail: nicole.stoesser@ndm.ox.ac.uk

ABSTRACT

Background

Healthcare-associated wastewater reservoirs and asymptomatic gastrointestinal

patient colonisation by carbapenemase-producing Enterobacterales (CPE) can be

important in nosocomial CPE dissemination and infection. We characterised these

niches and within-niche diversity in a bla_{KPC}-associated CPE (KPC-E) endemic

healthcare setting, to better understand transmission potential.

Methods

We systematically sampled wastewater sites and patients across three units (six

wards) over 6-12 months in 2016 in a KPC-E endemic hospital. We used Illumina

sequencing to characterise up to five isolates per sample. Recombination-adjusted

phylogenies were used to define genetically related strains; assembly and mapping-

based typing approaches were used to characterise antimicrobial resistance gene and

insertion sequence profiles, and Tn*4401* types/target site sequences. The wider

accessory genome was evaluated in a subset of the largest clusters, and those crossing

niches.

Findings

Wastewater site KPC-E-positivity was substantial (101/349 sites [28.9%] positive,

319/4,488 [7.1%] sampling events positive); 183/4,425 (4.1%) of patients were CPE

culture-positive over the same timeframe. 13 species and 109 KPC-E strains were

observed across niches, and 24% of wastewater and 26% of patient KPC-E-positive

samples harboured ≥1 strain. Most diversity was explained by the individual niche,

suggesting highly localised factors are important in selection and spread.

Tn*4401*+target site sequence (TSS) diversity was greater in wastewater sites

(p<0.001), suggesting these might favour Tn*4401*-associated transposition/evolution

and dissemination. Shower/bath and sluice/mop-associated sites were more likely to

be KPC-E-positive (Adjusted Odds Ratio [95% CI]: 2.69 [1.44-5.01], p=0.0019 and

2.60 [1.04-6.52], p=0.0410, respectively). Different strains had different transmission

and bla_{KPC} dissemination dynamics.

Interpretation

There may be substantial KPC-E colonisation of wastewater sites and patients in

KPC-E-endemic healthcare settings. Niche-specific factors, and different strains with

different transmission dynamics influence carbapenemase gene dissemination. New

transmission models incorporating complex, multi-level dynamics are needed to

better quantify CPE dissemination to inform interventions and reduce transmission.

Funding

This study was supported by the National Institute for Health Research, UK.

RESEARCH IN CONTEXT

Evidence before this study

We searched PubMed to identify previous studies using whole genome sequencing

(WGS) to characterise in-hospital dissemination and persistence of carbapenemase-

producing Enterobacterales (CPEs) or the presence of CPEs in hospital wastewater

reservoirs. In our search (01/Jan/1996-30/Sep/2021) we used the terms ((((CPE) OR

(KPC)) AND (healthcare-associated OR nosocomial OR hospital-associated)) AND

(genom*) AND ((environment OR patient))). We had no restrictions on language. We

found sixty-six studies, but these were limited by: (i) evaluating single species or

lineages only (n=34), (ii) including patients or environmental outbreaks only, and not

both (n=23), (iii) sampling only small numbers (n<50) of patients or environmental

sites (n=26), or (iv) performing no evaluation of within-niche diversity (n=64). 13

studies were not relevant (e.g. investigating non-hospital locations, *ex vivo* or *in vivo*

evolution), two studies did not use WGS, and one study was not primary research.

Added value of this study

We provide a comprehensive assessment of healthcare-associated CPE dissemination

and persistence in hospital wastewater reservoirs through systematic sampling of a

large number of environmental sites (n=349 sites, n=4,488 samples) and patients

(n=4,425) over 6-12 months in a single hospital. For all CPE-positive environmental

samples (n=319) and a subset of CPE-positive patient samples (n=97/399 [24.3%]

samples from 76/183 [41.5%] CPE-positive patients), we sampled up to five CPE

isolates per sample to capture within-niche diversity, and used short-read Illumina

WGS (n=1,732 isolates successfully sequenced) to characterise within-niche

diversity, changes in colonisation over time, and genetic overlap between patient and

environmental niches.

Implications of all the available evidence

Through a detailed and resource-intensive study we captured the dynamics of seeding

and dissemination of important carbapenemase genes amongst patients and

environmental reservoirs within a hospital. We found differential colonisation and

dissemination dynamics for different species and lineages within and between niches.

Improved approaches incorporating variable within-niche diversity, accessory

distances and horizontal gene transfer in transmission evaluations are required to

better understand CPE dissemination and persistence, in order to direct interventions

limiting transmission.

INTRODUCTION

Carbapenemase-producing Enterobacterales (CPE) are a global health threat[1], and commonly associated with multi-drug resistance, rendering treatment of CPE infections difficult. Major transmissible global carbapenemases include the metallo-beta-lactamases (bla_{NDM}, bla_{VIM}, bla_{IMP}), some oxacillinases (blaOXA$_{-48/48-like}$ variants), and the *Klebsiella pneumoniae* carbapenemase (KPC, encoded by bla_{KPC})[2]. Intra- and inter-species horizontal transfer of these genes is facilitated by mobile genetic elements such as transposons and plasmids, resulting in rapid dissemination of carbapenem resistance[3, 4].

CPE have been isolated from human and animal gastrointestinal tracts, and from sewage, rivers and sink drains[5-7]. Recent studies have highlighted that hospital wastewater sites may act as a CPE reservoir[5], but the diversity within these niches, genetic overlap with patient isolates and likely directionality and rates of transmission remain to be elucidated. Studies evaluating the contribution of patient-to-patient transmission as the sole explanatory factor explaining transmission have generally been unable to robustly explain the majority of transmission events, suggesting that patient, staff and/or environmental reservoirs remain insufficiently sampled to account for the diversity observed[8]. Few studies have considered within-niche diversity of strains by sampling multiple isolates per individual and/or site[9-11].

The Manchester University NHS Foundation Trust (MFT; Manchester, UK), has experienced on-going transmission and patient acquisition of multi-species CPE, predominantly bla_{KPC}-positive Enterobacterales (KPC-E), since 2009. Following a KPC-*Escherichia coli* outbreak on two cardiac wards in the Manchester Royal

Infirmary (MRI; part of MFT), and subsequent ward closure and plumbing

infrastructure replacement[12], we undertook a prospective study systematically

sampling all wastewater sites on six wards for 6-12 months alongside patient rectal

screening/clinical sampling, and used an anonymised electronic database to

characterise patient admission, ward movement, sampling profiles and culture results.

Multiple colonies were isolated from KPC-E-positive samples to define within-niche

diversity, and whole genome sequencing (Illumina) of individual isolates was used as

to characterise this diversity and consider modes of evolution and transmission.

METHODS

Study setting and epidemiological data

MFT is a large, regional referral centre in northwest England, UK, managing >10,000

patients/year. In response to the regional emergence of KPC-E[13], the Trust

implemented an extensive Infection Prevention and Control (IPC) programme,

consistent with UK CPE guidelines (published in 2014)[14]. Despite this, in April 2015

there was a large bla_{KPC}-*E. coli* outbreak in the cardiac unit[12], leading to its closure

(September 2015-January 2016) and complete refurbishment, including plumbing

replacement. Subsequently systematic, prospective wastewater site sampling was

undertaken (described further below).

To evaluate patient-level microbiological data and hospital stays, MFT electronic

bacteriology records were linked, based on NHS numbers, to patient administration

data, and anonymised. For this study, we analysed anonymised patient data from 1st

January 2016-31st December 2016 inclusive, and included only those patients with

exposure to any one of the three units (i.e. six study wards) during that time.

Environmental sampling and laboratory processing

All sink/drain/wastewater sites on six wards in three units were sampled: the cardiac

unit (wards 3 [W3] and 4 [W4]); the geratology unit (wards 45 [W45] and 46 [W46]);

and the acute medicine unit (wards AM1 and AM2) (Table S1 shows sampling site

designations). W3/W4 wastewater sites were sampled fortnightly on rotation from 8th

January 2016-28th December 2016. W45/W46 and AM1/AM2 wastewater sites were

similarly sampled, but from 18th July 2016-31st December 2016. Wastewater site

types are described in Appendix Methods.

Environmental sampling was carried out by aspirating ~20mls of wastewater from

sink P-traps, shower drains or toilets and performing enrichment-based culture (see

Appendix methods). Multiplex, real-time, quantitative PCR (for bla_{NDM}, bla_{KPC},

bla_{OXA-48}[12]) was performed on broths following incubation. 10μl of any bla_{KPC}-

positive-broth were streaked onto CPE-selective agar plates (Chromid CARBA) and

re-incubated (aerobically, 37°C).

Patient sampling and laboratory processing

Routinely collected patient clinical samples were processed using standard operating

procedures in line with UK Standards[15]. Rectal swabs were screened for CPE using

the Cepheid Xpert Carba-R assay or an in-house multiplex PCR (identifying bla_{KPC},

bla_{NDM}, and bla_{OXA-48}) from November 2014. Species identification of isolates was

performed by MALDI-ToF (Bruker Biotyper, Bruker MD); antimicrobial

susceptibility testing was performed as per EUCAST guidelines[16].

Analyses of within-niche diversity

To evaluate within-niche diversity in the environment and patients, up to five

different Enterobacterales colonies were individually sub-cultured; bla_{KPC}-positivity

in sub-cultures was re-confirmed by PCR and species identification performed by

MALDI-ToF prior to storage in nutrient broth at -80°C.

DNA extraction and isolate sequencing

DNA was extracted from frozen sub-culture stocks with the QuickGene kit, as per the

manufacturer's instructions, with an additional mechanical lysis step post-chemical

lysis (6m/s at 40secs x2 FastPrep, MPBio). Isolates were sequenced on the Illumina

HiSeq 4000 generating 150bp paired-end reads.

Sequence read processing and analysis

Sequencing reads were initially trimmed/quality controlled[17]; species identification

was performed with Kraken[18]. If at least 50 sequenced isolates were of a single

species, reads were mapped to species-specific references (Appendix Methods).

Variant calling and filtering was performed, and consensus fasta sequences of variant

calls with respect to the reference generated. Recombination-adjusted phylogenies by

species were created from core chromosomal single nucleotide variants

(SNVs)(padded to the length of the reference genome prior to analysis) using IQtree[19]

(flags: -m GTR+G -blmin 0.00000001 -t PARS) and then ClonalFrameML[20] (default

parameters). A distance of ≤400 SNVs was used as a threshold to define strains

(threshold used similar to[21]; Fig.S2A-E for most common species identified). For

detailed sub-strain analysis, sub-strains were defined on the basis of accessory

genome clusters characterised using Panaroo (default parameters including --clean-

mode strict[22]) and clustered using the R heatmap package.

Isolates were assembled using SPAdes (v3.6)[23] with default parameters. Multi-locus

sequence types were derived using BLASTn to relevant species-specific databases

hosted at https://pubmlst.org (Achtman 7-gene MLST for *E. coli*). AMR genes were

identified using ARIBA[24] with the CARD database[25]. bla_{KPC} genes, any Tn*4401*

transposon context and 5-bp target site sequences reflecting Tn*4401* insertion

signatures were characterised using TETyper[26], using a Tn*4401*b reference sequence,

a 5bp flank length and the default structural and SNV profiles.

For the analysis of *K. pneumoniae* strain 9 (see Results), putative transmission

networks were inferred first using SCOTTI[27] (Appendix Methods), and then

heuristically, linking isolates identified as clusters on the basis of accessory

component profiles and geographic/temporal overlap (Appendix Methods).

Sequence data have been deposited in NCBI's GenBank (BioProject accession

PRJNA768622).

Statistical analyses and data visualisation

Descriptive statistics were calculated in R v3.6.2 or STATA v16.1. To conduct

permutational analyses of variance, we used the *adonis* function from the vegan R

package (v2.5-7)[28] on a matrix of pairwise Gower distances based on genetic features

(i.e. composite of species-strain type, AMR gene profile, IS profile, and

Tn*4401*+flanking sequence profiles), which was calculated using the *daisy* function in

the R cluster package. To evaluate the impact of unit location or environmental site

type on the odds of CPE positivity, we used logistic regression with robust standard

errors clustered by environmental site sampled, using the R rms package (v6.2-0)[29].

Data visualisation was using the ggplot2[30] and scatterpie[31] (v.0.1.5) R packages, and

BioRender (www.biorender.com).

Ethical approvals

This study was performed as part of a Trust board-approved and UK Health Security

Agency (formerly Public Health England) long-term outbreak response, and ethics

approval was not required under NHS governance arrangements; all analyses were

undertaken using anonymised data. This was reviewed and agreed by the Joint

Research Office of the Oxford University Hospitals NHS Foundation Trust and the

University of Oxford (who oversee research governance, ethics and assurance for our

institution), and who deemed that neither sponsorship nor research ethics review was

required. Of note, sample and patient numbers represented in the

figures/supplementary data are study-specific labels that are not patient-identifiable.

RESULTS

Environmental sampling revealed high prevalence of wastewater site KPC-E

colonisation

349 sites across the six study wards (i.e. three units) were sampled a total of 4,488

times (Fig.1): Of these, 23/76 (30%) sites and 111/1,564 (7.1%) sampling events on

the cardiac unit were positive for KPC-E, 38/129 (29%) sites and 86/1,309 (6.6%)

sampling events on the acute medicine unit and 40/144 [28%] and 122/1,615 (7.6%)

on the geratology unit (Fig.2, top three panels). 248/349 (71.1%) sites were never

identified as KPC-E-positive (Fig.S3); 7/349 (2.0%) sites appeared more likely to be

persistently KPC-E positive (≥10 consecutive KPC-E-positive sampling events) and

interestingly five of these were medicines/treatment room handwash basin drains

likely only accessible to healthcare staff (A59, F45, F113, T43, T44; Fig.S4). The

proportion of positive sites did not vary significantly over time (Fig.2, bottom panel).

Two utility room sink drains on the cardiac unit (across W3, W4) were positive for

KPC-E a day after W3 reopened to patient admissions (11[th] January 2016) following

closure and complete plumbing replacement over three months; thereafter the unit

was rapidly re-colonised (Fig.2)[12]. Shower/bath drain sites and sluices/sluice

sinks/mop sinks were significantly more likely to be positive for KPC-E than others

(Adjusted Odds Ratio [AOR] (95% CI): 2.69 (1.44-5.01), p=0.0019 and 2.60 (1.04-

6.52), p=0.041, respectively), and toilet water sampling less likely to be positive

(AOR (95% CI): 0.28 (0.10-0.78), p=0.015); there was no evidence of a unit-

dependent difference (Table 1).

Approximately one in twenty patients was affected by CPE, with most cases reflecting

rectal colonisation with KPC-producing K. pneumoniae, E. coli or E. cloacae

Of 2,172 Enterobacterales positive-samples (obtained from 943 patients) and 2,000

with an associated carbapenem susceptibility result (172 carbapenem susceptibility

results missing), 399/2,000 (20.0%) were carbapenem-resistant Enterobacterales

(CRE; i.e. had intermediate or resistant susceptibilities) (Figs.1, S5). For 344/431

(79.8%) CRE isolates cultured from these 399 CRE-positive samples carbapenemase

PCR data were available: All contained at least one known carbapenemase resistance

gene, and 274 (79.7%) were KPC-E (Table S2, henceforth denoted CPE). These CPEs

were cultured from 183 patients (4.1% of all patients overall, 8.4% of all patients with

an Enterobacterales-positive culture) during the study period (median=1, IQR: 1-2,

range: 1-17 CPE isolates). Amongst these patients, CPE-positive samples represented

a median of 10% (IQR: 5-20%) of all samples taken and cultured.

The majority of confirmed CPE-positive samples were rectal screens (n=318/399

[79.7%]); others were non-screening clinical specimens (81/399 [20.3%] including

blood cultures [n=4], urine [45], respiratory [20], wound/intra-abdominal [12]) (Table

S3). 16/183 individuals were positive only on clinical culture (9%), 141 only on rectal

screen (77%), and 26 (14%) on both. By routine laboratory methods, at least 16

species and eight genera were represented amongst CPEs, with *K. pneumoniae*, *E.*

coli and *E. cloacae* complex predominating (364/431 [84.5%]). Notably, the

proportions of carbapenem-resistant cases within these top species varied (n=239/464

[51%] vs 88/1,083 [8%] vs 37/86 [43%] respectively; Fisher's exact test, p<0.001,

Fig.S5).

The proportion of patient CPE-positive cultures was lower than that of environmental

site samples, typically ~3% (Fig.3). Occasional peaks in prevalence were observed,

consistent with outbreaks (e.g. week 31, acute medicine unit). CPE prevalence

amongst patient cultures on the refurbished cardiology ward was 0% (97.5% CI: 0-

0.02) for the first 10 weeks post-reopening.

Structuring of diversity by niche, with most diversity explained by the individual niche

Overall, amongst successfully sequenced isolates (n=1,732) across patients and

environmental reservoirs, at least 13 Enterobacterales species harboured bla_{KPC}, with

109 unique strains represented (and a further ten species not characterised at the strain

level because of small numbers, see Methods). Of the 109 strains, 61 (56%) were

found only in the environment, 27 (24%) only in patients and 21 (19%) in both

patients and the environment (Fig.4); the median strain cluster size was 5 isolates

(range: 1-122), and the median number of unique niches (i.e. unique patients or

wastewater sites) affected by a strain was 2 (range: 1-23) (Fig.4A, Fig.4B). A much

greater proportion of all identified Tn*4401*/TSS types was observed in the

environment versus patients (118/121 [98%%] of types versus 9/121 [7%]

respectively; Fig.S6; Fisher's exact test, p<0.001), and all the isolates with >1 set of

TSSs were seen in environmental sites, consistent with these representing a more

favourable niche for Tn*4401*-associated transposition/evolution and dissemination.

For colonised niches (i.e. patient or environmental), using a metric of genomic

diversity based on strain type, AMR gene profile, IS profile, and Tn*4401*+ 5bp target

site sequence profiles, most variance was explained by the specific individual niche

sampled (R^2=46.0%; p<0.001), versus small but significant contributions made by

reservoir type (patient vs specific environmental niche; R^2=5.9%; p<0.001) and unit

location (R^2=3.0%; p<0.001). For the fifteen examples where exactly the same strain,

AMR gene profile, IS profile, plasmid profile and Tn*4401*/TSS type were seen in

patient and environmental niches, the temporal/geographic relationship of these

suggested: patient>environment transmission in 14/31 (45%) events,

environment>patient transmission in 2/31 (7%) events, environment>environment

transmission in 4/31 (13%) events, and patient>patient transmission in 11/31 (35%)

events (test of equality of proportions, p<0.001).

Environmental and patient niches harbour diverse KPC-E within single samples and over time

Environmental sites showed evidence of major strain-level diversity (Fig.5): in the 101/349 sites from which KPC-E was successfully sequenced at any timepoint, 78/319 (24.4%) KPC-E-positive samples had >1 KPC-E strain, namely 52/319 (16.3%) with two strains, 16/319 (5.0%) with three, 9/319 (2.8%) with four and 1/319 (0.3%) with five strains. Over the study period, colonised environmental sites had a median of 3 (range: 1-17, IQR: 2-5) KPC-E strains. 49/101 sites (49%) were apparently colonised by a single KPC-E strain only, of which 38 were at single sampling-points, with 11 sites showing persistent colonisation (≥2 timepoints positive with the same strain).

For patient samples, based on routine laboratory microbiology, evidence of mixed-species KPC-E colonisation/infection was found in 34/399 (9%) samples from 27/183 (15%) patients; these were predominantly *E. coli/K. pneumoniae* mixtures (n=21 samples; maximum two species identified). To evaluate within-host diversity with genomic rather than species-level resolution, in a subset of 76 patients, strains were sequenced from at least one CPE-positive sample during the study period; for 97 samples, multiple colonies were picked. In total 463 sequenced colonies from 97 samples were available for evaluation of within-patient diversity; 25/97 samples (26%) had more than one strain per sample, namely 17/97 samples (18%) with two KPC-E strains, 6/97 (6%) with three strains and 2/97 (2%) with four strains (Fig.5) - i.e. considerably more diversity identified than based on routine laboratory species identification data (26% vs 9% samples; Fisher's exact test, p<0.001). Longitudinal switches in strain composition of KPC-E-positive samples was common, occurring in

11/21 (52%) KPC-E-positive longitudinal patient samples and 108/202 (53%) KPC-

E-positive longitudinal wastewater site samples.

Phylogenetic analysis reveals distinct population biology for different KPC-E strains

To investigate different modes of *bla*KPC dissemination, we characterised in detail

several clusters, including the largest environmental-only cluster (*E. cloacae* strain 10

[n=122 isolates]), a large cluster involving patients and the environment (*K.*

pneumoniae strain 9 [n=116 isolates], and a cluster in which signatures of Tn*4401*

transposition appeared highly frequent (K. *pneumoniae* strain 11 [n=106 isolates]).

Rapid churn of diversity in restricted environmental niches observed at the

chromosome, plasmid, IS, and AMR gene level - E. cloacae strain 10

E. cloacae strain 10 (eclo10, ST32) appeared a restricted but persistent coloniser of

three co-located sink drains on a geratology ward (in a drug room clinical handwash

basin drain [F45] and utility sink drain [F46], and a separate treatment room clinical

handwash basin drain [F47]) over ~5 months (18/Jul/2016-20/Dec/2016) (Fig.6A). At

the core chromosomal level, strains appeared largely structured by site, with 7 SNVs

separating the majority of isolates in F45 from those in F47, and 14 SNVs those in

F46 from F45 (Fig.6A). There was evidence of rapid churn in AMR gene and IS

content (111 and 36 different profiles respectively; Fig.6A), and all isolates in F47

had acquired an additional set of IncHI2/HI2A replicons on a background of FIB/FII,

except for 1954277_1_4, consistent with plasmid gain/loss (Fig.6B). Although the

Tn*4401* type and flanking sequences were largely identical (Tn*4401*a-ATTGA-

ATTGA in 105/122 (86%) isolates across sinks), there was evidence of unique

additional Tn*4401* deletions occurring in multiple isolates at four distinct timepoints

in the three sites (Fig.6C).

Rapid patient-patient transmission - K. pneumoniae strain 9

K. pneumoniae strain 9 (kpne9, ST252) was highly related (Fig.S7; maximum 6 SNVs

across strain, 68 isolates within 0 SNVs of each other), and spread amongst 21

patients across three wards on two units (acute medicine AM1 and AM2 and

geratology W46), with a dense outbreak in Aug 2016, most consistent, at least

initially, with direct/indirect patient-patient transmission (Fig.7). bla_{KPC} was

consistently nested in Tn*4401*-ATTGA-ATTGA, except for one isolate which

acquired a Tn*4401*-associated mutation (isolate: 2216698_14, C4620T), and another

in which the right target site sequence (TSS) could not be identified (2035791_11).

Amongst the 106 isolates there was evidence of four transient plasmid replicon

acquisitions on a background of the stable presence of a set of IncFIB/IncFII replicons

(Fig.S7). Transmission probabilities inferred by SCOTTI were weak (see Appendix;

Fig.S8), as previously[8], but using accessory cluster profiling to attempt further

discrimination amongst strains suggested that there were several major transmission

clusters (Fig.S9), with some individuals involved in several of these (Fig.7). A single

shower drain site was involved (A13, serving a particular bay with four beds), which

was likely a seeded bystander from affected patients, but may have contributed to

transmission of sub-strain 3g (Fig.7).

Cross-unit dissemination with evidence of significant Tn4401-associated bla$_{KPC}$

transposition - K. pneumoniae strain 11

Given that Tn*4401* transposition is associated with bla_{KPC} dissemination[21], we also

evaluated whether the number of different Tn*4401*-TSS types was different across

strains, noting that the observed frequency of different Tn*4401*-TSS combinations

was higher in *K. pneumoniae* strain 11 (kpne11, ST11) than in other strains (Fig.S10).

Kpne11 represented two sub-clusters, separated by ~40SNVs (Fig.7, designated

"cluster 1" and "cluster 2"), both of which were isolated from patients and wastewater

sites across ward settings, including the cardiac unit after plumbing replacement.

Within both clusters of isolates there were ten different Tn*4401*a flanking signatures

observed, only one of which was seen in the other >1700 isolates sequenced in the

study (Tn*4401*a-1_ATTGA_ATTGA, the most common flanking signature), and

therefore most strongly consistent with the occurrence of multiple transposition

events within this *K. pneumoniae* strain. Similarly plasmid replicon gain/loss events

were observed (Col, IncN, IncR and IncFIA); all isolates contained IncFIB and IncFII

replicons.

DISCUSSION

This work highlights the substantial CPE diversity and flux that can occur in both

human and wastewater niches in healthcare settings. Many CPE strains have unique

features likely to make generalising results from analyses of transmission, evolution

and gene sharing of bla_{KPC} amongst these species very challenging. Although some

sites are persistently colonised with individual strains over a period of months, many

appear to be only transiently colonised. Sites particularly associated with CPE

colonisation included sluices, sluice sinks, mop sinks and shower/bath drains,

supported by other studies[5], and consistent with these representing a major interface

between human faeces and premise plumbing. Toilets were less likely to be CPE-

positive, perhaps because of regular flushing or particular cleaning protocols;

consistent with this, several studies have demonstrated that daily bleach application

decontaminated wastewater sites[32, 33].

At the strain-level, likely transmissions occurred from patient-patient, environment-

patient, patient-environment and environment-environment; however, our study

highlights the difficulty in accurately tracking strains across niches, and the detailed

and dense sampling effort required to understand within-sample diversity and

longitudinal flux. This difficulty has also been demonstrated in other recent work

investigating bla_{OXA-48}, where only patient-based sampling was undertaken and most

bla_{OXA-48}-associated transmission events either remained non-delineated or were only

weakly supported by joint phylogenetic/epidemiological inference[8]. Our study clearly

shows that different species and lineages represent different epidemiological risks and

have their own behaviours, and a focus on single, clonal strains as mediators of AMR

gene transmission is not relevant in this and potentially many other settings where

important AMR genes have become endemic.

The importance of evaluating asymptomatic CPE carriage, characterising within-host

diversity of strains, considering rapid acquisition/loss of strains over short

timeframes, and including environmental sites in transmission networks has been

partly demonstrated in previous studies investigating ESBL-*E. coli*[10, 34] and

Enterococcus spp.[35], but generally these features are not jointly investigated because

doing so is highly resource-intensive. However, without this effort, such as in studies

which consider only single or infrequent sampling timepoints, and do not investigate

asymptomatic colonisation or consider within-site diversity[36], our study highlights

that relevant events may easily be missed.

There are several limitations to this work. We undertook IPC-led real-time collection

patient isolates; sample and data capture for this process was not 24/7, and we

therefore only evaluated a subset of all the patient-associated CPE isolates. We were

not resourced to perform long-read sequencing to enable plasmid sequence

reconstruction, and plasmid replicon or bla_{KPC}-contig profiling is relatively low-

resolution[21]; we therefore avoided a detailed plasmid analysis which would be highly

relevant to evaluating horizontal bla_{KPC} transmission. However, even with relatively

low-resolution approaches, the diversity in bla_{KPC} contexts was substantial,

supporting high rates of HGT. A further limitation is the absence of metadata

capturing relevant selection pressures in each niche, including cleaning protocols for

environmental reservoirs and drug prescription/antimicrobial stewardship data for

patients. Although we only investigated one healthcare setting, the polyspecies,

polyclonal phenomenon we observed has been seen in other centres and with other

carbapenemase genes[4,8]. We focused only on Enterobacterales and may have missed

wider HGT of carbapenemase genes to other bacterial orders. In line with this, at least

23 *Pseudomonas* spp., five *Stenotrophomonas maltophilia* and two *Acinetobacter* spp.

that were PCR-positive for bla_{KPC} or bla_{NDM} were inadvertently picked up on rectal

screening.

Further optimisation of methods to evaluate multi-level genetic transmission and

within-sample diversity without individual colony-level characterisation is needed.

Holistic approaches using fully reconstructed chromosome/plasmid assemblies to

quantify the diverse types of genomic transmission (including clonal transmission,

horizontal transfer at the gene, transposon and plasmid-level) are also required to

better understand what facilitates emergence, selection and persistence of different

lineages. This information will improve our understanding of how to intervene to

limit the transmission of drug-resistant Enterobacterales in healthcare settings, and

avoid incorrect inference.

Declaration of interests

Acknowledgements

We are grateful to the microbiology staff and infection prevention and control teams at Manchester University NHS Foundation Trust, and to Alvaro San Millán Ricardo León-Sampedro for helpful discussion.

This study was funded by the National Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance (NIHR200915), a partnership between the UK Health Security Agency (UKHSA) and the University of Oxford, and was supported by the NIHR Oxford Biomedical Research Centre (BRC). The computational aspects of this research were funded from the NIHR Oxford BRC with additional support from the Wellcome Trust Core Award Grant Number 203141/Z/16/Z. The views expressed are those of the author(s) and not necessarily those of the NHS, NIHR, UKHSA or the Department of Health and Social Care. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

DWE is a Robertson Foundation Fellow and an NIHR Oxford Biomedical Research Centre Senior Fellow. NS is an Oxford Martin Fellow and an NIHR Oxford Biomedical Research Centre Senior Fellow. DWC, TEAP and ASW are NIHR Senior Investigators.

The Transmission of Carbapenemase-producing Enterobacteriaceae (TRACE)

Investigators' Group is made up of the following (listed alphabetically, includes

several of the authors also listed by name in the main author list): Zoie Aiken,

Oluwafemi Akinremi, Aiysha Ali, Julie Cawthorne, Paul Cleary, Derrick W. Crook,

Valerie Decraene, Andrew Dodgson, Michel Doumith, Matthew J. Ellington, Ryan

George, John Grimshaw, Malcolm Guiver, Robert Hill, Katie L. Hopkins, Rachel

Jones, Cheryl Lenney, Amy J. Mathers, Ashley McEwan, Ginny Moore, Andrew

Mumford, Mark Neilson, Sarah Neilson, Tim E.A. Peto, Hang T.T. Phan, Mark

Regan, Anna C. Seale, Nicole Stoesser, Jay Turner-Gardner, Vicky Watts, A. Sarah

Walker, Jimmy Walker, David H. Wyllie, William Welfare and Neil Woodford.

REFERENCES

1.   Munoz-Price LS, Poirel L, Bonomo RA et al. Clinical epidemiology of the

global expansion of Klebsiella pneumoniae carbapenemases. *Lancet Infect Dis* 2013;

13: 785-96.

2.   Logan LK, Weinstein RA. The Epidemiology of Carbapenem-Resistant

Enterobacteriaceae: The Impact and Evolution of a Global Menace. *J Infect Dis* 2017;

215: S28-S36.

3.   Nordmann P, Dortet L, Poirel L. Carbapenem resistance in

Enterobacteriaceae: here is the storm! *Trends Mol Med* 2012; **18**: 263-72.

4.   Mathers AJ, Cox HL, Kitchel B et al. Molecular dissection of an outbreak of

carbapenem-resistant enterobacteriaceae reveals Intergenus KPC carbapenemase

transmission through a promiscuous plasmid. *MBio* 2011; **2**: e00204-11.

5.   Kizny Gordon AE, Mathers AJ, Cheong EY et al. Is the hospital water

environment a reservoir for carbapenem-resistant organisms causing hospital-acquired

infections? A systematic review of the literature. *Clin Infect Dis* 2017.

6.   Bleichenbacher S, Stevens MJA, Zurfluh K et al. Environmental

dissemination of carbapenemase-producing Enterobacteriaceae in rivers in

Switzerland. *Environ Pollut* 2020; **265**.

7.   Cahill N, O'Connor L, Mahon B et al. Hospital effluent: A reservoir for

carbapenemase-producing Enterobacterales? *The Science of the total environment*

2019; **672**: 618-24.

8.   León-Sampedro R, DelaFuente J, Díaz-Agero C et al. Pervasive transmission

of a carbapenem resistance plasmid in the gut microbiota of hospitalized patients. *Nat*

Microbiol 2021; **6**: 606-16.

9.   Weingarten RA, Johnson RC, Conlan S et al. Genomic Analysis of Hospital

Plumbing Reveals Diverse Reservoir of Bacterial Plasmids Conferring Carbapenem

Resistance. *MBio* 2018; **9**.

10.   Stoesser N, Sheppard AE, Moore CE et al. Extensive Within-Host Diversity in

Fecally Carried Extended-Spectrum-Beta-Lactamase-Producing Escherichia coli

Isolates: Implications for Transmission Analyses. *J Clin Microbiol* 2015; **53**: 2122-

31.

11.   Ludden C, Coll F, Gouliouris T et al. Defining nosocomial transmission of

Escherichia coli and antimicrobial resistance genes: a genomic surveillance study. *The*

Lancet Microbe 2021; **2**: e472-e80.

12.   Decraene V, Phan HTT, George R et al. A Large, Refractory Nosocomial

Outbreak of Klebsiella pneumoniae Carbapenemase-Producing Escherichia coli

Demonstrates Carbapenemase Gene Outbreaks Involving Sink Sites Require Novel

Approaches to Infection Control. *Antimicrob Agents Chemother* 2018; **62**.

13.   Public Health England. Carbapenemase-producing Enterobacteriaceae:

laboratory confirmed cases, 2003 to 2015. London: Public Health England, 2016.

14.   Public Health England. Carbapenemase-producing Enterobacteriaceae: early

detection, management and control toolkit for acute trusts. 2014.

15.   UK Standards for Microbiology Investigations.

https://www.gov.uk/government/collections/standards-for-microbiology-

investigations-smi. Last updated: 1st July 2021.

16.   The European Committee on Antimicrobial Susceptibility Testing. Breakpoint

tables for interpretation of MICs and zone diameters. 2017.

17.   Stoesser N, Giess A, Batty EM et al. Genome sequencing of an extended

series of NDM-producing Klebsiella pneumoniae isolates from neonatal infections in

a Nepali hospital characterizes the extent of community- versus hospital-associated

transmission in an endemic setting. *Antimicrob Agents Chemother* 2014; **58**: 7347-57.

18.   Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence

classification using exact alignments. *Genome Biol* 2014; **15**: R46.

19.   Nguyen LT, Schmidt HA, von Haeseler A et al. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*

2015; **32**: 268-74.

20.   Didelot X, Wilson DJ. ClonalFrameML: efficient inference of recombination

in whole bacterial genomes. *PLoS Comput Biol* 2015; **11**: e1004041.

21.   Sheppard AE, Stoesser N, Wilson DJ et al. Nested Russian Doll-Like Genetic

Mobility Drives Rapid Dissemination of the Carbapenem Resistance Gene blaKPC.

Antimicrob Agents Chemother 2016; **60**: 3767-78.

22.   Tonkin-Hill G, MacAlasdair N, Ruis C et al. Producing polished prokaryotic

pangenomes with the Panaroo pipeline. *Genome Biol* 2020; **21**: 180.

23.   Bankevich A, Nurk S, Antipov D et al. SPAdes: a new genome assembly

algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012; **19**: 455-

77.

24.   Hunt M, Mather AE, Sánchez-Busó L et al. ARIBA: rapid antimicrobial

resistance genotyping directly from sequencing reads. *Microb Genom* 2017; **3**:

e000131.

25.   Jia B, Raphenya AR, Alcock B et al. CARD 2017: expansion and model-

centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids*

Res 2017; **45**: D566-D73.

26.   Sheppard AE, Stoesser N, German-Mesner I et al. TETyper: a bioinformatic

pipeline for classifying variation and genetic contexts of transposable elements from

short-read whole-genome sequencing data. *Microb Genom* 2018; **4**.

27.   De Maio N, Wu CH, Wilson DJ. SCOTTI: Efficient Reconstruction of

Transmission within Outbreaks with the Structured Coalescent. *PLoS Comput Biol*

2016; **12**: e1005130.

28.   Oksanen JB, Blanchet, FG, Friendly M, Kindt R, Legendre P, McGlinn D,

Minchin PR, O'Hara RB, Simpson GL, Solymos P, Henry M, Stevens H, Szoecs E,

Wagner H. vegan: Community Ecology Package. 2020.

29.   Harrell Jr FE. rms: Regression Modeling Strategies. 2021.

30.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016.

31.   Yu G. scatterpie: Scatter Pie Plot. 2020.

32.   Valentin AS, Santos SD, Goube F et al. A prospective multicentre surveillance

study to investigate the risk associated with contaminated sinks in the intensive care

unit. *Clin Microbiol Infect* 2021; **27**: 1347.e9-.e14.

33.   Heireman L, Hamerlinck H, Vandendriessche S et al. Toilet drain water as a

potential source of hospital room-to-room transmission of carbapenemase-producing

Klebsiella pneumoniae. *The Journal of hospital infection* 2020; **106**: 232-9.

34.   Kantele A, Kuenzli E, Dunn SJ et al. Dynamics of intestinal multidrug-

resistant bacteria colonisation contracted by visitors to a high-endemic setting: a

prospective, daily, real-time sampling study. *The Lancet Microbe* 2021; **2**: e151-e8.

35.   Gouliouris T, Coll F, Ludden C et al. Quantifying acquisition and transmission

of Enterococcus faecium using genomic surveillance. *Nat Microbiol* 2021; **6**: 103-11.

36.   Qiao F, Wei L, Feng Y et al. Handwashing Sink Contamination and

Carbapenem-resistant Klebsiella Infection in the Intensive Care Unit: A Prospective

Multicenter Study. *Clin Infect Dis* 2020; **71**: S379-s85.

TABLES

Table 1. Association of site type and unit location with bla_{KPC}-Enterobacterales (KPC-E) positivity in environmental samples. Outputs of a multivariable logistic regression model with robust standard errors clustered by environmental site type and unit location; no other variables were considered in the model. P-values <0.05 were considered significant and are highlighted in bold. No interactions with an interaction Wald p <0.05 were observed.

Environmental site	Number of sites	Number of sampling events KPC-E-positive (% of total sampling events)	Number of sampling events KPC-E-negative	OR (95% CI)	p
Environmental site type					
Clinical handwash basin drain	96	79 (5.9%)	1270	Reference	
Handwash basin drain	85	52 (4.6%)	1077	0.99 (0.51-1.93)	0.9735
Shower/bath drain	59	100 (19.2%)	421	2.69 (1.44-5.01)	**0.0019**
Sluice/sluice	16	47 (17.9%)	215	2.60 (1.04-	**0.0410**

sink/mop sink				6.52)	
Toilet	74	5 (0.5%)	931	0.28 (0.10-0.78)	**0.0152**
Utility sink	19	36 (12.4%)	255	1.32 (0.47-3.68)	0.5965
Environmental site location					
Acute medicine unit	129	86 (6.6%)	1223	Reference	
Cardiac unit	76	111 (7.0%)	1453	1.07 (0.57-2.00)	0.8412
Geratology unit	144	122 (7.6%)	1493	0.88 (0.52-1.48)	0.6243

	FIGURE LEGENDS

Figure 1. Sampling and sequencing flowchart for the study.

Figure 2. bla_{KPC}-Enterobacterales (KPC-E)-positive environmental sites by week

in 2016 and hospital unit. The number of environmental site samples performed and

KPC-E-positive samples by week of sampling, stratified by unit (top three panels),

and the proportion of KPC-E-positive sites by week of sampling, again stratified by

unit (bottom panel).

Figure 3. Carbapenem-susceptible Enterobacterales (CSE) and carbapenem-

resistant-Enterobacterales (CRE) patient cultures by week in 2016 and hospital

unit. Top three panels show counts of Enterobacterales culture-positive samples

stratified by unit and carbapenem susceptibility; bottom panel shows the proportion of

all cultures taken that were CRE, again stratified by unit.

Figure 4. Number of unique patient/environmental niches colonised by common

bla_{KPC} Enterobacterales species/strains. A) Includes species/strains identified in

both patients and wastewater niches during the study (if >50 isolates of a species were

identified [see Methods]). Orange circles denote wastewater niches, black circles

unique patients. B) Includes species/strains identified only in patients (left panel) and

only in wastewater sites (right panel). For *Raoultella ornithinolytica*, six strains were

identified and these were observed in wastewater sites only (data not shown).

Figure 5. Top panel: Within-patient KPC-E diversity. Pies reflect the proportion of

strains within a sample that were identified using whole genome sequencing, with

evaluation of diversity at the species-strain level. "cfre" = *Citrobacter freundii*, "eclo"

= *Enterobacter cloacae*, "ecol" = *Escherichia coli*, "evul" = *Escherichia vulnificus*,

"koxy" = *Klebsiella oxytoca*, "kpne"= *Klebsiella pneumoniae*, "lecl" = *Leclercia*

adecarboxylata. Numbers after species abbreviations denote unique strains. Bottom

panel: Within-wastewater sample KPC-E diversity. Pies reflect the proportion of

strains within a sample that were identified using whole genome sequencing, with

evaluation of diversity at the species-strain level. "aher" = *Escherichia hermanii*,

"cfar" = *Citrobacter farmeri*, "easb" = *Enterobacter asburiae*, "pant" = *Pantoea* spp.,

"plur" = *Pluralibacter gergoviae*; other assignations as for top panel.

Figure 6. A) Recombination-adjusted phylogeny for *E. cloacae* strain 10. Main

annotations as described in the legends as part of the figure. In the panel to the right

of the phylogeny, AMR profile, IS profile, plasmid replicon profile and Tn*4401*/target

site sequences (TSSs) are represented, with different colours distinguishing different

versions of each. Three isolates in F47 and one isolate in F46 clustered with the F45-

associated isolates; sampling dates suggest that these represented transmission events

from F45. The scale bar is in SNVs. B) Represents plasmid replicon profiles in the

isolates obtained from the environmental site samples over time, and C) similarly

represents the Tn*4401*/TSS profiles with additional deletions observed annotated as

an overlay for isolates typed as containing the "unknown-1_ATTGA_ATTGA

variants" using TETyper.

Figure 7. Putative transmission network for sub-lineages of *K. pneumoniae* strain 9,

defined by accessory genome clustering (clusters 1, 2, 3a-h). Horizontal bars

represent admission episodes to study wards for each patient (pale blue=W45,

geratology; pale pink=AM1, acute medicine; pale purple=AM2, acute medicine; A13

represents the only shower site involved. Empty black triangles denote CPE-negative

rectal screens on patients. Arrows denote possible transmission links (dashed arrows

denote events that have no geographic overlap or could be associated with two links

based on negative screen, geographic location and timing). For clarity, arrows are not

drawn for clusters 3g and 3h, which affected the most patients - these are shown as an

inset panel at the bottom right.

Figure 8. Left-hand panel shows a recombination-corrected phylogeny for *K.*

pneumoniae strain 11, designating the two identified clusters, 1 and 2, and annotated

with isolate names, unit, collection date, niche, and Tn*4401*-TSS type. Right-hand

panels reflect the plasmid replicon and Tn*4401*-TSS diversity within sequenced

isolates in given niches over time for each chromosomally genetically-related cluster,

highlighting features consistent with clonal transmission, Tn*4401* transposition events

within isolates (Tn*4401*a, change in inner dot colour with consistent outer ring

colour), deletions in Tn*4401*a (to "unknown-a"), plasmid replicon loss/gain events

(change in outer ring colour).

Wastewater sites cultured
n=349
Clinical handwash basin drains n=96
Handwash basin drains n 85=
Shower/bath drains n=59
Sluices/sluice sink/mop sinks n=16
Utility sinks n=19
Toilets n=74

Wastewater samples
Total cultures - n=4,488
CRE-positive -n=319 (7.1%)

Patients cultured
n=4,425
Patients ever CRE-positive 183 (4.1%)

Patient samples
Total cultures n=41,203
Culture-positive n=5,154 (12.5%)
Enterobacterales-positive n= 2,172
(2,000 with carbapenem susceptibility results) (42.1%)
CRE-positive n=399 (20.0% of Enterobacterales; 1.0% of total)

Total isolates sequenced
n=1452

Total isolates sequenced
n=482

Excluded
n=177
Not Enterobacterales
Mixed/poor quality
Replicates

Excluded
n=25
Not Enterobacterales
Mixed/poor quality
Replicates

Total wastewater isolate sequences available for analysis
n=1275

Enterobacter cloacae - n=536 isolates
Klebsiella pneumoniae - n=463 isolates
Citrobacter freundii - n=133 isolates
Raoultella ornithinolytica - n=63 isolates
Enterobacter asburiae - n=15 isolates
Citrobacter farmeri - n=14 isolates
Klebsiella oxytoca - n=13 isolates
Escherichia coli - n=12 isolates
Pluralibacter gergoviae - n=10 isolates
Leclercia adecarboxylata - n=9 isolates
Kluyvera intermedia - n=6 isolates
Pantoea spp. - n=1 isolate

Total patient isolate sequences available for analysis
n=457

Klebsiella pneumoniae - n=329 isolates
Escherichia coli - n=64 isolates
Enterobacter cloacae - n=28 isolates
Klebsiella oxytoca - n=22 isolates
Escherichia vulneris - n=7 isolates
Citrobacter freundii - n=4 isolates
Leclercia adecarboxylata - n=3 isolates

A *E. cloacae* strain 10 Geratology ward

F46

F47

F45

AMR profile
IS profile
Plasmid profile
Tn4401 type

location ● CHWB ● Utility_sink

B Probable IncHI2/HI2A plasmid gain/loss events

Sample location code: F47, F46, F45

Sample collection date: Aug, Sep, Oct, Nov, Dec

Plasmid replicon profile
a IncFIB|IncFII
a IncFIB|IncFII|IncHI2|IncHI2A

C Multiple deletion events in the Tn4401 transposon (denoted by black arrows)

Sample location code: F47, F46, F45

Sample collection date: Aug, Sep, Oct, Nov, Dec

tn4401type
a Tn4401a-1_ATTGA_ATTGA
a unknown-1_ATTGA_ATTGA