

1 **Employment status and its associated factor for**
2 **patients 12 months after intensive care: Secondary**
3 **analysis of the SMAP-HoPe-study**

4

5 Short title: Employment status 12 months after Intensive Care

6

7 Takeshi Unoki^{1*}, Mio Kitayama², Hideaki Sakuramoto³, Akira Ouchi³, Tomoki

8 Kuribara¹, Takako Yamaguchi⁴, Sakura Uemura⁵, Yuko Fukuda⁶, Junpei

9 Haruna⁷, Takahiro Tsujimoto⁸, Mayumi Hino⁹, Yuko Shiba¹⁰, Takumi

10 Nagao¹¹, Masako Shirasaka¹², Yosuke Sato¹³, Miki Toyoshima⁵, Yoshiki

11 Masuda¹⁴, on behalf of the SMAP-HoPe Study Project ^

12

13 1. Department of Acute and Critical Care Nursing, School of Nursing, Sapporo

14 City University, Chuo-ku, Sapporo, Hokkaido, Japan

15 ² Nursing Department Heart Center, Kanazawa Medical University Hospital,

16 Uchinada, Ishikawa, Japan

17 ³ Department of Adult Health Nursing, College of Nursing, Ibaraki Christian

18 University, Hitachi, Ibaraki, Japan

19 ⁴ Intensive Care Unit, Nippon Medical School Musashikosugi Hospital,

20 Kawasaki, Kanagawa, Japan

21 ⁵ Emergency and Critical Care Medical Center, Osaka City General Hospital,

22 Osaka, Japan

23 ⁶ Intensive Care Unit, Jichi Medical University Hospital, Yakushiji Shimotsuke-
24 shi, Tochigi, Japan

25 ⁷ Intensive Care Unit, Sapporo Medical University Hospital, Sapporo, Hokkaido,
26 Japan

27 ⁸ Nursing Practice and Career Support Center, Nara Medical University
28 Hospital, Kashihara City, Nara, Japan

29 ⁹ Intensive Care Unit, Tohoku Medical and Pharmaceutical University Hospital,
30 Sendai, Miyagi, Japan

31 ¹⁰ Intensive Care Unit, University of Tsukuba Hospital, Tsukuba, Ibaraki, Japan

32 ¹¹ Intensive Care Unit, Sakakibara Heart Institute, Fuchu-shi, Tokyo, Japan

33 ¹² Intensive Care Unit & Cardiac Care Unit, Japanese Red Cross Fukuoka
34 Hospital, Fukuoka, Japan

35 ¹³ Intensive Care Unit, Naha City Hospital, Naha, Okinawa, Japan

36 ¹⁴ Department of Intensive Care Medicine, Sapporo Medical University Hospital,
37 Sapporo, Hokkaido, Japan

38

39 *Corresponding author

40 E-mail: t.unoki@scu.ac.jp (TU)

41

42

43 [^]Membership of the SMAP-HoPe Study Project member is provided in the
44 Acknowledgements.

45

46 **Competing Interests:** The authors have declared that no competing interests
47 exist.

48

49 **Financial Disclosure:** TU received JSPS KAKENHI Grant Number 19K10929.
50 The funders had no role in study design, data collection and analysis, decision
51 to publish, or preparation of the manuscript.

52

53 **Data availability statement:** All relevant data are within the manuscript and
54 Supporting Information files.

55

56

57

58 **Abstract**

59 Returning to work is a serious issue that affects patients who are being
60 discharged from the intensive care unit (ICU). This study aimed to clarify the
61 employment status and the perceived household financial status of ICU patients
62 12 months following discharge from the ICU. Additionally, a hypothesis of
63 whether depressive symptoms were associated with subsequent unemployment
64 status was tested. This study was a subgroup analysis using data from the
65 published Survey of Multicenter Assessment with Postal questionnaire for Post-
66 Intensive Care Syndrome (PICS) for Home Living Patients (the SMAP-HoPe
67 study) in Japan. The patients included those who had a history of staying in the
68 ICU for at least three nights and had been living at home for one year following
69 discharge, between October 2019 and July 2020. We assessed employment
70 status, subjective cognitive functions, household financial status, Hospital
71 Anxiety and Depression Scale scores, and EuroQOL-5 dimensions of physical
72 function at 12 months following intensive care. This study included 328 patients
73 who were known to be employed prior to ICU admission. The median age was
74 64 (Interquartile Range [IQR] 52-72), and males were predominant
75 (86%). Seventy-nine (24%) of those evaluated were unemployed. The number
76 of patients who reported worsened financial status was significantly higher in
77 the unemployed group. ($p < .01$) Multivariate analysis showed that higher age
78 (Odds Ratio [OR]: 1.06, 95% Confidence Interval [CI]: 1.03-1.08) and severity
79 of depressive symptoms (OR: 1.13 [95% CI: 1.05-1.23]) were independent
80 factors for employment status after 12 months from being discharged from the

81 ICU. These factors were determined to be significant even after adjusting for
82 sex, physical function, and cognitive function. We found that one-fourth of our
83 patients who had been employed prior to ICU admission were subsequently
84 unemployed 12 months following ICU discharge. Additionally, depressive
85 symptoms were associated with unemployment status. The government and the
86 local municipalities should provide medical and financial support to such
87 patients. Additionally, community support for such patients is warranted.

88

89 **Keywords:**

90 Intensive Care Units, Employment, Depression, Critical Care

91

92 Introduction

93

94 The problem of patients returning to work following discharge from the
95 intensive care unit (ICU) is a serious issue. A systematic review and meta-
96 analysis, including 52 studies, of returning to work among previously critically ill
97 patients indicated that delayed return to work and unemployment were common
98 and persistent problems[1]. These studies showed that 36% of patients were
99 subsequently unemployed at 12 months following ICU admission. This change
100 in employment status shall have a corresponding effect on the household
101 income. A study suggested that 30% of patients discharged from the ICU had a
102 decline in household income even after 6 months to one year following
103 intensive care admission in the United Kingdom[2].

104 National employment and disability policies could contribute to
105 resumption of work productivity among those discharged from the ICU. Su et
106 al. [3] revealed that the disability policies of each country were related to
107 resumption of work productivity. Thus, it is worth evaluating how those
108 discharged from the ICU fare in terms of employment in each country. Notably,
109 there was a prior study examining the employment status of those discharged
110 from the ICU in Japan.[4] They showed that among 33 patients discharged
111 from the ICU, 26 (83.9%) were able to return to work six months following ICU
112 admission. However, the precision was insufficient due to a small sample
113 size. For enhanced understanding of employment status among those

114 discharged from the ICU, a study with a larger sample size was therefore
115 warranted.

116 The factors associated with the unemployment of those discharged
117 from the ICU had been examined in previous studies. Higher age and female
118 [5], cognitive function [6], depressive symptoms [7], physical disability and
119 educational level [8] had been suggested to influence unemployment status in
120 those discharged from the ICU. However, these factors are not consistent
121 between studies[1]. Among the general population, absence due to all sickness
122 and depression are key factors influencing a longer duration until return to work
123 [9].

124 The primary objective of our study was to elucidate the employment
125 status and perceived household financial status among those discharged from
126 the ICU 12 months following discharge. The secondary objective was to test a
127 hypothesis of whether depressive symptoms were associated with subsequent
128 employment status following discharge from intensive care.

129

130 **Materials and methods**

131 **Study design**

132

133 The present study was a sub analysis of the the Survey of Multicenter
134 Assessment with Postal questionnaire for Post-Intensive Care Syndrome
135 (PICS) for Home Living Patients (SMAP-HoPe) study [10] and focused on

136 employment status following ICU discharge and the factors that affect it. This
137 analysis was nested in the SMAP-HoPe study [10]. In summary of SMAP-HoPe
138 study, we conducted an ambidirectional study for patients living at home 12
139 months following ICU discharge. We sent postal surveys on post-intensive care
140 syndrome and employment status. Data from when the patients were admitted
141 were obtained for retrospective analysis. Detailed methods were provided
142 elsewhere in a distinct publication [10].

143

144 **Setting**

145

146 Twelve ICUs in Japan were included in this study. The detailed
147 characteristics of each ICU included in the study were previously reported in the
148 original study [10].

149

150 **Participants**

151

152 The patients included those who had stayed in the ICU for at least
153 three nights and had been living at home for one year following ICU discharge
154 between October 2019 and July 2020. The number of subjects in the previously
155 published study of which this is a subgroup analysis was 754 patients; the
156 response rate was 91.1%. In this secondary analysis focusing on employment

157 status, only patients aged 18 years or over and who had been working prior to
158 admission were included.

159

160 **Variables/Instruments/Data source**

161

162 Patients' characteristics including age, sex, diagnosis, APACHE II
163 score, pre-existing disease, and length of ICU stay were retrospectively
164 collected by medical chart review. Status after 12 months following discharge
165 from intensive care was collected through postal questionnaires, which explored
166 work status, cognitive function, and Euro-QOL-5D-5L (EQ-5D-5L). [11][11] EQ-
167 5D-5L is a validated questionnaire [11] and Japanese version is available upon
168 request to the EuroQOL group (<https://euroqol.org>). Work status prior to ICU
169 admission and present working status were classified according to the following
170 scheme: unemployed, self-employed, employed part-time, and employed full-
171 time. Additionally, we inquired about whether household finances changed
172 compared with the period prior to ICU admission. The respondents selected
173 between a three-point scale of "worse," "no change," or "better." Cognitive
174 function was measured using two simple questions that we developed, due to
175 lack of valid instruments for self-administration. One was "Do you think your
176 memory function was impaired compared with before ICU admission?" Another
177 was "Do you think your concentration function was impaired compared with
178 before ICU admission?" Four-point Likert scales, from not at all (0), sometimes
179 (1), frequently (2), and very frequently (3) were the acceptable responses. If the

180 subject responds “very frequently” or “frequently” for either of the two questions,
181 we define them to have cognitive dysfunction. Usual activities were measured
182 by part of question from EQ-5D-5L. The response consisted of five levels, from
183 having “no problems” performing usual activities to being “unable to do” their
184 usual activities. We defined responses other than “no problems” or “slight
185 problems” as evidence of physical dysfunction. Severity of depression was
186 measured by the Hospital Anxiety and Depression Scale (HADS) [12]. HADS
187 consists of depression and anxiety subscales, and each subscale has seven
188 components, which were rated on a scale from 0 to 3. Subscales of depression
189 are graded from 0 to 21. Previous studies demonstrated that the Japanese
190 version of HADS had good reliability and validity [13]. We used Japanese
191 version of HADS [13].

192

193 **Bias**

194

195 This study was a postal survey and selection bias may exist; however,
196 selection bias was minimum, given the high rate of response from the original
197 study from which this one was nested. Additionally, a state of emergency was
198 declared by the government due to the coronavirus disease 2019 (COVID-19)
199 pandemic in April 2020, which overlapped with the study period. This
200 declaration may have had an impact on employment. This bias was assessed
201 through sensitivity analysis. Furthermore, in many Japanese companies, the
202 compulsory age of retirement was generally between 60 and 65 years. The

203 compulsory retirement, other than the episodes of intensive care admission,
204 may influence our results. Thus, we accordingly conducted sensitivity analysis
205 to evaluate its effect on our findings.

206

207 **Sensitivity Analysis**

208

209 We conducted sensitivity analyses to confirm whether our findings
210 were robust. First, we calculated the percentage of unemployment after
211 excluding the subjects who responded within the duration of the COVID-19
212 pandemic; employment status after 12 months following ICU discharge may be
213 influenced by the COVID-19 pandemic. Nevertheless employment status before
214 ICU admission was collected between September 2018 and July 2019. We
215 defined the period after April 2020 as the start of the COVID-19 pandemic
216 because a state of emergency had been declared across the country on April
217 16, 2020. We calculated the proportion of the unemployed after excluding from
218 the analysis the subjects who posted their responses during this
219 period. Second, 60 years old was the beginning of eligibility for retirement from
220 employment [14]. Thus, we performed multivariable analysis after excluding the
221 subjects who were 60 years old and above and reviewed the impact of
222 retirement age on our findings.

223

224 **Statistical analysis**

225

226 First, descriptive statistical analyses were performed. Continuous
227 variables were expressed as median [interquartile range (IQR)]. Wilcoxon rank-
228 sum test was used for the comparison of continuous variables. Chi-square test
229 was used for categorical variables. We calculated the proportion of unemployed
230 subjects stratified by age. For multivariable analysis, a multilevel generalized
231 linear model (GLM) with a binomial distribution and log link was performed. We
232 pre-defined covariates consisting of work status before ICU admission, age [5],
233 sex [5], cognitive dysfunction [6], physical dysfunction [7], and severity of
234 depression [8] based on previous studies. Statistical analysis was used for
235 STATA IC ver. 16 (Statacorp, TX) and R 4.0.2 (The R foundation for Statistical
236 Computing), and the significance difference was set $< .05$.

237

238 **Ethical considerations**

239

240 Human Research Ethics Committee approval was obtained. (Sapporo
241 City University, Sapporo, Hokkaido, Japan, approval number: 1927-1)
242 Additionally, ethical approval was obtained from all sites for the original studies.
243 Detailed informed consent process was provided in the original study [10].

244

245 **Results**

246

247 **Participants**

248

249 Fig 1 displays a flow diagram of study enrollment. The SMAP-HoPe
250 study had 754 subjects. We excluded 425 subjects who were unemployed
251 before ICU admission. One patient who did not provide an answer to the
252 question of cognitive function was excluded. Thus, 328 subjects were analyzed
253 in this study.

254

255 **Fig 1. Patient recruitment scheme**

256

257 **Population of the study**

258

259 The characteristics of the population are shown in Table 1. The
260 median age of the population was 64 [IQR: 52-72] years. The number of
261 subjects in each age group was shown in S1 Table. Male was dominant
262 (86%). Approximately 41.2% of the subjects were admitted to the ICU after
263 undergoing elective surgery. The number of subjects receiving mechanical
264 ventilation was 219 (66.8%), and the median duration of mechanical ventilation
265 were 2 days. Of 328 employed subjects before ICU admission, the most

266 frequent employment status was part-time employed in 125 (50.2%) following
267 full-time employed in 86 (34.5%)

268

269 **Characteristics between employed and unemployed** 270 **subjects**

271

272 Among 328 subjects who were previously employed, 79 (24.1%, 95%
273 CI 19.7-29.1) were unemployed at 12 months following ICU discharge. A
274 comparison of patient characteristics between the employed and the
275 unemployed subjects is shown in Table 1. The median age of those who were
276 employed was significantly lower than that of those who were unemployed (62
277 [50-71] vs 69 [62-75], $p < 0.01$, respectively). The median APACHE II score
278 among employed subjects was lower than that among unemployed subjects (13
279 [9-19] vs 16 [13-19], $p < 0.001$, respectively).

280

281 **Table 1. Demographic characteristics of employed and unemployed**
282 **subjects from the SMAP-HoPe study**

Variables	Participant s n=328	Employed n=249	Unemployed n=79	p-value
Age (years), median [IQR]	64 [52-72]	62 [50-71]	69 [62-75]	<.001
Female, n (%)	46 (14.0)	32 (12.9)	14 (17.7)	.271

Types of admission, n (%)				
Elective surgery	149 (41.2)	109 (43.8)	37 (46.2)	.697
Unscheduled admission	182 (55.5)	140 (56.2)	42 (53.2)	.697
Unscheduled surgery	49 (14.9)	35 (14.1)	14 (17.7)	.469
Reason for ICU admission, n (%)				
CV surgery	122 (37.2)	87 (34.9)	35 (44.3)	.183
CHF/AMI/Arrhy	57 (17.4)	46 (18.5)	11 (13.9)	
Sepsis	34 (10.4)	21 (8.4)	13 (16.5)	
Abdominal surgery	30 (9.1)	25 (10.0)	5 (6.3)	
Other surgery	14 (4.3)	9 (3.6)	5 (6.3)	
Trauma	14 (4.3)	13 (5.2)	1 (1.3)	
ENT surgery	12 (3.7)	10 (4.0)	2 (2.5)	
Aortic dissection (non-operative)	12 (3.7)	11 (4.4)	1 (1.3)	
Acute renal failure	12 (3.7)	9 (3.6)	3 (3.8)	
Others	21 (6.4)	18 (7.2)	3 (3.8)	
Employment status before ICU admission, n (%)				
Full time	101 (30.8)	86 (34.5)	15 (19.0)	<.001

Part time	156 (47.6)	125 (50.2)	31 (39.2)	
Self	71 (21.6)	38 (15.3)	33 (41.8)	
APACHE II , median [IQR]	14 [10-19]	13 [9-19]	16 [13-19]	.001
MV use, n (%)	219 (66.8)	157 (63.1)	62 (78.5)	.013
MV (days), median [IQR]	2.0 [0.0-3.0]	2.0 [0.0-3.0]	2.0 [1.0-3.0]	.041
Psychological history, n (%)	4 (1.2)	4 (1.6)	0 (0.0)	.576
Delirium (days), median [IQR]	0.0 [0.0-1.0]	0.0 [0.0-1.0]	0.0 [0.0-2.0]	.094
ICU LOS (days), median [IQR]	5.0 [4.0-7.0]	5.0 [4.0-7.0]	5.0 [4.0-7.0]	.293
Hospital LOS (days), median [IQR]	26.0 [18.0-36.0]	25.0 [18.0-34.0]	27.0 [19.0-46.5]	.192

283 IQR, interquartile range; CV, cardiovascular; CHF/AMI/arrhy, congestive heart
284 failure/ acute myocardial infarction/ arrhythmia; ENT, ear nose throat; APACHE
285 II , Acute Physiology and Chronic Health Evaluation II ; MV, mechanical
286 ventilation; ICU, intensive care unit; LOS, length of stay; SMAP-HoPe: Survey
287 of Multicenter Assessment with Postal questionnaire for Post-Intensive Care
288 Syndrome (PICS) for Home Living Patients
289

290 **Employment status**

291

292 Fig 2 shows the employment status between the period prior to ICU
293 admission and 12 months following ICU discharge. Prior to ICU admission,
294 the full-time employees comprised 101 subjects. Of those, 15 (14.9%) of them
295 were unemployed at 12 months following discharge from the ICU. Additionally,
296 of 156 subjects with part-time employment, 31 subjects were unemployed at 12
297 months following ICU discharge.

298

299 **Figure 2. Change in the work status between before ICU admission and 12**
300 **months after ICU discharge**

301

302 **Household financial status**

303

304 Fig 3 showed the proportion of respondents who perceived lower
305 household financial status ; this figure distinguishes between employed and
306 unemployed subjects at 12 months following ICU discharge. Among the
307 subjects with an unemployed status, over half of them perceived their financial
308 status to be worse compared with that in the period prior to ICU admission
309 (p<.001).

310

311 **Figure 3 Perceived change in household financial status before and after**
312 **ICU admission**

313 The number of patients who reported that their household financial status had
314 worsened was significantly higher among the unemployed 12 months after ICU
315 discharge. ($p < .001$)

316

317 **Multivariable analysis**

318

319 In multilevel GLM, it showed that higher age, part-time or self-
320 employed status prior to ICU admission, and higher depressive symptoms were
321 independent factors for being unemployed at 12 months following
322 admission. The results of the multivariable analysis are shown in Table 2.

323

324 **Table 2. Multivariable analysis of factors associated with unemployed**
325 **status 12 months after intensive care unit discharge**

Variable	Odds ratio	95% CI	p-value
Age	1.06	1.03-1.08	<.001
Male	0.70	0.31-1.58	0.393
Previous employment status			
Part-time employed†	3.76	1.66-8.52	0.001

Self-employed†	8.57	3.78-19.43	<.001
Cognitive impairment	1.08	0.39-2.95	0.886
Physical dysfunction	2.43	0.92-6.40	0.073
Severity of depression	1.13	1.05-1.23	0.003

326 † indicates full-time employment as a reference.

327

328 **Sensitivity Analysis**

329

330 First, we calculated the proportion of the unemployed after excluding
331 all subjects who responded within the duration of the COVID-19 pandemic. The
332 total subjects who were employed during the period prior to ICU admission was
333 178 and subjects who were in a state of unemployment was 40. The
334 unemployment rate was 22.4% (95% CI, 16.9-29.2). Second, we excluded
335 subjects who were 60 years old and above and conducted multivariable
336 analysis as well as primary analysis. Consequently, age was not determined to
337 be an independent factor; however, worse depressive symptoms still remained
338 to be an independent factor among the unemployed (S2 Table).

339

340 **Discussion**

341

342 We determined that one-fourth of the patients who were previously
343 employed prior to ICU admission were subsequently unemployed at 12 months
344 following discharge from the ICU. Additionally, over half of the unemployed
345 patients perceived worse household financial status compared to that in the
346 period prior to ICU admission. Severity of depressive symptoms was an
347 independent factor for unemployment, as well as higher age and part-time or
348 self-employed status.

349 On the other hand, a recent meta-analysis reported that 40% of
350 patients did not return to work at 12 months following discharge from intensive
351 care [1]. Second, compared with a meta-analysis [1], our findings indicate a
352 relatively lower rate of unemployment. There are some possible reasons for the
353 discrepancies in the findings.

354 First, the characteristics of patients admitted to the ICU could be
355 contributory. Prior studies comparing ICU admissions in Japan and the U.S.
356 found that 59.2% of patients in Japan entered the ICU after elective surgery,
357 compared to 42.3% in the U.S., and that fewer patients entered the ICU from
358 the emergency department in Japan [15]. Upon review, more than half of our
359 subjects following elective surgery were included in this study.

360 Second, although it has not always been the case in recent years,
361 there has been a “lifetime employment system”, which may be specific to
362 Japan, whereas employees in their 50s, in particular, tend to be protected from
363 redundancy [16]. This might influence our findings because a median age of 64
364 years old was included in our study.

365 Third, disability policy in each country should influence the proportion
366 of employees returning to work among patients discharged from intensive
367 care. A recent systematic review and meta-analysis using meta-regression
368 revealed that disability policies in each country positively impacted the ability of
369 ICU-discharged patients to return to work [3]. From OECD (Organization for
370 Economic Co-operation and Development) data, the integration index that was
371 calculated from ten criteria such as employment programs and government-
372 provided job training was higher in Japan than the median index of OECD. [17]

373 Depressive symptoms are likely to be associated with unemployment
374 status after ICU discharge. We found depressive symptoms were an
375 independent factor associated with unemployed despite adjusting for covariates
376 including physical function and cognitive function. These findings are consistent
377 with our hypothesis. A previous study suggested that depression was
378 associated with not returning to work in Australia; however, the study did not
379 conduct multivariable analysis [18]. The present study did not indicate cause-
380 outcome relationships between unemployment and subsequent
381 depression. Depression may lead to unemployment and vice versa from our
382 study design.

383

384 **Strength of the study**

385

386 To date, this study is the largest multi-center study of its kind in
387 Japan. Thus, our findings are reflective of the population admitted in Japanese
388 ICUs.

389

390 **Limitation of the study**

391

392 First, since we collected the employment status and depressive
393 symptoms at the same time, we were not able to clarify the causal relationship
394 between these two variables. Further research measuring depressive
395 symptoms at the discharge from hospital or enough before collecting
396 employment data.

397 Second, COVID-19 may affect our findings. For about half of the study
398 period, Japan was under the COVID-19 pandemic; however, the level of spread
399 was different in the region. Moreover, we conducted an ambidirectional study
400 design; there was no effect on employment and household finance before ICU
401 admission. Employment and household finances at 12 months following ICU
402 discharge was likely influenced by the COVID-19 pandemic. However, we
403 considered this influence minimal, since our sensitivity analysis showed similar
404 results despite exclusion of subjects who responded within the COVID-19
405 pandemic.

406 Third, we included patients of all ages; thus, retirement and retirement
407 age both affect our results. We consider this effect inconsequential to our
408 findings based on results from our sensitivity analysis.

409 Fourth, we did not use validated tools to measure cognitive
410 function. This may affect our findings; however, translated Japanese self-
411 administered questionnaires were limited in utility.

412 Fifth, we excluded patients with central nervous system disease due to
413 this study being based on self-administered questionnaires. Thus, our findings
414 may underestimate the proportion of the unemployed.

415

416 **Implications for future research**

417

418 It is likely that the proportion of those returning to work depends not
419 only on mental health or physical function but also the employment culture and
420 governmental policy. This study is sub-group analysis; thus, a large cohort
421 study focusing on the return to work following intensive care admission is
422 warranted for each country.

423

424 **Implications for clinical practice**

425

426 We need to carefully monitor patients discharged from the ICU, even if
427 they are at home. In particular, depression is associated with unemployment
428 status and needs to be observed. The government and the local municipalities
429 should provide medical and financial support to such patients. Additionally,
430 community support for such patients is warranted.

431

432 **Conclusion**

433

434 We found one-four patients after 12 months from intensive care were
435 unemployed. Additionally, depressive symptoms were associated with
436 unemployment status. employment status and mental health should be follow-
437 up and adequate support are warranted.

438

439 **Acknowledgements**

440 The following SMAP-HoPe Study Project investigators were involved in the
441 protocol: Ryuta Indo, Hiroomi Tatsumi, Atsuko Handa, Kazuyo Koori, Ayano
442 Kudo, Kayo Kitaura, Etsuko Moro, Shin Nunomiya, Akira Ouchi, Masako Sato,
443 Yoshiaki Inoue, Etsuko Tsukioka, Yasuhiro Kishi, Chiaki Fujii, Kohei Matsuba,
444 Hiroki Isonishi, Ikumi Kobashi, Miki Toyoshima, Masahiro Yamane, Yumi
445 Kajiyama, and Yoshifumi Heshiki

446

447

448 **S1 Table 1. Distribution of age group and proportion of unemployed**
449 **subjects stratified by age group**

450 **S2 Table 2. Sensitivity analysis:** Multivariable analysis of factors associated
451 with unemployed status 12 months after intensive care unit discharge among

452 patients after excluding patients aged 60 years or older

453

454

455 **References**

456

457 1. Kamdar BB, Suri R, Suchyta MR, Digrande KF, Sherwood KD, Colantuoni
458 E, et al. Return to work after critical illness: a systematic review and meta-
459 analysis. *Thorax*. 2020;75: 17–27. doi:10.1136/thoraxjnl-2019-213803

460 2. Griffiths J, Hatch RA, Bishop J, Morgan K, Jenkinson C, Cuthbertson BH,
461 et al. An exploration of social and economic outcome and associated
462 health-related quality of life after critical illness in general intensive care unit
463 survivors: a 12-month follow-up study. *Crit Care*. 2013;17: R100.
464 doi:10.1186/cc12745

465 3. Su H, Dreesmann NJ, Hough CL, Bridges E, Thompson HJ. Factors
466 associated with employment outcome after critical illness: Systematic
467 review, meta-analysis, and meta-regression. *J Adv Nurs*. 2021;77: 653–
468 663. doi:10.1111/jan.14631

469 4. Kawakami D, Fujitani S, Morimoto T, Dote H, Takita M, Takaba A, et al.
470 Prevalence of post-intensive care syndrome among Japanese intensive
471 care unit patients: a prospective, multicenter, observational J-PICS study.

- 472 Crit Care. 2021;25: 69. doi:10.1186/s13054-021-03501-z
- 473 5. Myhren H, Ekeberg Ø, Stokland O. Health-related quality of life and return
474 to work after critical illness in general intensive care unit patients: a 1-year
475 follow-up study. Crit Care Med. 2010;38: 1554–1561.
476 doi:10.1097/CCM.0b013e3181e2c8b1
- 477 6. Rothenhäusler H-B, Ehrentraut S, Stoll C, Schelling G, Kapfhammer H-P.
478 The relationship between cognitive performance and employment and
479 health status in long-term survivors of the acute respiratory distress
480 syndrome: results of an exploratory study. Gen Hosp Psychiatry. 2001;23:
481 90–96. doi:10.1016/s0163-8343(01)00123-2
- 482 7. Zisopoulos G, Roussi P, Mouloudi E. Psychological morbidity a year after
483 treatment in intensive care unit. Health Psychol Res. 2020;8: 8852.
484 doi:10.4081/hpr.2020.8852
- 485 8. Norman BC, Jackson JC, Graves JA, Girard TD, Pandharipande PP,
486 Brummel NE, et al. Employment Outcomes After Critical Illness. Crit Care
487 Med. 2016;44: 2003–2009. doi:10.1097/ccm.0000000000001849
- 488 9. Vlasveld MC, van der Feltz-Cornelis CM, Bültmann U, Beekman ATF, van
489 Mechelen W, Hoedeman R, et al. Predicting return to work in workers with
490 all-cause sickness absence greater than 4 weeks: a prospective cohort
491 study. J Occup Rehabil. 2012;22: 118–126. doi:10.1007/s10926-011-9326-
492 0

- 493 10. Unoki T, Sakuramoto H, Uemura S, Tsujimoto T, Yamaguchi T, Shiba Y, et
494 al. Prevalence of and risk factors for post-intensive care syndrome:
495 Multicenter study of patients living at home after treatment in 12 Japanese
496 intensive care units, SMAP-HoPe study. PLoS One. 2021;16: e0252167.
497 doi:10.1371/journal.pone.0252167
- 498 11. Herdman M, Gudex C, Lloyd A, Janssen M, Kind P, Parkin D, et al.
499 Development and preliminary testing of the new five-level version of EQ-5D
500 (EQ-5D-5L). Qual Life Res. 2011;20: 1727–1736. doi:10.1007/s11136-011-
501 9903-x
- 502 12. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. Acta
503 Psychiatr Scand. 1983;67: 361–370. doi:10.1111/j.1600-
504 0447.1983.tb09716.x PMID - 6880820
- 505 13. Hatta H, Higashi A, Yashiro H, Kotaro O, Hayashi K, Kiyota K, et al. A
506 Validation of the Hospital Anxiety and Depression Scale. Jpn J Psychosom
507 Med. 1998;38: 309–315. doi:10.15064/jjpm.38.5_309
- 508 14. Okamoto S, Okamura T, Komamura K. Employment and health after
509 retirement in Japanese men. Bull World Health Organ. 2018;96: 826–833.
510 doi:10.2471/BLT.18.215764
- 511 15. Sirio CA, Tajimi K, Taenaka N, Ujike Y, Okamoto K, Katsuya H. A Cross-
512 Cultural Comparison of Critical Care Delivery Japan and the United States.
513 Chest. 2002;121: 539–548. doi:10.1378/chest.121.2.539

- 514 16. Kambayashi R, Kato T. Long-Term Employment and Job Security over the
515 Past 25 Years: A Comparative Study of Japan and the United States. ILR
516 Review. 2017;70: 359–394. doi:10.1177/0019793916653956
- 517 17. Organisation for Economic Co-operation and Development. Sickness,
518 disability and work breaking the barriers : a synthesis of findings across
519 OECD countries. Paris: OECD; 2010. Available:
520 [https://www.worldcat.org/title/sickness-disability-and-work-breaking-the-](https://www.worldcat.org/title/sickness-disability-and-work-breaking-the-barriers-a-synthesis-of-findings-across-oecd-countries/oclc/1136220722?referer=di&ht=edition)
521 [barriers-a-synthesis-of-findings-across-oecd-](https://www.worldcat.org/title/sickness-disability-and-work-breaking-the-barriers-a-synthesis-of-findings-across-oecd-countries/oclc/1136220722?referer=di&ht=edition)
522 [countries/oclc/1136220722?referer=di&ht=edition](https://www.worldcat.org/title/sickness-disability-and-work-breaking-the-barriers-a-synthesis-of-findings-across-oecd-countries/oclc/1136220722?referer=di&ht=edition)
- 523 18. Hodgson CL, Haines KJ, Bailey M, Barrett J, Bellomo R, Bucknall T, et al.
524 Predictors of return to work in survivors of critical illness. J Crit Care.
525 2018;48: 21–25. doi:10.1016/j.jcrc.2018.08.005
- 526
- 527

SMAP-HoPe study participants

N = 754

Unemployed before ICU admission

n = 425

Employed before ICU admission


n = 329

Exclude due to missing data

n = 1

Analysis

n = 328



**Financial status at 12 months
after ICU discharge**

