Health inequities in SARS-CoV-2 infection, seroprevalence, and COVID-19 vaccination:

Results from the East Bay COVID-19 study

Authors: Cameron Adams, MPH[1], Mary Horton, MPH[1], Olivia Solomon, MPH, Marcus Wong,

Sean L. Wu, PhD, Sophia Fuller, MA, Xiaorong Shao, MA, Indro Fedrigo, Hong L. Quach,

Diana L. Quach, Michelle Meas, MPH, Luis Lopez, Abigail Broughton, Anna L. Barcellos,

MPH, Joan Shim, Yusef Seymens, Samantha Hernandez, Magelda Montoya, Darrell M. Johnson,

MS, Kenneth B. Beckman, PhD, Michael P. Busch, MD, Josefina Coloma, PhD, Joseph A.

Lewnard, PhD, Eva Harris, PhD[2], Lisa F. Barcellos, PhD[2]

Affiliations: University of California, Berkeley, California, USA (C. Adams, M. Horton, O.

Solomon, M. Wong, S. Fuller, X. Shao, I. Fedrigo, H.L. Quach, D.L. Quach, M. Meas, L. Lopez,

A. Broughton, A.L. Barcellos, J. Shim, Y. Seymens, S. Hernandez, M. Montoya, J. Coloma, J.A.

Lewnard, E. Harris, L.F. Barcellos); Institute for Health Metrics and Evaluation, University of

Washington, Seattle, Washington, USA (S.L. Wu); University of Minnesota Genomics Center,

Minneapolis, Minnesota, USA (D.M. Johnson, K.B. Beckman); Vitalant Research Institute, San

Francisco, California, USA (M.P. Busch)

[1]These first authors contributed equally to this article.

[2]These senior authors contributed equally to this article.

Corresponding author: Lisa F. Barcellos, 324 Stanley Hall, Berkeley, CA 94720,

lbarcellos@berkeley.edu, 510-642-7814

Abstract

Comprehensive data on transmission mitigation behaviors and both SARS-CoV-2 infection and serostatus are needed from large, community-based cohorts to identify COVID-19 risk factors and the impact of public health measures. From July 2020-March 2021, approximately 5,500 adults from the East Bay Area, California were followed over three data collection rounds to investigate the association between geographic and demographic characteristics and transmission mitigation behavior with SARS-CoV-2 prevalence. We estimated the populated-adjusted prevalence of antibodies from SARS-CoV-2 infection and COVID-19 vaccination, and self-reported COVID-19 test positivity. Population-adjusted SARS-CoV-2 seroprevalence was low, increasing from 1.03% (95% CI: 0.50-1.96) in Round 1 (July-September 2020), to 1.37% (95% CI: 0.75-2.39) in Round 2 (October-December 2020), to 2.18% (95% CI: 1.48-3.17) in Round 3 (February-March 2021). Population-adjusted seroprevalence of COVID-19 vaccination was 21.64% (95% CI: 19.20-24.34) in Round 3, with Whites having 4.35% (95% CI: 0.35-8.32) higher COVID-19 vaccine seroprevalence than non-Whites. No evidence for an association between transmission mitigation behavior and seroprevalence was observed. Despite >99% of participants reporting wearing masks, non-Whites, lower-income, and lower-educated individuals had the highest SARS-CoV-2 seroprevalence and lowest vaccination seroprevalence. Results demonstrate that more effective policies are needed to address these disparities and inequities.

Text

The first confirmed severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in the California Bay Area was reported on February 28, 2020.[1] Measures to prevent

transmission were implemented soon after and included shelter-in-place orders, mask mandates,

business and school closures, and social distancing recommendations. Despite these measures,

spikes in reported infections occurred from July to September 2020 and December 2020 to

February 2021.[2]

Local public health and healthcare systems experienced major challenges in preventing

infections, identifying COVID-19 cases, and ensuring adherence to transmission mitigation

strategies. Furthermore, disparities and inequities in COVID-19 have been observed in the

United States[3] and there are few studies investigating the association of risk behaviors with

infection outcomes at the individual level.[4–6] A detailed understanding of the effectiveness of

transmission mitigation behaviors and sociodemographic factors that contribute to the

disproportionate impact of COVID-19 in vulnerable communities is critical. Public health and

policy directives aimed at controlling ongoing transmission, developing future prevention

strategies, and targeting health disparities and inequities must be evidence-based. Large

population-representative cohorts with individual-level data on social and behavioral factors[7]

associated with COVID-19 and SARS-CoV-2 infection and serostatus[8–12] are needed.

To address this need, we investigated individual-level characteristics and mitigation

behaviors that contributed to SARS-CoV-2 seroprevalence, self-reported infection, and viral

infection, and other outcomes in a large, population-based sample of over 5,500 individuals from

12 East Bay cities in Northern California followed longitudinally. During three time periods

from July 2020 to April 2021, we estimated the population-adjusted prevalence of SARS-CoV-2

outcomes and differences by age, sex, race/ethnicity, ZIP code, and other demographic strata as

well as the effect of transmission mitigation behavior on SARS-CoV-2 prevalence.

Methods

Recruitment and participants

Recruitment and selection of study participants was completed using a screening phase

followed by a longitudinal study phase with three rounds of data collection. In the screening

phase, all residential addresses within the East Bay cities and communities of Albany, Berkeley,

El Cerrito, El Sobrante, Emeryville, Hercules, Kensington, Oakland, Piedmont, Pinole,

Richmond, and San Pablo (~307,000 residential households) were mailed an invitation to

complete a consent form and screening questionnaire. In a household, the individual aged 18

years or older with the next birthday was eligible to participate. Additional eligibility criteria

included living within the study region, willingness to provide biospecimens and questionnaire

responses, ability to read and speak English or Spanish, and having internet access and an email

address.

The target sample size was 5,500 participants. The distribution of racial and ethnic

identification in the screening questionnaire responses was more White and non-Hispanic than

the region population. To obtain a sample that resembled the racial and ethnic proportions in the

2018 American Community Survey (ACS) for the study region, we ranked assigned ranks for

order of inclusion to all eligible individuals who responded to the screening questionnaire. Black

and/or Hispanic individuals were ranked the highest followed by other non-Whites. Order of

inclusion for Whites was randomly sampled. In Round 1, individuals ranked between 1 and

5,500 were offered study enrollment. Those who declined to enroll were replaced with next

highest ranked individuals who had not been offered study entry. In subsequent rounds,

individuals who had participated in the previous round(s) were offered participation in the next

round. If participation was declined, individuals from the pool of participants who had not

participated in a study round were invited. Approximate dates for each round were July-

September 2020, October-December 2020, and February-March 2021.

All participants provided their informed consent for the screening phase. All participants

in the study phase provided their informed consent for each study round. The study was

approved by the University of California, Berkeley Committee on Protection of Human Subjects

(Protocol #2020-03-13121).

Study procedures

At the start of each round, eligible participants were invited to participate. Those who

agreed to participate received a kit via FedEx containing materials for self-collection of

biospecimens, pre-paid return shipping labels, and instructions to complete an online-

administered questionnaire at the same time as biospecimen collection.

Questionnaire

The questionnaire addressed sex, gender, age, race/ethnicity, income, employment,

physical and mental health, as well as symptoms potentially related to COVID-19 within the

previous 2 weeks, and SARS-CoV-2 testing outside of the study. Participants were also asked

about transmission mitigation behaviors including physical distancing practices, close contacts

with others, and mask wearing. Questionnaires were available in English or Spanish.

SARS-CoV-2 viral and antibody testing

Anterior nasal nare swabs for viral RNA testing and dried blood spots (DBS) for antibody

testing were collected from participants at each study round. Quantitative reverse transcription

PCR (RT-qPCR) was used to identify SARS-CoV-2 viral infection. Three tests were used to

assess anti-SARS-CoV-2 antibodies in DBS: Ortho VITROS® Anti-SARS-CoV-2 Total IgG and

spike IgG ELISA targeted antibodies against the SARS-CoV-2 spike protein (indicating prior

natural infection or vaccination), and Roche-NC Total IgG targeted antibodies against the

nucleocapsid (NC) protein (indicating prior natural infection only).[13] Before COVID-19

vaccines were available in the study region during rounds 1 and 2, detection of antibodies against

the SARS-CoV-2 spike protein was considered evidence of SARS-CoV-2 infection. During

Round 3, vaccinations were widely available, therefore detection of antibodies against the NC

protein were considered evidence of SARS-CoV-2 infection, while detection of antibodies

against the spike protein were considered evidence of SARS-CoV-2 infection or COVID-19

vaccination (Supplement S-4.2, Figure S-1).

SARS-CoV-2 outcomes

The following outcomes were investigated in each round: (1) cumulative SARS-CoV-2

antibody positivity, (2) participants' self-reported history of SARS-CoV-2 positivity from RT-

qPCR testing outside the study, (3) and a surveillance definition of "probable COVID-19 case"

derived from self-reported symptoms and close contact with infected individuals[14], and (4)

viral SARS-CoV-2 positivity from RT-qPCR testing of nasal swabs (Supplement 4.2). We also

investigated antibodies induced by COVID-19 vaccination only in Round 3. SARS-CoV-2

antibody positivity was defined as having detectable antibodies in the current and/or previous

round(s). DBS samples that tested negative for anti-NC antibodies and positive for anti-spike

antibodies in Round 3 were considered to have antibodies from COVID-19 vaccination alone.

Self-reported COVID-19 viral positivity prevalence was defined as the proportion of participants

reporting a positive viral test among all participants who reported being tested within a study

round. Probable COVID-19 case prevalence was defined as the proportion of participants

identified as a probable COVID-19 case among all participants who provided valid responses

within a study round. Viral positivity prevalence was defined as testing positive by RT-qPCR

from nasal swab samples.

Statistical analysis

Population-adjusted seroprevalence and other SARS-CoV-2 outcomes

Bayesian multilevel regression and poststratification (MRP) was used to estimate

population-adjusted cumulative seroprevalence, self-reported SARS-CoV-2 viral positivity

prevalence at each study round, and "probable COVID-19" prevalence at each study round. MRP

is a regression-based method for estimating population and sub-population averages from survey

data that has been shown to perform better than survey weighting, particularly with sparse data

[15,16].

In addition to estimation of regional prevalence of our outcomes, we estimated

prevalence within demographic groups and geographic areas in the study region. Variables of

interest were gender, age, race, Hispanic ethnicity, income, education, household size, and ZIP

code. We used a method described by Leeman and colleagues to generate a synthetic population

for poststratification using data from the 2018 American Community Survey (ACS) and the

Public Use Microdata Sample.[16] Poststratification was done using binary sex because gender

is not reported by the ACS. Race and ethnicity were combined into a single variable to reduce

the number of poststratification strata.

At each study round, binary SARS-CoV-2 outcomes were modeled as a function of

geographic and demographic characteristics using multilevel logistic regression models.

Participant sex was included as a fixed effect. Vectors of random intercepts were defined for

each category of race/ethnicity, age, education, income, household size, and ZIP code and two-

way interactions between ZIP code, race/ethnicity, educational attainment, income, and age. To

improve estimation of geographic effects we allowed for spatial correlations using the modified

Besag-York-Mollié model and included the proportion of Spanish speaking households within

the ZIP code of residence from the ACS as a fixed effect (Supplement S-6).[17]

SARS-CoV-2 outcome prevalence and measures of association

We report populated-adjusted prevalence of SARS-CoV-2 outcomes across the study

region and within geographic and demographic groups of interest. To calculate prevalence

estimates, posterior distributions of the relevant poststratification stratum were aggregated. We

also estimated prevalence differences (PD) and prevalence ratios (PR) for the association

between populated-adjusted SARS-CoV-2 outcomes and race/ethnicity, education, and sex. For

each parameter of interest, the mean of the posterior distribution was the point estimate, and the

95% credible interval (CI) was the 2.5% and 97.5% quantiles of a posterior distribution.

SARS-CoV-2 test-kit bias corrected seroprevalence

169 We estimated cumulative SARS-CoV-2 seroprevalence at each study round adjusted for

170 the net sensitivity and specificity of the SARS-CoV-2 antibody testing algorithm in each study

171 round (Supplement S-6).

Transmission mitigation behavior analyses

In each round, participants were asked about physical distancing practices, recent close

contacts with others, mask wearing, and other behaviors and activities that might affect the risk

of SARS-CoV-2 infection. We classified participants into two behavior categories, "high-risk"

and "low-risk", with those responses using latent class analysis.[18] Crude associations between

behavior categories and characteristics such as sex, age, race/ethnicity, education, and income

were assessed with χ^2 tests. Associations between high-risk vs. low-risk behavior and within

round SARS-CoV-2 seroprevalence and self-reported test positivity were estimated using the

MRP model described above with random intercepts for behavior categories and interactions

between the behavior categories and ZIP code, age, race/ethnicity, education, and income

(Supplement S-5 and S-6).

Statistical analyses were completed in R 4.0.2. NIMBLE was used to implement MRP

models.[19] Detailed descriptions of MRP methods and code are provided in Supplementary

Methods and at github.com/adams-cam/ebcovid_prev.

Results

Enrollment and characteristics of study participants

Of the 16,115 residents who consented and completed the screening procedures between May-

July 2020, 1,777 did not satisfy inclusion criteria and were excluded (Fig 1). Characteristics of

participants are presented in Table 1 and Table S-1. Participation rates were high (Round 1:

76.8%, Round 2: 89.8%, and Round 3: 87.3%), and participants identified predominantly as

female (~63%). Those aged 45-64 years were the largest age group of participants across all

study rounds (ranging from 37.3% to 39.4%). Most participants identified as White (52.5% to

63.3%), followed by Asian/Pacific Islander (13.9% to 15.7%), Hispanic (11.0% to 15.6%), two

or more races (6.9% to 9.1%), African American or Black (3.0% to 4.9%), and Native

American/Alaska Native or other (1.7% to 2.2%).

Fig 1. Study flow chart.

Table 1. Characteristics of participants at each round of the study compared to study
region population.

	PS Strata[a] %	Round 1 no. (%)	Round 2 no. (%)	Round 3 no. (%)
Invited to round		7166	6242	5506
Participated in study round		5501 (76.8)	5603 (89.8)	4806 (87.3)
Gender				
Female		3451 (62.7)	3509 (62.6)	3040 (63.3)

	PS Strata[a] %	Round 1 no. (%)	Round 2 no. (%)	Round 3 no. (%)
Male		1996 (36.3)	2036 (36.3)	1721 (35.8)
Other		51 (0.9)	52 (0.9)	40 (0.8)
Missing		3 (0.1)	6 (0.1)	5 (0.1)
Sex				
Female	51.8	3502 (63.7)	3559 (63.5)	3080 (64.1)
Male	48.2	1996 (36.3)	2042 (36.4)	1725 (35.9)
Missing		3 (0.1)	2 (0)	1 (0)
Age				
18-29	24.2	476 (8.7)	378 (6.7)	286 (6)
30-44	29.0	1768 (32.1)	1656 (29.6)	1330 (27.7)
45-64	30.0	2052 (37.3)	2141 (38.2)	1895 (39.4)
65-74	10.2	928 (16.9)	1077 (19.2)	977 (20.3)
75+	6.6	275 (5)	345 (6.2)	313 (6.5)
Missing		2 (0)	6 (0.1)	5 (0.1)
Race/ethnicity				
African American/Black	17.4	268 (4.9)	180 (3.2)	146 (3)
American Indian/Alaska Native/Other	0.5	123 (2.2)	114 (2)	84 (1.7)
Asian/Pacific Islander	20.3	862 (15.7)	763 (13.6)	668 (13.9)
Hispanic	22.4	860 (15.6)	637 (11.4)	527 (11)
Two or more races	3.8	500 (9.1)	400 (7.1)	334 (6.9)
White	35.6	2886 (52.5)	3503 (62.5)	3041 (63.3)
Missing		2 (0)	6 (0.1)	6 (0.1)
Educational attainment				
College degree or more	47.0	4912 (89.3)	5101 (91)	4404 (91.6)
Less than college degree	53.0	552 (10)	496 (8.9)	399 (8.3)
Missing		37 (0.7)	6 (0.1)	3 (0.1)
Current annual household income, USD				
<$100K	62.9	2081 (37.8)	2075 (37)	1779 (37)
≥$100K	37.1	3192 (58)	3347 (59.7)	2866 (59.6)
Missing		228 (4.1)	181 (3.2)	161 (3.3)

Abbreviations: PS, Poststratification; USD, United States Dollar.

[a] Population percentages from synthetic poststratification tables generated from American Community Survey (ACS) and Public Use Microdata Sample data. Each cell contains the percentage of population age >18. Gender is not available in ACS data.

Of those who completed the questionnaire, 87.3%, 95.3%, and 96.6% provided DBS and

93.6%, 98.1% and 98.5% provided nasal swabs in rounds 1, 2, and 3, respectively (Table 2).

Antibodies against the SARS-CoV-2 spike protein were detected in 29 (0.6%) and 33 (0.6%) of

DBS in rounds 1 and 2, respectively. In Round 3, NC antibodies from natural infection alone

were detected in 84 participants (1.8% of 4,806) and spike antibodies from natural infection or

vaccination were detected in 1,452 participants (31.3% of 4,806). Viral infection was detected in

less than three nasal swabs in each round. The proportion of participants reporting both being

SARS-CoV-2 tested outside the study and testing positive increased over the study period:

10/1,030 (1.0%) participants reporting a positive COVID-19 result in Round 1, 19/2,059 (0.9%)

in Round 2, and 53/1,892 (2.8%) in Round 3. Few participants met the criteria for being a

COVID-19 probable case (<0.5%) (Table S-2).

Table 2. Prevalence of SARS-CoV-2 related outcomes among participants per study round.

	Round 1	**Round 2**	**Round 3**
Biospecimen collection period	Jul. 4, 2020 – Sep. 1, 2020	Oct. 10, 2020 – Dec. 16, 2020	Feb. 11, 2021 – Mar. 20, 2021
N	5501	5603	4806
Provided DBS, no. (%)	4801 (87.3)	5340 (95.3)	4641 (96.6)
Provided nasal swab, no. (%)	5148 (93.6)	5499 (98.1)	4735 (98.5)
SARS-CoV-2 spike antibody status[a], no. (%)			
Non-reactive	4641 (96.7)	5275 (98.8)	3157 (68.0)
QNS	131 (2.7)	31 (0.6)	32 (0.7)
Reactive	29 (0.6)	33 (0.6)	1452 (31.3)
SARS-CoV-2 nucleocapsid antibody status [b], no. (%)			
Non-reactive			4181 (90.1)
QNS			376 (8.1)
Reactive			84 (1.8)

SARS-CoV-2 viral status, no. (%)			
Invalid/Inconclusive	4 (0.1)	5 (0.1)	7 (0.1)
Not Detected	5143 (99.9)	5491 (99.9)	4725 (99.8)
Positive 2019-nCoV	1 (0.0)	3 (0.1)	3 (0.1)
Self-reported positive COVID-19 test, no. (%)			
No	1020 (18.5)	2059 (36.7)	1892 (39.4)
Yes	10 (0.2)	19 (0.3)	53 (1.1)
Not tested	4471 (81.3)	3525 (62.9)	2861 (59.5)
Probable COVID-19 case[c], no. (%)			
No	4800 (87.3)	4938 (88.1)	4419 (91.9)
Yes	21 (0.4)	10 (0.2)	8 (0.2)
Not reported	680 (12.4)	655 (11.7)	379 (7.9)

Abbreviations: DBS, dried blood spot; QNS, quantity not sufficient.

[a] Antibody assays specific to spike antibodies detect presence of antibodies induced by SARS-CoV-2 natural infection or vaccination.

[b] Antibody assays specific to nucleocapsid antibodies detect presence of antibodies induced by SARS-CoV-2 natural infection.

[c] Defined by the Council for State and Territorial Epidemiologists.

Population adjusted SARS-CoV-2 outcome prevalence in study region

Population adjusted SARS-CoV-2 seroprevalence, self-reported COVID-19 test

positivity, and probable COVID-19 cases are reported in Table 3. Overall, populated-adjusted

SARS-CoV-2 natural infection seroprevalence was low across the study region: Round 1 (July-

September 2020) 1.03% (95% CI 0.50-1.96), Round 2 (October-December 2020 1.37% (0.75-

2.39), and Round 3 (February-March 2021) 2.18% (95% CI 1.48-3.17). In Round 3 the

populated-adjusted seroprevalence of COVID-19 vaccination was 21.64% (95% CI: 19.2, 24.34).

Models incorporating sensitivity and specificity of the antibody assays yielded lower SARS-

CoV-2 seroprevalence estimates in Round 1, similar estimates in Rounds 2 and 3, and a higher

estimate of COVID-19 vaccine seroprevalence in Round 3. Population adjusted self-reported test

positivity to SARS-CoV-2 was similar in Rounds 1 (1.11%, 95 CI: 0.39-2.40) and 2 (1.29%,

95% CI: 0.55, 2.17) to seroprevalence estimates and increased to 4.58% (95% CI: 2.56-7.64) in

Round 3. Population-adjusted prevalence of being a COVID-19 probable case was <1% across

rounds (Figure S-2).

Table 3. Crude and population-adjusted prevalence (%) and 95% credible intervals of SARS-CoV-2 outcomes within the study region.

	Round 1	Round 2	Round 3
SARS-CoV-2 Outcome	**Prev.% (95% CI)**	**Prev. % (95% CI)**	**Prev. % (95% CI**
SARS-CoV-2 Ab+ NI[a]			
Crude	29/4670 = 0.62%	48/5949 = 0.81%	110/5991 = 1.83%
Pop-adjusted[b]	1.03 (0.50, 1.96)	1.37 (0.75, 2.39)	2.18 (1.48, 3.17)
Pop-adjusted + Test-bias adjusted[c]	0.63 (0.04, 2.00)	1.02 (0.17, 2.40)	2.19 (1.49, 3.18)
SARS-CoV-2 Ab+ V[d]			
Crude			1369/4608 = 29.7%
Pop-adjusted[b]			21.64 (19.2, 24.34)
Pop-adjusted + Test-bias adjusted[c]			27.56 (22.84, 33.55)
Self-reported test positivity[e]			
Crude	10/1030 = 0.97%	19/2078 = 0.91%	53/1945 = 2.73%
Pop-adjusted[b]	1.11 (0.39, 2.40)	1.29 (0.55, 2.71)	4.58 (2.56, 7.64)
Probable COVID-19 case[f]			
Crude	21/4821 = 0.44%	10/4948 = 0.2%	8/4427 = 0.18%
Pop-adjusted[b]	0.59 (0.26, 1.19)	0.64 (0.17, 1.75)	0.65 (0.17, 1.77)

Abbreviations: Ab, antibody; NI, natural infection; Prev., prevalence; V, vaccination.
[a] Cumulative antibody prevalence from natural infection by SARS-CoV-2.
[b] Population-adjusted prevalence estimated using multilevel regression and poststratification models.
[c] Population-adjusted prevalence estimated using multilevel regression and poststratification models adjusted for antibody assay sensitivity and specificity.
[d] Cumulative prevalence of antibodies from COVID-19 vaccination.
[e] Within round prevalence of self-reported COVID-19 test positivity.
[f] Within round prevalence of COVID-19 probable case.

SARS-CoV-2 seroprevalence and infection vary by geographic area

There was evidence for spatial differences in both populated-adjusted seroprevalence and self-reported test positivity (Fig 2). The northern areas (Richmond, San Pablo, Pinole, and Hercules) and southern areas (East Oakland) of the study region had higher seroprevalence than Berkeley, El Cerrito, and North/Downtown Oakland. Self-reported test positivity was also higher in the northern and southern areas. These trends were consistent across study rounds.

Fig 2. Population-adjusted prevalence of SARS-CoV-2 outcomes within study region ZIP codes.

(A) Cumulative seroprevalence of SARS-CoV-2 antibodies to natural infection. (B) Prevalence of self-reported COVID-19 test positivity across the study region. Data collected in three rounds: Round 1, July-September 2020; Round 2, October-December 2020), and Round 3, February-March 2021.

Population adjusted seroprevalence and self-reported test positivity prevalence within subgroups

Differences in populated-adjusted seroprevalence were observed among demographic groups (Fig 3). In Round 1, those aged less than 45 years had higher likelihood of antibody positivity than those aged 45 or older, but this relationship reversed over time. In general, non-Whites consistently had a higher likelihood of antibody positivity than Whites. The likelihood of antibody positivity also differed by household income and educational attainment, with those reporting a household income <$100,000 and less than a college degree having a higher likelihood of antibody positivity than their comparison groups. Those in households with more

than four people had a higher likelihood of antibody positivity than those in households with four

or less people. There was no clear relationship between sex and populated-adjusted

seroprevalence. Similar relationships were seen between demographic groups and self-reported

test prevalence (Figure S-3).

Fig 3. Cumulative seroprevalence of SARS-CoV-2 antibodies to natural infection among

demographic groups. A) Sex, B) Age, C) Race/ethnicity, D) Income, E) Educational

attainment, and F) Household size. Abbreviations: AA, African American; AMI, American

Indian or Alaskan Native; PI, Pacific Islander.

COVID-19 vaccination seroprevalence differs by race/ethnicity and age

In Round 3, populated-adjusted COVID-19 vaccination seroprevalence was lower in all

non-White groups compared to Whites, (White vs. Non-White, PD: -4.35%, 95% CI: (-9.32, -

0.35), Table 4, Figure S-4). The difference between racial/ethnic groups was largest among older

participants; non-Whites had a lower COVID-19 vaccination seroprevalence compared to Whites

among those aged between 65-74 (PD: -11.67%, 95% CI: (-20.98, -2.49)) and those aged 75 or

older (PD: -10.15%, 95% CI: (-22.25, 0.43)).

Table 4. Population-adjusted prevalence of antibodies from COVID-19 vaccination in
Round 3 within race/ethnicity and age groups and prevalence differences between non-
Whites and Whites.

	Within group	**Non-Whites compared to Whites**
	Prev. % (95% CI)[a]	**PD % (95% CI)[b]**
Race/ethnicity		
White	25.33 (23.14, 27.65)	Ref
African American/Black	19.69 (13.47, 26.13)	-5.65 (-12.17, 0.76)
American Indian/Alaska Native/Other	18.87 (12.32, 25.95)	-6.46 (-13.3, 0.51)
Asian/Pacific Islander	21.34 (17.24, 26.13)	-4 (-8.4, 0.93)

| | Within group | Non-Whites compared to Whites |
	Prev. % (95% CI)[a]	PD % (95% CI)[b]
Hispanic	18.64 (13.74, 24.4)	-6.69 (-11.83, -0.81)
Two or more races	15.74 (11.85, 19.94)	-9.6 (-13.88, -5.28)
All non-White	19.6 (16.16, 23.33)	-4.35 (-8.32, -0.35)
Age group, years		
18-29	15.04 (10.92, 19.81)	-3.2 (-9.14, 2.23)
30-44	16.53 (13.84, 19.57)	-1.94 (-6, 2.74)
45-64	16.6 (13.99, 19.61)	-1.04 (-5.44, 3.89)
65-74	45.74 (40.72, 50.75)	-11.67 (-20.98, -2.49)
75 +	54.01 (46.12, 61.79)	-10.15 (-22.25, 0.43)

Abbreviations: CI, credible interval; PD, prevalence difference; Prev, prevalence.

[a] Population adjusted seroprevalence in round 3 estimated using multilevel regression and poststratification (MRP); Seroprevalence and 95% CI are the mean and 2.5% and 97.5% quantiles of a posterior distribution respectively. Assay used detects antibodies from natural SARS-CoV-2 infection or from COVID-19 vaccination.

[b] Population adjusted COVID-19 seroprevalence difference in round 3 between each race/ethnicity group and Whites among race/ethnicity and between non-Whites and Whites within each age group. PDs estimated using MRP, and the PDs and 95% CIs are the mean and 2.5% and 97.5% quantiles of a posterior distribution, respectively.

Mask-wearing and association between high-risk vs. low-risk behavior and seroprevalence

More than 99% of participants reported ever wearing a mask 99% reported wearing a

mask during leisure and exercise activities, >91% reported wearing a mask at work, and >88%

reported wearing a mask while shopping (Table S-3). After clustering participants into "low-

risk" and "high-risk" groups according to self-reported mitigation behaviors, most participants

were considered low-risk across the study rounds (70%, 82%, and 77%, respectively; Figure

5A). Behaviors with the largest differences in high- versus low-risk behavior were reporting

"yes" to: left home for work, medical/healthcare, care of relative, or other; worked with potential

COVID-19 contact; attended gathering; and traveled to county outside of residence within last

two weeks (Figure 5B). Age, race/ethnicity, and education were all associated ($P<0.001$) with

high- vs. low-risk groups (Table S-4), but we did not observe strong evidence for an association

between mitigation behavior and either seroprevalence or self-reported test positivity (Table 5).

Although point estimates indicate that "low-risk" behavior was associated with lower

seroprevalence.

Table 5. Association of high-risk vs. low-risk mitigation behavior with seroprevalence and
self-reported test positivity and within each study round.

	MOA[a]	Round 1	Round 2	Round 3
SARS-CoV-2 antibody prevalence[b]	PD % (95% CI)	-0.21 (-1.4, 0.79)	0.02 (-1.59, 1.03)	-0.46 (-2.23, 1.00)
	PR (95% CI)	0.93 (0.34, 2.03)	1.4 (0.32, 3.93)	0.88 (0.45, 1.57)
Self-reported test positivity	PD % (95% CI)	0.53 (-0.73, 2.24)	0.11 (-2.12, 1.6)	-1.11 (-5.26, 2.18)
	PR (95% CI)	2.17 (0.54, 6.72)	1.56 (0.33, 4.65)	0.87 (0.41, 1.64)

Abbreviations: CI, credible interval; MOA, measure of association; PD, prevalence difference; PR, prevalence ratio
[a] Populated adjusted prevalence estimated using multilevel regression and poststratification models; point estimate is the mean of a posterior distribution of the parameter, 95% credible interval estimated from 2.5% and 97.5% quantiles of a posterior distribution.
[b] Defined as having detectable SARS-CoV-2 antibodies within a study round.

Discussion

In the current study, we investigated individual-level characteristics and behaviors that

contributed to SARS-CoV-2 related outcomes, including seroprevalence and self-reported

infection, in a large, population-based sample of over 5,500 participants from 12 East Bay

(Northern California) cities. During three time periods from July 2020 to March 2021, we

estimated the population-adjusted prevalence of SARS-CoV-2 outcomes across the study region

and within strata of age, sex, race/ethnicity, ZIP code, and household size. We then characterized

behaviors to mitigate transmission of SARS-CoV-2 and their association with related outcomes.

Overall, prevalence of SARS-CoV-2 outcomes for natural infection were low which may be

attributable to the high percentage of mask-wearing and other risk-mitigating behaviors among

our participants. COVID-19 vaccination seroprevalence estimates in Round 3 was greater than

20%, with non-Whites having much lower seroprevalence than Whites.

Despite the low overall SARS-CoV-2 seroprevalence and infection observed in our study,

ZIP code of residence, age, racial/ethnic identification, education, and income all contributed to

differences in seroprevalence. Specifically, non-Whites, those without a college degree, and

lower income households, had higher seroprevalence. Further, ZIP codes with higher proportions

of Spanish speakers had higher populated-adjusted seroprevalence estimates (Figure S-5). These

differences persisted despite the high rates of mask wearing reported by our study sample,

further adding to strong evidence that the risk of COVID-19 is distributed unequally and that

structural inequities play an important role in COVID-19 risk.[20–23] Moreover, during Round 3

(February-March 2021), COVID-19 vaccines were widely available across the study region. We

found that Whites had much higher prevalence of antibodies from COVID-19 vaccination in

Round 3, than non-Whites, as reported elsewhere.[3,24] Notably, this difference was largest

among those aged 65 or older. Vaccine hesitancy does not explain these findings. In Round 3,

<8% of participants reporting not receiving a COVID-19 vaccine within each racial/ethnic group

were hesitant to vaccination and of the 81 participants reporting hesitancy, 63 (77.8%) were

unsure of their vaccination plans. These findings demonstrate that in the first few months of

vaccine availability in the Bay Area, large disparities in vaccination rates by race/ethnicity were

observed among older persons. Furthermore, Whites, the group with the lowest prevalence of

SARS-CoV-2 infection, were more likely to be vaccinated, underscoring the inequities that exist

surrounding the coronavirus pandemic.

Another key finding was that almost all participants reported wearing masks. This

contrasts with models of mask usage reported by the Institute for Health Metrics and Evaluation

which reported mask usage ranging between 75-82% from December 2020 through March 2021

in California.[25,26] Mask wearing is one the most effective behaviors for controlling

community spread of SARS-CoV-2 infection.[27] The high rate of mask usage by study

participants may partially explain why we did not detect a differences between high-risk and

low-risk mitigation behavior and SARS-CoV-2 prevalence, and partially explain why our

estimates of SARS-CoV-2 seroprevalence and self-report test positivity were lower than public

case reports. However, a recent study estimating SARS-CoV-2 seroprevalence from blood

donors reported seroprevalence in the Bay Area region of California that were similar to our

estimates through December 2020.[28]

A major strength of this study was the longitudinal design and collection of individual-

level data, including biospecimens for antibody and virus testing, which is challenging but much

needed in current studies of the pandemic. Comprehensive data on social distancing, self-

quarantine, mask wearing, working from home, and other transmission mitigation efforts are also

needed to inform current and future prevention strategies. At-home collection of biospecimens,

including DBS for antibody testing, made regular testing without in-person interaction possible.

This was a critical feature, particularly early in the pandemic, when recommendations were to

travel only for essential purposes and to limit in-person interactions. At-home sample collection

was used to obtain more than 30,000 biospecimens and is a feasible approach for large

populations and geographic regions.

One limitation of this study was the under-representation of certain demographics in our

sample. Supplementary mailings of recruitment postcards in Spanish were sent to residences in

ZIP codes with high proportions of Spanish speaking households. We also placed recruitment

flyers in local grocery stores and conducted outreach to community organizations, local

government officials, and school districts in the study region. Despite these efforts, non-Whites,

males, lower income households, and individuals with lower education and from lower

socioeconomic ZIP codes were underrepresented in our sample. Unhoused individuals were also

not captured in our sample. This was important given evidence that individuals who identify as

Hispanic or Black, and other underrepresented groups, are at the highest risk for COVID-19.[23]

These discrepancies may have led to underestimation of SARS-CoV-2 seroprevalence, although

the MRP models may have partially minimized this by pooling information across similar

observed characteristics in the sample data. Our seroprevalence estimates were consistent with

COVID-19 case prevalence reported by public health agencies in ZIP codes with high response

rates (Figure S-6).[29] Additionally, there may be unmeasured confounding from variables not

included in the analyses and self-selection bias from some participants in COVID-19 research

who may be more fervent adherers to COVID-19 public health measures. Finally, some

individuals do not seroconvert after infection or vaccination.

Our results underscore the substantial and persistent inequities that exist surrounding the

coronavirus pandemic. Non-Whites, lower-income, and lower-educated individuals had the

highest SARS-CoV-2 seroprevalence. This disparity in seroprevalence was observed despite near

universal rates of mask wearing in our sample. We also observed large differences in COVID-19

vaccination seroprevalence between racial and ethnic groups. More work must be done to

address these disparities and inequities, such as allocation of resources for high-risk communities

and strategies to mitigate the structural barriers posed by social and structural determinants of

health.

Acknowledgments: We want to acknowledge the contributions of the following individuals to

this study: Ella Parsons, Jordan Keen, Janine Solomon, Jose Salinas, Kevin Duong, Joseph

Egbunikeokye, Maya Talavera, Riya Shrestha, Colin Warnes, José Victor Zambrana, Nicholas

Lo, Parnal Narvekar, Fausto Bustos, Gregorio Dias, Reinaldo Mercado-Hernandez, Julia

Huffaker, Raymond Montes, Alexandra Zermeno, Alejandra Zeiger, William Dow, Michael Lu,

Lila Krop, Kelly Lam, Yan Zhang, Sarah Folkmanis, Sophie Zhai, Dingjun Chen, Ruben Vargas

Ethan Garcia, Oliver Li, Manisha Sahoo, Raina Walencewicz, Sophia Wang, Antonia Gibbs,

Amrita Ramanathan, Catherine Livelo, Taylor Worley, Amanda Tanaka, Savinnie Ho, Jane Liu

Ryan Allen, Sofia Soltero, Victoria Van Metter, Madeleine Fraix, Allie Coyne, Subeksha

Sharma Lydia Yu, Shreeya Garg, Sanjeet Paluru, Malika Saxena, Talia Panadero, Ayra Rahman,

Joshua Calangian, Dharaa Upadhyaya, Sophia Kemp, Ruhi Parikh, Amy Rich, Sophie

Manoukian, Nola Vu, Crystal Nguyen, Jordyn Pinochi, Alma Kuc, Siri Ylenduri, Manvir Kaur,

Angikaar Chana, and Sannidhi Sarvadhavabhatla, Benjamin T. Auch, Dinesha Walek, Evan

Forsberg, Jerry Daniel, Veronica Tonnell, Ji Hyun (Jay) Kim, Mary Nieuwenhuis. Creative

Testing Solutions: Valerie Green, Sherri Cyrus, Phillip Willamson, Brett Hirsch, Paul

Contestable, Mars Stone, Joe Derisi, Emily Crawford, Emily Ahlvin, Armando Diaz, and

Favianna Rodriquez.

Funding: This study was supported by Open Philanthropy Projects, Fast Track grants from

Emergent Ventures, the UC Berkeley Innovative Genomics Institute, and the UC Berkeley

School of Public Health Center for Population Health.

Conflict of interest: JAL reports receipt of grants unrelated to this study from Pfizer and Merck,

Sharpe & Dohme, and consulting fees from Pfizer, Merck, Sharpe & Dohme, VaxCyte, and

Kaiser Permanente. All other authors report no disclosures.

Author Contributions: LB, EH, JC, MB, JAL, KB, CA, and MH conceived of the study. IF,

XS, CA, and MH were responsible for data curation. Formal data analysis was performed by CA,

OS, and SF. Financial support for the work presented in this study was acquired by LB and EH.

Data acquisition, including designing of surveys, design and construction of biospecimen

collection kits, FedEx mailings, receiving of biospecimens mailings from participants, and

communication with participants was performed by LL, IF, MH, HQ, DQ, AB, JS, XS, LB and

MH. Viral and antibody testing was performed and/or supervised by KB, BA, MW, DM, MM,

SM, YS, HQ, and DQ. Sampling design was conceived of by JAL and CA, and models for

prevalence analyses were developed by CA and SW. Management and coordination for

EBCOVID study was primarily the responsibility of LF and EH, with support from JC and MH.

LF and EH research facilities were the primary source of study and laboratory materials.

Computing resources were provided by LF. Implementation of NIMBLE models was performed

by CA with support from SW and SF. IF was responsible for management and implementation of

REDCap surveys and overall system administration of computing resources necessary for the

research. LF and EH supervised all research performed as part of the EBCOVID study. LB, EH,

CA, MH, and IF had full access to and verified the study data. CA, OS, and MH created all

figures in the manuscript. CA and MH wrote and prepared the initial draft of this manuscript. All

authors contributed to interpreting the results of analyses and critically editing and revising the

manuscript.

References

1. Eby K. Coronavirus Timeline: Tracking major moments of COVID-19 pandemic in San
Francisco Bay Area. In: ABC7 San Francisco [Internet]. 27 Mar 2020 [cited 29 Nov 2021].
Available: https://abc7news.com/6047519/

2. United States. Agency for Toxic Substances and Disease Registry. Geospatial Research A and
SP, editor. CDC COVID data tracker. 2020. Available:
https://stacks.cdc.gov/view/cdc/96118

3. Njoku A, Joseph M, Felix R. Changing the Narrative: Structural Barriers and Racial and
Ethnic Inequities in COVID-19 Vaccination. International Journal of Environmental
Research and Public Health. 2021;18: 9904. doi:10.3390/ijerph18189904

4. Qualls N. Community Mitigation Guidelines to Prevent Pandemic Influenza — United States,
2017. MMWR Recomm Rep. 2017;66. doi:10.15585/mmwr.rr6601a1

5. CDC. Interim pre-pandemic planning guidance□: community strategy for pandemic influenza
mitigation in the United States□: early, targeted, layered use of nonpharmaceutical
interventions. [cited 10 Nov 2021]. Available: https://stacks.cdc.gov/view/cdc/11425

6. Bell DM, World Health Organization Working Group on International and Community
Transmission of SARS. Public health interventions and SARS spread, 2003. Emerg Infect
Dis. 2004;10: 1900–1906. doi:10.3201/eid1011.040729

7. Hutchins HJ. COVID-19 Mitigation Behaviors by Age Group — United States, April–June
2020. MMWR Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6943e4

8. Havers FP, Reed C, Lim T, Montgomery JM, Klena JD, Hall AJ, et al. Seroprevalence of
Antibodies to SARS-CoV-2 in 10 Sites in the United States, March 23-May 12, 2020.
JAMA Internal Medicine. 2020;180: 1576–1586. doi:10.1001/jamainternmed.2020.4130

9. Bendavid E, Mulaney B, Sood N, Shah S, Ling E, Bromley-Dulfano R, et al. COVID-19
Antibody Seroprevalence in Santa Clara County, California. medRxiv. 2020;
2020.04.14.20062463-2020.04.14.20062463. doi:10.1101/2020.04.14.20062463

10. Sood N, Simon P, Ebner P, Eichner D, Reynolds J, Bendavid E, et al. Seroprevalence of
SARS-CoV-2–Specific Antibodies Among Adults in Los Angeles County, California, on
April 10-11, 2020. JAMA. 2020;323: 2425–2427. doi:10.1001/jama.2020.8279

11. Self WH. Seroprevalence of SARS-CoV-2 Among Frontline Health Care Personnel in a
Multistate Hospital Network — 13 Academic Medical Centers, April–June 2020. MMWR
Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6935e2

12. Rosenberg ES, Tesoriero JM, Rosenthal EM, Chung R, Barranco MA, Styer LM, et al.
Cumulative incidence and diagnosis of SARS-CoV-2 infection in New York. Ann
Epidemiol. 2020;48: 23-29.e4. doi:10.1016/j.annepidem.2020.06.004

13. Wong MP, Meas MA, Adams C, Hernandez S, Green V, Montoya M, et al. Development
and Implementation of Dried Blood Spot-based COVID-19 Serological Assays for
Epidemiologic Studies. medRxiv. 2021; 2021.11.25.21266786. doi:10/gnnsxg

14. CSTE Interim Position Statement: Update to COVID-19 Case Definition - Council of State
and Territorial Epidemiologists. [cited 13 Oct 2021]. Available:
https://www.cste.org/news/520707/CSTE-Interim-Position-Statement-Update-to-COVID-
19-Case-Definition.htm

15. Gelman A, Little TC. Poststratification Into Many Categories Using Hierarchical Logistic
Regression. Survey Research. 1997;23: 127–135.

16. Leemann L, Wasserfallen F. Extending the Use and Prediction Precision of Subnational
Public Opinion Estimation: EXTENDING USE AND PRECISION OF MrP. American
Journal of Political Science. 2017;61: 1003–1022. doi:10.1111/ajps.12319

17. Simpson D, Rue H, Riebler A, Martins TG, Sørbye SH. Penalising Model Component
Complexity: A Principled, Practical Approach to Constructing Priors. Statist Sci. 2017;32.
doi:10.1214/16-STS576

18. Linzer DA, Lewis JB. poLCA: An R Package for Polytomous Variable Latent Class
Analysis. Journal of Statistical Software. 2011;42: 1–29. doi:10.18637/jss.v042.i10

19. de Valpine P, Turek D, Paciorek CJ, Anderson-Bergman C, Lang DT, Bodik R.
Programming with models: writing statistical algorithms for general model structures with
NIMBLE. Journal of Computational and Graphical Statistics. 2017;26: 403–413.
doi:10.1080/10618600.2016.1172487

20. Reitsma MB, Claypool AL, Vargo J, Shete PB, McCorvie R, Wheeler WH, et al.
Racial/Ethnic Disparities In COVID-19 Exposure Risk, Testing, And Cases At The
Subcounty Level In California. Health Affairs. 2021;40: 870–878.
doi:10.1377/hlthaff.2021.00098

21. Paremoer L, Nandi S, Serag H, Baum F. Covid-19 pandemic and the social determinants of
health. BMJ. 2021;372: n129. doi:10.1136/bmj.n129

22. Abrams EM, Szefler SJ. COVID-19 and the impact of social determinants of health. Lancet
Respir Med. 2020;8: 659–661. doi:10.1016/S2213-2600(20)30234-4

23. Magesh S, John D, Li WT, Li Y, Mattingly-app A, Jain S, et al. Disparities in COVID-19
Outcomes by Race, Ethnicity, and Socioeconomic Status: A Systematic-Review and Meta-
analysis. JAMA Network Open. 2021;4: e2134147.
doi:10.1001/jamanetworkopen.2021.34147

24. Agarwal R, Dugas M, Ramaprasad J, Luo J, Li G, Gao G (Gordon). Socioeconomic privilege
and political ideology are associated with racial disparity in COVID-19 vaccination. PNAS.
2021;118. doi:10.1073/pnas.2107873118

25. Reiner RC, Barber RM, Collins JK, Zheng P, Adolph C, Albright J, et al. Modeling COVID-
19 scenarios for the United States. Nat Med. 2021;27: 94–105. doi:10.1038/s41591-020-
1132-9

26. IHME | COVID-19 Projections. In: Institute for Health Metrics and Evaluation [Internet].
[cited 1 Oct 2021]. Available: https://covid19.healthdata.org/

27. Brooks JT, Butler JC. Effectiveness of Mask Wearing to Control Community Spread of
SARS-CoV-2. JAMA. 2021;325: 998–999. doi:10.1001/jama.2021.1505

28. Jones JM, Stone M, Sulaeman H, Fink RV, Dave H, Levy ME, et al. Estimated US Infection-
and Vaccine-Induced SARS-CoV-2 Seroprevalence Based on Blood Donations, July 2020-
May 2021. JAMA. 2021 [cited 30 Sep 2021]. doi:10.1001/jama.2021.15161

29. The Los Angeles Times' open-source archive of California coronavirus data. Los Angeles
Times Data and Graphics Department; 2021. Available:
https://github.com/datadesk/california-coronavirus-data

A.

B.