Underlying co morbidity reveals unique immune signatures in Type II diabetes patients infected with SARS-CoV2

Soumya Sengupta [1,2†], Gargee Bhattacharya [1,2†], Sanchari Chatterjee[1,2], Ankita Datey[1,3], Shubham K Shaw[1,2], Sandhya Suranjika[1], Paritosh Nath[1], Prakash K Barik[1], Punit Prasad[1,2], Soma Chattopadhyay[1,2], Rajeeb K Swain[1,2]* Ajay Parida[1,2]*, Satish Devadas[1,2]*

1 Institute of Life Science, Nalco Square, Bhubaneswar, Odisha, India-751023.

2 Regional Centre for Biotechnology (RCB), 3rd Milestone, Faridabad-Gurgaon, Haryana-121001, India

3 School of Biotechnology, Kalinga Institute of Industrial Technology (KIIT) University, Bhubaneswar, Odisha, India-751024.

†These authors contributed equally

*Address of Corresponding authors:

Dr Rajeeb K Swain (RS)

Cancer Biology,

Institute of Life Sciences

(Autonomous Institute of Dept of Biotechnology, Govt. of India),

Nalco Square, Bhubaneswar-751023. Odisha, India

Phone No: 0091 0674 2304 301, Fax No: 0091 674 2300728

E-mail: rkswain@ils.res.in

Dr. Ajay Parida (AP)

Plant and Microbial Biotechnology,

Institute of Life Sciences,

(Autonomous Institute of Dept of Biotechnology, Govt. of India),

Nalco Square, Bhubaneswar-751023. Odisha, India

Phone No: 0091 674 2304324; Fax No: 0091 674 2300728

E-mail: drajayparida@gmail.com, ajayparida@ils.res.in

Dr. Satish Devadas (SD)

Infectious Disease Biology,

29   Institute of Life Sciences Institute of Life Sciences,

30   (Autonomous Institute of Dept of Biotechnology, Govt. of India),

31   Nalco Square, Bhubaneswar-751023. Odisha, India

32   Phone No: 0091 674 2300701; Fax No: 0091 674 2300728 E-mail: satdevs@ils.res.in

Abstract

Background: SARS-CoV2 infection in patients with comorbidities, particularly T2DM has been a major challenge globally. Here, we did whole blood immunophenotyping along with plasma cytokine, chemokine, antibody isotyping and viral load determination from oropharyngeal swab to understand the immune pathology in the T2DM patients infected with SARS-CoV2.

Methods: Blood samples from 25 Covid-19 positive patients having T2DM, 10 Covid-19 positive patients not having T2DM and 10 Covid-19 negative, non-diabetic healthy controls were assessed for various immune cells by analyzing for their signature surface proteins in mass cytometry. Circulating cytokines, chemokines and antibody isotypes were determined from plasma. Viral copy number was determined from oropharyngeal swabs. All our representative data corroborated with laboratory findings.

Results: Our observations encompass T2DM patients having elevated levels of both type I and type II cytokines and higher levels of circulating IgA, IgM, IgG1 and IgG2 as compared to NDM and healthy volunteers. They also displayed higher percentages of granulocytes, mDCs, plasmablasts, Th2-like cells, $CD4^+$ EM cells, $CD8^+$ TE cells as compared to healthy volunteers. T2DM patients also displayed lower percentages of pDCs, lymphocytes, $CD8^+$ TE cells, $CD4^+$, $CD8^+$ EM.

Conclusion: Our study demonstrated that patients with T2DM displayed higher inflammatory markers and a dysregulated anti-viral and anti-inflammatory response when compared to NDM and healthy controls.

Contribution to the field: Covid-19 infection in people with comorbidities, particularly T2DM has been a cause of mortality in several nations and they represent an extremely vulnerable population to Covid-19. This study is one of the most comprehensive study from India, to understand the interplay between immune response and viremia occurring in these T2DM patients infected with SARS-CoV2 and will help in designing public health response and vaccination priorities.

Keywords- SARS-CoV2, Type II Diabetes, Whole blood Immunophenotyping, Innate Immune Response, Adaptive Immune Response

Introduction

Since the outbreak of the novel coronavirus disease (Covid-19) in late 2019, India has recorded over 34 million cases and 4,54,745 deaths as of 25th October, 2021 and now has the second highest number of cases in the world, after the United States of America. Essentially, SARS-Cov2 is a betacoronavirus, belonging to Coronaviridae family and is closely related to SARS and MERS, which were responsible for earlier disease outbreaks in 2003 and 2011 respectively[1]. Broadly, Covid-19 cases were and are still classified into symptomatic and asymptomatic, based on the presence or absence of symptoms and eventual severity based on immune response and/or its failure. With respect to the symptoms of Covid-19 infection, symptomatic patients have fever, dry cough, shortness of breath, acute respiratory distress, loss of taste and smell and in certain cases diarrhoea [2, 3]. Based on the intensity of these symptoms, the patients are classified into mild and severe, during diagnosis. Prognosis of the severe patients is critical. However, the major challenge arises from the huge population of asymptomatic cases, as they are responsible for the undetected spread of infection.

In general, disease severity in Covid-19 is associated with lymphopenia, cytokine storm, blood coagulation, drop in pO_2 levels, etc. [4, 5]. However, with respect to the Indian population, these clinical parameters and the associated immune response were not definitely present in most Covid-19 infected patients. Interestingly, one of the major attributes of Covid-19 in Indian population, especially in the first wave, was quick recovery, but the underlying immunological mechanism was and is yet to be understood. However, similar to the first wave, 70% of India's mortality in Covid-19 in the second wave is still attributed to comorbidities, specifically type II diabetes (T2DM). Although vaccinations drive initiated by the government aims to protect the population in general, the overwhelming number of these patients poses a major challenge to recovery and recuperation of the individual. In that respect, co-relation between comorbidities including T2DM and viral load, T2DM and glucose levels, etc. in the patient can play a major role in dictating the immune response and eventual outcome of Covid-19 infection. Therefore, it is crucial to understand the interplay of these factors along with immunological parameters to give us a comprehensive idea about the status of T2DM patients having Covid-19.

Immune response can be divided into three types namely, type I (antiviral), type II (anti helminthic), type III (antifungal). Type I response mainly constitutes of T-bet and Interferon gamma (IFN-γ) mediated response, which is against intracellular pathogens, including viruses.

On the other hand, type II response is mediated by GATA-3 and is mainly against helminthic

worms, which are carried out by effector molecules such as IL-4, IL-9, and IgE. The third type

of response (type III response) is mediated by ROR-γt and effector molecules such as IL-17A,

IL-17F, 1L-22 etc. are responsible for controlling fungal infections (6, 7). Longitudinal

analysis has shown an immune dysregulation in Covid-19 patients and people with T2DM

contracting Covid-19 Infection had higher innate immune cells, lower T lymphocytes, a

sustained increase in antiviral, anti-fungal response and higher type 2 response such as IL-5,

IL-13, IgE and eosinophils (8).

Altogether, the immunological response in Covid-19 patients with co-morbidity is an

evolving area of study and these patients represents higher vulnerability in comparison to the

ones without any comorbidity or other comorbidities. Our study aims to elucidate the

differences between T2DM patients having Covid-19 as compared to those devoid of T2DM,

based on multiple parameters such as viral load, cytokine *milieu* and immune cells, which will

give us a comprehensive idea about the role played by T2DM in Covid-19 pathogenesis.

Materials and Methods

Population and Samples

A total of 25 samples with T2DM and 10 samples without T2DM (NDM) (Non-diabetes mellitus) were collected. These samples were collected from hospital within 4 days of patient's admission. Type II diabetes in these patients was confirmed by previous clinical history and for this particular study was assessed by fasting plasma glucose level and glycated haemoglobin and other clinical and biochemical parameters as requested by their consultant medical practitioner. Amongst these, 22 patients with T2DM had hypertension. The T2DM or NDM patients neither reported nor were assessed for other comorbidities during the infection. The detailed laboratory findings are given in Table 1. All the T2DM patients had severe symptoms, as assessed by the clinician. The NDM patients either had mild symptoms or were asymptomatic. Additionally, 10 healthy volunteers who tested negative for Covid-19 were included in the study to provide basal and or steady state biochemical and cytokine profile. Briefly, 5ml blood was collected in BD Vacutainer EDTA tubes (BD 367863) and 270 μl of whole blood from 10 T2DM patients, 5 NDM patients and 5 healthy volunteers was used for deep immune profiling by mass cytometry. The tubes were spun at 1600g for 20 minutes at room temperature. Plasma was then collected and stored in -80 °C till further analysis. The details of all the reagents and software used are given in the supplementary table 1 (S1).

SARS-CoV2 Viral load detection in respiratory specimen and plasma

300 μl each of oropharyngeal swab and plasma were taken for viral RNA extraction using TAN Bead Maelstrom 4800 as per manufacturer's instructions. The extracted viral RNA was stored at-80°C until further use (9).

qRT-PCR

The qRT-PCR was performed using 5μl of the extracted RNA from samples using the TRUPCR SARS-CoV-2 RT qPCR Kit V-2.0. The human RNase P served as an internal control whereas envelope (E) and nucleocapsid (N) genes were targeted for SARS-CoV-2 amplification (9).

Viral copy number determination

The viral copy number was determined for the above-mentioned samples by generating the standard curve of SARS CoV-2 N (nucleocapsid) gene. The N gene was cloned into pBiEx vector and 10-fold serial dilutions of the plasmid were done to obtain the standard curve. The

percentage of copy number/ml was calculated from the corresponding Ct values of all the

samples. For obtaining Ct values, cDNA was prepared from the extracted RNA using random

hexamers by TAKARA primescript 1st strand cDNA synthesis kit (Kusatsu, Japan). The

cDNA was subjected to qPCR (Mesagreen SYBR Green-No ROX, Eurogentec, Belgium)

using nucleocapsid gene specific primers. (FP: GTAACACAAGCTTTCGGCAG and RP:

GTGTGACTTCCATGCCAATG) (9).

Plasma Cytokine and Chemokine detection assay

148   Neat plasma from Covid-19 and controls was used to measure 41 cytokines and

149   chemokines using human Milliplex map cytokine assay kit (Millipore, Billerica, MA, USA).

150   The samples were acquired in a Bio-Plex 200 system (Bio-Rad, Hercules, CA, USA) and

151   cytokine concentrations were calculated using Bio-Plex manager software with a five-

152   parameter (5PL) curve-fitting algorithm applied for standard curve calculation (10).

Isotyping of circulating antibody from blood

For analysis of the isotype composition of antibodies in circulation, plasma of T2DM,

NDM patients and healthy volunteers, were analysed by ProcartaPlex Human Antibody

Isotyping Panels (Cat. No EPX070-10818-901, Invitrogen, Vienna, Austria), based on the

manufacturer's instructions. Briefly, antibody-coated magnetic bead mixtures were incubated

with 25μl of assay buffer, kit standards or diluted plasma (1:20000) samples in a ProcartaPlex

96-wells plate at room temperature for 1 hour. Detection antibodies (25 μl) were then added

and the plates were incubated on an orbital shaker at 500 rpm for 30 min. Next, the wells were

incubated with 50μl of diluted Streptavidin-Phycoerythrin for 30 min. Plates were then washed

using a hand-held magnetic plate washer. All incubations were performed at room temperature

in the dark. Afterwards, samples were suspended in 120 μl reading buffer. The samples were

acquired in a Bio-Plex 200 system (Bio-Rad, Hercules, CA, USA) and cytokine concentrations

were calculated using Bio-Plex manager software with a five-parameter curve-fitting algorithm

(5PL) applied for standard curve calculation (11).

Whole blood Immunophenotyping by Mass Cytometry

168   For immunophenotyping, 270μl of whole blood was collected, added to pre-coated

169   Maxpar Direct Immune Profiling tubes and incubated for 30 minutes in room temperature. For

170   lysis, 250μl of 1x BD FACs Lyse was added, followed by 10 min incubation. There were two

171   consecutive washes with Maxpar water and Cell staining buffer. The cells were then fixed

with 4% formaldehyde and incubated for 10 minutes followed by washing at 800g for 5 minutes. Eventually, the cells were suspended in 1ml Iridium solution and stored in -80 °C till acquisition. For acquisition, the cells were thawed, washed with Cell Staining buffer and Cell Acquisition Solution. The cell density was adjusted to 1 million cells/ml in Cell acquisition Solution with 0.1% EQ Beads and acquired in Helios Mass cytometer. FCS files were then normalised with CyTOF software V.7 (Fluidigm) and then exported and analysed by FlowJo software V10.7 (BD Biosciences)(12, 13).

Statistics

Statistical analysis was performed using the GraphPad Prism software, version 8.0.1. Data was presented as Mean ± Standard Deviation of Mean (SEM). Non-parametric Kruskal Wallis Test with post hoc Dunn's multiple comparison test was used to compare the levels of cytokines, antibodies and percentage of different immune cells among the three groups. Mann-Whitney U test was used to compare laboratory parameters between T2D and NDM patients. *P* values less than 0.05 were considered significant (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$).

High dimensional analysis

For high dimensional analysis, t-SNE analysis was performed using FlowJo V.10.7 (BD Biosciences). The gating for analysis was done on live, intact, $CD45^{high}CD66b^{low}$ lymphocytes. Amongst a total of 100,000 events, 5000 events in the lymphocyte population were used per sample for this analysis. 5 samples of T2D and of Healthy controls were used to compute t-SNE and a total of 25000 events per group were concatenated and exported for t-SNE analysis. The default parameters in the software, iterations-1000, perplexity-30, eta-675, KNN algorithm-Exact (vantage point tree), gradient logarithm-Barnes-Hut were used to compute the t-SNE plot.

Ethics Statement

The studies involving human participants were reviewed and approved by the Institutional Human Ethics Committee, Institute of Life Sciences. The Institutional Ethics Committee (IEC)/ Institutional Review Board (IRB) reference number is 106/HEC/2021. Written consent informed to participate in this study was provided by the participants' spouse/next of kin.

Data Availability Statement

The human clinical data is stored at KIMS Hospital, samples are stored at ILS
Biorepository, Ethical clearance with the corresponding authors and on request for this, ethical,
legal, and privacy consent will be requested from both the doctor and patient to share data.

Author Contributions

Experimental Design and Conceptualization – SD, SS and GB

Sample Collection and processing- RS, PP, SaS, SS, GB, SaC, PB and PN

Mass Cytometry Experiments- SS, GB and SKS

Chemokine and Cytokine Multiplexing- SS, SaC and GB

Antibody Isotyping- SS, SaC and AD

RT-PCR and Virological assays- SaC, AD and SC

Data analysis- SS, GB, SaC, AD, SC and SD

Manuscript Drafting- SS, GB, SaC, SC and SD

Manuscript Editing and Review- GB, SS, RS, PP, AP, SC and SD

Conflict of Interest.

The authors declare no conflict of interest. The funding agency had no role in the design
of the study; in the collection, analyses, or interpretation of data; in the writing of the
manuscript, or in the decision to publish the results.

Funding.

This study was supported by the core funding of Institute of Life Sciences,
Bhubaneswar, Dept of Biotechnology, India. SS was funded by the DBT fellowship. GB, SaC
and SKS were funded by the CSIR fellowship.

Acknowledgements.

We would like to acknowledge ILS Biorepository for providing the SARS-CoV2
samples. We would like to thank Dr. Gajendra Motiram Jogdand for valuable suggestions in
multiplexing. We would also like to thank Mr. Deepak Singh, Mr. Satyajit Behera and Ms.
Jyotsna Priyadarshini for data entry.

Results

Viral load does not corroborate with inflammation in T2DM

The study was executed on NDM and T2DM Covid-19 patients, who had been detected and admitted to a local tertiary COVID hospital. To examine for viral load, oropharyngeal (OP) samples were collected from various patients as per established protocol and processed in the BSL3 facility at ILS, Bhubaneswar. In most of the NDM and T2DM patients, there was no significant difference in the viral copy number or $\Delta\Delta$ Ct values of OP samples at the time of sampling **(Fig. 1A and B).** This suggests that the viral load in OP samples does not and did not correlate with disease severity in this population of study. This probably indicates that the viral load present in the host does not dictate the severity of Covid-19 infection.

Altered cytokine and chemokine profiles in Covid-19 patients with Type II Diabetes

To investigate the status of systemic inflammation concurrent with SARS-Cov2 infection, we did multiplexing analysis of 41 cytokines and chemokines from the plasma of T2DM and NDM patients to delineate the altered immune microenvironment in these patients. For comparison and for basal level expression of these cytokines and chemokines, we added ten healthy volunteers and segregated the proteins into significant, moderate and mild based on the level of their expression in T2DM as compared to NDM and healthy volunteers. Accordingly, the levels of IL-6, TNF-α, G-CSF, GM-CSF, IFN-α2, IL-10, VEGF, IL-1Rα, IL-12p40, IL-15, IL-1α, MIP-1β, were found to be significantly elevated (p<0.0001, Kruskal Wallis Test), indicating the simultaneous release of both Type I and Type II cytokines in severity **(Fig. 2A)**. In contrast, there was moderate elevation in the levels of IFN-γ, EGF, IL-8, indicative of the ongoing antiviral reaction in these patients (**Fig. 2B).** Amongst the slightly elevated ones, IP-10 (p=0.0365), IL-9, IL-4 (p=0.0146) showed higher significance **(Fig. 2C)**. Although IP-10 has been used as a real time marker for mortality in Covid-19 patients, we did not do a longitudinal patient tracking, therefore cannot ascertain its role in mortality. In addition, IL-4 and IL-9 which are both type II cytokines have also been linked to severity of Covid-19 patients. Although our study could not find any significant difference in levels of type III signature cytokines such as IL-17 and IL-1β, the presence of type I and type II immune signatures were overlapping in multiple patients (data not shown). However T2DM patients showed elevation of both type I and type II cytokines indicating a dysregulated immune response, prominent in these patients as earlier reported by Lucas et al **(Fig. 2)**.

Interestingly, the levels of chemokines as opposed to cytokines, were higher in NDM
and healthy volunteers as opposed to T2DM patients. For example, the levels of Eotaxin were
found to be lower in T2DM than in NDM patients and healthy controls (**Fig 2D**). In general,
Eotaxin is associated with chemoattraction of eosinophils, basophils and neutrophils. Previous
studies have reported its increase with severity but in our study, the population indicated a
reverse trend; decreasing Eotaxin levels in the T2DM patients. This can be indicative of an
impaired healing process as previous reports have suggested a positive correlation between
increasing number of eosinophils and healing. In addition, sCD40L was elevated in NDM
patients as compared to T2DM and healthy volunteers (**Fig. 2E**). Since sCD40L has been
associated with immunosuppression, it is indicative of an anti-inflammatory immune response
in NDM patients, which is clearly absent in T2DM cases. Similarly, the levels of MDC
(macrophage derived chemokine) were also higher in NDM as compared to T2DM patients,
demonstrating its potential role in immune suppression (**Fig. 2E**). In addition, GRO levels
were higher in both NDM and T2DM patients as compared to healthy controls (**Fig. 2F**). As
GRO serves as a chemoattractant for neutrophils, it is indirectly indicative of the status of
underlying inflammation in NDM and T2DM patients. Collectively, our cytokine and
chemokine profile of T2DM patients showed an increasing trend in the inflammatory cytokine
panel and augmented levels of chemokines that were primarily responsible for repair and
angiogenesis.

Circulating antibody isotype profiling showed a skewed inflammatory profile in Covid-19
patients with Type II Diabetes

Various antibody isotypes correlate with the clinical phenotype of an infection and is
therefore an important indicator of the type of immune response. Since cytokine *milieu* in
T2DM and NDM patients was skewed, we wanted to define the circulating antibody isotypes
in these patients as opposed to healthy controls. We did Luminex based multi plex assay from
their plasma and quantified the amount of IgA, IgM, IgE, IgG1, IgG2, IgG3 and IgG4. Here,
we found IgG1 and IgG2 to be significantly elevated in T2DM patients as compared to healthy
controls (p=<0.0001) and NDM patients (p=0.0024 for IgG1 and p=0.0033 for IgG2),
correlating with the higher levels of type I cytokines in these patients. IgE, being a hallmark
of type II immune response was also elevated in T2DM as compared to NDM (p=0.0453) and
healthy control (p=0.0347). Alongside, IgA being associated with mucosal immunity was also
higher in T2DM patients as compared to NDM (p=0.0012) and healthy controls (p=0.0140).
As IgM is the first antibody isotype to appear during the course of infection, accordingly we

found its levels to be higher in T2DM patients as compared to NDM (p=0.0178) and healthy

controls (p=0.0017) **(Fig. 3).** In summary, the antibody isotype profile was concurrent to the

cytokine levels, thus confirming and furthering the status of disease severity in Covid-19

patients.

Dysregulated innate immune system in Covid-19 patients with Type II Diabetes

With the cytokine multiplexing and antibody isotyping indicating an altered immune

response, our next objective was to delineate the alteration in the cells of innate and adaptive

immune system during Covid-19. Based on the surface markers analysed, we found T2DM

patients showing significant increase in $CD45^{low}CD66b^{high}$ granulocytes as compared to

healthy controls (p=0.0074), which is concurrent with the increase in the levels of IL-8 **(Fig.**

4A). The bulk population of these granulocytes were neutrophils, though we did not find any

significant difference between the levels of neutrophils in T2DM, NDM and healthy controls

(data not shown). Also, dendritic cell populations were variable, with their numbers decreasing

in T2DM as compared to NDM and healthy controls **(Fig. 4B)**. This was in contrast to the

increase in IFN-α2 levels, indicating towards dysregulated dendritic cell population in T2DM

patients **(Fig. 2A).** Within the dendritic cell population, plasmacytoid dendritic cells (pDCs)

decreased in T2DM patients as compared to NDM and healthy controls **(Fig. 4B).** However,

percentage of myeloid dendritic cells (mDCs) increased in T2DM patients as compared to

NDM and healthy controls **(Fig. 4C)**. The percentage of basophils was also reduced in T2DM

patients as compared to healthy volunteers, which was correlating with the decrease in levels

of chemokine, Eotaxin (CCL11) **(Fig. 4D).** Although, we did not observe any variation

between T2DM, NDM and healthy controls in the populations of gamma delta cells and natural

killer cells there was reduction in the percentage of MAIT/NKT cells in T2DM patients as

compared to healthy and NDM patients **(Fig. 4E).** As reported earlier (data not shown), there

were no changes in the overall or subsets of monocytes and natural killer cells. Altogether,

alterations in the innate immune cell population is suggestive of an aberrant immune response

incapable of controlling the infection.

Heterogenicity in Lymphocyte population in Covid-19 patients with Type II Diabetes, NDM
and healthy controls

Lymphocytes are instrumental in fighting infections and earlier reports indicated

towards its decreasing trend in T2DM patients. There was a moderate decrease in the

percentage of lymphocyte in T2DM as compared as to the healthy controls **(Fig. 4A).** Amongst

lymphocytes, percentage of total B cell population was expanding in T2DM patients as
compared to healthy controls **(Fig. 5A)** and was is in direct correlation with increased IgG1
and IgG2 in the former **(Fig. 3).** Further analyses revealed a significant increase in plasmablast
population in T2DM patients as compared to healthy controls. Additionally, we found a
significant decrease in the percentage of $CD3^+$ T cells in T2DM patients as compared to healthy
controls suggesting lymphopenia associated with increased severity **(Fig. 5B).** Within the T
cell population, there was an increase in the percentage $CD4^+$ T cells (statistically insignificant)
but decrease of $CD8^+$ T cells **(Fig. 5B).** Further, in depth analysis of $CD4^+$ T cells revealed
that T2DM patients had an increase of the Th2 subtype as compared to NDM or healthy
volunteers **(Fig. 5C).** This can be correlated with the increased levels of IL-4 and IL-9, found
in the plasma of these patients **(Fig. 2C).** However, Th1, Th17, Treg and Tfh cells did not
show any statistically significant difference, which can be attributed to the time of sampling or
be a consequence of dysregulated immune response. There was also no significant difference
in Naïve $CD4^+$ T cells or Central memory $CD4^+$ T cells within the groups. But terminal effector
(TE) cells were decreased and effector memory (EM) cells were increased in T2DM patients
as compared to healthy controls **(Fig. 5D**). This was in contrast to the $CD8^+$ compartment,
where TE cells were increased in T2DM patients as compared to NDM but there was no
difference between T2DM and healthy controls suggesting a robust response in NDM patients
as compared to T2DM and a certain degree of cross-reactive reaction occurring in these T2DM
patients **(Fig. 5E).** Similarly, there was significant decrease of EM cells in T2DM as compared
to NDM while no difference was observed in healthy controls when compared to T2DM or
NDM. Collectively, a bias was found towards terminal effector $CD8^+$ T cells in T cell immune
response which was suggestive of an effective ongoing anti-viral response.

High Dimensional analysis reveals immune perturbations in T2DM patients

We also performed t-distribution stochastic neighbour embedding (t-SNE) analysis to
understand how different markers reported previously, show variability in Covid-19 patients
in our population of study. t-SNE analysis revealed a decrease in the expression of CD3 and
CD8 in T2DM patients as compared to healthy volunteers **(Fig. 6A).** On the contrary, there
was increase in the expression of CD14, CD38 and HLA-DR in T2DM patients as compared
to healthy volunteers **(Fig. 6B).**

Discussion

The Covid-19 pandemic has brought an unprecedented global devastation on lives and livelihood of the common people, breakdown of the best healthcare system and economic collapse all across the globe. Till October 2021, India reported around 34 million cases, being second in place to the United States of America. Unfortunately, a bulk population of Covid-19 cases in India who had succumbed to the infection had comorbidities and amongst these, Type II Diabetes was the most common. At present, there are no significant reports from India providing an insight into the variations in immune response, extent of inflammation and outcome in T2DM patients as compared to NDM patients and healthy volunteers. In this study, we attempted to describe and compare immune response, viral loads and clinical parameters amongst T2DM, NDM and healthy volunteers. In that respect, 25 patients with T2DM, 10 NDM patients and 10 healthy volunteers were studied for cytokine, chemokine, viral loads, clinical parameters and antibody isotyping assays for understanding the immunopathology of the virus. Ten T2DM, five NDM and five healthy volunteers were studied for whole blood mass cytometry assay to determine the difference in immune cells amongst the groups. While it is understood that this is a very small group, some clear patterns are established here suggesting that the chronic Meta inflammation in Type II Diabetes might play a role in further precipitating the inflammatory signatures in Covid-19.

It is known that mild inflammation serves as one of the major initiator mechanisms of the immune system that is extremely crucial in fighting infection (14). However elevated levels of circulatory cytokines and eventual cytokine storm and dysregulated inflammatory responses have been evinced in Covid-19 infection (2, 4). Additionally, previous reports suggest that patients with T2DM have a higher vulnerability for SARS-CoV2 infection (15, 16) where persistent low grade inflammation associated with T2DM was evinced but not co related to be causal. Our data taken together with biochemical, cytokine and antibody profile strongly indicate that chronic low grade inflammation and the consequential dysregulated immune response might be responsible for the upsurge of inflammatory responses in Covid-19 patients leading to severe infection and or mortality.

Our preliminary objective in the study was to assess the differences in viremia in Oropharyngeal swab between the two groups. However, viral load of OP samples did not show any significant difference between the two groups. This was in accordance with previous studies, where it was observed that the viral load in nasal and throat swab samples were similar

in the NDM and T2DM patients (17, 18). Here, the T2DM are considered as severe Covid-19
patients, based on the severity of their symptoms. And, the NDM patients either had NDM
symptoms (n=4) or were asymptomatic (n=6). This is evident from the clinical parameters as
T2DM patients had higher levels of CRP which is a non-specific inflammatory marker,
dysregulated glucose metabolism as evident from higher plasma glucose, glycated
haemoglobin and impaired blood coagulation as interpreted from higher d-dimer levels(19-22).

Cytokines and chemokines are soluble mediators that control the migration of immune
cells to the site of inflammation, providing a pro or anti-inflammatory environment which
essentially shapes the type of immune response during a disease. In our T2DM patients who
were infected with SARS-CoV2, higher expression of both type I and type II cytokines were
evinced. Previous reports suggest the involvement of both the types of cytokines in aggravating
the severity of the disease. Increase in type I cytokines such as IFN-γ TNF-α, IFN-α2, IL-6,
GMCSF, IL-8 indicated a robust antiviral and inflammatory response occurring in these T2DM
patients. Additionally, these patients also showed an increase in type II cytokines such as IL-4
and IL-9, representing the anti-inflammatory response. Here, we understand that the presence
of both inflammatory and anti-inflammatory cytokines in T2DM patients as indicators of a
dysregulated immune response where the anti-inflammatory response was clearly incapable of
shutting the cytokine storm in these patients(23).

In diabetic patients, there is an increase in the inflammatory cytokines, CRP levels and
the presence of blood clots. As similar symptoms are associated with Covid-19 i.e. elevated
CRP levels, D-dimer and inflammatory cytokines, this suggests towards a synergistic effect
found in T2DM patients infected with SARS-Cov2, which was reduced in NDM patients.
Additionally, our study suggested an increase in IL-15 and 1L-7 in T2DM patients when
compared to NDM and healthy volunteers. As it is known that IL-15 is crucial for promoting
cytotoxic activity of both, NK and CD8[+] T cells and is also involved in memory CD8[+] T cell
differentiation, it should have led to increase in the CD8[+] population(24). However, we found
a decrease in the CD8[+] cell population, specifically in the EM compartment, again indicating
a dysregulated or non-responsive immune response in these patients. This might be a
consequence of reduced receptors for IL-15 in these cells. As IL-7 is also required for T cell
development and maintenance(25), its increased numbers in T2DM patients partially
corroborated with the increase in CD8[+] TE cells and CD4[+] EM cells. However, as a
consequence of immune dysregulation, neither the elevated levels of the above mentioned
cytokines nor the CD8[+] T cells, are capable of controlling the inflammation.

The sCD40L is one of the major co-stimulatory molecules on activated T cells that
interact with CD40 on B cells and is responsible for immunoglobulin isotype switching in the
membrane-bound form(26). In our study, levels of sCD40L were elevated in NDM as
compared to the T2DM patients indicating that the NDM patients may be undergoing an
immuno suppressive reaction, as elevated sCD40L is also associated to immunosuppression.
GRO (CXCL1) is a chemokine that attracts variety of immune cells, particularly neutrophils
and is also implicated in wound healing process(27). Similar to sCD40L levels, GRO levels
were also higher in NDM as compared to T2DM patients indicating that both inflammation
and healing were occurring simultaneously in these patients. However, the healing mechanism
was definitely poorer in T2DM patients. The levels of Macrophage-derived chemokine
(MDC/CCL2) were also higher in NDM patients than T2DM patients. As MDC is known to
be elevated in lung inflammation and hemorrhage but reduced in T2DM symptomatic patients,
this is also suggesting towards a possible immunosuppressive function of MDC in this
population(28).

To understand how different antibody isotypes influence the clinical phenotype in
T2DM and NDM patients, we did an antibody isotyping from the plasma of these patients.
Here, we found significant difference in the levels of IgA, IgM, IgG1, IgG2 and IgE between
the two groups but IgG3 and IgG4 did not show any significant difference and we understand
that the structure and function of different antibody isotypes vary from one another. For
example, IgG1, IgG2 and IgG3 can fix complement(29). However, IgG4 cannot fix
complement and is unable to induce antibody mediated cell cytotoxicity (ADCC). Similar to
previous reports, we found an increase in the levels of IgE in T2DM patients(8, 16). This
corroborated with the increase in the Th2 subset along with increase in cytokines, IL-4 and IL-
9 in T2DM patients, as it is known that IgE is associated with Type II response. In addition,
there was also an increase in IgG1 and IgG2 in these patients, indicative of the inflammatory
response. However, as IgE was high in these T2DM patients, we could see no significant
difference in IgG4 levels. This could be accredited to the competition between IgE and IgG4
against each other, for fixation sites in basophils and mast cells, as suggested by previous
reports(30). IgA levels were also found to be higher in T2DM, indicating towards viral immune
response at mucosal surfaces(31). With respect to IgM, its increased levels indicate towards
an ongoing infection. However, the lack of any longitudinal profiling restricts us from
understanding its specificity to antigen(32).

With respect to our whole blood analysis by mass cytometry, we found an increase in
the percentage of granulocytes in T2DM patients when compared to healthy controls indicating
underlying inflammatory responses in these patients. This is consistent with other reports
which suggested an increase in granulocytes with increasing severity(15, 33). On the other
hand, we found a decreasing percentage of monocytes in T2DM patients as compared to NDM
patients **(Table 1)**, although the difference was not significant. This has also been reported in
a previous study showing the absence of any significant changes in the total monocytes and its
subsets in T2DM patients with various comorbidities, T2DM being one of them(15).
Additionally, we also found that T2DM patients showed a decreasing trend in lymphocyte
percentage as compared to NDM and healthy volunteers, as reported in previous studies (4, 8,
15). One of the significant findings from our study is that, there is a drastic reduction in the
percentage of total dendritic cells in T2DM patients when compared to both, NDM and healthy
controls. Within the dendritic cell compartments, there is a decrease of pDCs but mDCs have
increased in T2DM patients when compared with healthy controls. In general, pDCs are known
to secrete type I interferons in response to viral infection(34). However, our study has shown
a decrease in the percentage of pDCs, but increase in the cytokine IFN-α2. This is suggesting
that although these cells are secreting high levels of type I interferon, their numbers are depleted
in T2DM patients and thus, are unable to control the infection. With respect to mDCs, they are
known to secrete the cytokines IL-12 and TNF-α, which polarizes the T cell towards a type I
response, crucial for controlling the viruses(35, 36). Our study also observed an increase in
the population of mDCs along with increase in the cytokines, IL-12p40, TNF-α in T2DM.
Although certain reports have suggested a decrease of mDCs along with pDCs, we observed
the presence of high numbers of mDCs in diabetic patients, indicating an ongoing low grade
inflammation, which is further augmented with Covid-19 infection. However, since we did
not do a longitudinal profiling, we could not delineate the mDC dynamics in these patients(37,
38).

With respect to basophils, literature suggested a decrease in its population with
increasing severity(39). We found a similar trend, as the basophil population decreased in
T2DM patients as compared to healthy controls. Similarly, MAIT/iNKT cells were
significantly decreased in T2DM patients as compared to NDM and healthy controls.
However, no significant differences were observed in the percentage of gamma delta T cells.
A decrease in MAIT/iNKT could indicate towards a migration to other inflamed tissues
including the lungs, as suggested by the previous reports. With respect to NK cell population,

it is known that they play a crucial role in fighting viral infections and previous reports

suggested a decrease in their numbers in T2DM cases. However, our population of study did

not show any significant difference amongst the groups and as mentioned earlier may be a

consequence of the time of sampling of patients(40, 41).

In case of B cells, we observed a slight elevation in its percentage in T2DM patients,

when compared to NDM and healthy controls, but statistical difference was found only

between T2DM patients and healthy volunteers. Although no difference was observed in naive

or memory B cell compartment, plasmablasts were significantly elevated in T2DM patients as

compared to healthy controls. The increase in plasmablasts in T2DM Covid-19 has also been

reported in other studies (8, 15)but unlike our study, they could not correlate it with any

comorbidity. Interestingly, there are reports suggesting an increase in extrafollicular B cell

response in Covid-19 and other inflammatory diseases such as Systemic Lupus Erythematous.

Since most of the Covid-19 patients had type II diabetes in our study, we hypothesize that these

two factors might have augmented the plasmablast production. Although we did not analyze

the clonality of B cells, but earlier study has suggested an oligoclonal expansion of B cells in

T2DM Covid-19, in turn correlating with increased plasmablast production. Increased B cell

population in T2DM patients also corroborated with the fact that most of the immunoglobulins

(IgM, IgA, IgG and IgE) were elevated in these patients as opposed to NDM and healthy

volunteers. Altogether, we found an increase in the plasmablast population in T2DM as

compared to NDM patients and healthy volunteers, suggesting that the low grade inflammation

in Diabetic patients resulted in increase in plasmablast population, which augmented further

with Covid-19 infection in these patients.

T2DM patients also displayed a lower percentage of $CD45^+CD3^+$ T cells as compared

to healthy controls. Within $CD3^+$ T cell population, there was a decrease in the percentage of

$CD8^+$ T cells in T2DM as compared to healthy volunteers. However, $CD4^+$ T cells did not

show any statistically significant difference amongst the groups. Earlier reports have suggested

the loss of $CD8^+$ T cells being greater than $CD4^+$ T cells. Since our population of study was

small with heterogenous manifestation, this could explain why $CD4^+$ T cells did not show any

difference amongst the groups. Additionally, several reports indicate towards T cell apoptosis

or migration to other tissues in SAR-CoV2 infection, leading to decrease in T cell population

in the periphery and could account for the decrease in T cells in the T2DM patients(42-44). In

general, there was decrease in the $CD3^+$ T cell population, where the $CD8^+$ compartment was

affected more significantly than $CD4^+$.

Type II diabetes has been reported to have various aberrancies such as an impaired

differential potential and secretion of multiple proinflammatory cytokines such as TNF-α and

IFN-γ(37, 45). Although we could not find any significant difference amongst the groups in

the percentage of Th1 like, Th17 like, Treg or cTfh subtypes, there was an increase in the

proinflammatory cytokines. We understand that the absence of difference amongst the groups

with respect to T cell subsets can be accounted to the time of sampling. On the other hand, our

study found an increase in the percentage of Th2 cells secreting type II cytokines such as IL-4

and IL-9. The high numbers of Th2 cells partially explained the augmented levels of

immunoglobulins in these patients, specifically IgE which is induced by cytokines such as IL-

4. Taken together, this indicated that the Diabetic patients already had low grade inflammation.

When encountered with Covid-19, dysregulation in the immune system escalated to such levels

that even with the increase in Th2 population secreting anti-inflammatory cytokines,

inflammation could not subdue.

We also analyzed for the naïve, effector and memory compartments in both $CD4^+$ and

$CD8^+$ T cells. In $CD4^+$ T cells, we did not see any difference in the overall percentage of naive

($CCR7^+CD45RA^+CD45RO^-$) and central memory cells ($CCR7^+CD45RA^-CD45RO^{+)}$ amongst

the various groups. However, effector memory cells ($CCR7^-CD45RA^-CD45RO^+CD27^+$) were

elevated in T2DM patients when compared to healthy controls, suggesting cross reactivity with

other family of coronavirus, as reported by a previous study from India(46, 47). An important

point to note here is that central memory cells did not show any significant difference and this

may be due to the fact that CM cells are lymph node residents as opposed to EM cells, which

circulate in blood and thus, respond faster to the antigen (data not shown). However, since we

did not analyze for the antigen specific cells, therefore we were unable to find any significant

difference in the percentage of CM cells. Terminal effector ($CCR7^-CD45RA^-CD45RO^+CD27^-$

) cells were decreased in T2DM patients with Covid-19 as compared to healthy controls,

corroborating with the observation that these cells are first to react with the virus and the

subsequent proinflammatory cytokine milieu is responsible for the apoptosis of these cells.

In $CD8^+$ T cell compartment, EM ($CCR7^-CD27^+$) cells were elevated in NDM as

compared to T2DM patients, indicating an effective immune response in these patients but less

effective in T2DM patients. TE ($CCR7^-CD27^-$) cells increased in T2DM patients indicating

that these cells are highly inflammatory and might be insensitive to AICD (47-49).

Altogether, our data suggests that Meta inflammation present in these T2DM patients is responsible for aggravation of the inflammation occurring due to Covid-19. With T2DM being one of the most common morbidity presents in India, these patients remain most vulnerable and were susceptible to secondary infections like mucormycosis. Though vaccinations are lowering the severity of the disease, reported waning antibody response in 3-6 months is a major concern apart from how long the protective immunity from T cells will work in these patients is also yet to be ascertained. Our study is essentially aimed at understanding how a low grade chronic inflammatory disorder such as Type II Diabetes dictates the immune response during the pathogenesis of Covid-19. We understand that our study is not devoid of limitations such as a small sample size, absence of longitudinal study or follow up, but nevertheless it will help in understanding the vulnerability of these patients and subsequent planning of vaccine coverage in these patients or future vaccines booster dosages to them.

References

1. Chen Y, Liu Q, Guo D. Emerging coronaviruses: Genome structure, replication, and pathogenesis. J Med Virol. 2020;92(10):2249.
2. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497-506.
3. Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. N Engl J Med. 2020;383(25):2451-60.
4. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of Immune Response in Patients With Coronavirus 2019 (COVID-19) in Wuhan, China. Clin Infect Dis. 2020;71(15):762-8.
5. Iba T, Levy JH, Levi M, Thachil J. Coagulopathy in COVID-19. J Thromb Haemost. 2020;18(9):2103-9.
6. Annunziato F, Romagnani C, Romagnani S. The 3 major types of innate and adaptive cell-mediated effector immunity. J Allergy Clin Immunol. 2015;135(3):626-35.
7. O'Shea JJ, Paul WE. Mechanisms underlying lineage commitment and plasticity of helper CD4+ T cells. Science. 2010;327(5969):1098-102.
8. Lucas C, Wong P, Klein J, Castro TBR, Silva J, Sundaram M, et al. Longitudinal analyses reveal immunological misfiring in severe COVID-19. Nature. 2020;584(7821):463-9.
9. Chatterjee S, Datey A, Sengupta S, Ghosh A, Jha A, Walia S, et al. 2021.
10. Jogdand GM, Sengupta S, Bhattacharya G, Singh SK, Barik PK, Devadas S. Inducible Costimulator Expressing T Cells Promote Parasitic Growth During Blood Stage Plasmodium berghei ANKA Infection. Front Immunol. 2018;9:1041.
11. Prodjinotho UF, von Horn C, Debrah AY, Batsa Debrah L, Albers A, Layland LE, et al. Pathological manifestations in lymphatic filariasis correlate with lack of inhibitory properties of IgG4 antibodies on IgE-activated granulocytes. PLoS Negl Trop Dis. 2017;11(7):e0005777.
12. Bagwell CB, Hunsberger B, Hill B, Herbert D, Bray C, Selvanantham T, et al. Multi-site reproducibility of a human immunophenotyping assay in whole blood and peripheral blood mononuclear cells preparations using CyTOF technology coupled with Maxpar Pathsetter, an automated data analysis system. Cytometry B Clin Cytom. 2020;98(2):146-60.

13.   Geanon D, Lee B, Gonzalez-Kozlova E, Kelly G, Handler D, Upadhyaya B, et al. A streamlined
whole blood CyTOF workflow defines a circulating immune cell signature of COVID-19. Cytometry A.
2021;99(5):446-61.
14.   Tay MZ, Poh CM, Renia L, MacAry PA, Ng LFP. The trinity of COVID-19: immunity, inflammation
and intervention. Nat Rev Immunol. 2020;20(6):363-74.
15.   Kuri-Cervantes L, Pampena MB, Meng W, Rosenfeld AM, Ittner CAG, Weisman AR, et al.
Comprehensive mapping of immune perturbations associated with severe COVID-19. Sci Immunol.
2020;5(49).
16.   Zhao R, Sun Y, Zhang Y, Wang W, Wang S, Wang C, et al. Distinguishable Immunologic
Characteristics of COVID-19 Patients with Comorbid Type 2 Diabetes Compared with Nondiabetic
Individuals. Mediators Inflamm. 2020;2020:6914878.
17.   Sohn Y, Jeong SJ, Chung WS, Hyun JH, Baek YJ, Cho Y, et al. Assessing Viral Shedding and
Infectivity of Asymptomatic or Mildly Symptomatic Patients with COVID-19 in a Later Phase. J Clin
Med. 2020;9(9).
18.   Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in Upper
Respiratory Specimens of Infected Patients. N Engl J Med. 2020;382(12):1177-9.
19.   Li G, Chen Z, Lv Z, Li H, Chang D, Lu J. Diabetes Mellitus and COVID-19: Associations and
Possible Mechanisms. Int J Endocrinol. 2021;2021:7394378.
20.   Chen L, Deng H, Cui H, Fang J, Zuo Z, Deng J, et al. Inflammatory responses and inflammation-
associated diseases in organs. Oncotarget. 2018;9(6):7204-18.
21.   Yaribeygi H, Sathyapalan T, Jamialahmadi T, Sahebkar A. The Impact of Diabetes Mellitus in
COVID-19: A Mechanistic Review of Molecular Interactions. J Diabetes Res. 2020;2020:5436832.
22.   Guo W, Li M, Dong Y, Zhou H, Zhang Z, Tian C, et al. Diabetes is a risk factor for the progression
and prognosis of COVID-19. Diabetes Metab Res Rev. 2020:e3319.
23.   Ozsurekci Y, Aykac K, Er AG, Halacli B, Arasli M, Oygar PD, et al. Predictive value of
cytokine/chemokine responses for the disease severity and management in children and adult cases
with COVID-19. J Med Virol. 2021;93(5):2828-37.
24.   Verbist KC, Klonowski KD. Functions of IL-15 in anti-viral immunity: multiplicity and variety.
Cytokine. 2012;59(3):467-78.
25.   Tani-ichi S, Shimba A, Wagatsuma K, Miyachi H, Kitano S, Imai K, et al. Interleukin-7 receptor
controls development and maturation of late stages of thymocyte subpopulations. Proc Natl Acad Sci
U S A. 2013;110(2):612-7.
26.   Huang J, Jochems C, Talaie T, Anderson A, Jales A, Tsang KY, et al. Elevated serum soluble CD40
ligand in cancer patients may play an immunosuppressive role. Blood. 2012;120(15):3030-8.
27.   Zaja-Milatovic S, Richmond A. CXC chemokines and their receptors: a case for a significant
biological role in cutaneous wound healing. Histol Histopathol. 2008;23(11):1399-407.
28.   Richter JR, Sutton JM, Belizaire RM, Friend LA, Schuster RM, Johannigman TA, et al.
Macrophage-derived chemokine (CCL22) is a novel mediator of lung inflammation following
hemorrhage and resuscitation. Shock. 2014;42(6):525-31.
29.   van der Zee JS, van Swieten P, Aalberse RC. Inhibition of complement activation by IgG4
antibodies. Clin Exp Immunol. 1986;64(2):415-22.
30.   Dodev TS, Bowen H, Shamji MH, Bax HJ, Beavil AJ, McDonnell JM, et al. Inhibition of allergen-
dependent IgE activity by antibodies of the same specificity but different class. Allergy.
2015;70(6):720-4.
31.   Sterlin D, Mathian A, Miyara M, Mohr A, Anna F, Claer L, et al. IgA dominates the early
neutralizing antibody response to SARS-CoV-2. Sci Transl Med. 2021;13(577).
32.   Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody Responses to SARS-CoV-2 in
Patients With Novel Coronavirus Disease 2019. Clin Infect Dis. 2020;71(16):2027-34.
33.   Vitte J, Diallo AB, Boumaza A, Lopez A, Michel M, Allardet-Servent J, et al. A Granulocytic
Signature Identifies COVID-19 and Its Severity. J Infect Dis. 2020;222(12):1985-96.

34.   Cervantes-Barragan L, Zust R, Weber F, Spiegel M, Lang KS, Akira S, et al. Control of
coronavirus infection through plasmacytoid dendritic-cell-derived type I interferon. Blood.
2007;109(3):1131-7.

35.   van der Molen RG, Sprengers D, Binda RS, de Jong EC, Niesters HG, Kusters JG, et al. Functional
impairment of myeloid and plasmacytoid dendritic cells of patients with chronic hepatitis B.
Hepatology. 2004;40(3):738-46.

36.   Cella M, Salio M, Sakakibara Y, Langen H, Julkunen I, Lanzavecchia A. Maturation, activation,
and protection of dendritic cells induced by double-stranded RNA. J Exp Med. 1999;189(5):821-9.

37.   Yao K, Lu H, Huang R, Zhang S, Hong X, Shi H, et al. Changes of dendritic cells and fractalkine
in type 2 diabetic patients with unstable angina pectoris: a preliminary report. Cardiovasc Diabetol.
2011;10:50.

38.   Sun Y, Zhou J, Ye K. White Blood Cells and Severe COVID-19: A Mendelian Randomization
Study. J Pers Med. 2021;11(3).

39.   Jouan Y, Guillon A, Gonzalez L, Perez Y, Boisseau C, Ehrmann S, et al. Phenotypical and
functional alteration of unconventional T cells in severe COVID-19 patients. J Exp Med. 2020;217(12).

40.   Parrot T, Gorin JB, Ponzetta A, Maleki KT, Kammann T, Emgard J, et al. MAIT cell activation
and dynamics associated with COVID-19 disease severity. Sci Immunol. 2020;5(51).

41.   Zhai X, Qian G, Wang Y, Chen X, Lu J, Zhang Y, et al. Elevated B Cell Activation is Associated
with Type 2 Diabetes Development in Obese Subjects. Cell Physiol Biochem. 2016;38(3):1257-66.

42.   Shrotri M, van Schalkwyk MCI, Post N, Eddy D, Huntley C, Leeman D, et al. T cell response to
SARS-CoV-2 infection in humans: A systematic review. PLoS One. 2021;16(1):e0245532.

43.   Mazzoni A, Salvati L, Maggi L, Capone M, Vanni A, Spinicci M, et al. Impaired immune cell
cytotoxicity in severe COVID-19 is IL-6 dependent. J Clin Invest. 2020;130(9):4694-703.

44.   Gil-Etayo FJ, Suarez-Fernandez P, Cabrera-Marante O, Arroyo D, Garcinuno S, Naranjo L, et al.
T-Helper Cell Subset Response Is a Determining Factor in COVID-19 Progression. Front Cell Infect
Microbiol. 2021;11:624483.

45.   Ansari A, Arya R, Sachan S, Jha SN, Kalia A, Lall A, et al. Immune Memory in Mild COVID-19
Patients and Unexposed Donors Reveals Persistent T Cell Responses After SARS-CoV-2 Infection. Front
Immunol. 2021;12:636768.

46.   Lipsitch M, Grad YH, Sette A, Crotty S. Cross-reactive memory T cells and herd immunity to
SARS-CoV-2. Nat Rev Immunol. 2020;20(11):709-13.

47.   Harty JT, Tvinnereim AR, White DW. CD8+ T cell effector mechanisms in resistance to
infection. Annu Rev Immunol. 2000;18:275-308.

48.   Reiser J, Banerjee A. Effector, Memory, and Dysfunctional CD8(+) T Cell Fates in the Antitumor
Immune Response. J Immunol Res. 2016;2016:8941260.

49.   Martin MD, Badovinac VP. Defining Memory CD8 T Cell. Front Immunol. 2018;9:2692.

Fig 1. Analysis of Viral load in Oropharyngeal samples in Covid-19 patients.(A) Bar diagram depicting the viral copy number of Oropharyngeal (OP) samples collected from T2DM(n=25) and NDM(n=10) COVID-19 patients.**(B)** Bar diagram representing the **ΔΔ** CT of OP samples of NDM and T2DM COVID-19 patients. The Mann- Whitney (non parametric, two tailed) test was performed. ns, not significant. All error bars were SEM.

Fig 2.Analysis of the cytokines and chemokines in T2DM ,NDM patients infected with SARS-CoV2

The bar diagrams representing cytokines and chemokines which were evaluated from COVID -19 positive plasma samples of T2DM(n=25) , NDM(n=10) and Covid-19 negative and non-diabetic healthy controls(n=10).**(A)**IL-6.TNF-α,GCSF,IL-12p40, IL-15,IL-1α,MIP-1β,GMCSF,IFN-α2,IL-10,VEGF and IL-Rα were highly elevated in T2DM patients when compared with healthy controls and NDM patients**. (B)** IFN-γ, EGF and IL-8 were moderately elevated in T2DM patients when compared to healthy controls and NDM.**(C)** IP-10, IL-9 and IL-4 were slightly elevated in T2DM patients when compared to healthy controls and NDM**(D)** Eotaxin was moderately elevated in healthy controls and NDM when compared to T2DM patients.**(E)** sCD40L was elevated in both NDM and T2DM when compared to healthy controls, MDC was elevated in NDM patients when compared to T2DM patients.**(F)**GRO was elevated in NDM patients when compared to healthy controls. The Kruskal Wallis Test (non parametric) with post hoc Dunn's multiple comparison test was performed p<0.05 was considered statistically significant (*), p<0.01 was considered to be very significant (**) , $P < 0.001$ was considered to highly significant(***) , $P < 0.0001$ was considered extremely significant(****) ns, not significant. All error bars were SEM.

Fig 3.Analysis of the circulating antibodies in T2DM ,NDM patients infected with SARS-CoV2
The bar diagram representing circulating antibody isotypes (IgA, IgM, IgG1, IgG2, IgG3,IgG4 and IgE) which were evaluated from COVID -19 positive plasma samples of T2DM(n=25) , NDM(n=10) and Covid-19 negative and non-diabetic healthy controls(n=10). The Kruskal Wallis Test (non parametric) withwith post hoc Dunn's multiple comparison test was performed p<0.05 was considered statistically significant (*), p<0.01 was considered to be very significant (**) , *P* < 0.001 was considered to highly significant(***) , *P* < 0.0001 was considered extremely significant(****) ns, not significant. All error bars were SEM.

Fig 4.Innate Immune Cells in T2DM and NDM patients. Whole blood from Covid-19 positive T2DM(n=10), NDM(n=5) and Covid-19 negative and non-diabetic healthy controls (n=5)was stained heavy metal tagged antibody and analyzed in mass cytometer.**(A)**Representative Gating strategy to identify CD45⁺CD66b^low lymphocytes and CD45⁻CD66b^high Granulocytes. Statistical analysis of percentage of Granulocytes and lymphocytes.**(B)** Gating strategy for CD123⁺CD11c⁻ pDCs . Statistical analysis for the frequency of total DCs and pDCs.**(C)** Gating strategy for CD11c⁺CD38⁺ mDCs.Statistical analysis for the percentage of mDCs.**(D)**Gating strategy for CD123⁺CD294⁺ Basophils. Statistical analysis for the frequency of basophils.**(E)** Gating strategy for MAIT/NKT cells.Statistical analysis for the frequency of MAIT/NKT cells.). The Kruskal Wallis Test (non parametric) withwith post hoc Dunn's multiple comparison test was performed p<0.05 was considered statistically significant (*), p<0.01 was considered to be very significant (**) , *P* < 0.001 was considered to highly significant(***) , *P* < 0.0001 was considered extremely significant(****) ns, not significant. All error bars were SEM.

Fig 5.Adaptive Immune Cells in T2DM and NDM patients. Whole blood from Covid-19 positive T2DM(n=10), NDM(n=5) and Covid-19 negative and non-diabetic healthy controls (n=5)was stained heavy metal tagged antibody and analyzed in mass cytometer.**(A)**Representative Gating strategy to identify CD38+CD20- Plasmablasts. Statistical analysis of percentage of plasmasblasts .**(B)** Gating strategy to identify CD45+CD3+CD4+ and CD45+CD3+CD8+ T cells. Statistical analysis for the frequency of CD3+,CD4+ and CD8+ T cells.**(C)** Gating strategy for CD4+CXCR3-CCR6-Th2 like cells.Statistical analysis for the percentage of Th2 like cells.. **(D)** Gating strategy to identify CD4+ TE(CD45RO+CD27-)and CD4+ EM cells(CD45RO+CD27+). Statistical analysis for the percentages of CD4+ TE and EM cells.**(E)** Gating strategy to identify CD8+ TE (CCR7- CD27-) and EM(CCR7-CD27+) cells. Statistical analysis for the percentages of the CD8+TE and EM cellsThe Kruskal Wallis Test (non parametric) with post hoc Dunn's multiple comparison test was performed p<0.05 was considered statistically significant (*), p<0.01 was considered to be very significant (**) , *P* < 0.001 was considered to highly significant(***) , *P* < 0.0001 was considered extremely significant(****) ns, not significant. All error bars were SEM.

Fig.6 t-SNE analysis of lowest and highest expressed markers in T2DM patients. **(A)** CD8 and CD3 showing decreased expression in T2DM patients.**(B)** HLA-DR and CD 14 showing increased expression in T2DM patients when compared to controls.

Table 1- Demographic details and laboratory findings of T2DM and NDM patients infected with SARS-CoV2

	T2DM n=25	NDM n=10	P VALUE
Age, median (IQR), years	46 (36-59)	49(40.25-55.50)	N. A
Sex, male/female, n (%)	17 (68%)/8(32%)	7 (70%)/3(10%)	N. A
Haemoglobin, gm/dl median (IQR)	12.2 (11.35-13.60)	14.25 (13.7-14.9)	0.0475
Platelets 10^3/μL median (IQR)	193 (154.5-235)	165 (159.5-197.5)	0.4027
White Blood Cells 10^3/μL, median (IQR)	6.1 (5.00-57.65)	5.8 (5.2-5.575)	0.4172
Neutrophils %, median (IQR)	75 (65.5-84)	55 (44.75-62)	0.0004
Lymphocytes%, median (IQR)	22 (15.5-29.75)	36 (26.75-39.75)	0.0349
Eosinophils %, median (IQR)	0	3 (3-4.5)	<0.0001
Monocytes%, median (IQR)	7 (5-9)	9 (8.5-10)	0.0373
D-dimer μg/mL, median (IQR)	0.45 (0.29-1.61)	0.19 (0.15-0.25)	0.0048
Ferritin ng/mL, median (IQR)	224.7 (145.9-464.1)	115 (110-289.5)	0.4975
Fasting Plasma Glucose mg/dL, median (IQR)	215 (142.72-244)	95.5 (88-101)	<0.0001
Hb1AC %, median (IQR)	6.9 (6-9)	N. T	N. A
CRP mg/mL, median (IQR)	69.62 (34.95-211)	13.28 (4.44-18.65)	0.0013
Serum SGOT U/L, median (IQR)	46.9 (30.2-64.6)	38 (30.75-43.25)	0.4432
Serum SGPT U/L, median (IQR)	27.6 (20-42)	40.5 (24.5-76.5)	0.1335

Hb1AC- Glycated haemoglobin, CRP- C-Reactive Protein, Serum SGOT-serum glutamic-oxaloacetic transaminase, Serum SGPT- serum glutamic-pyruvic transaminase, N.T-Not Tested, N.A-Not Applicable.

692