Increasing trends in mental health problems among Chinese young adolescents: results from repeated cross-sectional data in Changsha 2016-2020

Zhipeng Wu[1], Biao Wang[1], Zhulin Zou[2], Zhening Liu[1], Xudong Chen[1,*] and Yicheng Long[1,*]

[1]National Clinical Research Center for Mental Disorders, and Department of Psychiatry, The Second Xiangya Hospital, Central South University, Changsha, Hunan, China

[2]The High School Attached to Hunan Normal University and Bocai Experimental School, Changsha, Hunan, China

* Correspondence:

Xudong Chen

xudongchen@csu.edu.cn

and

Yicheng Long,

yichenglong@csu.edu.cn

Abstract

This study performed a repeated cross-sectional analysis to explore possible trends in mental health problems among Chinese adolescents during 2016-2020. A total of 2837 seventh-grade students from Changsha city were surveyed in three waves in 2016, 2019 and 2020. The results showed that obsessive–compulsive tendencies, interpersonal sensitivity, depression, anxiety, academic stress and emotional disturbance problems were significantly increased in surveyed adolescents during 2016-2020. Moreover, positive rates of most of these problems were higher in females than males, and were significantly increased in only females. These results highlight the importance of focusing on mental health among Chinese adolescents, especially girls.

Keywords: mental health; adolescents; depression; anxiety

1. Introduction

Childhood and early adolescence are crucial periods for mental health development with particularly high risk for mental health problems (Garcia-Carrion et al., 2019; Kessler et al., 2005). These problems can persist into adulthood and even lead to serious mental disease if they are undetected or not treated appropriately (Pine et al., 1998; Zeng et al., 2019; Zhang et al., 2019).

Over the past decades, China has witnessed great developments in economy and society; however, multiple studies have reported a dramatic decrease of mental health in China, especially in Chinese adolescents during the same period (Feng et al., 2020; Su and Liu, 2020; Wang and Tapia Granados, 2019; Xin et al., 2020). For example, a recent cross-temporal meta-analysis showed that Chinese adolescents' depression was significantly increased from 1989 to 2018 (Su and Liu, 2020). Another cross-temporal meta-analysis, which was based on data between 1992 and 2017, also reported that Chinese adolescents' anxiety level was significantly increased since 1992 (Xin et al., 2020). To improve public mental health, it is thus important to assess the prevalence and changes over time in prevalence of mental health problems in Chinese adolescents (Su and Liu, 2020).

Although having gained important insights into trends in mental health problems in Chinese adolescents in recent years, the above-mentioned published literatures are limited in several ways. First, reports on newly trends after the year 2018 are sparse. Second, most of them only focused on a single dimension of mental health such as depression or anxiety, while a more comprehensive survey in multiple dimensions is rare. Third, most of these studies are cross-temporal meta-analyses, which pooled previous studies using the same mental health assessing instruments (e.g., Self-rating Depression Scale (Su and Liu, 2020)) together. However, such an approach may not be able to fully address the potential bias caused by regional diversity (e.g., the pooled studies were across different provinces in China with different economic and educational levels (Su and Liu, 2020)).

In this study, we performed a repeated cross-sectional analysis to overcome the above limitations. Repeated cross-sectional studies utilize the same instruments and

methodologies in the same population over multiple time timepoints; therefore, they

were thought to be most suitable for investigating trends of mental health statuses over

time (Myhr et al., 2020; Richter et al., 2019). In specific, a total of 2837 adolescents

from a sample high school in Changsha city, China, were surveyed in three waves from

2016 to 2020. Mental health status was assessed using a validated multi-dimensional

mental health scale, Mental Health Inventory of Middle School Students (MMHI-60)

(Wang et al., 1997; Wu et al., 2021a) and compared between different years.

2.   Methods

This study uses data collected from 2016 to 2020 in a total of 2837 Chinese junior

high school students in three waves (978 students in September 2016; 949 students in

September 2019; and 910 students in September 2020). All participants were new

seventh-grade students from the same campus of a high school (Tianding campus of

The High School Attached to Hunan Normal University and Bocai Experimental

School in Changsha city, Hunan province, China) when they completed the survey.

Note that participants with missing data for any item of the MMHI-60 in the surveys

have been excluded from the study. All participants and their supervisors provided

written informed consents and the study was approved by the Ethics Committees of the

Second Xiangya Hospital of Central South University, Changsha.

All participants completed the MMHI-60 in the classroom to assess their mental

health statuses. MMHI-60 is a validated and widely used self-report measure of mental

health problems, which includes 10 subscales of distinct dimensions (60 items in total,

and 6 items for each subscale): obsessive–compulsive tendencies, paranoid ideation,

hostility, interpersonal sensitivity, depression, anxiety, academic stress, maladaptation,

emotional disturbance and psychological imbalance (Du et al., 2019; Luo et al., 2020;

Wang et al., 1997; Wu et al., 2021a; Xiong et al., 2021). All items were scored from 1

to 5, and higher scores indicate more severe mental health problems. According to

Wang et al. (Wang et al., 1997), a cutoff of average points of a subscale >=2 is identified

as having a mental health problem (positive for that subscale). This cutoff has shown

good specificity and sensitivity in previous research (Du et al., 2019; Luo et al., 2020;

Zhao and Liao, 2016). Positive rates of each subscale were calculated based on such a

cutoff, and Cochran–Armitage trend tests were used to determine if there are increasing

or decreasing trends of positive rates during the period of 2016-2020 (Yang et al., 2017).

Moreover, considering that sex differences in mental health among adolescents have

been widely reported (Kim et al., 2018; Luo et al., 2020; Myhr et al., 2020), positive

rates of all subscales were further compared between males and females using Chi-

square test; significance of trends were also tested in males and females separately.

3. Results

Demographic and mental health characteristics of students in each wave were

shown in **Table 1**. There was no significant difference in sex ratio among the three

waves ($\chi^2 = 1.873$, $p = 0.392$). Data on participants' age was only available for the

second wave and could not be compared. Cochran–Armitage trend tests showed that

positive rates of the obsessive–compulsive tendencies ($p < 0.001$), interpersonal

sensitivity ($p = 0.015$), depression ($p < 0.001$), anxiety ($p = 0.001$), academic stress (p

< 0.001) and emotional disturbance ($p = 0.006$) problems were significantly increased

during the period of 2016 to 2020, while no significant increasing or decreasing trends

were found in positive rates of the paranoid ideation, hostility, maladaptation and

psychological imbalance problems (all $p > 0.05$). The trends in positive rates of each

mental health problem were also visualized in **Figure 1**.

The females had significantly higher positive rates of obsessive–compulsive

tendencies, interpersonal sensitivity, depression, anxiety and academic stress, but a

significantly lower positive rate of psychological imbalance than males (**Figure 2 and**

Supplementary Table 1). During the period of 2016-2020, positive rate of obsessive–

compulsive tendencies was significantly increased in both males and females, while

positive rates of interpersonal sensitivity, depression, anxiety, academic stress,

maladaptation, emotional disturbance and psychological imbalance were significantly

increased in only females (**Figure 3 and Supplementary Tables 2-3**).

4. Discussion

In this study, we performed a repeated cross-sectional analysis to explore the
possible trends in mental health problems among Chinese adolescents over the years of
2016-2020. Compared with most published studies for similar purposes, we had
advantages in methodologies (uniformity of surveyed populations) and data timeliness
(until the most recent years).

Our results showed that positive rates of multiple mental health problems
including obsessive–compulsive tendencies, interpersonal sensitivity, depression,
anxiety, academic stress and emotional disturbance problems were all significantly
increased during September 2016 to September 2020 in surveyed participants. Thus, as
a supplement to previous studies analyzing data until 2018, our study suggests that
mental health in Chinese adolescents could have been deteriorating in the most recent
years. It is noteworthy that the COVID-19 outbreak in 2020, which has been widely
reported to have negative mental effects on worldwide population (Ahmed et al., 2020;
Rajkumar, 2020; Roy et al., 2020; Wu et al., 2021b). Therefore, the pandemic-related
effects may play an important role in these trends besides other reported social and
psychological factors (e.g., increasing divorce rate in China (Mo, 2017)).

Further analyses showed that positive rates of most of the above mental health
problems were significantly higher in females than males and were significantly
increased in only females during 2016-2020. Although being somewhat surprising,
similar results have been reported among adolescents in other countries (Eyuboglu et
al., 2021; Holik et al., 2021; Thorisdottir et al., 2021). It highlights the importance of
focusing on girls since they may be more affected by mental problems.

Our study has several limitations. First, this survey was not performed in years of
2017 and 2018. Second, data on age was lacking for most participants. However, since
children were required to enter school until 6 years old in China (Zhang et al., 2017),
all participants should had a close age around 12 when they were surveyed. Third, the
survey was performed in only a single school in Changsha city, and in only seventh-
grade students in early adolescence. Thus, it should be cautious to interpret our results
as trends in nationwide populations. Further studies conducted in a larger sample, in
wider regions including rural areas, and in a wider range of age would be necessary to
expand our knowledge in mental health among Chinese adolescents.

5. Conclusion

In this study, we found that positive rates of multiple mental health problems were significantly increased during 2016-2020 in a group of surveyed adolescents in Changsha city, China. Moreover, positive rates of most of these problems were found to be significantly higher in females than males, and significantly increased during 2016-2020 in only females. Such results highlight the importance of focusing on mental health problems among Chinese adolescents, especially among girls.

Funding

This work was supported by the Natural Science Foundation of Hunan Province, China (2021JJ40851 to YL, 2020JJ5800 to XC) and the National Natural Science Foundation of China (82071506 to ZL).

Acknowledgments

We would like to thank all the subjects who served as research participants.

Declaration of competing interest

The authors declare no conflict of interest.

Supplementary data

Supplementary material related to this article can be found in the online version.

References

Ahmed, M.Z., Ahmed, O., Aibao, Z., Hanbin, S., Siyu, L., Ahmad, A., 2020.
Epidemic of COVID-19 in China and associated Psychological Problems. Asian
J. Psychiatr. 51. https://doi.org/10.1016/j.ajp.2020.102092

Du, J., Li, Z., Jia, G., Zhang, Q., Chen, W., 2019. Relationship between mental health
and awareness of the knowledge on mental health in left-behind middle school
students. Med. (United States) 98.
https://doi.org/10.1097/MD.0000000000014476

Eyuboglu, M., Eyuboglu, D., Pala, S.C., Oktar, D., Demirtas, Z., Arslantas, D., Unsal,
A., 2021. Traditional school bullying and cyberbullying: Prevalence, the effect
on mental health problems and self-harm behavior. Psychiatry Res. 297.
https://doi.org/10.1016/j.psychres.2021.113730

Feng, B., Zhang, Y., Zhang, L., Xie, X., Geng, W., 2020. Change in the level of
depression among Chinese college students from 2000 to 2017: A cross-temporal
meta-analysis. Soc. Behav. Pers. 48. https://doi.org/10.2224/sbp.8832

Garcia-Carrion, R., Villarejo, B.C., Villardón-Gallego, L., 2019. Children and
adolescents mental health: A systematic review of interaction-based
interventions in schools and communities. Front. Psychol. 10.
https://doi.org/10.3389/fpsyg.2019.00918

Holik, D., Kribl, M., Milostić-Srb, A., Nujić, D., 2021. Epidemiology of youth mental
health problems from the area of osijek using strengths and difficulties
questionnaire (Sdq). Arch. psychiatry Res. 57, 61–68.
https://doi.org/10.20471/may.2021.57.01.06

Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R., Walters, E.E.,
2005. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in
the national comorbidity survey replication. Arch. Gen. Psychiatry.
https://doi.org/10.1001/archpsyc.62.6.593

Kim, S., Colwell, S.R., Kata, A., Boyle, M.H., Georgiades, K., 2018. Cyberbullying
Victimization and Adolescent Mental Health: Evidence of Differential Effects by
Sex and Mental Health Problem Type. J. Youth Adolesc. 47, 661–672.

214 https://doi.org/10.1007/s10964-017-0678-4

215 Luo, Y., Cui, Z., Zou, P., Wang, K., Lin, Z., He, J., Wang, J., 2020. Mental health

216  problems and associated factors in chinese high school students in henan

217  province: a cross-sectional study. Int. J. Environ. Res. Public Health 17, 1–13.

218  https://doi.org/10.3390/ijerph17165944

219 Mo, L., 2017. Trends in the divorce rate and its regional disparity in China. J. Comp.

220  Fam. Stud. 48, 383–394. https://doi.org/10.3138/jcfs.48.4.383

221 Myhr, A., Anthun, K.S., Lillefjell, M., Sund, E.R., 2020. Trends in Socioeconomic

222  Inequalities in Norwegian Adolescents' Mental Health From 2014 to 2018: A

223  Repeated Cross-Sectional Study. Front. Psychol. 11.

224  https://doi.org/10.3389/fpsyg.2020.01472

225 Pine, D.S., Cohen, P., Gurley, D., Brook, J., Ma, Y., 1998. The risk for early-

226  adulthood anxiety and depressive disorders in adolescents with anxiety and

227  depressive disorders. Arch. Gen. Psychiatry 55, 56–64.

228  https://doi.org/10.1001/archpsyc.55.1.56

229 Rajkumar, R.P., 2020. COVID-19 and mental health: A review of the existing

230  literature. Asian J. Psychiatr. 52. https://doi.org/10.1016/j.ajp.2020.102066

231 Richter, D., Wall, A., Bruen, A., Whittington, R., 2019. Is the global prevalence rate

232  of adult mental illness increasing? Systematic review and meta-analysis. Acta

233  Psychiatr. Scand. 140, 393–407. https://doi.org/10.1111/acps.13083

234 Roy, D., Tripathy, S., Kar, S.K., Sharma, N., Verma, S.K., Kaushal, V., 2020. Study

235  of knowledge, attitude, anxiety & perceived mental healthcare need in Indian

236  population during COVID-19 pandemic. Asian J. Psychiatr. 51.

237  https://doi.org/10.1016/j.ajp.2020.102083

238 Su, Q., Liu, G., 2020. Depression in Chinese adolescents from 1989 to 2018: An

239  increasing trend and its relationship with social environments. Curr. Psychol.

240  https://doi.org/10.1007/s12144-020-01181-6

241 Thorisdottir, I.E., Asgeirsdottir, B.B., Kristjansson, A.L., Valdimarsdottir, H.B.,

242  Jonsdottir Tolgyes, E.M., Sigfusson, J., Allegrante, J.P., Sigfusdottir, I.D.,

243  Halldorsdottir, T., 2021. Depressive symptoms, mental wellbeing, and substance

use among adolescents before and during the COVID-19 pandemic in Iceland: a

longitudinal, population-based study. The Lancet Psychiatry 8, 663–672.

https://doi.org/10.1016/S2215-0366(21)00156-5

Wang, J., Li, Y., He, E., 1997. Development and Standardization of Mental Health

scale for Middle School students in China. Psychosoc. Sci 4, 15–20.

Wang, Q., Tapia Granados, J.A., 2019. Economic growth and mental health in 21st

century China. Soc. Sci. Med. 220, 387–395.

https://doi.org/10.1016/j.socscimed.2018.11.031

Wu, Z., Liu, Z., Zou, Z., Wang, F., Zhu, M., Zhang, W., Tao, H., Ross, B., Long, Y.,

2021a. Changes of psychotic-like experiences and their association with

anxiety/depression among young adolescents before COVID-19 and after the

lockdown in China. Schizophr. Res. 237, 40–46.

https://doi.org/10.1016/j.schres.2021.08.020

Wu, Z., Zou, Z., Wang, F., Xiang, Z., Zhu, M., Long, Y., Tao, H., Palaniyappan, L.,

Liu, Z., 2021b. Family functioning as a moderator in the relation between

perceived stress and psychotic-like experiences among adolescents during

COVID-19. Compr. Psychiatry 111.

https://doi.org/10.1016/j.comppsych.2021.152274

Xin, S., Wang, Y., Sheng, L., 2020. Impact of social changes and birth cohort on

anxiety in adolescents in mainland China (1992–2017): A cross-temporal meta-

analysis. Child. Youth Serv. Rev. 116.

https://doi.org/10.1016/j.childyouth.2020.105159

Xiong, R., Xia, Y., Li, S.D., 2021. Perceived Discrimination and Aggression Among

Chinese Migrant Adolescents: A Moderated Mediation Model. Front. Psychol.

12. https://doi.org/10.3389/fpsyg.2021.651270

Yang, J.Z., Kang, C.Y., Li, J.H., Li, P.K., Zhao, X.D., 2017. A three-decade repeated

cross-sectional survey on mental health of the Chinese Jino minority. Aust. N. Z.

J. Psychiatry 51, 1134–1141. https://doi.org/10.1177/0004867416682599

Zeng, W., Chen, R., Wang, X., Zhang, Q., Deng, W., 2019. Prevalence of mental

health problems among medical students in China: A meta-analysis. Med.

(United States) 98. https://doi.org/10.1097/MD.0000000000015337

Zhang, S., Zhong, R., Zhang, J., 2017. School starting age and academic achievement:

Evidence from China's junior high schools. China Econ. Rev. 44, 343–354.

https://doi.org/10.1016/j.chieco.2017.03.004

Zhang, W., Zhu, Y., Sun, M., Guo, R., Wu, G., Wang, Z., Xue, Z., Shi, J., Ouyang,

X., Pu, W., Liu, Z., Chiu, H.F.K., Rosenheck, R., 2019. Longitudinal trajectories

of psychotic-like experiences and their relationship to emergent mental disorders

among adolescents: A 3-year cohort study. J. Clin. Psychiatry 80.

https://doi.org/10.4088/JCP.18m12437

Zhao, S., Liao, C., 2016. Mental Health and Life Events of Overseas Children Left-

Behind in China. Psychology 07, 1934–1946.

https://doi.org/10.4236/psych.2016.714176

 Table 1 Sample characteristics of the participants surveyed in each wave.

	Wave 1 (September 2016, $n = 978$)	Wave 2 (September 2019, $n = 949$)	Wave 3 (September 2020, $n = 910$)	Statistics
Sex (male/female)	544/434	503/446	481/429	$\chi^2 = 1.873$, $p = 0.392$
Age (years)	Unavailable	12.027 ± 0.432	Unavailable	/
MMHI-60 results (positive/negative/positive rate)				
Obsessive–compulsive tendencies	484/494/49.5%	596/353/62.8%	605/305/66.5%	$z = 7.566$, $p < 0.001$
Paranoid ideation	274/704/28.0%	272/677/28.7%	257/653/28.2%	$z = 0.115$, $p = 0.909$
Hostility	257/721/26.3%	265/684/27.9%	247/663/27.1%	$z = 0.437$, $p = 0.662$
Interpersonal sensitivity	305/673/31.2%	326/623/34.4%	332/578/36.5%	$z = 2.435$, $p = 0.015$
Depression	254/724/26.0%	294/655/31.0%	322/588/35.4%	$z = 4.437$, $p < 0.001$
Anxiety	354/624/36.2%	367/582/38.7%	396/514/43.5%	$z = 3.241$, $p = 0.001$
Academic stress	348/630/35.6%	362/587/38.2%	392/518/43.1%	$z = 3.326$, $p < 0.001$
Maladaptation	209/769/21.4%	235/714/24.8%	224/686/24.6%	$z = 1.683$, $p = 0.092$
Emotional disturbance	338/640/34.6%	350/599/36.9%	370/540/40.7%	$z = 2.731$, $p = 0.006$

Psychological imbalance	167/811/17.1%	168/781/17.7%	184/726/20.2%	$z = 1.753$, $p = 0.080$

Psychological imbalance 167/811/17.1% 168/781/17.7% 184/726/20.2% $z = 1.753$, $p = 0.080$

Figure Legends

Figure 1 Trends in positive rates of each mental health problem in surveyed participants during the period of 2016-2020. The "*" indicates a significant increasing trend with $p < 0.05$.

Figure 2 Comparisons on positive rates of each mental health problem between the female and male participants. The "*" indicates a significant sex difference with $p < 0.05$.

Figure 3 Trends in positive rates of each mental health problem during the period of 2016-2020 in surveyed females and males, respectively. The "*" indicates a significant increasing trend with $p < 0.05$.