Trust in the research community predicts intent to comply with COVID-19 prevention

measures: An analysis of a large-scale international survey dataset

Hyemin Han [1*]

[1*] Educational Psychology Program, University of Alabama

Tuscaloosa, Alabama, 35487, United States of America

hyemin.han@ua.edu

ORCID: https://orcid.org/0000-0001-7181-2565

Word Count (main text): 3,998 words

1   **Abstract**

In the present study, I explored the relationship between people's trust in different agents

related to prevention of spread of COVID-19 and their compliance with pharmaceutical and non-

pharmaceutical preventive measures. The COVIDiSTRESSII Global Survey dataset, which was

collected from international samples, was analysed to examine the aforementioned relationship

across different countries. For data-driven exploration, network analysis and Bayesian

generalized linear model (GLM) analysis were performed. The result from network analysis

demonstrated that trust in the scientific research community was most central in the network of

trust and compliance. In addition, the outcome from Bayesian GLM analysis indicated that the

same factor, trust in the scientific research community, was most fundamental in predicting

participants' intent to comply with both pharmaceutical and non-pharmaceutical preventive

measures. I briefly discussed the implications of the findings, the importance of trust in the

scientific research community in explaining people's compliance with measure to prevent spread

of COVID-19.

Introduction

Since the onset of the current COVID-19 pandemic, different agents, including but not

limited to, governments, organizations, and scientific communities, have been developing,

implementing, and enforcing measures to prevent spread of COVID-19. Such measures embrace

both pharmaceutical and non-pharmaceutical means. Since late 2020, there have been several

COVID-19 vaccines approved for public use [1]. Even before approval of the first COVID-19

vaccine, diverse non-pharmaceutical measures, such as mask use, social distancing, mandatory

self-isolation, stay-at-home order, have been implemented and enforced [2]. Although the

pandemic has not concluded, data collected so far suggests that implementation of such

preventive measures have significantly contributed to prevention and mitigation of severe

COVID-19 outbreaks [3,4].

Given the importance of preventive measures in prevention of spread of COVID-19,

whether public is compliant with such measures would be critical in the current pandemic

situation [5]. Even if diverse preventive measures that have been found to be effective are

planned and implemented by agents, without people's compliance with the measures, successful

control of the pandemic could not be achieved [6]. For instance, rejection of and incompliance

with the r recommended and required preventive measures associated with political debates

resulted in the recent drastic increase in COVID-19 cases and deaths caused by the Delta variant

in multiple countries across the globe [5,7]. Hence, it would be important to understand which

factors are involved in people's compliance as well as incompliance with preventive measures.

Previous research has suggested that trust in agents addressing pandemic-related matters

is one of the most fundamental factors predicting compliance with preventive measures [8]. For

instance, several researchers have examined and reported significant association between trust in

governmental agents and organizations in the domain of health care (e.g., World Health

Organization), and vaccination intent and compliance with non-pharmaceutical preventive

measures [9–11]. Furthermore, trust in science and scientific research communities, which play

fundamental roles in developing preventive measures and proposing guidelines based on

evidence, has also been considered as a central factor in predicting compliance [12,13]. This

would be particularly important within the context of the current pandemic, because spread of

misinformation and conspiracy theories, which are closely associated with distrust in science and

particularly problematic in recent days, drives people's tendency to disobey mandatory

preventive measures and vaccination requirement [14].

Although the aforementioned previous studies have examined the importance of trust in

compliance with preventive measures, several limitations would warrant further investigations.

First, the majority of the previous studies was conducted with datasets collected from single or a

limited number of countries. Given the current COVID-19 pandemic is a global issue [15], it

would be necessary to collect data across diverse countries in examining the mechanism of

compliance tendency. Such relatively small-scale research based on data from a small number of

countries might not be sufficient to draw conclusions that can be well generalizable across the

globe.

Second, in terms of methodology, the previous studies employed conventional analysis

methods, which are based on frequentist perspective; such conventional methods are suitable to

test one specific null hypothesis and/or model, but not ideal for model exploration [16]. For

instance, if we are primarily interested which trust factor is central in prediction of compliance

tendency, the previous studies employing conventional methods might not be able to address our

interest in a complete manner. In fact, exploration of the best prediction model among multiple

competing candidate models requires analysis methods specialized in data-driven analysis, in

lieu of conventional hypothesis-driven analysis [17]. Thus, the findings from the previous studies

that primarily focused on trust in specific agents and used conventional methods would not show

us the full picture of how trust in different agents is associated with compliance with different

types of preventive measures.

66   **Current Study**

67   In the current study, how people's trust in different agents predicts their intent to

compliance with preventive measures and get vaccinated within the context of the COVID-19

pandemic will be examined in a data-driven manner with a large-scale international survey

dataset to address the aforementioned limitations in the prior research. Unlike the previous

studies employing conventional analysis methods, which are suitable for one null-hypothesis

testing, I intend to explore which trust factor is particularly important in predicting compliance

by exploring the large-scale dataset, the COVIDiSTRESSII Global Survey dataset [18], with

data-driven analysis methods.

75   To conduct the data-driven exploration, I plan to implement two novel analysis methods.

First, network analysis will be performed to explore how trust in different agents and compliance

with different types of preventive measures are associated with each other. In this exploration, I

intend to examine which factor is positioned in the most central and influential position in the

network [19]. Second, I will explore the best model predicting compliance with different types of

preventive measures with Bayesian model exploration [20]. Through this process, all possible

candidate regression models in terms of all possible combinations of trust in different agents will

be tested, and the most probable model given data will be identified. Finally, based on results

from the aforementioned processes employing data-driven methods, I will examine which trust

factors are relatively more important in predicting their compliance with preventive measures

across different countries.

Methods

Dataset

In the present study, I analysed the COVIDiSTRESSII Global Survey dataset, which was

collected by the COVIDiSTRESS Global Survey Consortium and is available to public via the

Open Science Framework (https://osf.io/36tsd). Originally, the data was collected from 20,601

participants from 62 countries. However, as I employed mixed-effects model analysis to include

the between-country effect in analysis, to prevent potential convergence issue [8,21], only data

collected from countries where 100 or more participants completed the survey was used in the

present study. As a result, I analyzed a subset of the data collected from 14,349 participants from

35 countries. Demographics of the participants included in the subset is presented in Table 1.

Further details about data collection and cleaning procedures are explained in the project

page (https://osf.io/36tsd). All procedures regarding data collection and informed consent were

reviewed and approved by Institutional Review Board at University of Salford where the project

manager of the consortium was affiliated during the data collection period. The author asserts

that all procedures contributing to this work comply with the ethical standards of the relevant

institutional committee on human experimentation and with the Helsinki Declaration of 1975, as

revised in 2008.

Measures

The employed items were developed by the COVIDiSTRESS Global Survey Consortium

members. They were translated and back translated by the consortium members from different

countries. Further details about the measures are described in the survey project page

(https://osf.io/36tsd).

Trust Items

Trust in seven different agents related to development, implementation, and/or

enforcement of preventive measures against COVID-19 was surveyed. Participants were asked to

what extent they trust each agent based on their general impression. The seven agents were: a

parliament or government (Trust 1); police (Trust 2); civic service (Trust 3); health system (Trust

4); WHO (Trust 5); government's effort to handle Coronavirus (Trust 6); and scientific research

community (Trust 7). Participants' responses were anchored to an eleven-point Likert Scale (0:

no trust—10: complete trust).

Compliance Intent Items

Participants' intent to comply with eight different types of preventive measures was also

surveyed. First, in the domain of pharmaceutical measures, one item, "How willing are you to

get the vaccine if one becomes available to you?" was presented to assess their intent to get

vaccinated (Compliance 1). Participants' responses to this item were anchored to five-point

Likert scale (1: not willing at all—5: very willing).

Second, in the case of compliance with non-pharmaceutical preventive measures,

compliance with seven different types of measures was surveyed. Participants were asked to

what extent they were compliant with each measure during the last month. The seven surveyed

measures were: washing hands regularly (Compliance 2); wearing a face covering in public

when indoors (Compliance 3); wearing a face covering in public when outdoors (Compliance 4);

staying at least the recommended distance (Compliance 5); staying at home unless going out for

essential reasons (Compliance 6); self-isolating if you suspected that you had been in contact

with the virus (Compliance 7); staying away from crowded places generally (Compliance 8).

Answers to the items were anchored to a seven-point Likert scale (1: strongly disagree—7:

strongly agree).

Demographics

Following previous studies examining behavioural and psychological responses to

COVID-19 using international survey datasets [8,15,20], several demographic variables were

also employed as control variables in the present study. I used participants' age, gender, and

education level in analysis. Participants' gender was surveyed by presenting three options:

female; male; other or would rather not say. The survey presented seven options to ask

participants' education level: PhD or doctorate; university degree (e.g., MA, MSc, BA, BSc);

some university or equivalent (still ongoing, or completed a module or more, but did not

graduate); up to 12 years of school; up to 9 years of school; up to 6 years of school; none.

Analysis Plan

Network Analysis

To examine the overall association between responses to the seven trust and eight

compliance items, I conducted network analysis with *bootnet R* package. The main purpose of

network analysis is to demonstrate associations between nodes, trust and compliance in the case

of the present study. A connection between two specific node is defined as an edge, which has a

weight representing the strength of the association [22]. An edge weight is quantified in term of

partial correlation between two nodes by *bootnet*. As an illustrative example, in the case of the

edge between Trust 1 and Compliance 1, the edge weight can be understood in terms of

correlation between Trust 1 and Compliance 1 after controlling for correlation with all other

items in the same network (i.e., Trust 2 … Compliance 8). In a network plot, which visualizes

the result of network analysis, an edge between two nodes is presented in the format of a line

with a specific thickness, which represents its edge weight, the strength of the association.

154   While exploring a partial correlation network, *bootnet* employs one technique, graphical

LASSO (GLASSO), to identify a regularized partial correlation network through penalizing

spurious edge weights [19]. Implementation of GLASSO is required to minimize false positives

that may exist in a network of interest. For instance, we can imagine that there is no true non-

zero partial correlation between two specific nodes. In the reality, possibly due to noise and/or

measurement error, even after controlling for association with other nodes, the edge weight

between the two nodes could not exactly become zero, although that is a false positive [19]. Such

spurious edge weights can be excluded by GLASSO. Moreover, use of such a penalization

method can contribute to prevention of model overfitting [17,23]. Hence, in the present study

bootnet identified the best network model with the smallest extended Bayesian Information

Criterion (EBIC) value to penalize unnecessarily complex and spurious network edge structures.

165   Once a partial correlation network model was identified with GLASSO, I performed

centrality analysis to examine which node located at the most central and influential position in

the network. For this purpose, three indicators resulting from centrality analysis, i.e., strength,

closeness, and betweenness, were examined for each node [24]. Strength is calculated by

summing the absolute values of association strengths, edge weights, of a specific node.

Closeness is defined in terms of the inverse of summed distances from one specific node to the

other nodes in the same network. Finally, betweenness is estimated in terms of how many times

one specific node is in the shortest path between two other nodes in the whole network. In the

present study, I examined which node reported the highest strength, closeness, and betweenness

values.

Bayesian Model Exploration

To examine the best regression model predicting each compliance variable with trust

variables, I conducted Bayesian model selection with the Bayesian generalized linear model

(GLM) implemented in *BayesFactor* R package. Unlike conventional regression analysis based

on frequentist perspective, Bayesian regression analysis enables us to examine to what extent

evidence supports a regression model of interest [25]. In the case of conventional regression

analysis, only one model can be tested each time, and the resultant *p*-values can only inform us

whether a null hypothesis (e.g., whether a null model is the case) shall be rejected [26]. Thus, if

our interest is exploration of the best model among all possible candidate models, conventional

regression analysis could not be an ideal solution.

Bayesian analysis can provide us with more direct information about whether a specific

hypothesis of interest is likely to be accepted given evidence [26]; in the same vein, we can also

learn about to what extent a specific model is more likely to be the case compared with other

candidate models given evidence as well [27]. Once Bayesian GLM is performed with

BayesFactor, we can examine the Bayes Factor (BF_{M0}) of each model quantifying to what extent

the model of interest, Model M, is more strongly supported by data compared with a null model

(Model 0) [25]. In the present study, $2\log(BF_{Mo})$ was used for result interpretation. Statistical

guidelines suggest that $2\log(BF_{M0}) \geq 3$ indicates presence of positive evidence supporting Model

M against Model 0, $2\log(BF_{M0}) \geq 6$ presence of strong evidence, and $2\log(BF_{M0}) \geq 10$ presence of

very strong evidence. When $2\log(BF_{M0}) < 3$, I concluded that evidence is merely trivial or

anecdotal [26].

Given the methodological and epistemological benefits of Bayesian analysis, I conducted

Bayesian GLM analysis for each compliance dependent variable to identify which trust

predictors shall be included in the best regression model [20]. For each dependent variable, I

used seven trust variables as candidate predictors, the country as a random effect, and

demographic variables as control variables. All trust and compliance variables were standardized

at the country level for better convergence, and more straightforward interpretation and

comparison of estimated coefficients. Through the process, all possible 128 candidate models,

which were created in terms of all possible combinations of seven trust predictors ($2^{(7+1)}$), were

estimated and their BF_{M0} were calculated [28]. I identified the best model with the highest BF_{M0}

value. Furthermore, I also compared the identified best model and the full model including all

seven trust predictors by calculating BF_{MF}, a BF value indicating to what extent evidence more

strongly supported the best model against the full model. In general, a full model including all

candidate predictors is tested and evaluated by resultant p-values in conventional regression

analysis [16], so I compared the full model with the best model suggested from Bayesian GLM

analysis. For both BF_{M0} and BF_{MF}, I calculated 2log(BF) values for interpretation.

Furthermore, I performed mixed-effects analysis with the indicated best model with

lmerTest and *brms* R packages. This additional analysis was conducted to examine the effect size

of each trust variable included in the best models. Although all predictors included in the best

models might be statistically significant in terms of p-values, such a significance is perhaps due

to a large sample size even if an actual effect is nearly zero or trivial in a practical manner [26].

Effect sizes were calculated in terms of Cohen's D values with *EMAtools* R package after

performing multilevel modelling with *lmerTest*. In this process, for each dependent compliance

variable, I employed trust variables that were identified to be included in the best models as

predictors, the country as a random effect, and demographic variables as control variables. Then,

the resultant D values were interpreted qualitatively as well as quantitatively. For qualitative

interpretation, following [29]'s guidelines, I assumed that a Cohen's D value within the range of

-.10 and +.10 as an indicator of a practically non or trivial effect.

In addition to the qualitative interpretation of effect sizes, I also conducted Bayesian

multilevel modelling with *brms* for quantitative interpretation. The same mixed-effects model

analysed with *lmerTest* was tested with *brms* for each compliance variable. In this process, I

employed the default Cauchy prior, Cauchy (0, 1), suggested by [30] following the previous

studies [8,21]. After conducting Bayesian multilevel modelling for each dependent variable, the

result was analysed with *bayestestR* package for Bayesian quantitative interpretation of effect

sizes. I estimated to what extent the 95% highest density interval (HDI) of the posterior

distribution of each trust predictor was within the region of practical equivalence (ROPE)

[31,32]. A 95% HDI means an interval that "any parameter value inside the HDI has higher

probability density than any value outside the HDI, and the total probability of values in the 95%

HDI is 95% [31] (p. 271)". If 100% of the HDI falls inside the defined ROPE, then the most

credible (95%) values of the effect size are completely within the regions of trivial effect, so

accepting a null hypothesis (e.g., the effect does not significantly differ from zero) becomes

practically reasonable. In this context, a ROPE means a range of a parameter of interest that shall

be considered practically near-zero or trivial [32]. Since the guideline that I employed stated that

$-.10 \leq D \leq .10$ indicates zero or trivial effect [29], I set [-.10 .10] as the ROPE for this

quantitative interpretation. By following these steps, I examined to what extent the estimated

posterior value of the effect size of each trust predictor in each best model was within the defined

ROPE.

242   **Results**

243   **Network Analysis**

244   Figure 1 is a network plot demonstrating connectivity between seven trust and eight

compliance items in the best network model with the lowest EBIC value identified by *bootnet*.

Figure 2 shows the result of centrality analysis. In this plot, three centrality indicators, strength,

closeness, and betweenness of each node, were presented. All three indicators unequivocally

suggest that Trust 7, trust in the scientific research community, is most central in the analysed

network.

Bayesian Model Exploration

251   Table 2 demonstrates the results from Bayesian model exploration via Bayesian GLM

252   analysis. The outcome of Bayesian GLM analysis with each compliance dependent variable was

253   presented in each row. Only the coefficients and effect sizes of trust predictors that were

254   included in each best model were presented in Table 2. In addition, the same table reports the

255   proportion of the 95% HDI of each survived trust predictor within the defined ROPE, [-.10 .10].

256   In terms of BF_{M0} and BF_{MF}, all best models identified by Bayesian GLM analysis, except

257   for the best model predicting vaccination intent (Compliance 1), were supported by very strong

258   evidence compared with both the null and full models. In the case of Compliance 1, the full

259   model including all seven trust predictors was identified as the best model.

260   **Discussion**

261   In the present study, first, I conducted network analysis to understand association

between participants' trust in seven different agents addressing the COVID-19 pandemic and

their intent to comply with eight different types of preventive measures. Second, Bayesian GLM

analysis was performed to explore the best model predicting each compliance intent variable

with trust predictors. From network analysis, robust connectivity between trust and compliance

variables even after penalizing unnecessary edge features via GLASSO was demonstrated in the

visualized network plot. Among all nodes, including both all seven trust and eight compliance

variables, Trust 7, trust in the scientific research community, was found to be most central in the

network according to all three centrality indicators, strength, closeness, and betweenness. The

similar trend was also reported from Bayesian GLM analysis to identify the best model

predicting each compliance variable. Although several other trust variables were included in the

identified best models, only Trust 7 was included in all eight best prediction models.

Furthermore, in terms of the Cohen's *D* and proportion of the 95% HDI within the [-.10 .10]

ROPE [29,31,32], Trust 7 reported the greatest effect size, which was most likely to be out of the

region of near-zero or trivial effect, compared with all six other trust predictors.

276   The findings suggest that in predicting people's intent to comply with both

pharmaceutical (e.g., vaccination) and non-pharmaceutical measures to prevent spread of

COVID-19 (e.g., hand washing, mask use, social distancing, self-isolation), trust in scientific

research and the community of scientists play the most fundamental role compared with trust in

other agents, e.g., government, healthcare system, health organization [12,13]. Given that such

measures were primarily tested and suggested by scientific studies with empirical evidence, even

if their implementation and enforcement are tasks to be done by other agents, trust in science is

expected to make the greatest, fundamental influence on people's compliance [33]. Hence, if

people do not have robust trust in scientific research regarding COVID-19, then they are unlikely

to abide by preventive measures implemented by governments and health-related organizations

[13].

Given the prevalence of distrust in science, which is being closely linked to widespread

of misinformation and conspiracy theories [34], within the current situation, the potential reason

of why such distrust contributing to incompliance with preventive measures would be worth

consideration. [35] argued that epistemological doubts and ontological insecurity about scientific

knowledge shared within the modern society has promoted and reinforced conspiracy theories

and then challenged trust in science among public. A trend related to widespread conspiracy

theories and distrust in science is also influential in the current pandemic situation [14].

Conspiracy theories and distrust in science regarding COVID-19 have been promoted by

political motive and ideology, authoritarianism, and extremism in particular, and resulted in

distrust in scientific evidence supporting preventive measures, and finally, rejection of and

incompliance with recommended preventive measures [34,36]. Furthermore, lack of rational

deliberation and reflection upon information and messages is also reported to relate to

acceptance of conspiracy theories and distrust in science, which eventually cause incompliance

[37].

Therefore, if researchers and policy makers are interested in promoting people's

compliance with preventive measures, which are suggested and supported by scientific research,

they need to consider how to promote people's trust in scientific research and scientists.

Although consideration of concrete solutions for promoting trust in science in public is out of the

scope of the current study, let me list a couple of possible starting points. Educators may start

with improving science education to educate science-informed citizens who are capable of

rationally evaluating and accepting knowledge and information around them in a scientific

manner [38]. Moreover, it would be possible to examine how to improve science communication

with public, which improve people's understanding and perception on science [39]. Because

many of the current social issues related to incompliance with COVID-19 preventive measures

have been emerged from and reinforced by misinformation shared through diverse forms of

media, improvement of science communication would be required to address the issues [40].

We may consider several strengths of the present study and how it could make significant

contributions to literature. First, a large-scale international survey dataset, the COVIDiSTRESSII

Global Survey dataset, was analysed instead of a relatively small-size dataset collected from a

limited number of countries. Because the COVID-19 pandemic is a global issue, findings from

the current study will be able to provide researchers and policy makers across the globe with

useful insights about how to promote people's compliance with preventive measures based on

generalizable evidence from a cross-national investigation. Second, I explored the overall

association between compliance and trust in different agents instead of testing specific

hypotheses. With novel quantitative methods, network analysis and Bayesian GLM analysis, I

was able to demonstrate that trust in scientific research is most influential and fundamental in

predicting compliance.

However, several limitations in the present study may warrant further investigations.

First, while collecting data regarding compliance, the project consortium employed the self-

report method for the feasibility of the global survey project. Hence, whether the reported

compliance intent predicts compliance behaviour in the reality could be questionable. Second,

only one item per trust in each specific agent or compliance with each specific preventive

measure was employed in the survey. Because the consortium was not able to use multiple items

per construct due to the feasibility issue, the psychometrical aspects of the trust and compliance

items could not be tested in a complete manner. Thus, future studies shall employ more direct

measures for compliance and conduct psychometrics tests by employing multiple items per

construct to address the limitations in the current study.

334   **Data Availability Statement**

335   The data and codes that support the findings of this study are openly available in the

336   Open Science Framework project page (https://doi.org/10.17605/OSF.IO/Y4KGH).

337   **Financial support**

338   This research received no specific grant from any funding agency, commercial or not-for-

339   profit sectors.

340   **Conflicts of Interest**

341   Conflicts of Interest: None.

342   **References**

1.   **Tregoning JS, *et al.*** Progress of the COVID-19 vaccine effort: viruses, vaccines and

variants versus efficacy, effectiveness and escape. *Nature Reviews Immunology* 2021;

21Published online: October 9, 2021.doi:10.1038/s41577-021-00592-1.

2.   **Perra N**. Non-pharmaceutical interventions during the COVID-19 pandemic: A review.

Physics Reports 2021; **913**Published online: May

2021.doi:10.1016/j.physrep.2021.02.001.

3.   **Dagan N, *et al.*** BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination

Setting. *New England Journal of Medicine* 2021; **384**Published online: April 15,

2021.doi:10.1056/NEJMoa2101765.

4.   **Flaxman S, *et al.*** Estimating the effects of non-pharmaceutical interventions on COVID-

19 in Europe. *Nature* 2020; **584**Published online: August 13, 2020.doi:10.1038/s41586-

020-2405-7.

5.   **Gao J, Radford BJ**. Death by political party: The relationship between COVID-19 deaths

and political party affiliation in the United States. *World Medical & Health Policy* 2021;

13Published online: June 5, 2021.doi:10.1002/wmh3.435.

6.   **Borchering RK, *et al.*** Modeling of Future COVID-19 Cases, Hospitalizations, and

Deaths, by Vaccination Rates and Nonpharmaceutical Intervention Scenarios — United

States, April–September 2021. *MMWR. Morbidity and Mortality Weekly Report* 2021;

70Published online: May 14, 2021.doi:10.15585/mmwr.mm7019e3.

7.   **Freira L, *et al.*** The interplay between partisanship, forecasted COVID-19 deaths, and

support for preventive policies. *Humanities and Social Sciences Communications* 2021;

8Published online: December 3, 2021.doi:10.1057/s41599-021-00870-2.

8.   **Lieberoth A, *et al.*** Stress and worry in the 2020 coronavirus pandemic: Relationships to

trust and compliance with preventive measures across 48 countries. *Royal Society Open*

Science 2021; **8**: 200589.

9.   **Bargain O, Aminjonov** U. Trust and compliance to public health policies in times of

COVID-19. *Journal of Public Economics* 2020; **192**Published online: December

2020.doi:10.1016/j.jpubeco.2020.104316.

10.   **Lalot F, *et al.*** The dangers of distrustful complacency: Low concern and low political

trust combine to undermine compliance with governmental restrictions in the emerging

Covid-19 pandemic. *Group Processes & Intergroup Relations* 2020; Published online:

October 30, 2020.doi:10.1177/1368430220967986.

11.   **Chan HF, *et al.*** How confidence in health care systems affects mobility and compliance

during the COVID-19 pandemic. *PLOS ONE* 2020; **15**Published online: October 15,

2020.doi:10.1371/journal.pone.0240644.

12.   **Plohl N, Musil B**. Modeling compliance with COVID-19 prevention guidelines: the

379   critical role of trust in science. *Psychology, Health & Medicine* 2021; **26**Published online:

380   January 2, 2021.doi:10.1080/13548506.2020.1772988.

381   13.   **Bicchieri C, *et al.*** In science we (should) trust: Expectations and compliance across nine

382   countries during the COVID-19 pandemic. *PLOS ONE* 2021; **16**Published online: June 4,

383   2021.doi:10.1371/journal.pone.0252892.

384   14.   **Rutjens BT, van der Linden S, van der Lee R**. Science skepticism in times of COVID-

385   19. *Group Processes & Intergroup Relations* 2021; **24**Published online: February 4,

386   2021.doi:10.1177/1368430220981415.

387   15.   **Yamada Y, *et al.*** COVIDiSTRESS Global Survey dataset on psychological and

388   behavioural consequences of the COVID-19 outbreak. *Scientific Data* 2021; **8**: 3.

389   16.   **Han H, Dawson KJ**. Improved model exploration for the relationship between moral

390   foundations and moral judgment development using Bayesian Model Averaging. *Journal*

391   *of Moral Education* 2021; Published online: 2021.doi:10.1080/03057240.2020.1863774.

392   17.   **McNeish DM**. Using Lasso for Predictor Selection and to Assuage Overfitting: A Method

393   Long Overlooked in Behavioral Sciences. *Multivariate Behavioral Research* 2015; **50**:

394   471–484.

395   18.   **Lieberoth A, *et al.*** *COVIDiSTRESSII Global Survey*. 2021(https://osf.io/36tsd/).

396   Accessed September 26, 2021.

397   19.   **Epskamp S, Fried EI**. A tutorial on regularized partial correlation networks.

398   *Psychological Methods* 2018; **23**Published online: December

399   2018.doi:10.1037/met0000167.

20. **Han H**. Exploring the association between compliance with measures to prevent the

spread of COVID-19 and big five traits with Bayesian generalized linear model.

Personality and Individual Differences 2021; **176**: 110787.

21. **Rachev NR, *et al.*** Replicating the Disease framing problem during the 2020 COVID-19

pandemic: A study of stress, worry, trust, and choice under risk. *PLOS ONE* 2021;

16Published online: September 10, 2021.doi:10.1371/journal.pone.0257151.

22. **Dalege J, *et al.*** Network Analysis on Attitudes. *Social Psychological and Personality*

Science 2017; **8**Published online: July 10, 2017.doi:10.1177/1948550617709827.

23. **Han H, Dawson KJ**. Applying elastic-net regression to identify the best models

predicting changes in civic purpose during the emerging adulthood. *Journal of*

Adolescence 2021; **93**Published online: December

2021.doi:10.1016/j.adolescence.2021.09.011.

24. **Opsahl T, Agneessens F, Skvoretz J**. Node centrality in weighted networks:

Generalizing degree and shortest paths. *Social Networks* 2010; **32**Published online: July

2010.doi:10.1016/j.socnet.2010.03.006.

25. **Wagenmakers E-J, *et al.*** Bayesian inference for psychology. Part II: Example

applications with JASP. *Psychonomic Bulletin & Review* 2018; **25**: 58–76.

26. **Han H, Park J, Thoma SJ**. Why do we need to employ Bayesian statistics and how can

we employ it in studies of moral education?: With practical guidelines to use JASP for

educators and researchers. *Journal of Moral Education* 2018; **47**: 519–537.

27. **Dawson KJ, Han H, Choi YR**. How are moral foundations associated with empathic

traits and moral identity? *Current Psychology* 2021; Published online: October 12,

2021.doi:10.1007/s12144-021-02372-5.

28.   **Gronau QF, *et al.** A Simple Method for Comparing Complex Models: Bayesian Model*

424   *Comparison for Hierarchical Multinomial Processing Tree Models using Warp-III Bridge*

425   *Sampling*. 2017.

426   29.   **Funder DC, Ozer DJ**. Evaluating Effect Size in Psychological Research: Sense and

427   Nonsense. *Advances in Methods and Practices in Psychological Science* 2019; **2**Published

428   online: June 8, 2019.doi:10.1177/2515245919847202.

429   30.   **Rouder JN, Morey RD**. Default Bayes Factors for Model Selection in Regression.

430   *Multivariate Behavioral Research* 2012; **47**: 877–903.

431   31.   **Kruschke JK**. Rejecting or Accepting Parameter Values in Bayesian Estimation.

432   *Advances in Methods and Practices in Psychological Science* 2018; **1**Published online:

433   June 8, 2018.doi:10.1177/2515245918771304.

434   32.   **Makowski D, Ben-Shachar M, Lüdecke D**. bayestestR: Describing Effects and their

435   Uncertainty, Existence and Significance within the Bayesian Framework. *Journal of Open*

436   *Source Software* 2019; **4**Published online: August 13, 2019.doi:10.21105/joss.01541.

437   33.   **Nicola M, *et al.*** Evidence based management guideline for the COVID-19 pandemic -

438   Review article. *International Journal of Surgery* 2020; **77**Published online: May

439   2020.doi:10.1016/j.ijsu.2020.04.001.

440   34.   **Milošević Đorđević J, *et al.*** Links between conspiracy beliefs, vaccine knowledge, and

441   trust: Anti-vaccine behavior of Serbian adults. *Social Science & Medicine* 2021;

442   **277**Published online: May 2021.doi:10.1016/j.socscimed.2021.113930.

443   35.   **Aupers S**. 'Trust no one': Modernization, paranoia and conspiracy culture. *European*

444   *Journal of Communication* 2012; **27**Published online: March 29,

445   2012.doi:10.1177/0267323111433566.

446   36.   **Tonković M, *et al.*** Who Believes in COVID-19 Conspiracy Theories in Croatia?

447   Prevalence and Predictors of Conspiracy Beliefs. *Frontiers in Psychology* 2021;

448   **12**Published online: June 18, 2021.doi:10.3389/fpsyg.2021.643568.

449   37.   **Sutton RM, Douglas KM**. Conspiracy theories and the conspiracy mindset: implications

450   for political ideology. *Current Opinion in Behavioral Sciences* 2020; **34**Published online:

451   August 2020.doi:10.1016/j.cobeha.2020.02.015.

452   38.   **Fensham PJ**. Scepticism and trust: two counterpoint essentials in science education for

453   complex socio-scientific issues. *Cultural Studies of Science Education* 2014; **9**Published

454   online: September 6, 2014.doi:10.1007/s11422-013-9560-1.

455   39.   **Bubela T, *et al.*** Science communication reconsidered. *Nature Biotechnology* 2009;

456   **27**Published online: June 2009.doi:10.1038/nbt0609-514.

457   40.   **Mheidly N, Fares J**. Leveraging media and health communication strategies to overcome

458   the COVID-19 infodemic. *Journal of Public Health Policy* 2020; **41**Published online:

459   December 21, 2020.doi:10.1057/s41271-020-00247-w.

461 **Tables**

462 **Table 1:**

463 **Demographics of the whole dataset and each country**

	N	Age		Gender			Education level						
		M	SD	Female	Male	Other/ Would rather not say	Doctorate	University	University or equivalent	≤ 12 years	≤ 9 years	≤ 6 years	None
All	14,349	36.56	14.48	67.52%	31.55%	.93%	6.03%	47.65%	26.71%	16.19%	2.33%	.49%	.60%
Bolivia	115	39.56	11.23	65.22%	34.78%	.00%	13.91%	72.17%	13.91%	.00%	.00%	.00%	.00%
Bosnia and Herzegovina	109	39.85	11.51	76.15%	23.85%	.00%	13.89%	65.74%	7.41%	12.04%	.00%	.93%	.00%
Brazil	448	38.56	13.22	72.32%	27.23%	.45%	15.18%	66.07%	15.40%	3.35%	.00%	.00%	.00%
Bulgaria	299	41.47	16.74	73.58%	25.75%	.67%	6.04%	50.67%	32.55%	10.40%	.00%	.00%	.34%
Colombia	548	40.04	12.63	67.88%	31.57%	.55%	5.67%	77.15%	11.52%	4.20%	.73%	.37%	.37%
Costa Rica	270	35.92	10.37	69.63%	29.26%	1.11%	1.11%	78.15%	14.81%	3.70%	1.85%	.37%	.00%
Czech Republic	365	34.14	11.31	70.68%	27.95%	1.37%	5.75%	46.03%	31.51%	16.44%	.00%	.27%	.00%
Denmark	127	42.30	10.59	61.42%	38.58%	.00%	11.81%	57.48%	23.62%	2.36%	1.57%	3.15%	.00%
Ecuador	291	31.78	10.80	66.32%	32.99%	.69%	2.75%	69.76%	22.68%	3.44%	.69%	.34%	.34%
Estonia	246	39.36	10.21	86.53%	13.47%	.00%	1.22%	56.33%	22.04%	18.37%	2.04%	.00%	.00%
Finland	963	46.07	14.38	78.40%	20.15%	1.45%	4.69%	54.95%	18.56%	16.68%	2.61%	1.88%	.63%

Country	N												
Germany	152	36.73	12.49	67.76%	29.61%	2.63%	8.55%	55.92%	20.39%	13.82%	1.32%	.00%	.00%
Guatemala	287	36.94	14.02	84.32%	15.68%	.00%	4.88%	65.85%	26.83%	1.74%	.35%	.35%	.00%
Honduras	429	25.86	8.13	66.90%	32.17%	.93%	1.40%	20.05%	62.00%	14.22%	.93%	.70%	.70%
Ireland	401	29.01	10.79	68.00%	31.00%	1.00%	4.74%	48.38%	45.89%	.50%	.50%	.00%	.00%
Italy	310	44.83	16.27	74.43%	25.24%	.32%	6.54%	45.75%	23.20%	22.22%	1.96%	.33%	.00%
Japan	2,133	45.49	11.08	41.87%	56.96%	1.17%	1.03%	31.77%	20.84%	36.46%	6.34%	.84%	2.72%
Kyrgyzstan	254	32.44	12.46	82.28%	14.96%	2.76%	2.77%	52.57%	14.23%	27.67%	2.77%	.00%	.00%
Lebanon	141	29.31	12.24	73.05%	26.24%	.71%	6.43%	43.57%	37.86%	9.29%	2.14%	.71%	.00%
Malaysia	225	27.22	8.83	69.33%	28.44%	2.22%	4.44%	53.78%	39.11%	2.67%	.00%	.00%	.00%
Norway	376	40.86	13.58	80.32%	19.68%	.00%	8.24%	61.17%	18.35%	9.57%	1.06%	.80%	.80%
Pakistan	157	24.33	6.54	97.45%	2.55%	.00%	8.92%	45.22%	38.22%	7.01%	.64%	.00%	.00%
Portugal	484	33.33	14.94	70.25%	28.51%	1.24%	23.55%	42.56%	21.90%	11.16%	.83%	.00%	.00%
Russian	2,260	26.05	10.49	70.93%	28.14%	.93%	.66%	28.98%	46.70%	19.58%	3.54%	.31%	.22%
Slovakia	313	34.49	13.64	88.82%	10.86%	.32%	8.09%	49.19%	26.21%	13.92%	.97%	.97%	.65%
Spain	575	40.44	13.83	64.52%	34.96%	.52%	20.70%	53.91%	21.39%	3.30%	.70%	.00%	.00%
Sweden	134	39.28	13.91	76.87%	18.66%	4.48%	12.88%	54.55%	22.73%	8.33%	1.52%	.00%	.00%
Switzerland	593	44.84	19.01	63.58%	36.09%	.34%	6.91%	45.36%	18.21%	24.28%	4.38%	.34%	.51%
Taiwan	221	34.94	9.81	61.99%	36.20%	1.81%	5.43%	89.14%	1.81%	3.62%	.00%	.00%	.00%
Turkey	200	23.79	8.26	68.50%	30.50%	1.00%	3.00%	34.50%	4.50%	57.50%	.00%	.00%	.50%
Uganda	135	24.37	4.17	32.59%	67.41%	.00%	.00%	23.70%	74.81%	1.48%	.00%	.00%	.00%

Ukraine	252	31.39	10.06	63.49%	35.71%	.79%	10.32%	77.38%	5.95%	5.16%	.79%	.40%	.00%
United Kingdom	134	36.58	12.88	76.12%	22.39%	1.49%	28.36%	52.99%	11.19%	6.72%	.75%	.00%	.00%
United States	114	34.74	11.49	64.04%	34.21%	1.75%	22.81%	47.37%	23.68%	3.51%	.88%	1.75%	.00%
Uruguay	288	41.97	12.94	88.15%	11.85%	.00%	5.90%	74.31%	13.19%	5.21%	1.04%	.00%	.35%

465 **Table 2:**

466 **Results from Bayesian GLM analysis**

	Trust 1 β (SE)	Trust 1 ES (ROPE %)	Trust 2 β (SE)	Trust 2 ES (ROPE %)	Trust 3 β (SE)	Trust 3 ES (ROPE %)	Trust 4 β (SE)	Trust 4 ES (ROPE %)	Trust 5 β (SE)	Trust 5 ES (ROPE %)	Trust 6 β (SE)	Trust 6 ES (ROPE %)	Trust 7 β (SE)	Trust 7 ES (ROPE %)	Null model (BF_{M0})	Full model (BF_{MF})
Comp. 1	.04 (.01)	.06 (100.00%)	-.04 (.01)	-.07 (100.00%)	-.04 (.01)	-.06 (100.00%)	.08 (.01)	.12 (99.63%)	.15 (.01)	.25 (.00%)	.05 (.01)	.08 (100.00%)	.27 (.01)	.46 (.00%)	1375.36	
Comp. 2	-.03 (.01)	-.04 (100.00%)	.04 (.01)	.06 (100.00%)			.07 (.01)	.10 (100.00%)					.14 (.01)	.25 (.00%)	256.86	5.91
Comp. 3							.03 (.01)	.05 (100.00%)	.07 (.01)	.11 (100.00%)	.04 (.01)	.08 (100.00%)	.17 (.01)	.28 (.00%)	480.29	5.57
Comp. 4	.06 (.01)	.11 (100.00%)							.07 (.01)	.12 (100.00%)			.11 (.01)	.19 (9.10%)	254.77	3.72
Comp. 5									.07 (.01)	.11 (100.00%)	.05 (.01)	.09 (100.00%)	.10 (.01)	.17 (30.44%)	225.27	8.82
Comp. 6									.06 (.01)	.10 (100.00%)	.03 (.01)	.05 (100.00%)	.09 (.01)	.16 (74.11%)	158.90	9.18
Comp. 7									.07 (.01)	.12 (100.00%)	.04 (.01)	.07 (100.00%)	.11 (.01)	.18 (24.57%)	177.70	7.67
Comp. 8									.06 (.01)	.09 (100.00%)	.03 (.01)	.05 (100.00%)	.13 (.01)	.22 (.00%)	227.54	10.53

467   Comp. 1—Comp. 8: Compliance 1—Compliance 8. β (SE): Standardized regression coefficient (standard error). ES (ROPE %): Effect

468   size in Cohen's *D* (% of 95% HDI within the defined region of practical equivalence).

469   **Figures**

470   **Figure 1:**

471   **Network plot**

473   Solid line: positive edge weight. Dashed line: negative edge weight.

474   **Figure 2:**

475   **Result from centrality analysis**

476