Effects of racial/ethnic disparities in healthcare utilization on antibiotic use, 2016/2018

Scott W. Olesen, PhD[1*], Sanjat Kanjilal, MD MPH[2], Stephen M. Kissler, PhD[1], Daphne S. Sun, BS[1], Yonatan H. Grad, MD PhD[1*]

1 Department of Immunology and Infectious Diseases, Harvard Chan School of Public Health, Boston, MA

2 Department of Population Medicine, Harvard Medical School & Harvard Pilgrim Healthcare Institute, Boston, MA

* Correspondence:

Harvard Chan School of Public Health

665 Huntington Ave

Building 1 Room 715

Boston, MA 02115

732.351.2311

olesen@hsph.harvard.edu

Word count: 1402

Keywords: antibiotics, race/ethnicity, equity, healthcare utilization

ABSTRACT

White Americans make more office-based and emergency department visits per capita than other races/ethnicities, but the proportion of visits during which antibiotics are administered or prescribed is similar across races/ethnicities. Racial/ethnic disparities in antibiotic use may be due more to disparities in healthcare access and utilization than to prescriber behavior.

INTRODUCTION

Antibiotics are an important therapy for treating infectious disease, but use of antibiotics carries risk of negative effects, like adverse events or antibiotic resistance. Avoiding antibiotic overuse through antibiotic stewardship helps avoid unnecessary and inappropriate antibiotic use. However, underuse of antibiotics, when a person has an infection for which antibiotics are medically recommended but the person does not receive the antibiotic, is also problematic [1]. Even for conditions like sore throat or otitis media, which typically resolve without complications, failure to use antibiotics can sometimes lead to substantial morbidity [2].

Disparities in antibiotic use rates, when one population uses more or less antibiotics than another group, are a likely indicator of antibiotic overuse or underuse [1,3]. Many studies have shown that prescribing practice varies by patient race/ethnicity, usually with whites more likely to receive antibiotics or broader-spectrum antibiotics [1,4–7].

However, studies of antibiotic prescribing per healthcare visit, while important, do not address all the factors that could lead to disparities in antibiotic use. First, rates of bacterial infections may vary between populations. Second, access to healthcare and healthcare seeking behavior may differ across populations. In other words, the probability that someone with an infection presents at a healthcare visit may vary by population. Third, different populations may be more or less likely to receive antibiotics when they present at a healthcare visit. Studies of prescribing

per visit only address the third step, but previous work suggests that geographical differences in

antibiotic use rates are due more to differences in healthcare access and utilization than to

prescribing practice [8] and that temporal trends in racial disparities in antibiotic use among

children are due more to changes in healthcare utilization than to changes in prescribing

practice [7].

Here, we use a nationally representative survey of healthcare visits to assess the relative

importance of healthcare utilization and prescribing practice on antibiotic use by race/ethnicity.

We hypothesized that any racial/ethnic differences in antibiotic use are due both to differences

in healthcare utilization, measured as per capita rates of ambulatory care visits, as well as to

differences in prescribing practice, measured as the probability that an antibiotic will be used

given that certain diagnoses are present.

METHODS

We analyzed data from the National Ambulatory Medical Care Survey (NAMCS) and National

Hospital Ambulatory Medical Care Survey (NHAMCS), two nationally representative samples

used to characterize antibiotic prescribing practice [9,10]. Each survey is a sample of office-

based and emergency department visits with associated patient demographics, diagnosis

codes, and prescriptions as provided by physicians or hospital staff. Missing race/ethnicity data

are imputed using a model developed by NAMCS/NHAMCS staff [10]. We used the two most

recent years with data from both surveys, 2016 and 2018. NHAMCS from 2017 data were not

available [11].

Each healthcare visit was classified as antibiotic-appropriate, potentially antibiotic-appropriate,

or antibiotic-inappropriate by a two-step process [12]. First, using previously established

categorizations of the ICD-10 diagnosis codes, each diagnosis code associated with each visit

was classified as always antibiotic-appropriate, sometimes antibiotic-appropriate, or never

antibiotic-appropriate. Second, each visit was classified based on its diagnoses. A visit with at

least one always-appropriate diagnosis was classified as antibiotic-appropriate. Remaining

unclassified visits with at least one sometimes-appropriate code were classified as potentially

antibiotic-appropriate. Remaining unclassified visits with at least one never-appropriate code

were classified as antibiotic-inappropriate. The remaining visits were left unclassified. Note that,

in this classification, most visits are considered antibiotic-inappropriate, not because patients

are seeking antibiotics for antibiotic-inappropriate conditions, but rather because most

healthcare visits are not made for the purpose of treating bacterial infections.

Separately, each visit was classified as a visit with antibiotics if any of the prescribed or

administered medications included oral antibiotics [9].

Differences in rates were assessed using t-tests, with $p < 0.01$ considered statistically

significant, as recommended in the NHAMCS documentation [10]. Differences in proportions

were assessed using χ^2 tests. All analyses were performed using R (version 4.0.5), with the

survey package (version 4.0) used to account for the complex survey design in variance

estimation. The population denominators for each race/ethnicity provided in the

NAMCS/NHAMCS documentation, which are drawn from US Census estimates, were used for

population-based rate estimates. Code to reproduce these analyses is available at GitHub (doi:

10.5281/zenodo.6233588).

RESULTS

During 2016 and 2018, rates of physician office and hospital emergency room visits varied by

race/ethnicity. Non-Hispanic whites made the highest rate of visits, 7.0 (95% CI 6.6 to 7.4)

annual visits per capita, followed by non-Hispanic Blacks (5.5, 95% CI 4.9 to 6.0), Hispanics

(5.5, 95% CI 4.7 to 6.3), and non-Hispanic people of multiple or other races (4.2, 95% CI 3.2 to

5.2; Figure 1a). The difference in rates of visits was mostly attributable to different rates of

antibiotic-inappropriate visits, that is, visits with no associated diagnoses for which antibiotics

were appropriate or sometimes appropriate (*t*-tests, Figure 1a).

The proportion of visits during which antibiotics were prescribed or administered (5.5%, 95% CI

4.9% to 6.0%) did not vary statistically significantly by race/ethnicity ($p = 0.57$, χ^2 test), even

when stratifying by the antibiotic-appropriateness of the diagnoses associated with the visit ($p >$

0.05, χ^2 tests; Figure 1b).

Non-Hispanics whites had a higher rate of visits with antibiotics administered or prescribed (0.40

annual visits per capita, 95% CI 0.34 to 0.46) compared to non-Hispanic people of multiple or

other races (0.20 annual visits per capita, 95% CI 0.14 to 0.27; $p < 0.001$, *t*-test), but differences

between whites and Blacks (who had 0.33 annual antibiotic visits per capita, 95% CI 0.27 to

0.39) and between whites and Hispanics (who had 0.29, 95% CI 0.23 to 0.36) were not

statistically significant ($p > 0.01$, *t*-tests).

DISCUSSION

In this analysis of nationally representative survey data, whites made more healthcare visits per

capita. However, there was no statistically significant difference across races/ethnicities in the

proportion of visits that included antibiotics, even when stratifying by whether antibiotics were

appropriate for the visit. These results suggest that differences in healthcare utilization are at

least as important as per-visit prescribing practice to disparities in antibiotic use. For example,

whites had a 27% higher rate of healthcare visits than Blacks (7.0 vs. 5.5 annual visits per

capita), but the point estimate in the proportion of visits with antibiotics was 6% lower for whites

(5.7% vs. 6.0% of visits with antibiotics prescribed). Thus, difference in the overall rate of

antibiotic use between whites and Blacks is likely more attributable to differences in the number

of opportunities for antibiotics to be prescribed rather than to the probability that an antibiotic will

be prescribed at a visit.

This study's primary strength is its use of a nationally representative survey. Its most important

limitation is that the survey's sampling frame is the healthcare visits. A survey of visits can

estimate visits by race/ethnicity and the proportions of visits with antibiotics, but it cannot

address the proportion of infections that are treated with antibiotics because not every infection

that merits antibiotics results in a visit. Thus, without knowing the underlying rates of disease,

we cannot determine whether patients of different races/ethnicities with antibiotic-treatable

infections are more or less likely to present for treatment [1]. For example, if non-whites had

higher underlying rates of bacterial infections, then whites' approximately equal rates of

antibiotic-appropriate visits actually represent a disparity in healthcare access or healthcare

seeking.

This study has other important limitations. First, these national surveys did not have sufficient

statistical power to characterize any differences across races/ethnicities in the probability of per-

visit antibiotic use. For example, the point estimate for the proportion of Blacks' antibiotic-

appropriate visits that had antibiotics (35%, 95% CI 27% to 43%) was higher than that for whites

(30%, 95% CI 25% to 35%), but the confidence intervals were wide and overlapping (Figure 1b).

Second, these surveys do not include every kind of visit type and exclude, notably, urgent care,

telephone or telehealth contact, hospital outpatient departments, Veterans Administration

hospitals, and inpatient care. People of different races/ethnicities seek care in different

healthcare contexts [7] and so these results are necessarily incomplete. Third, there may be

systematic biases in the diagnosis codes that prescribers assign to people of different races

with the same underlying disease [1]. Finally, the available data cannot assess the effect of the

Covid-19 pandemic on antibiotic prescribing patterns.

Despite these limitations, these results point to two important components for ensuring equity in

the quality of care for bacterial infections: first, characterizing racial/ethnic differences in

underlying rates of antibiotic-treatable infections, and second, mitigating barriers to healthcare

access and utilization. It may be that these components are more important to antibiotic equity

than prescriber behavior.

STATEMENTS

Conflicts of interest

SWO is an employee of Biobot Analytics, Inc. YHG has consulted for GSK, holds grants from

Pfizer and Merck, and serves on the scientific advisory board for Day Zero Diagnostics.

Funding

YHG: Wellcome Trust Award 219812/Z/19/Z and internal Harvard T. H. Chan School award.

Data availability

Source data are openly available from the National Center for Health Statistics

(https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets, https://www.cdc.gov/nchs).

Figure 1. (a) Annual office-based and emergency department visits per capita, stratified by
patient race/ethnicity and by antibiotic-appropriateness of the diagnoses recorded for the visit.
Error bars show 95% confidence intervals. *: *t*-test, $p < 0.01$. (b) Proportion of visits with
antibiotics administered or prescribed.

REFERENCES

1. Fleming-Dutra KE, Shapiro DJ, Hicks LA, Gerber JS, Hersh AL. Race, Otitis Media, and Antibiotic Selection. Pediatrics **2014**; 134:1059–1066.

2. Little P. Delayed prescribing of antibiotics for upper respiratory tract infection. BMJ **2005**; 331:301.

3. Cosgrove SE, Klein EY. Reassessing the Link Between Healthcare Access and Outpatient Antibiotic Prescribing. J Infect Dis **2021**; 223:2017–2019.

4. Olesen SW, Grad YH. Racial/Ethnic Disparities in Antimicrobial Drug Use, United States, 2014–2015 - Volume 24, Number 11—November 2018 - Emerging Infectious Diseases journal - CDC. Emerg Infect Dis **2018**; 24:2126–2128.

5. Goyal MK, Johnson TJ, Chamberlain JM, et al. Racial and Ethnic Differences in Antibiotic Use for Viral Illness in Emergency Departments. Pediatrics **2017**; :e20170203.

6. Steinman MA, Landefeld SC, Gonzales R. Predictors of Broad-Spectrum Antibiotic Prescribing for Acute Respiratory Tract Infections in Adult Primary Care. JAMA **2003**; 289:719–725.

7. Halasa NB, Griffin MR, Zhu Y, Edwards KM. Differences in antibiotic prescribing patterns for children younger than five years in the three major outpatient settings. J Pediatr **2004**; 144:200–205.

8. Kissler SM, Klevens RM, Barnett ML, Grad YH. Childhood respiratory outpatient visits correlate with socioeconomic status and drive geographic patterns in antibiotic prescribing. J Infect Dis **2021**;

9. Fleming-Dutra KE, Hersh AL, Shapiro DJ, et al. Prevalence of Inappropriate Antibiotic Prescriptions Among US Ambulatory Care Visits, 2010-2011. JAMA **2016**; 315:1864–1873.

10. Centers for Disease Control and Prevention. Ambulatory Health Care Data Datasets and Documentation. 2021; Available at: https://www.cdc.gov/nchs/ahcd/datasets_documentation_related.htm. Accessed 13 February 2022.

11. Centers for Disease Control and Prevention. Notices for NAMCS and NHAMCS Public Use Data File Users. 2021; Available at: https://www.cdc.gov/nchs/ahcd/notice.htm. Accessed 12 February 2021.

12. Chua K-P, Fischer MA, Linder JA. Appropriateness of outpatient antibiotic prescribing among privately insured US patients: ICD-10-CM based cross sectional study. BMJ **2018**; 364:k5092.