1

SARS-CoV-2 and endemic coronaviruses: Comparing symptom

presentation and severity of symptomatic illness among Nicaraguan

Children

Aaron M Frutos[1], John Kubale[1], Guillermina Kuan[2], Sergio Ojeda[3], Nivea Vydiswaran[1], Nery

Sanchez[3], Miguel Plazaola[3], May Patel[1], Roger Lopez[4], Angel Balmaseda[3¶], Aubree Gordon[1*¶]

[1] Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor,

MI, USA

[2] Health Center Sócrates Flores Vivas, Ministry of Health, Managua, Nicaragua

[3] Laboratorio Nacional de Virología, Centro Nacional de Diagnóstico y Referencia, Ministry of

Health, Managua, Nicaragua.

[4] Sustainable Sciences Institute, Managua, Nicaragua.

*Corresponding author

E-mail: gordonal@umich.edu (AG)

¶ These authors contributed equally.

Abstract

It has been proposed that as SARS-CoV-2 transitions to endemicity, children will represent the greatest proportion of SARS-Co-V-2 infections as they currently do with endemic coronavirus infections. While SARS-CoV-2 infection severity is low for children, it is unclear if SARS-CoV-2 infections are distinct in symptom presentation, duration, and severity from endemic coronavirus infections in children. We compared symptom risk and duration of endemic coronavirus infections from 2011-2016 with SARS-CoV-2 infections from March 2020-September 2021 in a Nicaraguan pediatric cohort. Respiratory samples were collected from participants that met testing criteria and blood samples were collected annually. Respiratory samples were tested for each of the endemic coronaviruses from 2011-2016 and for SARS-CoV-2 from 2020-2021 via rt-PCR. 2021 blood samples were tested for SARS-CoV-2 antibodies and a subset of 2011-2016 blood samples from four-years-old participants were tested for endemic coronavirus antibodies. By April 2021, 854 (49%) active participants were ELISA positive for SARS-CoV-2 antibodies. Most participants had antibodies against one alpha and one beta coronavirus by age four. We observed 595 symptomatic endemic coronavirus infections from 2011-2016 and 121 symptomatic with SARS-CoV-2 infections from March 2020-September 2021. Symptom presentation of SARS-CoV-2 infection and endemic coronavirus infections were very similar, and SARS-CoV-2 symptomatic infections were as or less severe on average than endemic coronavirus infections. This suggests that, for children, SARS-CoV-2 may be just another endemic coronavirus. However, questions about the impact of variants and the long-term effects of SARS-CoV-2 remain.

Introduction

As severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) transitions to global endemicity, there are many questions about how that will occur. Over time, children will represent the greatest proportion of primary SARS-CoV-2 infections as adults gain immunity from natural infection or vaccination. (1) It is well established that pediatric risk of severe illness and death is much lower than that for adults. (2, 3) Differences in immune response between adults and children likely provide children better protection against severe SARS-CoV-2 infections. (3-5) Previous research found no differences in severity between SARS-CoV-2 infections and influenza A and B among children. (6, 7) However, it is unknown if, in children, SARS-CoV-2 infections are distinct in disease presentation and severity from endemic human coronavirus (HCoV) infections. If SARS-COV-2 infections are more severe, routine pediatric vaccination will be necessary to reduce excess mortality. If not, vaccine-induced immunity should prevent severe disease while allow for transmission to facilitate frequent immune boosting. (1) To determine if SARS-CoV-2 infections have distinct disease presentation from endemic HCoV infections in children, we compare symptomatic SARS-CoV-2 and endemic HCoVs infections symptomology and severity in a prospective, community-based pediatric cohort in Managua, Nicaragua from 2011-2016 and 2020-2021

Methods

61   Institutional review boards at the Nicaraguan Ministry of Health and the University of

62 Michigan approved this study. Parents/guardians of participants provided written informed

63 consent and participants aged ≥6 years provided verbal assent. The Nicaraguan Pediatric

64 Influenza Cohort Study (NPICS) is a prospective cohort study that began in 2011 and continues

65 today. Participants included in this analysis were members of the NPICS located in District 2 of

66 Managua, Nicaragua between 2011-2016 or March 2020 to September 2021. Children ages 0-14

67 years who live within the catchment area of Health Center Sócrates Flores Vivas were eligible to

68 participate.

69   Blood samples were collected from each study participant between February and April

70 each year. To confirm our assumption that endemic HCoV infection rates are high in the cohort,

71 we tested a random subset of 100 blood samples from four-year-old's from 2011-2016; using

72 protocols for an enzyme-linked immunosorbent assay (ELISA) developed at Mount Sinai (8), we

73 tested samples for antibody response to the spike protein for each of the four endemic HCoV

74 (alpha: NL63, and 229E, beta: OC43 and HKU1). To confirm that SARS-CoV-2 infections rates

75 were also high, we tested the 2021 blood samples for SARS-CoV-2 antibodies as described

76 previously. (9)

77   Parents/guardians agreed to bring participants to the health center at the first signs of

78 fever. Study personnel collected a respiratory swab from participants if they met the testing

79 criteria: feverishness for participants under two years old; measured fever/feverishness and

80 cough, sore throat, or rhinorrhea; severe respiratory symptoms such as apnea or chest indrawing;

81 hospitalization with respiratory symptoms or sepsis. Testing criteria were expanded in June 2020

to capture mild symptomatic SARS-CoV-2 cases; however, this analysis is limited to those

meeting the original testing criteria to ensure comparability. We used real-time reverse

transcription polymerase-chain reaction (RT-PCR) to test respiratory samples from 2011-2016

for each of the four endemic HCoV, influenza, respiratory syncytial virus (RSV) and human

metapneumovirus (HMpV) and samples from March 2020-September 2021 for SARS-CoV-2

and influenza. (10, 11)

Study clinicians record all participant symptoms and diagnoses during each clinic visit.

Parents/guardians also report participant symptoms at the initial clinic visit, and each day

thereafter until symptom resolution. We considered symptoms to be associated with an infection

if they occurred within 28 days of symptom onset. Symptoms that were included for follow-up in

2011-2016 and 2020-2021 were feverishness, cough, rhinorrhea, nasal congestion, loss of

appetite, myalgia, arthralgia, and rapid breathing. We considered participants who presented to

the clinic with rapid breathing, rhonchi, indrawing, wheezing, or shortness of breath as having

abnormal breathing. We defined acute lower respiratory infections (ALRI) as physician

diagnosed cases of pneumonia, bronchiolitis, bronchitis, or bronchial hyperreactivity.

To compare risk of symptoms, we calculated symptom specific risk differences between

SARS-CoV-2 infections and endemic HCoV infections, overall and stratified by the following

age groups: 0-4, 5-9, and 10-14 years. We also stratified the results by sex and endemic HCoV

species. Using upset plots, we explored which signs/symptoms tended to present together. We

also assessed symptom duration by comparing the time between each specific symptom onset

and the last day participants presented with that symptom. We plotted symptom duration using

boxplots and compared the distribution of symptom duration between SARS-CoV-2 and

endemic HCoV using the Mann Whitney U test.

105   We used SAS version 9.4 (SAS Institute Inc.) to calculate risk differences and ratios and

106   R version 4.1.0 to create figures and conduct all other analyses.

Results

109   There were 3,220 participants active in NPICS during the included years: 2,576 from

2011-2016 and 1,942 from March 2020-September 2021 with an average of 1,792 per year. Our

assumption that detected symptomatic infections consist of only a small proportion of total

SARS-CoV-2 and endemic HCoV infections that occurred in the cohort children. Specifically, of

the 1,942 children in the cohort from March 2020-September 2021, 1,743 had a blood sample

collected in early to mid- 2021 that was tested via ELISA for SARS-CoV-2. 854 (49%) tested

positive for SARS-CoV-2 antibodies indicating that SARS-CoV-2 infections were very high in

the cohort. We found that 94% of our randomly selected subset of four-year-olds from 2011-

2016 had an antibody response to at least one alpha and one beta HCoV, confirming our

assumption of high endemic HCoV infection rates in the community. Antibody response

prevalence was highest for OC43 (99%) followed by HKU1 (86%), NL63 (83%), and then 229E

(74%) (Table 1).

Table 1: Endemic Coronavirus ELISA Results, % Positive

	Total (n=100)
Alpha	94%
NL63	83%
229E	74%
Beta	100%
OC43	99%
HKU1	86%

Within this cohort we observed high infection rates of both endemic HCoVs and SARS-

CoV-2. That there were 595 RT-PCR+ symptomatic endemic HCoV infections from 2011-2016

and 121 RT-PCR+ symptomatic SARS-CoV-2 infections from March 2020- September 2021

again suggests that symptomatic cases represent only a small proportion of overall infections.

Most endemic HCoV, 432 (73%), and SARS-CoV-2, 59 (49%), infections occurred in

participants <5 years. Fever, cough, rhinorrhea, and congestion were the most common

symptoms for both endemic HCoVs and SARS-CoV-2 symptomatic infections. Among SARS-

CoV-2 symptomatic infections, 4 (3%) were acute lower respiratory infections compared to 98

(16%) endemic HCoV cases (p=<0.0001; Table 2).

Table 2: Study Participants and Symptom Prevalence

		Endemic (n=595)	SARS-CoV-2 (n=121)	p-value
Age Group (%)				<.0001
	0-4	432 (73)	59 (49)	
	5-9	110 (18)	29 (24)	
	10-14	53 (9)	33 (27)	
Symptoms (%)				
Measured Fever		272 (46)	68 (56)	0.035
Cough		524 (88)	85 (70)	<0.001

Rhinorrhea	505 (85)	93 (77)	0.030
Congestion	276 (46)	45 (37)	0.064
Sore Throat	145 (24)	36 (30)	0.221
Headache	81 (14)	38 (31)	<0.001
Loss of Appetite	142 (24)	30 (25)	0.828
Diarrhea	64 (11)	14 (12)	0.793
Pneumonia	40 (7)	4 (3)	0.154
Hospitalized	23 (4)	6 (5)	0.578
Abnormal Breathing	108 (18)	8 (6)	0.002
Acute Lower Respiratory Infection	98 (16)	4 (3)	<0.001

* p-value from chi-square test

Across age groups, participants with symptomatic endemic HCoV infections displayed

greater risk of cough compared to those with symptomatic SARS-CoV-2 infections. Among

participants aged <5 years, symptomatic endemic HCoV infections showed greater risk of

rhinorrhea, congestion, abnormal breathing, and ALRI than symptomatic SARS-CoV-2

infections, while SARS-CoV-2 exhibited greater risk of measured fever (Fig 1, Table 3).

Notably, symptomatic endemic HCoV infection was associated with greater risk of ALRI even

after excluding participants that also tested positive for influenza A, influenza B, RSV, or

HMpV. We observed no difference in risk of pneumonia or hospitalization between SARS-CoV-

2 and endemic HCoV symptomatic infections. These results were consistent after stratifying by

sex (S1-S2 Fig), or endemic HCoV species (S3-S6 Fig).

Fig 1: Symptom Risk Difference between Endemic HCoV and SARS-CoV-2.

A: All participants. B: Ages 0-4 years. C: Ages 5-9 years. D Ages: 10-14 years. Positive risk difference indicates higher risk for endemic HCoVs.

155 **Table 3: Symptom Risk Overall and by Age Group**

All

	Endemic (%)	SARS-CoV-2 (%)	Risk Difference (95% CI)	Risk Ratio (95% CI)
Measured Fever	272 (46)	68 (56)	-0.10 (-0.20, -0.01)	0.81 (0.68, 0.97)
Cough	524 (88)	85 (70)	0.18 (0.09, 0.26)	1.25 (1.11, 1.41)
Runny nose	505 (85)	93 (77)	0.08 (-0.00, 0.16)	1.10 (1.00, 1.22)
Congestion	276 (46)	45 (37)	0.09 (-0.00, 0.19)	1.24 (0.97, 1.60)
Sore throat	145 (24)	36 (30)	-0.05 (-0.14, 0.03)	0.82 (0.60, 1.11)
Headache	81 (14)	38 (31)	-0.18 (-0.27, -0.09)	0.43 (0.31, 0.60)
Loss of appetite	142 (24)	30 (25)	-0.01 (-0.09, 0.07)	0.96 (0.68, 1.35)
Diarrhea	64 (11)	14 (12)	-0.01 (-0.07, 0.05)	0.93 (0.54, 1.60)
Pneumonia	40 (7)	4 (3)	0.03 (-0.00, 0.07)	2.03 (0.74, 5.58)
Hospitalized	23 (4)	6 (5)	-0.01 (-0.05, 0.03)	0.78 (0.32, 1.87)
Abnormal Breathing	108 (18)	8 (7)	0.12 (0.06, 0.17)	2.75 (1.38, 5.48)
ALRI	98 (16)	4 (3)	0.13 (0.09, 0.18)	4.98 (1.87, 13.28)
ALRI[†]	77 (15)	4 (3)	0.12 (0.07,0.16)	4.55 (1.70, 12.19)

0-4

	Endemic (%)	SARS-CoV-2 (%)	Risk Difference (95% CI)	Risk Ratio (95% CI)
Measured Fever	218 (50)	44 (75)	-0.24 (-0.36, -0.12)	0.68 (0.57, 0.81)
Cough	382 (88)	42 (71)	0.17 (0.05, 0.29)	1.24 (1.05, 1.47)
Runny nose	378 (88)	42 (71)	0.16 (0.04, 0.28)	1.23 (1.04, 1.45)
Congestion	241 (50)	21 (36)	0.14 (0.09, 0.27)	1.39 (0.97, 1.99)
Sore throat	66 (15)	6 (10)	0.05 (-0.03, 0.14)	1.50 (0.68, 3.31)
Headache	23 (5)	4 (7)	-0.01 (-0.08, 0.05)	0.79 (0.28, 2.19)
Loss of appetite	116 (27)	18 (31)	-0.04 (-0.16, 0.09)	0.88 (0.58, 1.33)
Diarrhea	58 (13)	9 (15)	-0.02 (-0.12, 0.08)	0.88 (0.46. 1.68)
Pneumonia	38 (9)	3 (5)	0.04 (-0.03, 0.10)	1.73 (0.54, 6.03)
Hospitalized	17 (4)	4 (7)	-0.03 (-0.10, 0.04)	0.58 (0.20, 1.67)
Abnormal Breathing	95 (22)	3 (5)	0.17 (0.10, 0.24)	4.34 (1.42, 13.21)
ALRI	88 (20)	3 (5)	0.15 (0.9, 0.22)	4.00 (1.31, 12.26)
ALRI[†]	69 (19)	3 (5)	0.13 (0.07,0.20)	3.65 (1.19, 11.21)

5-9

	Endemic (%)	SARS-CoV-2 (%)	Risk Difference (95% CI)	Risk Ratio (95% CI)
Measured Fever	39 (35)	12 (41)	-0.06 (-0.26, 0.14)	0.86 (0.52, 1.41)
Cough	99 (90)	21 (72)	0.18 (000, 0.35)	1.24 (0.98, 1.57)
Runny nose	87 (79)	25 (86)	-0.07 (-0.22, 0.08)	0.92 (0.77, 1.09)

Congestion	42 (38)	14 (48)	-0.10 (-0.10, 0.30)	0.79 (0.51, 1.23)
Sore throat	52 (47)	12 (41)	0.06 (-0.14, 0.26)	1.14 (0.71, 1.84)
Headache	35 (32)	14 (48)	-0.16 (-0.37, 0.04)	0.66 (0.41, 1.04)
Loss of appetite	19 (17)	7 (24)	-0.07 (-0.24, 0.10)	0.72 (0.33, 1.54)
Diarrhea	4 (4)	3 (10)	-0.07 (-0.18, 0.05)	0.35 (0.08, 1.48)
Pneumonia	2 (2)	1 (3)	-0.02 (-0.9, 0.05)	0.53 (0.05, 5.61)
Hospitalized	5 (5)	2 (7)	-0.02 (-0.12, 0.08)	0.66 (0.13, 3.23)
Abnormal Breathing	10 (9)	2 (7)	0.02 (-0.08, 0.12)	1.32 (0.31, 5.69)
ALRI	8 (7)	1 (3)	0.04 (-0.04, 0.12)	2.11 (0.27, 16.19)
ALRI[†]	6 (7)	1 (3)	0.03 (-0.05, 0.11)	1.89 (0.23, 16.93)

10-14

	Endemic (%)	SARS-CoV-2 (%)	Risk Difference (95% CI)	Risk Ratio (95% CI)
Measured Fever	15 (28)	12 (36)	-0.08 (-0.28, 0.12)	0.78 (0.42, 1.45)
Cough	43 (81)	22 (67)	0.14 (-0.05, 0.34)	1.22 (0.93, 1.60)
Runny nose	40 (75)	26 (79)	-0.03 (-0.21, 0.15)	0.96 (0.76, 1.21)
Congestion	20 (38)	10 (30)	0.07 (-0.13, 0.28)	1.25 (0.67, 2.32)
Sore throat	27 (51)	18 (55)	-0.04 (-0.25, 0.18)	0.93 (0.62, 1.41)
Headache	23 (43)	20 (61)	-0.17 (-0.39, 0.04)	0.25 (0.47, 1.08)
Loss of appetite	7 (13)	5 (15)	-0.02 (-0.17, 0.13)	0.87 (0.30, 2.52)
Diarrhea	2 (4)	2 (6)	-0.02 (-0.12, 0.07)	0.62 (0.09, 4.21)
Pneumonia	0	0	-	-
Hospitalized	1 (2)	0	0.02 (-0.02, 0.06)	-
Abnormal Breathing	3 (6)	3 (9)	-0.03 (-0.15, 0.08)	0.62 (0.13, 2.90)
ALRI	2 (4)	0	0.04 (-0.01, 0.09)	-
ALRI[†]	2 (4)	0	0.04 (-0.01, 0.10)	-

156 *Positive risk difference, risk ratio > 1 indicates higher risk for endemic HCoV infection.
157 † Excluding influenza A, influenza B, RSV, and HmpV coinfections
158

For both endemic HCoV and SARS-CoV-2, we found that cough, rhinorrhea, and sore

throat frequently presented together. Loss of appetite appeared in common symptom groupings

for endemic HCoV, while headache was part of more common groupings for SARS-CoV-2 (Fig

2). We did find that among participants aged 0-4 years decreased appetite lasted longer and

among participants aged 5-9 and 10-14, cough lasted longer for SARS-CoV-2 infections., (Fig 2,

Supplemental Table 4).

Fig 2. Comparison of Common Symptom Groupings and Duration between Symptomatic

Endemic Coronavirus and SARS-CoV-2 Infections.

A: Upset plot of symptom groupings for symptomatic SARS-CoV-2 infections. B. Upset plot of

symptom groupings for symptomatic endemic coronavirus infections. C. Boxplots of symptom

duration

Table 4: Symptom Duration Comparison: SARS-CoV-2, Endemic Coronaviruses

Mean Symptom Duration in Days (SD)		
All		
SARS-CoV-2	**Endemic Coronaviruses**	**p-value***
Feverish	3.8 (6.1)	3.8 (7.1)	0.137
Cough	6.7 (6.4)	5.5 (8.0)	0.003
Runny nose	4.4 (5.7)	4.6 (7.3)	0.354
Congestion	2.8 (4.7)	2.8 (5.6)	0.997
Decreased appetite	2.2 (2.8)	1.4 (3.4)	0.032
0-4		
SARS-CoV-2	**Endemic Coronaviruses**	**p-value***
Feverish	5.2 (7.3)	4.4 (7.6)	0.069
Cough	6.4 (6.0)	6.4 (8.6)	0.178
Runny nose	5.1 (5.8)	5.4 (7.8)	0.278
Congestion	2.6 (3.8)	3.0 (5.8)	0.875
Decreased appetite	2.5 (2.9)	1.6 (3.7)	0.039
5-9		
SARS-CoV-2	**Endemic Coronaviruses**	**p-value***
Feverish	2.0 (4.0)	2.4 (5.4)	0.397
Cough	8.1 (7.5)	3.2 (5.8)	0.002
Runny nose	3.9 (5.5)	2.3 (4.7)	0.125
Congestion	3.4 (6.3)	1.9 (4.4)	0.690
Decreased appetite	2.1 (26)	1.0 (1.2)	0.351
10-14		
SARS-CoV-2	**Endemic Coronaviruses**	**p-value***
Feverish	2.6 (4.7)	1.7 (3.8)	0.434
Cough	6.0 (6.2)	2.3 (4.3)	0.015
Runny nose	3.5 (6.0)	2.2 (5.0)	0.440
Congestion	2.5 (3.9)	2.6 (5.6)	0.479
Decreased appetite	1.2 (2.2)	0.4 (0.8)	0.698

To assess the potential impact of variants on symptoms, we compared symptoms for

SARS-CoV-2 infections from 2020 prior to the global emergence of variants and 2021 when

delta, gamma, and lambda strains circulated in the cohort area and found no difference in

presentation by year (Fig 3). All observed cases of ALRI associated with SARS-CoV-2 occurred

in 2020. However, we did find that rhinorrhea lasted longer in SARS-CoV-2 cases from 2021

compared to 2020 (Table 5).

Fig 3. Comparison of Common Symptom Groupings between Symptomatic SARS-CoV-2

Infections in 2020 and 2021

A: 2020. B. 2021

Table 5: Symptom Duration Comparison: SARS-CoV-2 by Year

	Mean Symptom Duration in Days (SD)		p-value*
	2020	**2021**	
Feverish	4.1 (7.4)	3.46(5.0)	0.129
Cough	6.8 (8.0)	6.7 (5.5)	0.378
Runny nose	2.7 (5.0)	5.4 (6.0)	0.004
Congestion	1.6 (2.3)	3.3 (5.2)	0.382
Decreased appetite	3.5 (3.8)	1.6 (1.9)	0.172

*From Mann-Whitney U test

Discussion

This is, to our knowledge, the first study that assesses differences in symptom

presentation, duration, and severity between SARS-CoV-2 and endemic HCoV symptomatic

infections among children. Understanding how SARS-CoV-2 infections compare to endemic

HCoV infections is important as SARS-CoV-2 becomes endemic. While our study does not

evaluate asymptomatic or very mild illness, we provide comparisons of medically attended

illness associated with endemic HCoV and SARS-CoV-2 symptomatic infections in a large

cohort of children with high infection rates.

In this pediatric cohort, we found that with 854 (49%) participants tested positive for

SARS-CoV-2 antibodies in 2021 with only 121 PCR confirmed infections that met the original

testing criteria, like results from our community-based household cohort study. (9) We also

found that most participants had at least one alpha and one beta HCoV infection by age four

suggesting that in this cohort, participants have had at least two endemic HCoV exposures by the

age of four. This is similar to previous studies that showed that by age six, most children have

had infections with each of the four endemic HCoVs with a majority being asymptomatic

infections. (12) Thus for SARS-CoV-2 and endemic HCoV, symptomatic infections also

represent only a small proportion of all pediatric infections.

Comparing these symptomatic infections, we found that, overall, disease presentation is

very similar between endemic HCoV and SARS-CoV-2, with each frequently presenting with

"common cold" symptoms. Consistent with other studies, we found differences in symptom

presentation by age perhaps in part because older children have had more endemic HCoV

exposures (2, 4, 13). We also found great variability in symptom duration for SARS-CoV-2 and

endemic HCoV infections with symptoms lasting from 1-28 days or more. (14) There was a

difference in duration of decreased appetite for the youngest participants and cough for those

aged 5-14 years suggesting that some symptoms may last longer for SARS-CoV-2 infections.

We also found a difference in rhinorrhea duration between SARS-CoV-2 infections in 2020 and

2021, suggesting that SARS-CoV-2 variants may increase symptom duration among children.

Across age groups, risk of ALRI associated with SARS-CoV-2 was the same or lower

compared to ALRI associated with endemic HCoV infection. Even when excluding endemic

HCoV co-infections with pathogens commonly associated with ALRI, the conclusions did not

change. These results show that, for children, the risk of ALRI and severe illness from SARS-

CoV-2 infections is comparable to the risk from endemic HCoV infections at the community

level.

The main strength of this community-based study is its size and duration. Consistent viral

surveillance and symptom evaluation within the same population allow for year-to-year

comparisons and facilitates our comparisons of endemic HCoV and SARS-CoV-2. The high

number of participants under the age of five (about 36% of participants during these years),

allows us to evaluate SARS-CoV-2 in an age group with little representation in current literature.

Additionally, this cohort was already well established when SARS-CoV-2 began circulating in

Nicaragua allowing us to quickly incorporate questions regarding its effects on this population.

However, this study does have some limitations. First, using data from this community-

based cohort study we were not powered to detect the most severe manifestations of SARS-CoV-

2 including death or Multisystem Inflammatory Syndrome in Children (MIS-C) and other rare

outcomes. (15, 16) Second, our analysis did not include genetic sequencing preventing us from

assessing the importance of variants in presentation and severity of SARS-CoV-2 illness. We did

compare SARS-CoV-2 symptom presentation, severity, and duration by year and found little or

no difference. Finally, while our study does not evaluate asymptomatic illness for endemic

HCoVs, our results were consistent with other research, showing that childhood HCoVs are

ubiquitous and that symptomatic cases represent only a small proportion of infections, as with

SARS-CoV-2. (12)

In this study, we observed that symptomatic SARS-CoV-2 infections at the community

level are very similar to symptomatic endemic HCoV infections in symptom presentation.

Among children in a setting with a high SARS-CoV-2 infection rate, symptomatic SARS-CoV-2

infections are on average as or less severe as endemic HCoV infections. These findings support

the hypothesis that SARS-CoV-2 may be like another endemic HCoV for children—most

children will be asymptomatic with rare cases of severe symptomatic illness. This does not mean

SARS-CoV-2 in children is not important. There are many unknowns about the long-term effects

and impact of repeat infections among children. (1) Increased transmissibility of emerging

variants is also a cause for concern, as it will lead to increased frequency of severe

manifestations. Future mutations in the virus needed to be monitored as they may also increase

illness severity in children. Despite relatively low risk of severe illness among children, pediatric

vaccination that mirrors natural induced immunity against SARS-CoV-2 would further lower

individual risk and reduce the number of severe cases and deaths due to SARS-CoV-2.

References

1.   Lavine JS, Bjornstad ON, Antia R. Immunological characteristics govern the transition of

COVID-19 to endemicity. Science. 2021;371(6530):741-5.

2.   Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiology of COVID-19 Among

Children in China. Pediatrics. 2020;145(6).

3.   Do LAH, Anderson J, Mulholland EK, Licciardi PV. Can data from paediatric cohorts

solve the COVID-19 puzzle? PLoS Pathog. 2020;16(9):e1008798.

4.   Dhochak N, Singhal T, Kabra SK, Lodha R. Pathophysiology of COVID-19: Why

Children Fare Better than Adults? Indian J Pediatr. 2020;87(7):537-46.

5.   Khan T, Rahman M, Ali FA, Huang SSY, Ata M, Zhang Q, et al. Distinct antibody

repertoires against endemic human coronaviruses in children and adults. JCI Insight. 2021;6(4).

6.   Siddiqui M, Gultekingil A, Bakirci O, Uslu N, Baskin E. Comparison of clinical features

and laboratory findings of coronavirus disease 2019 and influenza A and B infections in

children: a single-center study. Clin Exp Pediatr. 2021.

7.   Pokorska-Spiewak M, Talarek E, Popielska J, Nowicka K, Oldakowska A, Zawadka K, et

al. Comparison of clinical severity and epidemiological spectrum between coronavirus disease

2019 and influenza in children. Sci Rep. 2021;11(1):5760.

8.   Stadlbauer D, Amanat F, Chromikova V, Jiang K, Strohmeier S, Arunkumar GA, et al.

SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a Serological Assay, Antigen

Production, and Test Setup. Curr Protoc Microbiol. 2020;57(1):e100.

9.   Maier HE, Kuan G, Saborio S, Bustos Carrillo FA, Plazaola M, Barilla C, et al. Clinical

spectrum of SARS-CoV-2 infection and protection from symptomatic re-infection. Clin Infect

Dis. 2021.

10.   Chu DKW, Pan Y, Cheng SMS, Hui KPY, Krishnan P, Liu Y, et al. Molecular Diagnosis

of a Novel Coronavirus (2019-nCoV) Causing an Outbreak of Pneumonia. Clin Chem.

2020;66(4):549-55.

11.   Real-Time RT-PCR Assays for Non-Influenza Respiratory Viruses. In: Diseases NCfIaR,

editor.: Centers for Disease Control and Prevention; 2015. p. 1-16.

12.   Zhou W, Wang W, Wang H, Lu R, Tan W. First infection by all four non-severe acute

respiratory syndrome human coronaviruses takes place during childhood. BMC Infect Dis.

2013;13:433.

13.   Ogimi C, Kim YJ, Martin ET, Huh HJ, Chiu CH, Englund JA. What's New With the Old

Coronaviruses? J Pediatric Infect Dis Soc. 2020;9(2):210-7.

14.   Molteni E, Sudre CH, Canas LS, Bhopal SS, Hughes RC, Antonelli M, et al. Illness

duration and symptom profile in symptomatic UK school-aged children tested for SARS-CoV-2.

Lancet Child Adolesc Health. 2021.

15.   Children and COVID-19: State-Level Data Report. American Academy of Pediatrics

2021 August 12, 2021.

16.   Payne AB, Gilani Z, Godfred-Cato S, Belay ED, Feldstein LR, Patel MM, et al.

Incidence of Multisystem Inflammatory Syndrome in Children Among US Persons Infected

With SARS-CoV-2. JAMA Netw Open. 2021;4(6):e2116420.

Supporting information

302 S1 Fig. Symptom Risk Difference between Endemic HCoVs and SARS-CoV-2 for Females.

A: All participants. B: Ages 0-4. C: Ages 5-9. D Ages: 10-14. Positive risk difference indicates

higher risk for endemic HCoV infection

S2 Fig. Symptom Risk Difference between Endemic HCoVs and SARS-CoV-2 for Males.

A: All participants. B: Ages 0-4. C: Ages 5-9. D Ages: 10-14. Positive risk difference indicates

higher risk for endemic HCoV infection

S3 Fig. Symptom Risk Difference NL63 and SARS-CoV-2.

A: All participants. B: Ages 0-4. C: Ages 5-9. D Ages: 10-14. Positive risk difference indicates

higher risk for endemic HCoV infection

S4 Fig. Symptom Risk Difference between 229E and SARS-CoV-2.

A: All participants. B: Ages 0-4. C: Ages 5-9. D Ages: 10-14. Positive risk difference indicates

higher risk for endemic HCoV infection

S5 Fig. Symptom Risk Difference between OC43 and SARS-CoV-2.

A: All participants. B: Ages 0-4. C: Ages 5-9. D Ages: 10-14. Positive risk difference indicates

higher risk for endemic HCoV infection

S6 Fig. Symptom Risk Difference between HKU1 and SARS-CoV-2.

A: All participants. B: Ages 0-4. C: Ages 5-9. D Ages: 10-14. Positive risk difference indicates

higher risk for endemic HCoV infection