Title. Proteomic Characterization of Acute Kidney Injury in Patients Hospitalized with SARS-CoV2 Infection

Running Title. Proteomic Characterization of AKI in COVID-19

Authors. Ishan Paranjpe[1,2], Pushkala Jayaraman[4], Chen-Yang Su[11,13], Sirui Zhou[11,12], Steven Chen[3], Ryan Thompson[1†], Diane Marie Del Valle[1†], Ephraim Kenigsberg[3,4,5] , Shan Zhao[6], Suraj Jaladanki[1,2], Kumardeep Chaudhary[1,2,4], Steven Ascolillo[1], Akhil Vaid[1], Arvind Kumar[1], Manish Paranjpe[7], Ross O'Hagan[1,2], Samir Kamat[1], Faris F. Gulamali[1,2], Hui Xie[8], Joceyln Harris[8], Manishkumar Patel[8], Kimberly Argueta[8], Craig Batchelor[8], Kai Nie[8], Sergio Dellepiane[9], Leisha Scott[8], Matthew A Levin[1,6], John Cijiang He[9], Steven G Coca[9], Lili Chan[9], Evren U Azeloglu[9], Eric Schadt[4], Noam Beckmann[1,4], Sacha Gnjatic[10], Miram Merad[3], Seunghee Kim-Schulze[3,8], Brent Richards[11,12,14,15], Benjamin S Glicksberg[1,2,4], Alexander W Charney[1,4,11], Girish N Nadkarni[1,2,9,12]

† These authors contributed equally to the work.

Affiliations:

1. The Mount Sinai Clinical Intelligence Center (MSCIC), Icahn School of Medicine at Mount Sinai, New York, NY, USA

2. The Hasso Plattner Institute for Digital Health at Mount Sinai, Icahn School of Medicine at Mount Sinai, New York, NY, USA

3. The Precision Immunology Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

4. Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

5. Icahn Institute for Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

6. Department of Anesthesiology, Perioperative and Pain Medicine, Icahn School of Medicine at Mount Sinai, New York, New York, United States of America

7. Division of Health Sciences and Technology, Harvard Medical School, Boston, MA, USA

8. Human Immune Monitoring Center, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

9. Department of Medicine, Division of Nephrology, Icahn School of Medicine at Mount Sinai, New York, New York, United States of America

10. Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

11. Lady Davis Institute, Jewish General Hospital, McGill University, Montréal, Québec, Canada

12. Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montréal, Québec, Canada

13. Department of Computer Science, McGill University, Montréal, Québec, Canada

14. Department of Human Genetics, McGill University, Montréal, Québec, Canada

15. Department of Twin Research, King's College London, London, United Kingdom

16. The Pamela Sklar Division of Psychiatric Genomics, Icahn School of Medicine at Mount Sinai, New York, New York, United States of America

17. The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, New York, United States of America

*Corresponding Author:

Girish N Nadkarni, MD

Department of Medicine, Division of Nephrology

Icahn School of Medicine at Mount Sinai, New York, NY, USA

girish.nadkarni@mountsinai.org

Abstract

Acute kidney injury (AKI) is a known complication of COVID-19 and is associated with an increased risk of in-hospital mortality. Unbiased proteomics using longitudinally collected biological specimens can lead to improved risk stratification and discover pathophysiological mechanisms. Using longitudinal measurements of ~4000 plasma proteins in two cohorts of patients hospitalized with COVID-19, we discovered and validated markers of COVID-associated AKI (stage 2 or 3) and long-term kidney dysfunction. In the discovery cohort (N= 437), we identified 413 upregulated and 40 downregulated proteins associated with COVID-AKI (adjusted p <0.05). Of these, 62 proteins were validated in an external cohort (p <0.05, N =261). We demonstrate that COVID-AKI is associated with increased markers of tubular injury (NGAL) and myocardial injury. Using estimated glomerular filtration (eGFR) measurements taken after discharge, we also find that 25 of the 62 AKI-associated proteins are significantly associated with decreased post-discharge eGFR (adjusted p <0.05). Proteins most strongly associated with decreased post-discharge eGFR included desmocollin-2, trefoil factor 3, transmembrane emp24 domain-containing protein 10, and cystatin-C indicating tubular dysfunction and injury. Using longitudinal clinical and proteomic data, our results suggest that while both acute and long-term COVID-associated kidney dysfunction are associated with markers of tubular dysfunction, AKI is driven by a largely multifactorial process involving hemodynamic instability and myocardial damage.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a novel coronavirus that has caused the coronavirus disease 2019 (COVID-19) pandemic. Although effective vaccines are available, novel variants that may evade neutralizing antibodies exist in the population and have led to high case counts and periodic case surges. COVID-19 most commonly presents with fever, cough, and dyspnea[1,2] and is associated with acute respiratory distress syndrome (ARDS). However, the clinical syndrome resulting from SARS-CoV-2 infection is broad, ranging from asymptomatic infection to severe disease with extrapulmonary manifestations[3], including acute kidney injury[4], acute myocardial injury[5,6] and thrombotic complications[7–11].

Acute kidney injury (AKI) is a particularly prominent complication. The rates of AKI vary greatly based on patient population, but evidence suggests that at least 30% of hospitalized patients and 50% of patients in the intensive care unit (ICU) develop AKI[1,4,12–15]. Although the rate of AKI in hospitalized COVID-19 patients has decreased since the initial surge in 2020, the incidence remains high[16]. Like community-acquired pneumonia[17], AKI is increasingly recognized as a common complication of COVID-19 in the hospitalized setting and confers significantly increased morbidity and mortality[18].

However, unlike AKI secondary to sepsis or other viral organisms, there remains a limited understanding of the pathophysiology of COVID-19-associated AKI. Histopathological reports from autopsy specimens have provided conflicting insights into the pathological changes in the kidney in COVID-19. A report of 26 patients who died with COVID-19 AKI revealed acute tubular injury as a prominent mechanism[19]. Additionally, the presence of viral particles in the tubular epithelium and podocytes in

autopsy specimens has been reported[19,20], which is evidence of direct viral invasion of the kidney. In addition, coagulopathy and endothelial dysfunction are hallmarks of COVID-19[21] and may also contribute to AKI. Finally, SARS-CoV-2 may directly activate the complement system[22]. In addition to these mechanisms, systemic effects of critical illness (hypovolemia, mechanical ventilation) and derangements in cardiac function and volume may also contribute to COVID-19 AKI.

In addition to morbidity and mortality in the acute setting, COVID-19 is also associated with long term manifestations i.e., the post-acute sequelae of SARS-CoV2 (PASC)[23]. Kidney function decline is a major component of PASC and a study of more than 1 million individuals found that survivors of COVID-19 had an elevated risk of post-acute eGFR decline[24], suggesting long term kidney dysfunction may occur following the acute infection.

Given the high incidence of COVID-19 associated kidney dysfunction, the unknown pathophysiology, and the urgent need for better approaches for risk stratification for long term kidney function decline we aimed to characterize the proteomic changes associated with COVID associated AKI and long-term kidney function. We measured protein expression of more than 4000 proteins from serum samples collected in a diverse large cohort of hospitalized patients with COVID-19 and validated significant results in an independent cohort and identified proteins that are significantly different between patients with and without AKI. We then determined whether these proteomic perturbations also characterize post-discharge kidney function decline as measured by estimated glomerular filtration rate (eGFR).

MATERIALS AND METHODS

Patient cohort

An overview of the discovery cohort selection process is provided in **Fig 1**. We prospectively enrolled patients hospitalized with COVID-19 between March 24, and August 26, 2020, at five hospitals of a large urban, academic hospital system in New York City, NY into a cohort as previously described[25]. The cohort enrolled patients who were admitted to the health care system with a COVID-19 infection and had broad inclusion criteria without specific exclusion criteria. The Mount Sinai Institutional Review Board approved this study under a regulatory approval allowing for access to patient level data and biospecimen collection. Peripheral blood specimens were collected at various points during the hospital admission for each patient.

The validation cohort included a prospective biobank from Quebec, Canada that enrolled patients hospitalized with COVID-19, as previously described[26]. Patients were recruited from the Jewish General Hospital and Centre Hospitalier de l'Université de Montréal. Peripheral blood specimens were collected at multiple time points after admission.

We defined an AKI cohort using proteomic data acquired at the last available timepoint during the hospital course for all individuals. Patients who developed AKI after the last specimen collection timepoint were excluded. Controls were defined as individuals who developed AKI stage 1 or did not develop AKI during their hospital course.

Serum collection and Processing

Blood samples were collected in Serum Separation Tubes with a polymer gel for serum separation. Samples were centrifuged at 1200 g for 10 minutes at 20°C. After centrifugation, serum was pipetted to a 15 mL conical tube. Serum was then aliquoted into cryovials and stored at -80°C.

Definition of Acute Kidney Injury

We defined AKI (stage 2 or 3) as per Kidney Disease Improving Global Outcomes (KDIGO) criteria: an increase in serum creatinine of at least 2.0 times the baseline creatinine[27]. For patients with previous serum creatinine measurement available in the 365 days prior to admission, the minimum value in this period was considered the baseline creatinine. For patients without a baseline creatinine in this period, a baseline value was calculated based on an estimated glomerular filtration rate (eGFR) of 75 ml/min per 1.73 m^2 as per the KDIGO AKI guidelines.

Clinical data collection

We collected demographic and laboratory data collected as part of standard medical care from an institutional electronic health record (EHR) database. We defined clinical comorbidities using diagnostic codes recorded in the EHR before the current hospital admission. To account for disease severity at the time of specimen collection, we defined supplemental oxygen requirement as 0 if the patient was not receiving supplemental oxygenation or on nasal cannula, 1 if the patient was receiving non-invasive mechanical ventilation (CPAP, BIPAP), or 2 if the person was receiving invasive mechanical ventilation.

Somalogic proteomic assay

We used the *SomaScan* discovery platform to quantify levels of protein expression. The *SomaScan* platform is a highly multiplexed aptamer based proteomic assay based on modified single-stranded DNA aptamers capable of simultaneously detecting 4497 proteins in biological samples. The assay was run using the standard 12 hybridization normalization control sequences to assess for variability in the Agilent plate quantification process, five human calibrator control pooled replicates, and 3 quality control pooled replicates to control for batch effects. Standard preprocessing as per Somalogic's guidelines were applied. Briefly, the data was first normalized using the 12 hybridization controls to remove hybridization variation within a run. Then, median signal normalization is performed with calibrator samples across plates to remove variation in sample-to-sample differences attributable to variations due to pipetting, reagent concentrations, assay timings and other technical aspects. Data was then calibrated to remove assay differences between runs. Standard Somalogic acceptance criteria for quality control metrics were used (plate scale factor between 0.4 and 2.5 and 85% of QC ratios between 0.8 and 1.2). Samples with intrinsic issues such as reddish appearance or low sample volume were also removed as part of the Somalogic quality control protocol. After quality control and normalization procedures, the resulting relative fluorescence unit (RFU) values were log2 transformed.

Dimensionality reduction

Principal component analysis (PCA) was performed using log2 transformed RFU values of all proteins. Pairwise plots of the top three principal components were plotted.

Differential expression analysis for prevalent AKI

Using data from the AKI cohort, linear regression models were fit independently for each protein using the Limma package in R using the $\log_2$ transformed protein values as the dependent variable. Models were adjusted for age, sex, history of chronic kidney disease (CKD), and supplemental oxygen requirement (0,1, or 2 [see above]) at the time of specimen collection. P values were adjusted using the Benjamin-Hochberg procedure to control the false discovery rate (FDR) at 5%.

Proteomic characterization of long-term kidney function in discovery cohort

Outpatient creatinine values measured after discharge were used to compute estimated glomerular filtration rate (eGFR) values the CKD-EPI equation. To determine whether AKI associated protein expression correlated with post-discharge kidney function, we fit a mixed effects linear regression model with random intercept and slope. Using the discovery cohort, protein expression of AKI-associated proteins measured at the last available timepoint during admission was used. The dependent variable was eGFR and the model was adjusted for age, sex, baseline creatinine, history of CKD, maximum AKI stage during the hospital admission, and day of eGFR measurement after hospital discharge. Models included a random effect of patient ID to adjust for correlation between eGFR values taken from the same individuals. Significance was evaluated using a F-test with Sattherwaite degrees of freedom implemented in the lmerTest[28] R package. P values were adjusted using the Benjamin-Hochberg procedure to control the false discovery rate (FDR) at 5%.

Data analysis and visualization

We performed all statistical analysis using R version 4.0.3. Protein-protein interaction (PPI) network was constructed using the Network X package in Python to display a Minimum Spanning Tree (MST) using Prim's algorithm. Network clustering was conducted using the MCL cluster algorithm and functional enrichment was carried out using the STRING Database in Cytoscape[29].

Data and Code Availability

Data is available by contacting the senior author, Girish Nadkarni (girish.nadkarni@mountsinai.org).

Code is available at https://github.com/Nadkarni-Lab/aki_covid_proteomics

RESULTS
Discovery and Validation Cohort Overview

To discover proteins associated with COVID-AKI, we enrolled a prospective cohort of patients hospitalized with COVID-19 admitted between March 24, 2020 and August 26, 2020 into a biobank as previously described[25]. Cases were defined as patients who developed AKI (stage 2 or 3) during their hospital admission and controls included all other patients (**Fig 1**). Characteristics of cases and controls in the discovery cohort are provided in **Table 1**. Patients who developed AKI were significantly older (67 vs. 60 years, p <0.001), more commonly Hispanic/Latino (48% vs 37%, p =0.012), and had a greater prevalence of atrial fibrillation (17% vs 8%, p =0.008), diabetes (37% vs 20%, p <0.001), and chronic kidney disease (20% vs 4%, p< 0.001). We then validated these associations in an external cohort from Quebec, Canada. Characteristics of the

validation cohort are provided in **Supplementary Table 1.** In the validation cohort, compared to controls, AKI (stage 2 or 3) cases had a significantly higher prevalence of CKD (29% vs 11%, p = 0.01) and a higher rate of intubation at the time of blood draw (49% vs 13%).

Identification of proteins associated with prevalent AKI

In the discovery cohort, serum levels of 4496 proteins were quantified using the *SomaScan* platform using samples collected at multiple timepoints during the hospital course (**Supplementary Table 2**) as previously described[25]. We first identified proteins associated with prevalent AKI using measurements taken after the onset of AKI in cases and the last available measurement in controls (**Fig 1**, 71 cases and 366 controls). The top three principal components (PCs) distinctly separate samples by case status (**Fig 2A**). We fit a multivariate linear regression model adjusted for age, sex, history of chronic kidney disease (CKD), and maximum oxygen requirement at the time of blood draw. We identified 413 upregulated and 30 downregulated proteins (**Supplementary Table 3**).

Validation of AKI-associated proteins

We then performed an external validation of AKI-associated proteins in a prospective biobank cohort from Quebec, Canada. Of the 443 proteins significantly associated with AKI in the discovery cohort, 62 were also associated with AKI in the validation cohort (p <0.05, **Table 2**). All validated proteins were associated with an increased risk of AKI. The fold changes of validated proteins in the discovery and validation cohort were highly correlated with a Pearson correlation of 0.71 (**Fig 2B**). The 62-protein signature distinctly separated AKI cases from cohorts in the discovery cohort

(**Fig 2C**). Protein-protein interaction (PPI) network analysis revealed enrichment of several highly connected proteins, including LCN2 (alternative name: NGAL), REG3A, and MB (**Fig 4A**). The AKI-associated protein network also included a cluster of cardiac structural proteins (**Fig 4B**), TNNT2, TTN, MYL3, SRL, and NPPB (alternative name: BNP).

Proteomic characterization of post-acute kidney dysfunction

Given the previously reported association of COVID-19 AKI with long-term eGFR decline[30], we hypothesized that significant proteomic markers associated with COVID-19 AKI are also associated with post-discharge eGFR. We included all outpatient eGFR measurements taken after discharge from patients in the Mount Sinai biobank cohort. Of the 437 patients in the cohort, 181 patients had at least one outpatient post-discharge eGFR measurement. The median number of eGFR measurements was 4 with an interquartile of 9.

We used a mixed effects linear model accounting for baseline creatinine, AKI stage during the COVID admission and repeated eGFR measurements to associate the 62 protein AKI signature with long-term eGFR trend. Of the 62 AKI-associated proteins, 25 were significantly (adjusted P <0.05) associated with eGFR trend (**Fig 3, Fig 5A**). All 25 eGFR-associated proteins were negatively correlated with eGFR (**Table 3**). However, the strength of association with AKI was not significantly associated with the strength of association with post-discharge eGFR. Proteins most strongly associated (by P value) with decreased post-discharge eGFR included desmocollin-2, trefoil factor 3, transmembrane emp24 domain-containing protein 10, and cystatin-C (**Fig 5B**).

DISCUSSION

Using longitudinal proteomic profiling in two large groups of patients hospitalized with COVID-19, we report several observations. First, we identified specific protein markers of AKI and post-discharge kidney dysfunction, both well-documented sequelae of COVID-19[4,30]. Second, in the acute phase, tubular injury and hemodynamic perturbation seem to be likely mechanisms. Third, tubular injury is a predominant mechanism of long-term kidney function decline. Thus, proteomic characterization reveals potentially different mechanisms of AKI and long-term eGFR decline with implications for more specific prognostic models and targeted therapeutic development.

Our data suggest that in the acute setting, COVID AKI is characterized by two broad mechanisms: 1) tubular injury and 2) hemodynamic perturbation. First, we found significant upregulation of NGAL (LCN2), a canonical marker of tubular injury. NGAL is secreted by circulating neutrophils and kidney tubular epithelium in response to systemic inflammation or ischemia. Since renal tubular epithelial cells express the angiotensin-converting enzyme 2 (ACE2) receptor which enables SARS-CoV2 viral entry into cells, direct tubular infection may cause the release of NGAL into the serum and urine. This potential mechanism is supported by our results and remains a testable hypothesis. Although NGAL is a known marker for intrinsic AKI accompanied by tubular injury, it is relatively insensitive to pre-renal AKI caused by hemodynamic disturbance[31,32]. However, our results demonstrate upregulation of BNP, a protein

released in the setting of volume overload as well as several cardiac structural proteins (cardiac troponin T, titin, myosin light chain 1, and sarcalumenin). This proteomic signature may represent an impaired crosstalk between the cardiovascular system and kidney in which myocardial injury leads to decreased renal perfusion and eventual AKI. Myocardial injury has been previously reported in patients hospitalized with COVID-19[6] and thus may contribute to the multifactorial nature of COVID-AKI.

Since COVID-AKI increases the risk of long-term eGFR decline[30], we then sought to determine whether these two phenomena shared common proteomic markers. Surprisingly, we found that although almost half of the AKI-associated proteins were also significantly associated with post-discharge eGFR decline, the strengths of associations were not correlated. While COVID-AKI is likely caused by a combination of intrinsic tubular injury and hemodynamic disturbance in the setting of critical illness, long term eGFR decline was associated with increased expression of trefoil factor 3 (TFF3), a known prognostic marker for incident CKD[33]. Trefoil factors are a class of small peptides expressed in colonic and urinary tract epithelia that play essential roles in regeneration and repair of epithelial tissue[34,35]. Immunohistochemistry reveals TFF3 expression is localized to the tubular epithelial cells in kidney specimens from patients with CKD[33], suggesting that long term eGFR decline may be associated with renal tubular epithelial damage. The exact pathological role of TFF3 in the renal tubules is unclear but it has been hypothesized to play a role in repair of kidney damage[36]. Additionally, TFF3 release from the renal interstitium has also been hypothesized to direct the epithelial-to-mesenchymal transition (EMT) in renal interstitial fibrosis, a main pathway that leads to ESKD[33]. Our results implicate tubular damage in both AKI and

long term eGFR decline suggesting that SARS-CoV2 may preferentially target this region of the nephron. While AKI in the acute setting may be a result of ischemia and decreased renal perfusion associated with critical illness, the specific elevation of TFF3 associated with eGFR decline implicates a more general pattern of tubular injury that underlies COVID mediated kidney dysfunction. Since the ACE2, is preferentially expressed in the tubular epithelial cells of the kidney[37,38], the elevation of markers of tubular damage in the plasma may represent direct viral invasion of tubular epithelia cells. However, again this would need to be tested using biopsy/autopsy specimens or other mechanistic studies

Our study should be interpreted in the context of certain limitations. First, samples were collected during the hospital course of patients with confirmed COVID-19. However, the timepoints were not systematic due to logistical challenges during the peak of the COVID-19 pandemic and thus are not standardized between patients. Since a subset of patients had AKI at the time of admission, these patients were excluded from our analysis since specimens were collected after admission. Thus, our AKI cases may be biased towards less severe presentations. Second, since kidney injury is usually not an isolated phenomenon in critically ill patients, the protein expression changes observed may have been partially due to damage to other organs, such as the lung, liver, and heart. However, we accounted for non-kidney damage by adjusting for the highest level of ventilatory support and thus our results are likely a reflection of kidney injury. However, our results do show the importance of crosstalk between the cardiac system and the kidneys. Finally, our cohort did not include autopsy or kidney biopsy

specimens. Histopathological analysis of kidney specimens is necessary to determine the mechanism of AKI and whether viral particles are present in the kidney.

In conclusion, we provide the first comprehensive characterization of the plasma proteome of AKI and long term eGFR decline in hospitalized COVID-19 patients. Our results suggest that COVID-AKI and post-discharge kidney dysfunction may be caused by tubular damage but that in the acute setting, several factors including hemodynamic disturbance and myocardial injury also play a role.

Author contributions:

Conceptualization: IP, GNN, AWC, SZ, NB, ES, EK, LC, SC, BM, SD
Methodology: IP, RT, KC, SA, AV, SJ, MP, RO, SK, SZ, EUA, PJ, FFG
Investigation: IP, RT, EK, SZ, SJ, KC, AK, SG, HX, JH, MP, KA, CB, KN, PJ, RO
Visualization: IP, PJ
Funding acquisition: GNN, AWC, BSG, MM, NB
Supervision: GNN, AWC, BSG, MM, SK, NB, ES, JCH
Writing: IP, GNN, SK

Disclosures: GNN and SGC reports grants, personal fees, and non-financial support from Renalytix. GNN reports non-financial support from Pensieve Health, personal fees from AstraZeneca, personal fees from BioVie, personal fees from GLG Consulting, personal fees from Siemens Healthineers from outside the submitted work.

Funding: EUA, GNN, JCH and SGC are partially funded by R01 DK118222. GNN also

is supported by R01DK127139, R01HL155915 and R56DK126930.

Figure Captions

Fig 1: Overview of the discovery cohort selection process.

Fig 2: A. Top 3 Principal Components show separation of the sample by AKI (stage 2 or 3) case status. **B.** External validation of AKI associated proteins in the discovery cohort shows high correlation with increased risk of AKI with significance of $p < 0.05$. **C.** Expression heatmap shows a distinct separation of the cases and controls using the 62 significant proteins identified from the validation cohort in the discovery cohort.

Fig 3: Nested Venn diagram of the analyses performed.

Fig 4: A. Protein–protein interaction (PPI) network (Minimum Spanning Tree) of the 62 overlapping AKI associated proteins with a score >0.4. The size of each node corresponds to number of interactions and the thickness of the edges represent the weight of the interactions between the nodes. B. MCL algorithm was used to identify tightly connected cluster of proteins which was functionally enriched for cardiac structure proteins using the STRING database.

Fig 5: Proteomic characterization of long-term eGFR decline. A. Comparison of strengths of association with AKI and long term eGFR for proteins associated with AKI in both the discovery and validation cohorts. B. Trend in eGFR values separated by protein expression for tertiles for proteins most significantly (by P value) associated with eGFR trend.

Tables

Table 1: Demographic and clinical comorbidities of patients hospitalized with COVID-19 in the discovery cohort separated by the development of AKI (stage 2 or 3) during their hospital course.

Characteristic	Developed AKI (stage 2 or 3) during hospitalization (N = 71)	AKI stage 1 or no AKI during hospitalization (N = 366)	P Value
Age, mean (SD)	63 (14.7)	61.8 (18.2)	0.80
Male, n (%)	46 (65%)	218 (60%)	0.43
Race, n (%)			0.74
Caucasian	23 (32%)	136 (37%)	
Black	17 (24%)	81 (22%)	
Other	31 (44%)	149 (41%)	
Ethnicity, n (%)			0.23
Hispanic	34 (48%)	144 (39%)	
Not Hispanic or Latino	37 (52%)	213 (58%)	
Unknown / Not Reported	0 (0%)	9 (2%)	
Comorbidities, n (%)			
Atrial Fibrillation	9 (13%)	38 (10%)	0.53
Coronary Artery Disease	10 (14%)	50 (14%)	1
Arterial Hypertension	30 (42%)	142 (39%)	0.60
Diabetes	30 (42%)	81 (22%)	<0.001
Chronic Kidney Disease	22 (31%)	20 (5%)	<0.001
Highest respiratory support, n (%)			<0.001
Intubation	33 (46%)	39 (11%)	
Non-invasive ventilation (CPAP, BIPAP, high-flow cannula)	3 (4%)	19 (5%)	
Nasal cannula	21 (30%)	184 (50%)	
None of the above	14 (20%)	124 (34%)	

Table 2: Validation of AKI-associated proteins. Strength of association of proteins significantly associated with AKI (stage 2 or 3) in both the discovery (adjusted P <0.05) and validation cohort (P <0.05) are provided. Significance was determined by fitting a

linear model adjusted for age, sex, history of CKD, and maximum oxygen requirement at the time of blood draw.

Table 3: Association of AKI-associated proteins with post-discharge eGFR. A mixed effects linear model adjusted for age, sex, history of CKD, baseline creatinine, AKI stage during the COVID admission and number repeated eGFR measurements was fit. Proteins significantly associated with eGFR (adjusted P <0.05) are provided.

Gene name	Coefficient	P Value	Adjusted P Value	Uniprot ID
BolA-like protein 1	-18.022031	1.38E-05	0.00086933	Q9Y3E2
Cystatin-C	-17.63355	3.40E-07	2.14E-05	P01034
Desmocollin-2	-16.597692	1.63E-09	1.03E-07	Q02487
Transmembrane emp24 domain-containing protein 10	-15.819807	1.36E-08	8.54E-07	P49755
Ephrin-B1	-15.167647	0.00020447	0.01288142	P98172
Serine/threonine-protein kinase receptor R3	-13.939695	7.64E-05	0.00481422	P37023
Ganglioside GM2 activator	-13.060806	1.17E-05	0.00073632	P17900
Matrix-remodeling-associated protein 7	-11.88046	0.0002809	0.01769682	P84157
PILR alpha-associated neural protein	-11.803896	3.23E-05	0.00203468	Q8IYJ0
Trefoil factor 3	-11.738116	1.60E-08	1.01E-06	Q07654
Copper transport protein ATOX1	-11.216693	4.38E-06	0.00027586	O00244
Serine protease inhibitor Kazal-type 7	-11.157036	5.77E-06	0.00036339	P58062
IGF-like family receptor 1	-11.007485	4.79E-07	3.02E-05	Q9H665
Complexin-2	-10.73936	1.33E-06	8.36E-05	Q6PUV4
Collagen alpha-1(XXVIII) chain	-10.047217	0.00010393	0.00654748	Q2UY09
CD59 glycoprotein	-9.9917438	1.74E-05	0.00109912	P13987
Ribonuclease K6	-9.9898356	0.00010554	0.00664925	Q93091

Transgelin	-9.9393974	0.00015956	0.01005208	Q01995
DnaJ homolog subfamily B member 12	-8.2638876	0.00049523	0.03119973	Q9NXW2
Protein delta homolog 2	-7.7406286	0.00029449	0.01855277	Q6UY11
Discoidin domain-containing receptor 2	-7.6524963	0.00067262	0.04237511	Q16832
Ribonuclease pancreatic	-7.40997	8.01E-07	5.05E-05	P07998
Ephrin-B2	-7.2943084	0.00015596	0.00982541	P52799
Peroxidasin homolog	-7.2282085	1.75E-05	0.00109962	Q92626
Trefoil factor 2	-7.2179491	1.20E-05	0.0007563	Q03403

References.

1. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting Characteristics, Comorbidities, and Outcomes among 5700 Patients Hospitalized with COVID-19 in the New York City Area. JAMA - J Am Med Assoc 2020;

2. Goyal P, Choi JJ, Pinheiro LC, et al. Clinical characteristics of COVID-19 in New York City. N. Engl. J. Med. 2020;

3. Gupta A, Madhavan M V., Sehgal K, et al. Extrapulmonary manifestations of COVID-19. Nat. Med. 2020;

4. Chan L, Chaudhary K, Saha A, et al. AKI in Hospitalized Patients with COVID-19. J Am Soc Nephrol [Internet] 2020;ASN.2020050615. Available from: http://jasn.asnjournals.org/content/early/2020/09/02/ASN.2020050615.abstract

5. Sala S, Peretto G, Gramegna M, et al. Acute myocarditis presenting as a reverse Tako-Tsubo syndrome in a patient with SARS-CoV-2 respiratory infection. Eur. Heart J. 2020;

6. Lala A, Johnson KW, Januzzi JL, et al. Prevalence and Impact of Myocardial Injury in Patients Hospitalized With COVID-19 Infection. J Am Coll Cardiol 2020;76(5):533–46.

7. Cui S, Chen S, Li X, Liu S, Wang F. Prevalence of venous thromboembolism in patients with severe novel coronavirus pneumonia. J Thromb Haemost 2020;

8. Bilaloglu S, Aphinyanaphongs Y, Jones S, Iturrate E, Hochman J, Berger JS. Thrombosis in Hospitalized Patients with COVID-19 in a New York City Health System. JAMA - J. Am. Med. Assoc. 2020;

9. Lodigiani C, Iapichino G, Carenzo L, et al. Venous and arterial thromboembolic

complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. Thromb Res 2020;

10. Helms J, Tacquard C, Severac F, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. Intensive Care Med 2020;

11. Paranjpe I, Fuster V, Lala A, et al. Association of Treatment Dose Anticoagulation With In-Hospital Survival Among Hospitalized Patients With COVID-19. J. Am. Coll. Cardiol. 2020;

12. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet 2020;

13. Argenzian MG, Bruc SL, Slate CL, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: Retrospective case series. BMJ 2020;

14. Mohamed MMB, Lukitsch I, Torres-Ortiz AE, et al. Acute Kidney Injury Associated with Coronavirus Disease 2019 in Urban New Orleans. Kidney360 2020;

15. Cao J, Tu WJ, Cheng W, et al. Clinical features and short-term outcomes of 102 patients with coronavirus disease 2019 in Wuhan, China. Clin Infect Dis 2020;

16. Dellepiane S, Vaid A, Jaladanki SK, et al. Acute Kidney Injury in Patients Hospitalized With COVID-19 in New York City: Temporal Trends From March 2020 to April 2021. Kidney Med 2021;

17. Murugan R, Karajala-Subramanyam V, Lee M, et al. Acute kidney injury in non-severe pneumonia is associated with an increased immune response and lower

survival. Kidney Int 2010;

18. Nadim MK, Forni LG, Mehta RL, et al. COVID-19-associated acute kidney injury: consensus report of the 25th Acute Disease Quality Initiative (ADQI) Workgroup. Nat. Rev. Nephrol. 2020;

19. Su H, Yang M, Wan C, et al. Renal histopathological analysis of 26 postmortem findings of patients with COVID-19 in China. Kidney Int 2020;

20. Farkash EA, Wilson AM, Jentzen JM. Ultrastructural evidence for direct renal infection with sars-cov-2. J Am Soc Nephrol 2020;

21. Klok FA, Kruip MJHA, van der Meer NJM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. Thromb Res 2020;

22. Noris M, Benigni A, Remuzzi G. The case of complement activation in COVID-19 multiorgan impact. Kidney Int. 2020;

23. Al-Aly Z, Xie Y, Bowe B. High-dimensional characterization of post-acute sequelae of COVID-19. Nature 2021;

24. Bowe B, Xie Y, Xu E, Al-Aly Z. Kidney Outcomes in Long COVID. J Am Soc Nephrol 2021;

25. Charney AW, Simons NW, Mouskas K, et al. Sampling the host response to SARS-CoV-2 in hospitals under siege. Nat. Med. 2020;

26. Su C-Y, Zhou S, Gonzalez-Kozlova E, et al. Circulating proteins to predict adverse COVID-19 outcomes. medRxiv [Internet] 2021;2021.10.04.21264015. Available from:

http://medrxiv.org/content/early/2021/10/05/2021.10.04.21264015.abstract

27. Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. Nephron -

Clin. Pract. 2012;

28. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed Effects Models . J Stat Softw 2017;

29. Doncheva NT, Morris JH, Gorodkin J, Jensen LJ. Cytoscape StringApp: Network Analysis and Visualization of Proteomics Data. J Proteome Res 2019;

30. Nugent J, Aklilu A, Yamamoto Y, et al. Assessment of Acute Kidney Injury and Longitudinal Kidney Function after Hospital Discharge among Patients with and without COVID-19. JAMA Netw Open 2021;

31. Basu RK, Wong HR, Krawczeski CD, et al. Combining functional and tubular damage biomarkers improves diagnostic precision for acute kidney injury after cardiac surgery. J Am Coll Cardiol 2014;

32. Nickolas TL, Schmidt-Ott KM, Canetta P, et al. Diagnostic and prognostic stratification in the emergency department using urinary biomarkers of nephron damage: A multicenter prospective cohort study. J Am Coll Cardiol 2012;

33. Du TY, Luo HM, Qin HC, et al. Circulating serum trefoil factor 3 (TFF3) is dramatically increased in chronic kidney disease. PLoS One 2013;

34. Madsen J, Nielsen O, Tornøe I, Thim L, Holmskov U. Tissue localization of human trefoil factors 1, 2, and 3. J Histochem Cytochem 2007;

35. Kjellev S. The trefoil factor family - Small peptides with multiple functionalities. Cell. Mol. Life Sci. 2009;

36. Astor BC, Köttgen A, Hwang SJ, Bhavsar N, Fox CS, Coresh J. Trefoil factor 3 predicts incident chronic kidney disease: A case-control study nested within the Atherosclerosis Risk in Communities (ARIC) study. Am J Nephrol 2011;

37. Chen Z, Hu J, Liu L, et al. SARS-CoV-2 Causes Acute Kidney Injury by Directly Infecting Renal Tubules. Front Cell Dev Biol 2021;

38. Lin W, Fan J, Hu LF, et al. Single-cell analysis of angiotensin-converting enzyme II expression in human kidneys and bladders reveals a potential route of 2019 novel coronavirus infection. Chin Med J (Engl) 2021;

Fig 1

Fig 2

A

Principal component analysis of protein expression in discovery cohort

PC1,
12.62%

PC2,
11.22%

■ AKI stage 2 or 3

■ AKI stage 1 or no AKI

PC3,
6.56%

PC4,
4.37%

B

Validation of AKI associated proteins

C

Expression of 62 protein AKI signature in discovery cohort

■ AKI stage 2 or 3

■ AKI stage 1 or no AKI

Fig 3

Results Overview

443 proteins associated with AKI stage 2 or 3 in discovery cohort (adjusted P <0.05)

62 proteins associated with AKI stage 2 or 3 in validation cohort (P <0.05)

25 proteins associated with post discharge eGFR trend (adjusted P <0.05)

Fig 4

AKI associated protein network

Cardiac structure protein network

Fig 5

Proteomic characterization of post-discharge kidney function

A

B Protein expression ▮ Top tertile ▮ Middle tertile ▮ Bottom tertile