Deficiency of U4atac snRNA secondarily results in ciliary defects

Deepak Khatri,[1,16] Audrey Putoux,[1,2,16] Audric Cologne,[1,3] Sophie Kaltenbach,[4] Alicia Besson,[1] Eloïse Bertiaux,[5] Justine Guguin,[1] Adèle Fendler,[1] Marie A. Dupont,[6] Clara Benoit-Pilven,[1,3] Sarah Grotto,[7] Lyse Ruaud,[8,9] Caroline Michot,[10,11] Martin Castelle,[12] Agnès Guët,[13] Laurent Guibaud,[14] Virginie Hamel,[5] Rémy Bordonné,[15] Anne-Louise Leutenegger,[8] Tania Attié-Bitach,[4,11] Patrick Edery,[1,2] Sylvie Mazoyer,[1,17] and Marion Delous[1,17]*

[1]Genetics of Neurodevelopment Team, Lyon Neuroscience Research Centre, UMR5292 CNRS, U1028 Inserm, University of Claude Bernard Lyon 1, Bron, France.

[2]Clinical Genetics Unit, Department of Genetics, Centre de Référence Anomalies du Développement et Syndromes Polymalformatifs, Hospices Civils de Lyon, University of Claude Bernard Lyon 1, Bron, France.

[3]INRIA Erable, LBBE, UMR5558 CNRS, University of Claude Bernard Lyon 1, Villeurbanne, France.

[4]Department of Histology Embryology and Cytogenetics, APHP, Necker-Enfants Malades Hospital, University of Paris, Paris, France.

[5]Department of Cell Biology, Sciences III, University of Geneva, Geneva, Switzerland.

[6]Laboratory of hereditary kidney diseases, Imagine Institute, UMR1163 Inserm, University of Paris, Paris, France.

[7]Clinical Genetics Unit, Maternité Port-Royal, APHP, Cochin Broca Hôtel-Dieu Hospitals, Paris, France.

[8]NeuroDiderot, UMR1141 Inserm, Université de Paris, Paris, France.

[9]Departement of Genetics, APHP, Robert Debré Hospital, Paris, France.

[10]Clinical Genetics Department, Centre de Référence Maladies Rares - Maladies Osseuses Constitutionnelles, APHP, Necker-Enfants Malades Hospital, Paris, France.

[11]Imagine Institute, UMR1163 Inserm, University of Paris, Paris, France.

[12]Hematology-Immunology Unit, APHP, Necker-Enfants Malades Hospital, Paris, France.

[13]Neonatal and Pediatric Units, Louis-Mourier Hospital, Colombes, France

[14]Pediatric and Fetal Imaging, Hospices Civils de Lyon, University of Claude Bernard Lyon 1, Bron, France

[15]Institute of Molecular Genetics of Montpellier, UMR5535 CNRS, University of Montpellier, Montpellier, France

[16]DK and AP share the first author position.

[17]SM and MD share the last author position.

*Address correspondence to: Marion Delous, Institut IDEE, 59 Bd Pinel, 69500 BRON, France. Phone: (+33) 4 72 13 89 69; Email: marion.delous@inserm.fr

Abstract

In the human genome, about 700 genes contain usually one intron excised by the minor spliceosome. This spliceosome comprises its own set of snRNAs, among which U4atac. Its non-coding gene, *RNU4ATAC*, has been found mutated in Taybi-Linder (MOPD1/TALS), Roifman (RFMN) and Lowry-Wood syndromes (LWS). These rare developmental disorders, whose physiopathological mechanisms remain unsolved, associate ante- and post-natal growth retardation, microcephaly, skeletal dysplasia, intellectual disability, retinal dystrophy and immunodeficiency. Here, we report a homozygous *RNU4ATAC* mutation in the Stem II domain, n.16G>A, in two unrelated patients presenting with both typical traits of the Joubert syndrome (JBTS), a well-characterized ciliopathy, and of TALS/RFMN/LWS, thus widening the clinical spectrum of *RNU4ATAC*-associated disorders and indicating ciliary dysfunction as a mechanism downstream of minor splicing defects. This finding is supported by alterations of primary cilium function in TALS and JBTS/RFMN fibroblasts, as well as by *u4atac* zebrafish model, which exhibit ciliopathy-related phenotypes and ciliary defects. Altogether, our data indicate that alteration of cilium biogenesis is part of the physiopathological mechanisms of TALS/RFMN/LWS, secondarily to defects of minor intron splicing.

Introduction

Pathogenic variants of the *RNU4ATAC* gene, transcribed into the U4atac small non-coding RNA involved in minor splicing, are associated with three rare autosomal recessive syndromes: Taybi-Linder (TALS, also named microcephalic osteodysplastic primordial dwarfism type 1, MOPD1) (Edery *et al*, 2011; He *et al*, 2011), Roifman (RFMN) (Merico *et al*, 2015) and Lowry-Wood (LWS) (Farach *et al*, 2017) syndromes. TALS, RFMN and LWS patients are characterized by distinct patterns of skeletal dysplasia and different degrees of severity of ante- and post-natal growth retardation, microcephaly, intellectual disability, retinal dystrophy and immunodeficiency, as well as variable accompanying traits including syndrome-specific facial dysmorphism, dermatitis, hepatomegaly (RFMN), endocrine abnormalities and reduced survival (TALS) (Hagiwara *et al*, 2021; Putoux *et al*, 2016). The most serious cases, i.e. TALS patients, display severe cerebral malformations including abnormal gyral pattern, corpus callosum abnormalities and hypoplasia of cerebellar vermis. For those who carry a specific recurrent *RNU4ATAC* mutation (n.51G>A), death usually occurs within the two first years of life, rapidly after an apparently trivial infectious episode, for yet unknown reasons. The U4atac snRNA is a component of the minor spliceosome involved in the splicing of 935 so-called minor or U12-type introns found in 748 genes, in an up-to-date analysis of the human genome, recognized through their consensus splice site sequences (Turunen *et al*, 2013). U4atac/U6atac shares with U4/U6, despite sequence differences, a common secondary structure consisting of intra- and inter-molecular base-pairing. Some of the resulting domains such as Stem II and 5' Stem-Loop bind essential splicing proteins. Most TALS patients carry homozygous or compound heterozygous mutations in the 5' Stem-Loop, while all RFMN patients carry one of the two variants in Stem II, never found mutated in TALS (Benoit-Pilven *et al*, 2020). The physiopathological links between *RNU4ATAC* variants and TALS/RFMN/LWS symptoms remain largely unknown. Although it has been shown that splicing defects affect most U12-containing genes (Dinur Schejter *et al*, 2017; Merico *et al*, 2015; Heremans *et al*, 2018) to an extent that depends on the cell type (Cologne *et al*, 2019), the resulting molecular perturbations causing the

developmental anomalies have not been investigated yet, due to the lack of animal models and biological hypotheses.

In the present work, we show for the first time that a homozygous pathogenic variant of *RNU4ATAC* causes a disorder with overlapping traits with Joubert syndrome (JBTS), a well-known ciliopathy characterized by developmental delay, dysregulation of the breathing pattern, hypoplasia of the cerebellar vermis and a distinctive neuroradiologic "molar tooth sign" (MTS). We thus describe an unsuspected link between minor splicing deficiency and cilium dysfunction in humans and zebrafish. The *RNU4ATAC*-associated pathologies hence join the newly proposed group of "disorders with ciliary contribution".

Results

Identification of a homozygous RNU4ATAC *mutation in patients with JBTS traits*

Two unrelated children suspected with having JBTS were referred to the genetic consultation. A multi-panel targeted exome sequencing was performed, but no bi-allelic pathogenic variants were found in known JBTS-related genes or other ciliary protein-encoding genes of the ciliome panels (Failler *et al*, 2014; Thomas *et al*, 2012). After modification of several bioinformatic parameters and the removal of a filter excluding non-coding genes, we identified a homozygous pathogenic variant, n.16G>A, in *RNU4ATAC* (GenBank: NR_023343) (Fig. 1A, Table S1). This variant was previously identified either at the compound heterozygous (n=4) (Merico *et al*, 2015) or homozygous (n=2) (Dinur Schejter *et al*, 2017; Heremans *et al*, 2018) state in patients with RFMN syndrome, for whom no brain investigation was reported. Patient 1 (F1:II-2, P1) exhibited a severe developmental delay with severe axial hypotonia, microcephaly and skeletal dysplasia, associated with facial dysmorphism, retinal coloboma, nystagmus, eczema, bilateral postaxial polydactyly, genital anomalies, growth hormone deficiency, small interventricular septal defect and immunodeficiency. Brain MRI showed thin but complete corpus callosum, vermis hypoplasia, a MTS and a small posterior arachnoid cyst below the left internal vein (Table 1, Fig. 1B). Patient 2 (F2:II-3, P2), who died before two weeks of age, showed microcephaly and skeletal dysplasia, associated with an atrial septal defect, hand and feet abnormalities and ocular nystagmus (and strabismus). Brain MRI showed a short and dysgenetic vermis with enlarged retro- and infra-cerebellar cisterns, and a MTS (Table 1, Fig. 1C).

Close examination of all clinical features showed that, in addition to the cerebellar and brainstem anomalies suggestive of JBTS, both patients presented with other ciliopathy traits, i.e. polydactyly and cardiac septal defects, as well as typical features of the RFMN/TALS spectrum, i.e. microcephaly, growth retardation, skeletal dysplasia and immunodeficiency (Table 1). Thus, we concluded that these patients exhibited phenotypic traits overlapping both ciliopathy and *RNU4ATAC*-disease spectra, suggesting that *RNU4ATAC* loss of function secondarily results in ciliary defects.

MTS and ciliopathy traits can be found in other RNU4ATAC-*mutated patients*

This finding led us to conduct a thorough review of all *RNU4ATAC*-mutated published cases. A MTS had been noted in a TALS patient carrying the most frequent *RNU4ATAC* n.51G>A mutation at the homozygous state (Abdel-Salam *et al*, 2013), even though the posterior interpeduncular fossa was not as deep as in the classic MTS (G. Abdel-Salam, personal communication). Also, other ciliopathy-related traits have been observed in TALS/RFMN/LWS patients, notably retinitis pigmentosa, a cardinal feature of RFMN, and less frequently, polydactyly (TALS8 case) (Edery *et al*, 2011) and cystic kidney (TALS10 case) (Edery *et al*, 2011). Atrial or ventricular septal defects, frequently observed in TALS/RFMN/LWS patients, are also a trait commonly found in ciliopathies (Burnicka-Turek *et al*, 2016), although not as specific as other phenotypes. Therefore, the presence of ciliopathy traits in other *RNU4ATAC* patients suggests that alteration of cilium structure/function is a possible mechanism downstream any mutations of *RNU4ATAC*, and is not restricted to n.16G>A pathogenic variant.

Cilium-related genes are over-represented among human U12 genes

To better apprehend the link between minor splicing and cilium function, we conducted a gene ontology (GO) enrichment analysis of U12 genes. Relatively few studies addressed the functions of U12 genes in recent years, since their full identification has been facilitated both by the completion of the human genome sequence and the development of computational predictive tools. To the best of our knowledge, only three GO enrichment analyses have been performed, one based on functional gene-sets and mouse phenotypes from the Mammalian Phenotype Ontology resource (Merico *et al*, 2015), one on the most differentially expressed U12 genes in adult human tissues (Olthof *et al*, 2019) and the last on a subset of U12 genes associated with a disease (Olthof *et al*, 2020). Here, we used a different approach and performed a GO term enrichment analysis of the U12 genes *versus* the U2 genes, using the entire set of genes with GO annotations (n=618 and 19,939 for U12 and U2 genes, respectively). An enrichment is observed if more than 3% (618/20,557) of U12 genes are present in a GO term. Hence, we found 243 enriched biological processes and 96

enriched cellular components with a p-value ≤0.05 (Fig. 2A, Table S2, S3). The most enriched biological process is the *membrane depolarization during action potential* (p-value = 5.2e-15, with 40% of U12 genes), which partially overlaps with the other GO terms *neuronal action potential* (p=5.5e-13, 40%), *regulation of ion transmembrane transport* (p=1.2e-12, 8%), *calcium ion transmembrane transport* (p=4.3e-6, 9%) and *sodium ion transmembrane transport* (p=5.2e-6, 10%). Interestingly, the *non-motile cilium assembly* process (p=6.8e-6, 18%) was the thirteenth most significant term (red in Fig. 2A, Table S2). Other significant terms explicitly related to cilia were identified, namely *intraciliary transport involved in cilium assembly* (p=1.2e-3, 15%), *ciliary basal body-plasma membrane docking* (p=1.6e-2, 7%) and *protein localization to ciliary transition zone* (p=2.8e-2, 22%). Similarly, GO term enrichment analysis of cellular components revealed an unexpected amount of U12 gene-encoding proteins that localize at the basal body (p=1.4e-5, 11%), the ciliary base (p=4.3e-3, 14%) and the ciliary tip (p=1.0e-2, 11%) (Table S3). Of note, the opposite comparison of U2 *vs* U12 genes did not reveal any cilium-related terms (Table S4).

We then examined cilium-related genes, bearing in mind that 2.0% of the genes are of the U12-type category in the human genome (748 U12 vs 37,173 U2 genes). Since we observed 7.6% (23 U12 genes vs 279 U2 genes, Table S5) of U12 genes among the curated ciliary "Gold_Standard" genes (Van Dam *et al*, 2013), we concluded that there is an enrichment of U12 genes among them (χ^2 p-value = 1.6e-11). By broadening this list through the compilation of several databases (1426 genes in total, see Materials and Methods), we identified 86 U12 genes (86/1426, 6%) within this in-house ciliome database (Fig. 2B, Table S5). Finally, among the 329 ciliopathy-related genes (Lovera & Lüders, 2021; Reiter & Leroux, 2017; Wheway *et al*, 2019) (Table S5), 23 (~7%) are U12 genes, which is more than the 2.0% expected by chance (χ^2 p-value = 4.1e-10). Strikingly, nearly a third of them are associated with JBTS (*KIAA0556, CEP41, TCTN3, ARMC9, PDE6D, TMEM107, TCTN1, TMEM231*) (Fig. 2B, Table S5), corresponding to 21.6% (8/37) of all known JBTS-associated genes.

Altogether, these enrichment analyses indicate that U12-intron mis-splicing is likely to impair the expression of many genes crucial for cilium formation and/or function, thus suggesting an explanation for the ciliary contribution of *RNU4ATAC*-associated disorders.

JBTS/RFMN RNU4ATAC *mutation results in U12 intron retentions*

We recently published an in-depth analysis of transcriptomes of three different cell types (fibroblasts, amniocytes, lymphocytes) from seven TALS patients, and showed a specific and global alteration of minor splicing (U12 intron retentions (IR) mostly) while gene expression profiles were unchanged (Cologne *et al*, 2019). We carried out a similar RNA-seq experiment on fibroblasts derived from the skin biopsy of patient P1 (Table S6), and observed, like previously in TALS samples, about the same amount of U12 expressed genes (~450), of which a small proportion (10%) showed differences of percent-spliced-in (PSI) values of the U12 intron greater than 10% compared to the control (Fig. 3A). Although biological replicates (i.e. fibroblasts issued from other patients with n.16G>A mutation) would be needed to draw conclusions, the levels of U12 intron retentions, globally weak (maximum deltaPSI of ~30%), are reminiscent of those seen in TALS samples, with most U12 genes (~60%) having a dPSI<10% (Cologne *et al*, 2019). We further performed qRT-PCR analyses to specifically analyse the U12 intron retention of some ciliary genes with known functions (*RABL2A, TMEM107, TMEM231, TCTN1, IFT80*, see Fig. 2B) and observed, for all of them, a higher intron retention level in patients' cells than in controls, with a stronger effect seen in P1 compared to a TALS patient's cells (Fig. 3B). The fact that qRT-PCR evidence stronger IR than RNA-seq, particularly for *RABL2A* and *TMEM107*, may be explained by the low coverage of some U12 intron boundaries by NGS, at least for *RABL2A*.

Altogether, these results confirmed that the n.16G>A mutation, similarly to other *RNU4ATAC* mutations associated with TALS, causes U12 intron retention in fibroblasts of patients, including in U12 ciliary genes.

Primary cilium function is altered in RNU4ATAC-*associated TALS and JBTS/RFMN fibroblasts*

To investigate whether intron retention in the cilium-related genes had an impact on ciliogenesis, we analysed cilium structure in patients' fibroblasts, a well described and favored ciliated cell model. This analysis was conducted in cell lines issued from skin biopsy of JBTS/RFMN patient P1, and from three patients carrying the n.51G>A homozygous mutation (Patients TALS2, TALS4, TALS6 described in Edery *et al*, 2011). Analysis of the percentage of ciliated cells and cilium length, based on immunostainings of cilia with an antibody anti-Arl13b, revealed slight ciliogenesis alterations in two (TALS4, TALS6) out of four patients' cell lines (Fig. S1). To evaluate cilium function, we explored the ciliary localisation of the protein adenylate cyclase III (AC3), which plays a crucial role in signal transduction by associating with G-protein coupled receptors (GPCR) and producing the second messenger cAMP upon their activation. We observed that in TALS6 and P1 fibroblasts, there is a dramatic loss of AC3 into the cilium compared to control and TALS4 cell lines. This result could be explained by the fact that six U12 genes encode proteins of the transition zone, which plays an essential role as a filter barrier selecting ciliary proteins going in or out of the cilium.

Hence, we conclude that in fibroblasts, loss of U4atac function impairs primary cilium function rather than ciliogenesis, perhaps as a result of global low level of U12 intron retentions.

Deficiency of u4atac in zebrafish leads to ciliary defects

To evaluate the importance of minor splicing during development, we turned to the zebrafish, a widely used animal model to study genetic diseases, including ciliopathies (Song *et al*, 2016). Its genome includes 710 U12 introns present in 655 genes in the latest genome assembly (GRCz11), as revealed by a computational scan we performed using U12 intron scoring matrices (Alioto, 2007). These figures are in the same range as those found in humans, and most of the U12 zebrafish genes (544/655) are orthologs of human genes. Zebrafish has two orthologs for *RNU4ATAC*, and by combining the analysis of sequence and secondary structure conservation, expression throughout development and phenotype of CRISPR/Cas9-mediated knock-out lines (see Supplementary Information, Fig. S2, S3), we

concluded that *rnu4atac* on chromosome 11 was the sole functional ortholog of human *RNU4ATAC*.

Pathogenic variants in humans are hypomorphic and full deletion of rnu4atac_chr11 in zebrafish precluded the analysis of phenotypes beyond 22 hpf due to growth arrest and lethality. We therefore opted for a morpholino-based approach and designed two morpholino oligonucleotides (MO) that targeted two distinct functional domains of u4atac, the 5' Stem Loop and Stem II (Fig. S4A). Both MO led to a wide range of developmental anomalies whose severity was dose-dependent and concomitant with an increasing level of U12 intron splicing deficiency (Fig. 4A-G, Fig. S4B-E). Among the most penetrant phenotypes (>50% of MO-injected embryos), we observed body axis curvature, pronephric cysts and otolith defects as well as cardiac dysfunction (Supplementary Information, Movies S1 and S2) and absence of touch response. To a lesser extent (<30%), u4atac morphants also exhibited microcephaly and brain hemorrhages (Fig. 4A-G, Fig. S4D). All these phenotypes were rescued by the co-injection of human WT U4atac snRNA (Fig. 4G, Fig. S4D), in accordance with the highly similar bi-dimensional predictions of human U4atac/U6atac, zebrafish u4atac/u6atac and human U4atac/zebrafish u6atac duplexes (Fig. S2D). Strikingly, the three most penetrant phenotypes (i.e. body axis curvature, pronephric cysts and otolith defects) are hallmarks of cilium dysfunction in zebrafish, thus strengthening the link between u4atac, minor splicing and cilium function. For further evidence, we analysed cilium structure in the central canal, pronephros and otic vesicle. Cilium dysfunction in the central canal results in cerebrospinal fluid flow alteration and Reissner's fiber mis-aggregation, shown to be the origin of the ventral body curvature phenotype (Cantaut-Belarif *et al*, 2018). In the pronephros, motile cilia, and notably those of multiciliated cells, allow fluid flow in renal tubules; their impairment leads to fluid accumulation, tubule luminal expansion and cyst formation in glomeruli (Kramer-Zucker *et al*, 2005). In the otic vesicle, motile cilia participate in otolith formation, favoring the binding of otolith precursor particles to the tip of kinocilia (Wu *et al*, 2011). We analysed motile cilia in the central canal by performing confocal live imaging of *Tg(bactin:arl13bGFP)* transgenic line. We observed a drastic decrease of cilium number in morphants (Fig. 4H), associated with defective cilium beating (Fig. 4H, I). In

pronephros, we observed tubule dilations, mostly of the proximal straight tubule, that were accompanied by a lack of multiciliated tufts (Fig. 4J). Similarly, the number of cilia lining the otic vesicle was decreased (Fig. 4K, L). Finally, analysis of cilium formation at the nasal pit, at 48 hpf, revealed a significant alteration of the ciliary tuft in u4atac morphants compared to control embryos (Fig. 4M). All these ciliary defects were partially or totally rescued by the expression of human WT U4atac snRNA (Fig. 4G-M).

Altogether, these data indicate that loss of function of u4atac is associated *in vivo* with ciliary defects, which are likely secondary to minor splicing alterations*.*

JBTS/RFMN and TALS U4atac mutations lead to ciliopathy-related phenotypes in zebrafish

To have a better assessment of the impact of human U4atac variants on embryonic development processes linked to cilium function, we took advantage of the zebrafish model and conducted rescue experiments. We co-injected u4atac 5'SL MO with human U4atac carrying either the n.16G>A, n.51G>A or n.55G>A mutation. These mutations have been identified at the homozygous state in 4, 31 and 8 patients or fetuses respectively (Benoit-Pilven *et al*, 2020). They are associated with slightly different presentations (Table S7) and large differences in life expectancies: mean age at death is 10 months for patients homozygous for n.51G>A (three surviving patients, ≤ 2 years old), while 3/4 and 4/8 of those with n.16G>A or n.55G>A were still alive at the time of publication, their age varying from 2 to 40 and 2 to 20 years old, respectively.

Stability of injected human snRNA was validated by Northern blot (Supplementary Information, Fig. S4F), and the above-mentioned phenotypes were recorded. Regarding the body curvature, we observed that 16A, 51A and 55A snRNAs had little capacity to rescue the phenotype, with nevertheless a gradation between their effect, 55A snRNA co-injected-embryos having a slightly milder phenotype than those co-injected with 16A, themselves a little better than those co-injected with 51A snRNAs (Fig. 5A, B). Considering all other phenotypic features, the 51A snRNA seemed to have the lowest ability to rescue the anomalies compared to 16A and 55A, the latest corresponding to the less severe mutation (Fig. 5C). We also explored the impact of the mutations at the molecular level by analysing

in zebrafish the U12 intron retention of six cilium-related genes (*rabl2, tmem107l, tmem231, ift80, tctn1, pde6d*), among which the five genes tested in TALS and P1 patients' cells. For all of them, the U12 intron was strongly retained in 5'SL MO-injected embryos, and mis-splicing could be rescued to the level of control by the co-injection of WT U4atac. Consistent with the observed phenotype, n.51G>A had a more severe impact on minor splicing than n.16G>A, and then n.55G>A (Fig. 5D).

In conclusion, JBTS/RFMN and TALS *RNU4ATAC* pathogenic variants all lead to ciliary defects in zebrafish as the result of deficient minor splicing, with a gradation in the effect, sustaining a phenotype/genotype correlation that will need to be further explored.

Discussion

We report here for the first time the identification, in two independent cases suspected of having JBTS, of a homozygous pathogenic variant of *RNU4ATAC*, a gene previously involved in three rare syndromes (TALS, RFMN, LWS) characterized by microcephaly, short stature and other specific features. In line with this finding, we found that cilium-related genes are over-represented among human U12 genes, that several ciliary genes are impacted by defective minor splicing and that primary cilium function is altered in TALS and JBTS/RFMN fibroblasts. Furthermore, the zebrafish u4atac models present with ciliopathy-related phenotypes and ciliary defects that could be rescued by WT but not by U4atac carrying TALS and JBTS/RFMN variants.

Quite remarkably, the two infants we describe here have a complex syndromic disorder combining both traits seen in JBTS patients (MTS, hypotonia, nystagmus/strabismus, septal defects, polydactyly/clinosyndactyly) and in TALS/RFMN/LWS patients (microcephaly, skeletal dysplasia, growth retardation, immunodeficiency, eczema). They harbour a homozygous *RNU4ATAC* pathogenic variant, n.16G>A, the only one previously found at the homozygous state in Stem II in two RFMN patients while all others harboured compound heterozygous variants, one located in Stem II and the other elsewhere in U4atac. These two RFMN cases demonstrated that the disease could occur without the presence of a *RNU4ATAC* variant in TALS-implicated regions, as the six year old girl (Dinur Schejter *et al*, 2017) and the 38 year old male (Heremans *et al*, 2018) presented with typical RFMN phenotypes, albeit quite mild in the latter patient. To explain the phenotypic variability associated with n.16G>A homozygotes, one could consider that other genetic alterations are responsible for the ciliopathy-related traits seen in the JBTS/RFMN patients reported here, even if all of them were found at the heterozygous state (Table S1). Nevertheless, it is noteworthy that one of the two published n.16G>A homozygous RFMN patients presented with "eye motility problems" (Table 1) (Heremans *et al*, 2018), a feature previously unseen in *RNU4ATAC*-associated pathologies but common in JBTS. As for the MTS, most RFMN patients present with a mild microcephaly and borderline-to-mild intellectual impairment, so that brain MRI is not systematically performed. Based on our findings, we suggest that the

two RFMN previously published patients should have a thorough investigation of their brain morphology to explore the possible presence of a MTS. The overlap with ciliopathies is clearly not restricted to *RNU4ATAC*-associated JBTS as ciliopathy-related traits have been observed in some TALS/RFMN/LWS patients (including a MTS in one of them), even if it might be greater for n.16G>A homozygotes or more largely for carriers of two pathogenic variants in U4atac Stem II. In agreement with this, a recent disease scoring system based on the occurrence of ciliopathy-like phenotypes produced high scores for RFMN (267[th] highest score among 6058 analysed disorders) and TALS (428[th]), making them likely candidates for disorders with ciliary contribution (Lovera & Lüders, 2021). Moreover, we bring up the identification of the homozygous c.2953A>G variant in the *RTTN* gene, encoding the centrosome/cilium protein rotatin, in a child presenting with TALS features but *RNU4ATAC*-negative (Guguin et al., manuscript in preparation). Our finding correlates with other published cases (Grandone *et al*, 2016; Shamseldin *et al*, 2015), and strongly supports the notion that many of the developmental abnormalities seen in TALS patients could be due to ciliary/centrosome dysfunctions.

Additional evidence is provided by the zebrafish u4atac model. As seen in patients, MO-mediated deficiency of u4atac led to phenotypes that recapitulate features both associated with TALS/RFMN/LWS (microcephaly, brain hemorrhages (Abdel-Salam et al, 2013, 2011), cardiac defects), and with ciliopathies (body axis curvature, pronephric cysts and otolith defects). Brain hemorrhages could even fall in this last category as they have been shown to result from ciliary defects of endothelial cells in zebrafish (Eisa-Beygi et al, 2018; Kallakuri et al, 2015). Our results demonstrate that the ciliopathy-related phenotypes are indeed due to ciliary defects in the central canal, pronephros, otic vesicle and olfactive bulb. These phenotypes could be mostly rescued by co-injection of the human WT U4atac snRNA but not, or to a lesser extent, by snRNA molecules carrying JBTS/RFMN or TALS-associated mutations. These results thus strongly support the fact that several types of pathogenic variants of *RNU4ATAC* can result in ciliopathy-related phenotypes.

By identifying a homozygous *RNU4ATAC* mutation in two patients with emblematic ciliopathy traits, we shed light on a new "second-order" mechanism leading to ciliary defects,

i.e. splicing of minor introns. Second-order ciliopathies are caused by mutations in genes that have an indirect role in cilium formation or function (Reiter & Leroux, 2017), among which some are involved in different aspects of RNA metabolism. As examples, we can cite *DDX59* encoding a RNA helicase or *BICC1* coding for a RNA-binding protein involved in Oro-Facial-Digital (OFD) syndrome and cystic renal dysplasia, respectively (Shamseldin et al, 2013; Kraus et al, 2012). More recently, a subunit of the snRNA-processing complex Integrator, INTS13, has been linked to a complex developmental ciliopathy (Mascibroda *et al*, 2020) while other subunits were shown *in vitro* to be required for ciliogenesis (Jodoin et al, 2013). In all the second-order ciliopathies linked to the deficiency of an RNA-processing function, including minor splicing unravelled here, it remains to be explained why ciliary genes are specifically affected and how they drive the phenotype. In the case of minor splicing, the high proportion of U12 genes encoding ciliary components (at least 86, i.e. more than one tenth of the total number of U12 genes) is likely to be an important factor, combined to the fact that the cilium is a complex sensory machine involved in transducing a wide range of extracellular stimuli into cellular responses essential for organogenesis. Further investigations are needed to address these questions, including global transcriptomic analyses of splicing events in the zebrafish model.

To conclude, our work is of significant impact for both JBTS and TALS/RFMN/LWS patients, notably regarding the genetic diagnosis and counselling. Due to its non-coding status, *RNU4ATAC* is still under-considered among the clinically relevant genes. Not only should it be evaluated in the JBTS cases lacking a molecular diagnosis, but given the pleiotropy of the clinical signs associated with this gene, one should remain open when assessing its potential pathogenic variants in patients with rare diseases. Also, our work widens the fields of research on the physiopathological mechanisms underlying the *RNU4ATAC*-associated diseases.

Materials and Methods

Targeted next-generation sequencing

Sequencing of genes involved in ciliary diseases or cilia function was performed using a custom SureSelect capture kits (Ciliome V3c (1339 genes) or CilDiag (170 genes)) (Agilent Technologies, Santa Clara, CA, USA). Libraries were then sequenced using a HiSeq or NextSeq sequencers (Illumina, San Diego, CA, USA), respectively. Resulting paired-end reads were mapped on the GRCh37 reference genome using Burrows-Wheeler Aligner (Illumina). Bioinformatic analysis was performed using the Genome Analysis Toolkit, SAMTools and Picard Tools, in accordance to best practices issued by the Broad Institute. Variants were annotated using a software developed by the Paris Descartes University Bioinformatics platform. As a first screen, only nonsense, frameshift, splice site and missense variants, leading to an abnormal protein sequence, were retained; variants in non-exonic regions, pseudogenes, UTRs or with a frequency above 1% in public databases (dbSNP, 1000 genomes project, ExAC), or frequent in an in-house database of previously sequenced patients, were excluded. The functional consequence of missense variants was predicted using in silico tools, namely PolyPhen-2, SIFT, Mutation Taster, and CADD. Since this first screen turned out to be negative, i.e. identification of bi-allelic variants in one gene, variants in non-coding genes were thus analysed. Variants identified by NGS were then confirmed by Sanger sequencing.

Bioinformatic analyses

All the sequences were analyzed using the Ensembl Genome Browser. Nucleotide and amino acid sequences were compared to the non-redundant sequences present at the NCBI (National Center for Biotechnology Information) using the Basic Local Alignment Search Tool (BLAST). Secondary structure prediction for snRNA duplexes was performed by using RNAstructure (Reuter & Mathews, 2010).

U12 genes were identified by running the U12db (Alioto, 2007) annotation pipeline on the GRCh38 version of the human genome, with the gencode38 version of the annotations, or the GRCz11 version of the zebrafish genome. GO term analysis was performed with the

TopGO (v2.38.1) R tool (Alexa *et al*, 2006), with the default "weight01" algorithm, the Fisher-test and the org.Hs.eg.db (v3.10.0) R database of the genome wide annotation for humans. Genes without any GO annotation were discarded. Two analyses were run: i- U12 (618) vs U2 (19,939) genes; ii- U2 (19,939) vs U12 (618) genes.

To constitute a list of cilium-related genes, we opted for the combination of three existing databases, and thus defined a gene as cilium-related if 1) it was part of the Gene_Ontology (GO=Cilium), of the Gold_Standard (SysCilia, a curated list of known ciliary components) (Van Dam *et al*, 2013) or of the Predicted candidates based on a bayesian integration in CiliaCarta (van Dam *et al*, 2017) (last updated version of March, 18th 2018); 2) it was in the CilDB table v3.0 and had the maximum human ciliary evidence stringency level and at least 3 ciliary evidence (Arnaiz *et al*, 2015, 2009) (last update June 2014); 3) it was a ciliopathy-related gene based on three reviews, (Reiter & Leroux, 2017), (Wheway *et al*, 2019) and (Lovera & Lüders, 2021).

cDNA library preparation and high-throughput sequencing
One to two micrograms of RNA were sent for RNA-sequencing to IntegraGen Genomics (Evry, France), where a DNA library was generated with the tTruSeq Stranded mRNA Sample Prep" kit (Illumina) that comprises a step of mRNA purification using oligo(dT) beads. RNA-seq experiments have been performed on a HiSeq 4000 sequencer (Illumina), yielding approximately 2670 million of stranded two time 75 bp paired-end reads. Raw RNA-seq data are available upon request. IRFinder v1.2.0 was used to compute the PSI values for U12 introns.

Cell culture and immunofluorescence
Primary human fibroblasts were cultivated in HAM F10 media (Eurobio, CM1H1000-01) supplemented with 12% fetal bovine serum and 1% penicillin-streptomycin. On Day 1, $4x10^4$ cells were plated onto coverslips in 24-wells plates in order to have them at confluence the next day. On Day 2, cells were starved using HAM F10 media supplemented with 0.5% fetal bovine serum for two days to induce ciliogenesis. Then, cells were fixed in 4% PFA or with

cold methanol during 20 minutes at room temperature, and saturated with blocking buffer (PBS 1X, goat serum 10%, bovine serum albumin 1%, Triton 0.1%) for an hour. The following primary antibodies were used: Arl13b (1:500, Proteintech, 17711-1 AP), gamma-tubulin (1:250, Abcam, ab11316), acetylated alpha-tubulin (1:1000, Sigma, T6793) and adenylate cyclase III (1:500, Invitrogen, PA5-35382), diluted in blocking buffer and incubated overnight at 4°C. The following day, coverslips were washed with PBS before being stained with anti-rabbit Alexa 488 (1:1000, Invitrogen, A32731) and anti-mouse Alexa 647 (1:1000, Invitrogen, A32728) for an hour and half, and with DAPI for 10 minutes. Coverslips were finally mounted in FluorPreserveTM Reagent (EMD Millipore, 3457870) and z-stack images were taken using a Zeiss LSM880 confocal microscope. Measurement of the cilium length was performed manually using ImageJ software.

Zebrafish husbandry and microinjection of embryos

Zebrafish (*Danio rerio*) adults and embryos were maintained at 28.5°C under standard protocols. Breeding and staging were performed as per (Kimmel *et al*, 1995). To avoid pigmentation of embryos dedicated to imaging, 0.003% 1-phenyl-2-thiourea (PTU) was added to embryo medium. The following strains were used: the mixed wild-type ABxTU, and the transgenic lines *Tg(wt1b:EGFP)li1* ((44), *Tg(cmlc2:GFP)* and *Tg(bactin:arl13bGFP)hsc5Tg* (45) to label proximal pronephros, heart and cilia, respectively. Live embryos were photographed using a Leica M165FC stereoscope or Zeiss Axiozoom V16. Morpholinos, designed to target the 5' Stem Loop (5'SL) (5'-GATGTTCTCAGTTAACCTTCATTGA-3') or the Stem II (SII) (5'-AAACCACCCCCAGACAAGGAAGGTT-3') regions of u4atac_chr11, were provided by GeneTools, LCC and injected into the yolk at the one-cell stage (quantity ranging from 0.016 to 0.066 pmol (i.e. 135 to 556 pg) per embryo for 5'SL MO, and 0.002 to 0.006 pmol (i.e. 17 to 51 pg) per embryo for SII MO). A five-mismatch morpholino (5'-GATcTTgTCAcTTAACCTTgATTgA-3') was used as a negative control. For rescue experiments, constructions of human U4atac snRNA of WT or mutated (c.16G>A, c.51G>A and c.55G>A) sequences were previously described (Benoit-Pilven *et al*, 2020). snRNA

molecules were synthesized using a PCR amplicon as template (forward primer containing the T7 promoter (in italics): 5'-*TAATACGACTCACTATAGGG*AACCATCCTTTTCTT<u>G</u>GGGGTTG-3' (the underlined G being replaced by A for c.16G>A mutation), reverse primer: 5'-ATTTTTCCAAAAATTGCACCAAAATAAAGC-3') and the mMESSAGE mMACHINE™ T7 Transcription Kit (ThermoFisher Scientific), followed by a purification using RNA Clean and Concentrator kit (Zymo Research). Seventy-five picograms of snRNA were co-injected with u4atac-targeted morpholinos per embryo.

RNA extraction, RT-PCR and qRT-PCR

Total RNA was extracted from cultured fibroblasts or a pool of 20 zebrafish embryos and treated with DNase I using NucleoSpin RNA plus kit (Macherey-Nagel). 1.5 µg of total RNA was retro-transcribed into cDNA using random hexamers or oligodT and GoScript Reverse Transcriptase (Promega) following the manufacturer's protocol. RT-PCR were performed using the GoTaq Green Master Mix (Promega) according to the manufacturer's instructions (primer sequences in Table S8). PCR products were separated by electrophoresis on 1.5% agarose gel using GelRed nucleic acid gel stain (Biotium). Quantitative real time PCR was performed using the Rotor-gene Q (Qiagen). Reactions were performed in a 20 µl volume, with 0.5 µM of each primer, 10 µl of RotorGene SYBR Green Mix (Qiagen) and 25 ng of cDNA. The amplification was carried out using a denaturation program (95°C for 1 min), 45 cycles of two-steps amplification (95°C for 15s and 60°C for 30s) followed by a melting cycle. Reactions were performed in triplicate, and *ACTB* for fibroblasts or *gapdh* for zebrafish were used as a reference gene (see primer sequences in Table S8). Relative levels of expression of *rnu4atac* genes were calculated using the 2–ΔΔCt method. As for the quantification of intron retention by qRT-PCR, two pairs of primers were designed to target specifically the spliced and unspliced transcripts (see sequences in Table S8). After normalisation to the level of expression of ACTB/*gapdh* gene, the percentage of intron retention was calculated as the ratio of unspliced transcript level to that of the total of unspliced and spliced forms.

Zebrafish embryo O-Dianisidine staining

O-Dianisidine staining is used to detect hemoglobin in embryos. Staining was performed following a previously described method (Detrich *et al*, 1995). Briefly, embryos were dechorionated at 48 hpf and incubated for 15 minutes in staining solution containing O-Dianisidine 0.6 mg/mL, 0.01 M sodium acetate (pH 4.5), 0.65% H_2O_2, and 40% ethanol. Embryos were post-fixed in 4% PFA for 20 minutes, and dehydrated in methanol before being bathed in a clearing solution of benzyl benzoate/ benzyl alcohol (2:1, vol/vol) for imaging.

Zebrafish whole mount immunostaining

Embryos were dechorionated, washed twice in PBS, and then fixed in 4% PFA overnight. The next day, embryos were washed in PBS with 0.1% Tween 20 (PBST) and yolks were removed with clean forceps. After incubation for 3 hours in blocking solution PBST+3% BSA, embryos were washed thrice in PBST and incubated overnight at 4°C with anti-acetylated alpha-tubulin antibody in PBST (1:500; Sigma-Aldrich T6793). The next day, following three PBST washes, samples were incubated with Phalloidin-tetramethylrohdamine B isothiocyanate (1:100, Sigma-Aldrich P1951) for 20 min at room temperature, washed several times in PBST and incubated overnight at 4°C with anti-mouse secondary antibody in PBST (1:2000, Invitrogen A32728). Finally, following several PBST washes, embryos were mounted on glass slides using FluoPreserveTM Reagent (Merck) prior to confocal imaging.

Microscopy of zebrafish embryos

Live and fixed embryos were imaged using Zeiss LSM810 confocal microscope. For live imaging, embryos were mounted in 0.4% low melting agarose mixed with Tricain in glass-bottom Petri dishes (Ibidi). Live beating cilia were imaged by optimizing the signal to noise ratio and the scan speed, as previously described in ref.(45). Z-stacks were performed to

acquire the whole organs (nasal pit, central canal lumen, pronephros lumen, otic vesicle), and cilium number and range of motion were analysed using ImageJ software.

Statistics

All the data are reported as the mean of at least three independent experiments with s.e.m. All hypothesis tests were two-sided, and statistically significant differences (p<0.05) were calculated by one-way ANOVA or t-tests as indicated in figure legends. When sample size was too small or when normality was not reached (following Shapiro-Wilk test), a non-parametric test was used. Statistical analyses were performed using GraphPad Prism software.

Study approval

Animal studies (zebrafish) were conducted in accordance with the European regulations on animal use. Written informed consent was obtained from parents for the genetic study and for the publication of clinical information.

Acknowledgments and funding sources

We thank the families who participated in this study for their contribution to this project. We thank the CBC Biotec biobank for biosample management (Emilie Chopin, Isabelle Rouvet), the PRECI (Plateau de Recherche Expérimentale en Criblage In Vivo) aquatic core facility (Laure Bernard), the TEFOR facility, and the GENDEV team members for stimulating discussions. This work was supported by CNRS, Inserm, Université de Montpellier, Université Paris 7 and Université Lyon 1 through recurrent funding, the Fondation Maladies Rares ("Small Animal Models and Rare Diseases" program, no. 20161207), the Agence Nationale de la Recherche (no. ANR-18CE12-0007-01) grants.

References

Abdel-Salam GMH, Abdel-Hamid MS, Hassan NA, Issa MY, Effat L, Ismail S, Aglan MS & Zaki MS (2013) Further delineation of the clinical spectrum in RNU4ATAC related microcephalic osteodysplastic primordial dwarfism type I. *Am J Med Genet Part A*

Abdel-Salam GMH, Miyake N, Eid MM, Abdel-Hamid MS, Hassan NA, Eid OM, Effat LK, El-Badry TH, El-Kamah GY, El-Darouti M, *et al* (2011) A homozygous mutation in RNU4ATAC as a cause of microcephalic osteodysplastic primordial dwarfism type I (MOPD I) with associated pigmentary disorder. *Am J Med Genet Part A*

Alexa A, Rahnenführer J & Lengauer T (2006) Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. *Bioinformatics* 22: 1600–1607

Alioto TS (2007) U12DB: a database of orthologous U12-type spliceosomal introns. *Nucleic Acids Res* 35: D110–D115

Arnaiz O, Cohen J, Tassin AM & Koll F (2015) Remodeling Cildb, a popular database for cilia and links for ciliopathies. *Cilia*: 1–10

Arnaiz O, Malinowska A, Klotz C, Sperling L, Dadlez M, Koll F & Cohen J (2009) Cildb: A knowledgebase for centrosomes and cilia. *Database* 2009: 1–14

Benoit-Pilven C, Besson A, Putoux A, Benetollo C, Saccaro C, Guguin J, Sala G, Cologne A, Delous M, Lesca G, *et al* (2020) Clinical interpretation of variants identified in RNU4ATAC, a non-coding spliceosomal gene. *PLoS One* 15: 1–22

Burnicka-Turek O, Steimle JD, Huang W, Felker L, Kamp A, Kweon J, Peterson M, Reeves RH, Maslen CL, Gruber PJ, *et al* (2016) Cilia gene mutations cause atrioventricular septal defects by multiple mechanisms. *Hum Mol Genet* 25: 3011–3028

Cantaut-Belarif Y, Sternberg JR, Thouvenin O, Wyart C & Bardet P (2018) The Reissner Fiber in the Cerebrospinal Fluid Controls Morphogenesis of the Body Axis Report The Reissner Fiber in the Cerebrospinal Fluid Controls Morphogenesis of the Body Axis. *Curr Biol*: 1–8

Cologne A, Benoit-Pilven C, Besson A, Putoux A, Campan-Fournier A, Bober MB, Die-smulders CEMDE, Paulussen ADC, Pinson L, Toutain A, *et al* (2019) New insights into

minor splicing — a transcriptomic analysis of cells derived from TALS patients. *RNA* 25(9): 1130–1149

van Dam TJP, Kennedy J, der Lee R van, de Vrieze E, Wunderlich KA, Rix S, Dougherty GW, Lambacher NJ, Li C, Jensen VL, *et al* (2017) CiliaCarta: An integrated and validated compendium of ciliary genes

Van Dam TJP, Wheway G, Slaats GG, Huynen MA & Giles RH (2013) The SYSCILIA gold standard (SCGSv1) of known ciliary components and its applications within a systems biology consortium. *Cilia* 2: 1–5

Detrich HW, Kieran MW, Chan FY, Barone LM, Yee K, Rundstadler JA, Pratt S, Ransom D & Zon LI (1995) Intraembryonic hematopoietic cell migration during vertebrate development. *Proc Natl Acad Sci U S A* 92: 10713–10717

Dinur Schejter Y, Ovadia A, Alexandrova R, Thiruvahindrapuram B, Pereira SL, Manson DE, Vincent A, Merico D & Roifman CM (2017) A homozygous mutation in the stem II domain of RNU4ATAC causes typical Roifman syndrome. *npj Genomic Med* 2: 23

Edery P, Marcaillou C, Sahbatou M, Labalme A, Chastang J, Touraine R, Tubacher E, Senni F, Bober MB, Nampoothiri S, *et al* (2011) Association of TALS Developmental Disorder with Defect in Minor Splicing Component U4atac snRNA. *Science (80-)* 332: 240–243

Eisa-Beygi S, Benslimane FM, El-Rass S, Prabhudesai S, Khatib M, Abdelrasool A, Simpson PM, Yalcin HC, Burrows PE & Ramchandran R (2018) Characterization of endothelial cilia distribution during cerebral-vascular development in zebrafish (Danio rerio) HHS Public Access. *Arter Thromb Vasc Biol* 38: 2806–2818

Failler M, Gee HY, Krug P, Joo K, Halbritter J, Belkacem L, Filhol E, Porath JD, Braun DA, Schueler M, *et al* (2014) Mutations of CEP83 cause infantile nephronophthisis and intellectual disability. *Am J Hum Genet*

Farach LS, Little ME, Duker AL, Logan C V, Jackson A, Hecht JT & Bober M (2017) The expanding phenotype of RNU4ATAC pathogenic variants to Lowry Wood syndrome. *Am J Med Genet A*: 1–5

Grandone A, Torella A, Santoro C, Giugliano T, del Vecchio Blanco F, Mutarelli M, Cirillo M, Cirillo G, Piluso G, Capristo C, *et al* (2016) Expanding the phenotype of RTTN

variations: a new family with primary microcephaly, severe growth failure, brain malformations and dermatitis. *Clin Genet*

Hagiwara H, Matsumoto H, Uematsu K, Zaha K, Sekinaka Y, Miyake N, Matsumoto N & Nonoyama S (2021) Immunodeficiency in a patient with microcephalic osteodysplastic primordial dwarfism type I as compared to Roifman syndrome. *Brain Dev* 43: 337–342

He H, Liyanarachchi S, Akagi K, Nagy R, Li J, Dietrich RC, Li W, Sebastian N, Wen B, Xin B, *et al* (2011) Mutations in U4atac snRNA, a component of the minor spliceosome, in the developmental disorder MOPD i. *Science (80-)* 332: 238–240

Heremans J, Garcia-Perez JE, Turro E, Schlenner SM, Casteels I, Collin R, de Zegher F, Greene D, Humblet-Baron S, Lesage S, *et al* (2018) Abnormal differentiation of B cells and megakaryocytes in Roifman syndrome. *J Allergy Clin Immunol*: 1–17

Jodoin JN, Shboul M, Albrecht TR, Lee E, Wagner EJ, Reversade B & Lee LA (2013) The snRNA-processing complex, Integrator, is required for ciliogenesis and dynein recruitment to the nuclear envelope via distinct mechanisms. *Biol Open* 2: 1390–1396

Kallakuri S, Yu JA, Li J, Li Y, Weinstein BM, Nicoli S & Sun Z (2015) Endothelial cilia are essential for developmental vascular integrity in zebrafish. *J Am Soc Nephrol* 26: 864–875

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B & Schilling TF (1995) Stages of Embryonic Development of the Zebrafish. 10

Kramer-Zucker AG, Olale F, Haycraft CJ, Yoder BK, Schier AF & Drummond IA (2005) Cilia-driven fluid flow in the zebrafish pronephros, brain and Kupffer's vesicle is required for normal organogenesis. *Development* 132: 1907–1921

Kraus MRC, Clauin S, Pfister Y, Di Maïo M, Ulinski T, Constam D, Bellanné-Chantelot C & Grapin-Botton A (2012) Two mutations in human BICC1 resulting in wnt pathway hyperactivity associated with cystic renal dysplasia. *Hum Mutat* 33: 86–90

Lovera M & Lüders J (2021) The ciliary impact of nonciliary gene mutations. *Trends Cell Biol* 31: 876–887

Mascibroda LG, Shboul M, Elrod ND, Colleaux L, Hamamy H, Huang KL, Peart N, Singh MK, Lee H, Merriman B, *et al* (2020) INTS13 mutations causing a developmental

ciliopathy disrupt integrator complex assembly. *bioRxiv*

Merico D, Roifman M, Braunschweig U, Yuen RKC, Alexandrova R, Bates A, Reid B, Nalpathamkalam T, Wang Z, Thiruvahindrapuram B, *et al* (2015) Compound heterozygous mutations in the noncoding RNU4ATAC cause Roifman Syndrome by disrupting minor intron splicing. *Nat Commun* 6: 8718

Olthof AM, Hyatt KC & Kanadia RN (2019) Minor intron splicing revisited: Identification of new minor intron-containing genes and tissue-dependent retention and alternative splicing of minor introns. *BMC Genomics* 20: 1–19

Olthof AM, Rasmussen JS, Campeau PM & Kanadia RN (2020) Disrupted minor intron splicing is prevalent in Mendelian disorders. *Mol Genet Genomic Med* 8: 1–14

Putoux A, Alqahtani A, Pinson L, Paulussen ADC, Michel J, Besson A, Mazoyer S, Borg I, Nampoothiri S, Vasiljevic A, *et al* (2016) Refining the phenotypical and mutational spectrum of Taybi-Linder syndrome. *Clin Genet* 90: 550–555

Reiter JF & Leroux MR (2017) Genes and molecular pathways underpinning ciliopathies. *Nat Rev Mol Cell Biol* 18: 533–547

Reuter JS & Mathews DH (2010) RNAstructure: Software for RNA secondary structure prediction and analysis. *BMC Bioinformatics* 11

Shamseldin H, Alazami AM, Manning M, Hashem A, Caluseiu O, Tabarki B, Esplin E, Schelley S, Innes AM, Parboosingh JS, *et al* (2015) RTTN Mutations Cause Primary Microcephaly and Primordial Dwarfism in Humans. *Am J Hum Genet*

Shamseldin HE, Rajab A, Alhashem A, Shaheen R, Al-Shidi T, Alamro R, Al Harassi S & Alkuraya FS (2013) Mutations in DDX59 implicate RNA helicase in the pathogenesis of orofaciodigital syndrome. *Am J Hum Genet* 93: 555–560

Song Z, Zhang X, Jia S, Yelick PC & Zhao C (2016) Zebrafish as a Model for Human Ciliopathies. *J Genet Genomics* 43: 107–120

Thomas S, Legendre M, Saunier S, Bessières B, Alby C, Bonnière M, Toutain A, Loeuillet L, Szymanska K, Jossic F, *et al* (2012) TCTN3 mutations cause Mohr-Majewski syndrome. *Am J Hum Genet* 91: 372–378

Turunen JJ, Niemelä EH, Verma B & Frilander MJ (2013) The significant other: splicing by

the minor spliceosome. *Wiley Interdiscip Rev RNA* 4: 61–76

Wheway G, Mitchison HM, Ambrose JC, Baple EL, Bleda M, Boardman-Pretty F, Boissiere JM, Boustred CR, Caulfield MJ, Chan GC, *et al* (2019) Opportunities and challenges for molecular understanding of ciliopathies–the 100,000 genomes project. *Front Genet* 10: 1–21

Wu D, Freund JB, Fraser SE & Vermot J (2011) Mechanistic Basis of Otolith Formation during Teleost Inner Ear Development. *Dev Cell* 20: 271–278

Table 1 : Clinical features of patients with *RNU4ATAC* mutations

Disease	TALS	LWS	RFMN	RFMN		JBTS/RFMN		JBTS
***RNU4ATAC* genotype**	various[a]	various[a]	various[a]	16G>A		16G>A		none
# Patients	55 cases[a,b]	5 cases[a]	15 cases[a]	Case 1[c]	Case 2[d]	F1:II-2 (P1)	F2:II-3 (P2)	none
Growth retardation	severe	mild	mild	+	severe	+	mild	-
Neurological features								
Developmental delay	+	+	+	+	+	+	+	+
Intellectual deficiency	moderate to severe	mild	mild	NR	NE	NE	NE	+ (inconstant)
Microcephaly	severe	variable	mild	mild	NR	+	+	-
Abnormal gyral pattern	+	-	-	NR	NR	-	-	rare
Corpus callosum agenesis	+	-	-	NR	NR	-	-	rare
Molar tooth sign	1 case	-	-	NR	NR	+	+	+
Skeletal abnormalities								
Skeletal dysplasia	+	+	+	+	+	+	+	-
Polydactyly	1 case	-	-	-	-	+	-	occasional
Ophthalmological abnormalities								
Retinal dystrophy	occasional	+	frequent	+	+	-	NE	occasional
Nystagmus / EMA	-	-	-	NR	+	+	+	+
Coloboma	-	-	-	-	-	+	NE	occasional (COACH)
Cardiopathy	occasional	rare	occasional	-	-	+	+	occasional
Skin and skin appendage								
Eczema	frequent	-	+	+	+	+	NR	-
Sparse hair	+	-	-	NR	NR	-	NR	-
Endocrine abnormalities								
Genital abnormalities	occasional	-	-	NR	+	+	NR	-
Hypothyroidism	occasional	-	-	NR	+	NR	NR	rare
IGF1 deficiency	?	-	-	NR	+	+	NE	-
Hepatomegaly	-	-	+	-	+	+	NR	occasional (COACH)
Immunological features								
IgG deficiency	occasional	occasional	+	+	+	+	NR	-
IgA deficiency	-	-	+	-	+	+	NR	-
Lymphocyte B deficiency	1 case	-	+	+	+	-	NR	-
Specific Ab deficiency	1 case	-	+	+	+	+	NR	-

TALS, Taybi-Linder Syndrome; LWS, Lowry Wood Syndrome; RFMN, Roifman Syndrome; JBTS, Joubert Syndrome; NE, not evaluated; NR, not reported; Ab, antibody; EMA, eye movement abnormalities; COACH, Cerebellar vermis hypoplasia, Oligophrenia, Ataxia, Coloboma and Hepatic fibrosis. [a]Cases reported in the literature, reviewed in Benoit-Pilven et al, *PLoS One* 2020. [b]Hagiwara et al, *Brain Dev* 2020. [c]Case 1 is reported in Dinur Schejter et al., *NPJ Genom Med* 2017; [d]Case 2 is reported in Heremans et al., *J Allergy Clin Immunol* 2018.

Figure legends

Figure 1. *RNU4ATAC* is mutated in two independent cases of patients with JBTS traits

(A) Schema of the structure of the U4atac/U6atac bi-molecule, with the main interacting proteins, showing the nucleotides mutated in patients with *RNU4ATAC*-associated syndromes. The red arrowhead points to the guanine nucleotide in position 16, which is mutated (n.16G>A) in patients with JBTS/RFMN; the two green arrowheads point to the guanine nucleotides in positions 51 and 55, the most frequently mutated nucleotides in TALS patients. **(B)** Patient 1, F1:II-2: Cerebral examination performed within first year of age. Mid-sagittal T1 weighted MR image showing a complete but short vermis suggestive of vermian hypoplasia (vermian height 25 mm, double arrow) with normal corpus callosum (left panel). Infra (middle panel) and supra (right panel) tentorial T2-weighted images

demonstrating unilateral hypoplasia of the left cerebral peduncle (red arrow) as well as elongation and thickening of the superior cerebellar peduncle leading to the characteristic « molar tooth sign » (red asterisk), and a small arachnoid cyst below the left internal cerebral vein (black star). **(C)** Patient 2, F2:II-3: Cerebral examination performed in a premature baby at few days of age. Mid-sagittal T1 weighted MR image (left panel) showing a complete but short (vermian height 12 mm, 50th centile 18,5 mm) and dysgenetic (lack of fissures) vermis with enlarged retro and infra-cerebellar cyst suggestive of Blake pouch cyst (black asterisk), well demonstrated on infra-tentorial T2-weighted image (middle panel) (black asterisk). Axial T1-weighted image (right panel) showing abnormal elongated superior cerebellar peduncles (black stars) leading to typical molar tooth sign associated to abnormal fourth ventricle shape (antero-posterior diameter superior to transverse diameter) as previously reported as a prenatal feature for Joubert or Joubert-like syndrome.

Figure 2. The non-motile cilium assembly pathway is enriched among U12 genes

(A) GO term enrichment among U12 intron-containing genes. Comparison of 618 U12 and 19,939 U2 genes with Gene Ontology annotations, using the TopGO R package, allowed to identify 80 enriched biological processes with a p-value<0.05, of which the 13 most significant are shown. Numbers in bars indicate the ratio of U12 gene to the total gene number belonging to each GO term; in brackets the percentage of U12 genes among the

GO term. **(B)** Schema representing the localisation and/or function of the 86 cilium-related U12 genes. Underlined are the "Gold standard" genes (Van Dam *et al*, 2013), in bold the genes associated with a ciliopathy, and in red those associated with JBTS. Hh, hedgehog pathway; PCP, Planar Cell Polarity pathway.

A

deltaPSI	Number of U12 genes		% of expressed U12 genes	
	TALS	P1	TALS	P1
≥ 10%	48	18	10	4
1 - 10%	275	140	57,4	30,8
< 1%	156	296	32,6	65,2
Total	479	454	100	100

Figure 3. JBTS/RFMN-related *RNU4ATAC* mutation leads to U12 intron retention in ciliary genes and results in ciliary function alteration in patient's fibroblasts

(A) Table summarising the analysis of U12 intron retention in the transcriptomic data of Patient 1 (P1) *vs* ctl1 fibroblasts, in comparison to previously published TALS samples (5 patient *vs* 5 control) (Cologne *et al*, 2019). Indicated are the number and percentage of U12 genes presenting differences of percent-spliced-in (PSI) values greater than 10%, between 10 and 1%, or below 1% between patient and control cells (deltaPSI). **(B)** qRT-PCR analyses of inton retention (IR) of U12 ciliary genes in control, TALS6 and P1 fibroblasts. Graphs show the mean ± s.e.m. of at least 3 independent experiments. $*p<0.05$, $**p<0.005$, using Kruskal-Wallis test with Dunn's multiple comparisons. **(C)** Confocal images of age- and sex-matched control, TALS and P1 patients' fibroblasts derived from skin biopsy. Adenylate cyclase (AC3, green) is co-labelled with Arl13b (cilium marker, magenta). Scale bar, 10 µm. **(D)** Quantification of the percentage of AC-positive cilia seen in (C). Graphs show the mean ± s.e.m. of at least 6 fields of 5 independent experiments (total number of analysed cells >200). $****p<0.0001$, using Kruskal-Wallis test with Dunn's multiple comparisons.

Figure 4. Deficiency of u4atac leads to cilium alterations in zebrafish

(A-F) Global morphology of control (ctl) (upper panel) and u4atac 5'SL (lower panel) morpholino (MO)-injected embryos at 48 hpf. u4atac morphants are characterized by body curvature and cardiac edema (A', arrowhead); cysts (B', asterisks) in *Tg(wt1b:GFP)*-labelled glomeruli; anomalies of otolith morphology/number in otic vesicle (C'); alterations of cardiac morphology with atrial hypertrophy, as seen with *Tg(cmlc2:GFP)* transgene (D'); blood hemorrhages in brain, as visualised by O-dianisidine staining (E', arrowhead); and microcephaly. Dorsal views in B-B' and ventral views in D-D', with anterior to the top. **(G)** Percentage of embryos displaying each of the phenotypes shown in A-F, following injection of control MO, u4atac 5'SL MO alone or with human WT U4atac snRNA. Graph shows the mean ± s.e.m. of three independent experiments (total number of embryos : >300). $p < 0.01$, t-tests comparing 5'SL MO to ctl MO, and 5'SL MO + WT to 5'SL MO. **(H)** Live confocal imaging of motile cilia in central canal of MO-injected *Tg(bactin:arl13bGFP)* transgenic embryos, at 48 hpf. Slow acquisition speed of motile cilia produces imaging artefacts, delineating the range of motion of each cilium (yellow angles). **(I)** Quantification of the angle defined by the range of motion of motile cilia in control MO (n=22), u4atac 5'SL MO alone (n=22) or with human WT U4atac snRNA (n=20) injected embryos, as described in (H). Box-and-whisker plots show in the box the median and the 25th-75th percentiles, and in whiskers the minimum to the maximum values. Values are the mean ranges of motion of each single embryo. ****$p < 0.0001$, by Kruskal-Wallis test with Dunn's multiple comparisons test. **(J)** Immunostaining of cilia (acetylated α-tubulin) in the proximal straight tubule of control MO, u4atac 5'SL MO alone or with human WT U4atac snRNA injected embryos, at 48 hpf. In brackets, the number of embryos exhibiting the phenotype shown in the image over the total amount of analysed embryos. Dotted lines delineate the tubule lumen. **(K)** Immunostaining of cilia (acetylated α-tubulin) in the otic vesicle of control MO, u4atac 5'SL MO alone or with human WT U4atac snRNA injected embryos, at 24 hpf. **(L)** Quantification of the cilium number in otic vesicle of control MO (n=15), u4atac 5'SL MO alone (n=15) or with human WT U4atac snRNA (n=16) injected embryos, as described in (K). Box-and-whisker plots show in the box the median and the 25th-75th percentiles, and in whiskers the minimum to the maximum values. ****$p < 0.0001$, ***$p < 0.005$, **$p < 0.01$ by Kruskal-Wallis test with Dunn's

multiple comparisons test. **(M)** Immunostaining of cilia (acetylated α-tubulin) at the nasal pit of control MO, u4atac 5'SL MO alone or with human WT U4atac snRNA injected embryos, at 48 hpf. In brackets, the number of embryos exhibiting the phenotype shown in the image over the total amount of analysed embryos. Scale bars, 10 µm. a, atrium; v, ventricle; hem, hemorrhages.

Figure 5. Co-injected JBTS/RFMN and TALS U4atac mutants lead to ciliopathy-related phenotypes in zebrafish.

(A) Global morphology of embryos co-injected with 5'SL MO and JBTS/RFMN (16A) or TALS (51A, 55A) -related U4atac variants. A varying severity of body curvature could be observed (mild, curly, severe). **(B)** Quantification of embryos exhibiting the various severities

of body curvature shown in (A). **(C)** Percentage of embryos displaying each of the phenotypes shown in Fig. 4A-F, following co-injection of 5'SL MO with human WT or mutated U4atac snRNA. Graph shows the mean ± s.e.m. of three independent experiments (total number of embryos: 180). **(D)** qRT-PCR analysis of U12 intron retention (IR) in six cilium-related genes in embryos co-injected with 5'SL MO and human WT or mutated U4atac snRNA. Graphs show the mean ± s.e.m. of three independent experiments.