Rapid fluoroquinolone resistance detection in *Pseudomonas aeruginosa* using mismatch

amplification mutation assay-based real-time PCR

Danielle E. Madden[1,2,3], Kate L. McCarthy[4,5], Scott C. Bell[6,7,8], Olusola Olagoke[1,2,3], Timothy Baird[2,9],

Jane Neill[9], Kay A. Ramsay[8], Timothy J. Kidd[10,11], Adam G. Stewart[5,12], Shradha Subedi[3,12], Keat

Choong[3,12], Tamieka A. Fraser[1,2], Derek S. Sarovich[1,2,3*], and Erin P. Price[1,2,3*]

[1]GeneCology Research Centre, University of the Sunshine Coast, Sippy Downs, Queensland, Australia

[2]Sunshine Coast Health Institute, Birtinya, Queensland, Australia

[3]Infection Research Network Sunshine Coast, Birtinya, Queensland, Australia

[4]Infectious Diseases Unit, Royal Brisbane and Women's Hospital, Herston, Queensland, Australia

[5]University of Queensland Centre for Clinical Research, Herston, Queensland, Australia

[6]Adult Cystic Fibrosis Centre, The Prince Charles Hospital, Chermside, Queensland, Australia

[7]Translational Research Institute, Woolloongabba, Queensland, Australia

[8]Child Health Research Centre, The University of Queensland, Centre for Children's Health Research,

South Brisbane, QLD, Australia

[9]Respiratory Department, Sunshine Coast University Hospital, Birtinya, Queensland, Australia

[10]School of Chemistry and Molecular Biosciences, University of Queensland, St Lucia, Queensland,

Australia

[11]Central Microbiology, Pathology Queensland, Royal Brisbane and Women's Hospital, Herston,

Queensland, Australia

[12]Infectious Diseases Unit, Sunshine Coast University Hospital, Birtinya, Queensland, Australia

Keywords: Antibiotic, AMR, fluoroquinolone, ciprofloxacin, *Pseudomonas aeruginosa*, real-time PCR,

SYBR-MAMA

Running title: Rapid FQ resistance detection in *P. aeruginosa*

*Authors contributed equally

Corresponding author: Erin Price, eprice@usc.edu.au Ph: +61 7 5456 5568

Abstract

Background: Antimicrobial resistance (AMR) is an ever-increasing global health concern. One crucial facet in tackling the AMR epidemic is earlier and more accurate AMR diagnosis, particularly in the dangerous and highly multi-drug resistant ESKAPE pathogen, *Pseudomonas aeruginosa*.

Objectives: We aimed to develop two SYBR Green-based mismatch amplification mutation assays (SYBR-MAMAs) targeting GyrA T83I (*gyrA*248), and GyrA D87N, D87Y, and D87H (*gyrA*259). Together, these variants cause the majority of fluoroquinolone (FQ) AMR in *P. aeruginosa*.

Methods: Following assay validation, the *gyrA*248 and *gyrA*259 SYBR-MAMAs were tested on 84 clinical *P. aeruginosa* isolates from Queensland, Australia, 45 of which demonstrated intermediate/full ciprofloxacin resistance according to antimicrobial susceptibility testing.

Results: Our two SYBR-MAMAs correctly predicted an AMR phenotype in the majority (84%) of isolates with intermediate/full FQ resistance. Importantly, all FQ-sensitive strains were predicted to have a sensitive phenotype. Whole-genome sequencing confirmed 100% concordance with SYBR-MAMA genotypes.

Conclusions: Our GyrA SYBR-MAMAs provide a rapid and cost-effective method for same-day identification of FQ AMR in *P. aeruginosa.* An additional SYBR-MAMA targeting the GyrB S466Y/S466F variants would increase FQ AMR prediction to 91%. Clinical implementation of our assays will permit more timely treatment alterations in cases where decreased FQ susceptibility is identified, leading to improved patient outcomes and antimicrobial stewardship.

Background

The ESKAPE pathogen, *P. aeruginosa*, has a remarkable capacity to develop AMR towards all

clinically-relevant antibiotic classes (1). This bacterium can cause life-threatening infections,

particularly in people with wounds, cancer, or chronic respiratory diseases such as cystic fibrosis (CF)

or chronic obstructive pulmonary disease (COPD) (2).

Rapid, affordable, accessible, and accurate AMR diagnosis is crucial in the battle against ESKAPE

pathogens (1). However, few diagnostic tests exist (3) for rapidly and inexpensively characterising

AMR-conferring single-nucleotide polymorphisms (SNPs) in *P. aeruginosa*, a striking knowledge gap

given that SNPs confer AMR towards anti-pseudomonal drugs such as fluoroquinolones (FQs) (4).

FQs (predominantly ciprofloxacin [CIP]) have proven clinically useful for treating *P. aeruginosa*

infections (5). Yet, upon exposure, *P. aeruginosa* often develops FQ resistance (FQr). Codon-altering

mutations within the GyrA quinolone-resistance-determining region (QRDR) can confer an

intermediate (CIPi) or fully resistant (CIPr) CIP phenotype (6, 7); in contrast, *gyrB*, *parC*, *parE* and

nfxB typically require ≥2 mutations to impart CIPi/CIPr (8).

Due to the single-step nature of QRDR mutations in conferring CIPi/CIPr, and their prevalence in

clinical isolates (9), we targeted the two most common QRDR SNPs, *gyrA*248 (4) and *gyrA*259 (10),

for assay development. We chose SYBR Green-based mismatch amplification mutation assay (SYBR-

MAMA), an inexpensive (~AUD$1-2/assay when run in duplicate), simple, rapid (~1h turnaround

time) and scalable method (11-15) that exploits the differential efficiency of allele-specific

amplification for SNP interrogation; this efficiency disparity can be observed in real-time by

measuring the difference in cycles-to-threshold (ΔC_T) (12, 14, 15).

Methods

Eighty-four *P. aeruginosa* isolates from Southeast Queensland, Australia, were examined: 42 from sputum derived from adults with CF and chronic *P. aeruginosa* infection admitted to The Prince Charles Hospital between 2017 and 2019 (16); 35 bloodstream isolates retrieved from adults admitted to several public and private hospitals in Brisbane between 2008 and 2011 (17), three from COPD sputum, collected during in-home community nurse visits in the Sunshine Coast region (16); one from an adult with non-CF bronchiectasis collected in 2017, and one from an adult with urinary tract infection collected in 2018, both during admission to the Sunshine Coast University Hospital (Table 1). One ulcer and one ear infection isolate, both from Brisbane, were obtained from the 1,000 International *P. aeruginosa* Consortium collection (18). Ethics approvals were obtained as previously described (16, 17, 19).

Strains were isolated using MacConkey agar (Oxoid, VIC, Australia) incubated at 37°C for 24h and confirmed as *P. aeruginosa* by *ecfX* real-time PCR (20). Susceptibility towards CIP [5µg], levofloxacin (LEV; 5 µg), moxifloxacin (MFX; 5 µg), and ofloxacin (OFX; 5 µg) was determined by disc diffusion (Edwards Group, QLD, Australia) using CLSI M100S-Ed27:2017 guidelines (Table 2).

Isolates were DNA-extracted using the DNeasy kit (Qiagen, Chadstone Centre, VIC, Australia), followed by Illumina paired-end whole-genome sequencing (WGS) (19). Quality-filtered reads (21) were assembled with MGAP v1.1 (https://github.com/dsarov/MGAP---Microbial-Genome-Assembler-Pipeline) or SPAdes (22). AMR prediction was undertaken using a *P. aeruginosa*-specific ARDaP (23) database. Assemblies were deposited into the PubMLST database (https://pubmlst.org/organisms/pseudomonas-aeruginosa). Illumina data are available via BioProject PRJNA761496. We also tested rapid DNA extraction of 10 representative strains using the 5% chelex-100 rapid heat soak method (24), followed by a 1:10 dilution in molecular-grade H_2O prior to PCR.

SYBR-MAMA primers were assessed *in silico* for dimer formation and specificity as described

previously (25). For the *gyrA*248 SYBR-MAMA, gyrA248_T_AMR amplifies the CIPi/CIPr-conferring

T83I allele, whereas gyrA248_C_WT amplifies the wild-type allele (Table 3). For the *gyrA*259 SYBR-

MAMA, gyrA259_D_AMR amplifies mutant alleles at position 259 (D87N, D87Y, D87H), all of which

confer CIPi/CIPr (4, 8), whereas gyrA259_G_WT amplifies the wild-type allele (10). PCRs consisted of

1× SsoAdvanced Universal SYBR Green Supermix (Bio-Rad, NSW, Australia), 0.2µM primers, 1µL

template, and PCR-grade H_2O, to 5µL. Thermocycling comprised 95°C for 2min, followed by 40 cycles

of 95°C for 5s and 60°C for 5sec. Strains used to validate *gyrA* 248 and 259 alleles are listed in Table

2.

Results

SYBR-MAMAs were screened across the 85 genome-sequenced *P. aeruginosa* isolates, comprising 40

CIP-sensitive (including PAO1), 29 CIPi, and 16 CIPr strains (Table 1). Of these, only one,

SCHI0050.S.1, was derived from a participant receiving FQ (CIP) treatment, and this isolate was CIP-

sensitive (Table 1). Unexpectedly, none of the 45 CIPi/CIPr strains were from participants known to

be receiving contemporaneous FQ antibiotics (Table 1), although we were unable to investigate

historical FQ exposure due to ethical limitations on participant data collection. It is therefore

possible that some of our participants have previously received FQ antibiotics in the weeks or

months prior to our sample collection. Alternatively, given that most of our participants were

hospitalised, another possibility is that the CIPi/CIPr strains were nosocomially acquired, either from

other admitted patients who had or were being treated with FQs, or from the hospital environment.

The *gyrA*248 SYBR-MAMA robustly discriminated GyrA T83I from wild-type strains, with matched

alleles consistently amplifying earlier than mismatched counterparts (T83I ΔC_T=4.0±0.03 vs. wild-

type ΔC_T=7.6±0.1 [Figures 1A and 1B]). Four tested GyrA T83I strains also demonstrated

intermediate or full resistance towards LEV, MFX, and OFX (Table 2), confirming the importance of

this variant in broader FQr. T83I is considered the most common GyrA variant in CIPi/CIPr strains (4,

9, 26); for example, two Japanese studies reported 82% (60/73) (27) and 75% (112/150) (28) T83I

prevalence among CIPi/CIPr isolates, and a Vietnamese study reported 54% (76/141) prevalence (9).

In our dataset, T83I was detected in 29% (13/45) CIPi/CIPr strains and 0% (0/40) CIP-sensitive strains

(Table 1), suggesting that T83I is an important, but not dominant, cause of CIPi/CIPr in Australian

isolates, although testing across a broader isolate collection is required to confirm this observation.

Like *gyrA*248, there was clear discrimination between AMR and wild-type genotypes for the *gyrA*259

SYBR-MAMA, with AMR alleles amplifying earlier in AMR-encoding strains (D87Y ΔC_T=14.7 [Figure

2A]; D87N ΔC_T=9.6±0.04 [Figure 2B]; D87H ΔC_T=13.5±0.2 [Figure 2C]), and vice versa for wild-type

strains (ΔC_T=10.8; Figure 2D). *gyrA*259 AMR was detected in 56% (25/45) CIPi/CIPr and 0% (0/40)

CIP-sensitive strains. The degenerate nature of the *gyrA*259 AMR primer has the advantage of

enabling all four nucleotide variants to be detected using just two reactions; it does not require each

variant to be tested individually. We therefore used WGS to differentiate each of the *gyrA*259 AMR

variants to determine their prevalence. D87H was most common according to WGS, accounting for

88% (22/25) *gyrA*259 AMR strains (Table 1). In no instance did we observe a strain encoding both

*gyrA*248 and *gyrA*259 AMR variants.

Like T83I, seven tested strains encoding D87Y, D87N and D87H all exhibited intermediate or full LEV,

MFX, and OFX resistance (Table 2). The *gyrA* mutations had a larger impact on MFX and OFX (causing

AMR) compared with CIP and LEV (causing intermediate resistance) (Table 2). This observation

supports the hypothesis that additional mutation/s are sometimes required to confer AMR towards

CIP and LEV (10).

We tested our two SYBR-MAMAs against rapid chelex-extracted heat soak DNA. In all instances,

isolates yielded excellent, early amplification, and genotyped as expected. Although not tested in

this study, further time savings could be made by performing SYBR-MAMAs using colony PCR (29);

however, in our experience, this approach requires a high degree of skill due to the inhibitory nature

of total cellular extracts in PCR, and typically results in a proportion of PCR failures, even with skilled

operators. This issue is particularly acute when using low (e.g. 5 μL) reaction volumes, as was used in

our study to minimise costs. Therefore, we recommend that chelex extractions followed by 1:10

dilution be performed where rapid DNA extraction is desired to ensure 100% amplification success.

Of the seven CIPi/CIPr strains that did not encode *gyrA*248 or *gyrA*259 AMR variants, three had GyrB

Ser466Phe or Ser466Tyr missense mutations with or without other FQr-conferring mutations (ParE

Ala473Val, MexR Gln25Stop), whereas four had no known FQr determinants (Table 1). A possible

explanation for these latter strains is phenotype misclassification as 3/4 strains sat on the

CIPi/sensitivity breakpoint. Although not performed in this study, determining the minimum

inhibitory concentration of these strains using Etests or broth microdilution may help resolve these

borderline phenotypes.

Conclusions

Our two *gyrA*248 and *gyrA*259 SYBR-MAMAs detected FQ non-susceptibility in 84% Australian

CIPi/CIPr strains. Importantly, all FQ-sensitive strains yielded wild-type genotypes for both assays,

demonstrating 100% specificity. Our two *gyrA* SYBR-MAMAs thus provide a same-day, inexpensive,

simple, and accurate tool for detecting the two most prevalent causes of FQ non-susceptibility in *P.*

aeruginosa. An additional SYBR-MAMA targeting GyrB Ser466Phe and Ser466Tyr would increase

detection of FQ non-susceptibility to 91% without loss of specificity. Conversion of our assays to

single-tube Melt-MAMA format (11) would further decrease assay costs and PCR setup time.

Acknowledgements

We thank Patrick Harris at UQ Centre for Clinical Research for helpful discussions.

Funding

This work was supported by Advance Queensland (awards AQIRF0362018 and AQRF13016-17RD2),

an Australian Government Research Training Program Scholarship, the Wishlist Sunshine Coast

Health Foundation (award 2019-14), and the National Health and Medical Research Council (award

455919).

Transparency declaration

The authors declare no competing interests. The funders had no role in study design, data

acquisition, analysis, interpretation, writing or submission of the manuscript.

Tables

Table 1. *Pseudomonas aeruginosa* isolates used in this study, and their DNA gyrase A (GyrA) SYBR-MAMA PCR genotypes, whole-genome sequencing

(WGS) results, multilocus sequence types, and treatment details

Strain	Disease	*gyrA*248 allele (PCR)	*gyrA*259 allele (PCR)	GyrA AMR mutation (WGS)	Other CIP AMR-conferring mutations (WGS)	Sequence Type	Treatment at time of collection
CIP-resistant							
SCHI0004.S.5	CF	AMR	WT	T83I	None	ST649	Day 14 IV MEM, TOB
SCHI0008.S.3	CF	AMR	WT	T83I	None	ST649	Day 8 IV ATM, DOX
SCHI0010.S.1	CF	AMR	WT	T83I	None	ST4	Day 7 IV CAZ, TOB
SCHI0016.S.124	UTI	WT	AMR	D87N	MexR (Ile24fs; loss; upregulates MexAB efflux pump)	ST309	Unknown
SCHI0021.S.7	CF	WT	AMR	D87H	None	ST801	Day 13 IV CAZ, TOB
SCHI0025.S.15	CF	WT	AMR	D87H	None	ST801	Unknown
SCHI0027.S.1	CF	WT	AMR	D87H	None	ST801	Day 1 IV CAZ, TOB
SCHI0029.S.1	CF	WT	AMR	D87H	None	ST801	Day 1 unknown antibiotic/s
SCHI0030.S.4	CF	WT	AMR	D87H	None	ST801	None
SCHI0032.S.32	Ulcer	WT	AMR	D87N	None	ST143#	Unknown
SCHI0032.S.33	Ear infection	WT	AMR	D87Y	None	ST848^	Unknown
SCHI0033.S.2	BSI	AMR	WT	T83I	ParE (Ala473Val), NalD (Met1del; upregulates MexAB efflux pump)	ST847	Unknown

SCHI0033.S.4	BSI	AMR	WT	T83I	None	ST244	Unknown
SCHI0033.S.15	BSI	AMR	WT	T83I	ParC (Ser87Leu)	ST532	Unknown
SCHI0033.S.19	BSI	AMR	WT	T83I	ParC (Ser87Leu)	ST235	Unknown
SCHI0033.S.26	BSI	WT	WT	None	None	ST147	Unknown
CIP-intermediate							
SCHI0002.S.8	CF	WT	WT	None	GyrB (Ser466Phe)	ST4	Day 7 IV CAZ, MEM, TOB
SCHI0002.S.9	CF	WT	WT	None	None	ST683	Day 7 IV CAZ, MEM, TOB
SCHI0002.S.12	CF	WT	WT	None	None	ST683	Day 37 IV CAZ, MEM, TOB
SCHI0003.S.1	CF	AMR	WT	T83I	None	ST649	Day 4 IV ATM, MEM, TOB
SCHI0003.S.3	CF	AMR	WT	T83I	None	ST649	Day 4 IV ATM, MEM, TOB
SCHI0003.S.8	CF	WT	AMR	D87H	None	ST801	Day 4 IV ATM, MEM, TOB
SCHI0005.S.9	CF	WT	AMR	D87H	None	ST801	Day 10 IV MEM, TOB
SCHI0005.S.10	CF	WT	AMR	D87H	None	ST801	Day 10 IV MEM, TOB
SCHI0005.S.11	CF	WT	AMR	D87H	None	ST801	Day 10 IV MEM, TOB
SCHI0006.S.10	CF	WT	AMR	D87H	None	ST801	None
SCHI0008.S.2	CF	AMR	WT	T83I	None	ST649	Day 8 IV ATM, DOX
SCHI0008.S.4	CF	AMR	WT	T83I	None	ST649	Day 8 IV ATM, DOX
SCHI0013.S.12	CF	WT	WT	None	GyrB (Ser466Tyr), ParE (Ala473Val)	ST775	Day 4 FOF, MEM
SCHI0016.S.57	BE	WT	WT	None	GyrB (Ser466Phe), MexR (Gln25stop; upregulates MexAB efflux pump)	ST2601	>2 weeks MEM, GEN, CRO, FEP
SCHI0021.S.5	CF	WT	AMR	D87H	None	ST801	Day 5 IV CAZ, TOB

SCHI0021.S.6	CF	WT	AMR	D87H	None	ST801	Day 5 IV CAZ, TOB
SCHI0021.S.8	CF	WT	AMR	D87H	None	ST801	Day 13 IV CAZ, TOB
SCHI0021.S.10	CF	WT	AMR	D87H	None	ST801	Day 1 IV CAZ, TOB
SCHI0021.S.11	CF	WT	AMR	D87H	None	ST801	Day 1 IV CAZ, TOB
SCHI0021.S.12	CF	WT	AMR	D87H	None	ST801	Day 1 IV CAZ, TOB
SCHI0021.S.14	CF	WT	AMR	D87H	None	ST801	None
SCHI0025.S.9	CF	AMR	WT	T83I	None	ST649	Unknown
SCHI0027.S.2	CF	WT	AMR	D87H	None	ST801	Day 12 IV CAZ, TOB
SCHI0027.S.3	CF	WT	AMR	D87H	None	ST801	Day 9 IV CAZ, TOB
SCHI0027.S.5	CF	WT	AMR	D87H	None	ST801	None
SCHI0030.S.2	CF	WT	AMR	D87H	None	ST801	Day 6 IV ATM, TOB
SCHI0030.S.3	CF	WT	AMR	D87H	None	ST801	Day 13 IV ATM, TOB
SCHI0033.S.14	BSI	WT	WT	None	None	ST571	Unknown
SCHI0033.S.29	BSI	AMR	WT	T83I	None	ST244	Unknown
CIP-sensitive							
SCHI0005.S.8	CF	WT	WT	None	ParE (Ala473Val), AmgS (Val121Gly; upregulates MexXY efflux pump)	ST262	Day 10 IV MEM, TOB
SCHI0013.S.2	CF	WT	WT	None	GyrB (Ser466Tyr), ParE (Ala473Val)	ST775	Day 4 FOF, MEM
SCHI0020.S.4	CF	WT	WT	None	ParE (Ala473Val), NfxB (large deletion, upregulates MexCD efflux pump)	ST3828*	Day 1 IV CAZ, TOB
SCHI0020.S.5	CF	WT	WT	None	ParE (Ala473Val), NfxB (large deletion, upregulates MexCD efflux pump)	ST3828*	None

SCHI0020.S.6	CF	WT	WT	None	ParE (Ala473Val), NfxB (loss, upregulates MexCD efflux pump)	ST3828*	Day 1 IV CAZ, TOB
SCHI0027.S.4	CF	WT	WT	None	None	ST801	None
SCHI0030.S.1	CF	WT	WT	None	None	ST275	None
SCHI0030.S.5	CF	WT	WT	None	None	ST275	None
SCHI0033.S.1	BSI	WT	WT	None	None	ST3864*	Unknown
SCHI0033.S.3	BSI	WT	WT	None	None	ST3829	Unknown
SCHI0033.S.5	BSI	WT	WT	None	None	ST274	Unknown
SCHI0033.S.7	BSI	WT	WT	None	None	ST252	Unknown
SCHI0033.S.8	BSI	WT	WT	None	None	ST649	Unknown
SCHI0033.S.9	BSI	WT	WT	None	None	ST708	Unknown
SCHI0033.S.10	BSI	WT	WT	None	None	ST244	Unknown
SCHI0033.S.11	BSI	WT	WT	None	None	ST3865*	Unknown
SCHI0033.S.12	BSI	WT	WT	None	None	ST865	Unknown
SCHI0033.S.13A	BSI	WT	WT	None	None	ST395	Unknown
SCHI0033.S.13B	BSI	WT	WT	None	None	ST395	Unknown
SCHI0033.S.16	BSI	WT	WT	None	NfxB (Phe126fs; upregulates MexCD efflux pump)	ST909	Unknown
SCHI0033.S.17	BSI	WT	WT	None	None	ST274	Unknown
SCHI0033.S.18	BSI	WT	WT	None	None	ST348	Unknown
SCHI0033.S.20	BSI	WT	WT	None	None	ST274	Unknown
SCHI0033.S.23	BSI	WT	WT	None	None	ST3843*	Unknown
SCHI0033.S.24	BSI	WT	WT	None	None	ST471	Unknown
SCHI0033.S.25	BSI	WT	WT	None	None	ST815	Unknown
SCHI0033.S.27	BSI	WT	WT	None	None	ST395	Unknown

SCHI0033.S.28	BSI	WT	WT	None	None	ST3804	Unknown
SCHI0033.S.30	BSI	WT	WT	None	None	ST782	Unknown
SCHI0033.S.32	BSI	WT	WT	None	None	ST298	Unknown
SCHI0033.S.33	BSI	WT	WT	None	MexR (Arg91fs; upregulates MexAB efflux pump	ST17	Unknown
SCHI0033.S.35	BSI	WT	WT	None	None	ST348	Unknown
SCHI0033.S.36	BSI	WT	WT	None	None	ST3830*	Unknown
SCHI0033.S.37	BSI	WT	WT	None	None	ST298	Unknown
SCHI0033.S.38	BSI	WT	WT	None	None	ST235	Unknown
SCHI0033.S.39	BSI	WT	WT	None	None	ST1189	Unknown
SCHI0038.S.3	COPD	WT	WT	None	None	ST888	None
SCHI0039.S.1	COPD	WT	WT	None	None	ST3134	None
SCHI0050.S.1	COPD	WT	WT	None	ParC (Gln405Arg)	ST3323	Day 3 oral CIP
PAO1	Wound	WT	WT	None	None	ST549	Unknown

Abbreviations: AMR, antimicrobial-resistant; ATM, aztreonam; BE, bronchiectasis; BSI, bloodstream infection, CAZ, ceftazidime; CF, cystic fibrosis; CIP,

ciprofloxacin; COPD, chronic obstructive pulmonary disease; CRO, ceftriaxone; DOX, doxycycline; FEP, cefepime; FOF, fosfomycin; GEN, gentamicin; IV,

intravenous; MEM, meropenem; SYBR-MAMA, SYBR Green-based mismatch amplification mutation assay; TOB, tobramycin; UTI, urogenital tract infection;

WGS, whole-genome sequencing; WT, wild-type

*Novel multilocus sequence type (ST) identified in this study

#SCHI0032.S.32 submitted to PubMLST under isolate name AUS205 (ID:1023) (30)

^SCHI0032.S.33 submitted to PubMLST under the isolate name AUS134 (ID:952) (30)

Table 2. *Pseudomonas aeruginosa* isolates used for *gyrA*248 and *gyrA*259 SYBR-MAMA assay

validation, and associated fluoroquinolone-class antibiotic disc diffusion data

gyrA phenotype	Isolate ID	CIP	LEV	OFX	MFX	WT to AMR (codon change)
AMR (mutant)	SCHI0003.S.3	I	I	R	R	*gyrA*248 T→C (T83I)
	SCHI0008.S.4	I	I	R	R	
	SCHI0010.S.1	R	R	R	R	
	SCHI0025.S.9	I	R	R	R	
	SCHI0016.S.124	R	R	R	R	*gyrA*259 G→A (D87N)
	SCHI0032.S.32	R	R	R	R	
	SCHI0003.S.8	I	I	R	R	*gyrA*259 G→C (D87H)
	SCHI0005.S.10	I	I	I	R	
	SCHI0025.S.15	R	R	R	R	
	SCHI0029.S.1	R	R	R	R	
	SCHI0032.S.33	R	I	R	R	*gyrA*259 G→T (D87Y)
WT (control)	SCHI0038.S.3	S	S	S	S	No mutation at *gyrA*248 or *gyrA*259
	SCHI0039.S.1	S	S	S	S	
	SCHI0050.S.1	S	S	S	S	
	PAO1	S	S	S	S	

183 Abbreviations: AMR, antimicrobial-resistant; CIP, ciprofloxacin; I, intermediate; LEV, levofloxacin;

184 MFX, moxifloxacin; OFX, ofloxacin; R, resistant; S, sensitive; WT, wild-type.

185 **Table 3. SYBR Green-based mismatch amplification mutation assay (SYBR-MAMA) primers**

186 **designed in this study**

gyrA SNP	GyrA AMR variant	Primer name	Sequence (5'-3')*	Optimised concentration (µM)
***gyrA*248**	T83I	gyrA248_T_AMR	ACGATGGTGTCGTAGACCGt**G**a	0.30
		gyrA248_C_WT	CGATGGTGTCGTAGACCGt**G**g	0.30
		gyrA248_F	TGTGGTCGGCGACGTGATC	0.20
***gyrA*259**	D87N, D87Y, D87H	gyrA259_D_AMR	CATGCGCACGATGGTGa**d**	0.20
		gyrA259_G_WT	GCCATaCGCACGATGGTGa**c**	0.20
		gyrA259_F	AGCTGGGCAACGACTGGAA	0.20

187 Abbreviations: AMR, antimicrobial resistant (allele); SNP, single-nucleotide polymorphism; WT, wild-

188 type (allele)

189 *Bolded nucleotides indicate the SNP; underlined nucleotides indicate deliberately incorporated

190 antepenultimate/penultimate mismatches to enhance allele specificity

Figures

Figure 1. SYBR Green-based mismatch amplification mutation assay (SYBR-MAMA) interrogation of

the *gyrA*248 (T➔C) biallelic single-nucleotide polymorphism, resulting in a threonine to isoleucine

substitution at position 83 (T83I) in certain fluoroquinolone (FQ)-resistant *P. aeruginosa* isolates.

*gyrA*248 SYBR-MAMA performance in: **A**. SCHI0010.S.1 (FQ-resistant, encodes T83I); **B.** SCHI0038.S.3

(FQ-sensitive wild-type isolate). Orange, antimicrobial-resistant allele; blue, wild-type allele. No-

template controls did not amplify. All samples were run in duplicate.

Figure 2. SYBR Green-based mismatch amplification mutation assay (SYBR-MAMA) interrogation of

the *gyrA*259 (G→A/C/T) tetra-allelic single-nucleotide polymorphism, resulting in aspartate to

asparagine (D87N), histidine (D87H), or tyrosine (D87Y) substitutions, respectively, at position 87 in

certain fluoroquinolone (FQ)-resistant *Pseudomonas aeruginosa* isolates. *gyrA*259 SYBR-MAMA

performance in: **A.** SCHI0005.S.10 (intermediate resistance towards FQs, encodes D87H); **B.**

SCHI0032.S.32 (FQ-resistant, encodes D87N); **C.** SCHI0032.S.33 (FQ-resistant, encodes D87Y); and **D.**

SCHI0038.S.3 (FQ-sensitive wild-type isolate). Purple, antimicrobial-resistant allele; aqua, wild-type

allele. No-template controls did not amplify. All samples were run in duplicate.

References

1. Breidenstein EB, de la Fuente-Nunez C, Hancock RE. 2011. *Pseudomonas aeruginosa*: all roads lead to resistance. *Trends Microbiol* 19:419-26.
2. Botelho J, Grosso F, Peixe L. 2019. Antibiotic resistance in *Pseudomonas aeruginosa* - Mechanisms, epidemiology and evolution. *Drug Resist Updat* 44:100640.
3. Schwartz T, Volkmann H, Kirchen S, Kohnen W, Schon-Holz K, Jansen B, Obst U. 2006. Real-time PCR detection of *Pseudomonas aeruginosa* in clinical and municipal wastewater and genotyping of the ciprofloxacin-resistant isolates. *FEMS Microbiol Ecol* 57:158-67.
4. Higgins PG, Fluit AC, Milatovic D, Verhoef J, Schmitz FJ. 2003. Mutations in GyrA, ParC, MexR and NfxB in clinical isolates of *Pseudomonas aeruginosa*. *Int J Antimicrob Agents* 21:409-13.
5. Poole K. 2011. *Pseudomonas aeruginosa*: resistance to the max. *Front Microbiol* 2:65-65.
6. Lee JK, Lee YS, Park YK, Kim BS. 2005. Alterations in the GyrA and GyrB subunits of topoisomerase II and the ParC and ParE subunits of topoisomerase IV in ciprofloxacin-resistant clinical isolates of *Pseudomonas aeruginosa*. *Int J Antimicrob Agents* 25:290-5.
7. Yonezawa M, Takahata M, Matsubara N, Watanabe Y, Narita H. 1995. DNA gyrase *gyrA* mutations in quinolone-resistant clinical isolates of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 39:1970-1972.
8. Rehman A, Jeukens J, Levesque RC, Lamont IL. 2021. Gene-gene interactions dictate ciprofloxacin resistance in *Pseudomonas aeruginosa* and facilitate prediction of resistance phenotype from genome sequence data. *Antimicrob Agents Chemother* 65:e0269620.
9. Nguyen KV, Nguyen TV, Nguyen HTT, Le DV. 2018. Mutations in the gyrA, parC, and mexR genes provide functional insights into the fluoroquinolone-resistant Pseudomonas aeruginosa isolated in Vietnam. *Infect Drug Resist* 11:275-282.
10. Rehman A, Patrick WM, Lamont IL. 2019. Mechanisms of ciprofloxacin resistance in *Pseudomonas aeruginosa*: new approaches to an old problem. *J Med Microbiol* 68:1-10.
11. Birdsell DN, Pearson T, Price EP, Hornstra HM, Nera RD, Stone N, Gruendike J, Kaufman EL, Pettus AH, Hurbon AN, Buchhagen JL, Harms NJ, Chanturia G, Gyuranecz M, Wagner DM, Keim PS. 2012. Melt analysis of mismatch amplification mutation assays (Melt-MAMA): a functional study of a cost-effective SNP genotyping assay in bacterial models. *PLoS One* 7:e32866.
12. Germer S, Higuchi R. 1999. Single-tube genotyping without oligonucleotide probes. *Genome Res* 9:72-8.
13. Donà V, Smid JH, Kasraian S, Egli-Gany D, Dost F, Imeri F, Unemo M, Low N, Endimiani A. 2018. Mismatch amplification mutation assay-based real-time PCR for rapid detection of *Neisseria gonorrhoeae* and antimicrobial resistance determinants in clinical specimens. *J Clin Microbiol* 56:00365-18.
14. Sarovich DS, Price EP, Von Schulze AT, Cook JM, Mayo M, Watson LM, Richardson L, Seymour ML, Tuanyok A, Engelthaler DM, Pearson T, Peacock SJ, Currie BJ, Keim P, Wagner DM. 2012. Characterization of ceftazidime resistance mechanisms in clinical isolates of *Burkholderia pseudomallei* from Australia. *PLoS One* 7:e30789.
15. Germer S, Holland MJ, Higuchi R. 2000. High-throughput SNP allele-frequency determination in pooled DNA samples by kinetic PCR. *Genome Res* 10:258-66.
16. Webb KA, Olagoke O, Baird T, Neill J, Pham A, Wells TJ, Ramsay KA, Bell SC, Sarovich DS, Price EP. 2021. Genomic diversity and antimicrobial resistance of *Prevotella* spp. isolated from chronic lung disease airways. *MedRxiv* doi:https://doi.org/10.1101/2021.08.30.21262864.
17. McCarthy KL, Paterson DL. 2017. Long-term mortality following *Pseudomonas aeruginosa* bloodstream infection. *J Hosp Infect* 95:292-299.
18. Freschi L, Jeukens J, Kukavica-Ibrulj I, Boyle B, Dupont MJ, Laroche J, Larose S, Maaroufi H, Fothergill JL, Moore M, Winsor GL, Aaron SD, Barbeau J, Bell SC, Burns JL, Camara M, Cantin A, Charette SJ, Dewar K, Deziel E, Grimwood K, Hancock RE, Harrison JJ, Heeb S, Jelsbak L, Jia

B, Kenna DT, Kidd TJ, Klockgether J, Lam JS, Lamont IL, Lewenza S, Loman N, Malouin F,
Manos J, McArthur AG, McKeown J, Milot J, Naghra H, Nguyen D, Pereira SK, Perron GG,
Pirnay JP, Rainey PB, Rousseau S, Santos PM, Stephenson A, Taylor V, Turton JF, Waglechner
N, et al. 2015. Clinical utilization of genomics data produced by the international
Pseudomonas aeruginosa consortium. *Front Microbiol* 6:1036.

19.   Stewart AG, Price EP, Schabacker K, Birikmen M, Harris PNA, Choong K, Subedi S, Sarovich
DS. 2021. Molecular epidemiology of third-generation-cephalosporin-resistant
Enterobacteriaceae in Southeast Queensland, Australia. *Antimicrob Agents Chemother*
65:e00130-21.

20.   Anuj SN, Whiley DM, Kidd TJ, Bell SC, Wainwright CE, Nissen MD, Sloots TP. 2009.
Identification of *Pseudomonas aeruginosa* by a duplex real-time polymerase chain reaction
assay targeting the *ecfX* and the *gyrB* genes. *Diagn Microbiol Infect Dis* 63:127-131.

21.   Price EP, Viberg LT, Kidd TJ, Bell SC, Currie BJ, Sarovich DS. 2018. Transcriptomic analysis of
longitudinal *Burkholderia pseudomallei* infecting the cystic fibrosis lung. *Microb Genom*
4:e000194.

22.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner
PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell
sequencing. *J Comput Biol* 19:455-477.

23.   Madden DE, Webb JR, Steinig EJ, Currie BJ, Price EP, Sarovich DS. 2021. Taking the next-gen
step: comprehensive antimicrobial resistance detection from *Burkholderia pseudomallei*.
EBioMedicine 63:103152.

24.   de Lamballerie X, Zandotti C, Vignoli C, Bollet C, de Micco P. 1992. A one-step microbial DNA
extraction method using "Chelex 100" suitable for gene amplification. *Res Microbiol*
143:785-90.

25.   Fraser TA, Bell MG, Harris PNA, Bell SC, Bergh H, Nguyen T-K, Kidd TJ, Nimmo GR, Sarovich
DS, Price EP. 2019. Quantitative real-time PCR assay for the rapid identification of the
intrinsically multidrug-resistant bacterial pathogen *Stenotrophomonas maltophilia*. *Microb*
Genom 5:e000307.

26.   Farahi RM, Ali AA, Gharavi S. 2018. Characterization of *gyrA* and *parC* mutations in
ciprofloxacin-resistant *Pseudomonas aeruginosa* isolates from Tehran hospitals in Iran. *Iran J*
Microbiol 10:242-249.

27.   Takenouchi T, Sakagawa E, Sugawara M. 1999. Detection of *gyrA* mutations among 335
Pseudomonas aeruginosa strains isolated in Japan and their susceptibilities to
fluoroquinolones. *Antimicrob Agents Chemother* 43:406-409.

28.   Akasaka T, Tanaka M, Yamaguchi A, Sato K. 2001. Type II topoisomerase mutations in
fluoroquinolone-resistant clinical strains of *Pseudomonas aeruginosa* isolated in 1998 and
1999: role of target enzyme in mechanism of fluoroquinolone resistance. *Antimicrob Agents*
Chemother 45:2263-2268.

29.   Bergkessel M, Guthrie C. 2013. Colony PCR. *Methods Enzymol* 529:299-309.

30.   Kidd TJ, Ritchie SR, Ramsay KA, Grimwood K, Bell SC, Rainey PB. 2012. *Pseudomonas*
aeruginosa exhibits frequent recombination, but only a limited association between
genotype and ecological setting. *PLoS One* 7:e44199.